

BASELINE ECOLOGICAL RISK ASSESSMENT REPORT PATRICK BAYOU SUPERFUND SITE DEER PARK, TEXAS

Prepared for

Patrick Bayou Joint Defense Group
U.S. Environmental Protection Agency

Prepared by

Anchor QEA, LLC 614 Magnolia Avenue Ocean Springs, Mississippi 39564

March 2013



BASELINE ECOLOGICAL RISK ASSESSMENT REPORT PATRICK BAYOU SUPERFUND SITE DEER PARK, TEXAS

Prepared for

Patrick Bayou Joint Defense Group
U.S. Environmental Protection Agency

Prepared by

Anchor QEA, LLC 614 Magnolia Avenue Ocean Springs, Mississippi 39564

March 2013

RECEIVED

13 APR 16 PM 4: 45

SUPERFUND DIV. REMEDIAL BRANCH (65F-R)

TABLE OF CONTENTS

E	EXECUTIVE SUMMARY	ES-1
	Problem Formulation	1
	Wildlife Risk Assessment	3
	Fish Risk Assessment	3
	Benthic and Aquatic Invertebrate Risk Assessment	3
	Conclusions and Risk Management Recommendations	5
1	INTRODUCTION	1
	1.1 Overview of Document	2
2	2 SITE DESCRIPTION	
3		
	3.1 Identification of Ecological COPCs	
	3.2 Refined Ecological Conceptual Site Model	
	3.3 Assessment Endpoints and Measurement Endpoints	
	3.4 Analysis Plan	9
4	BERA DATASET	11
	4.1 Data Evaluation	11
	4.1.1 Data Quality and Management	11
	4.1.2 Data Treatment, Summation, and Reduction	
	4.1.2.1 General Data Treatment Rules	12
	4.1.2.2 Summation Rules	12
	4.1.2.3 Toxic Equivalency Quotients	13
	4.2 Sediment	14
	4.3 Surface Water	15
	4.4 Tissue	16
	4.5 Bioassay/Sediment Chemistry Data	17
	4.6 Porewater Data	18
	4.7 Benthic Community Data	18
5	ECOLOGICAL COPC SUMMARY	20
_	5.1 Benthic and Aquatic Invertebrate COPCs	
	3.1 Dettatio and riquate invertebrate 301 35	

_				
	5.2 I	Fish CC	DPCs	21
	5.3 A	Aquatic	e-Dependent Wildlife COPCs	22
6	WII.r	HIFE E	RISK ASSESSMENT	24
Ö			re Assessment	
	6.1.1	-	ary Exposure Parameters	
	6.1.2		osure Point Concentrations	
		2.1	Fish and Shellfish	
		.2.2	Sediment	
	6.1	.2.3	Terrestrial Animals and Plant Matter	33
	6.1.3	Area	a Use Factors	35
	6.2 I	Effects .	Assessment	36
	6.2.1	Dev	elopment of Inorganic Mercury TRV	37
	6.3 I		naracterization and Uncertainty Analysis	
	6.3.1	Cha	racterizing PCB Risks	38
	6.3.2	Risk	Estimates	39
	6.3	3.2.1	Belted Kingfisher	39
	6.3	3.2.2	Carnivorous Birds	39
	6.3	3.2.3	Raccoon	40
	6.3	3.2.4	Spotted Sandpiper	40
	6.3.3	Unc	ertainty Analysis	40
	6.3	3.3.1	Exposure Assessment	41
	6.3	3.3.2	Effects Assessment	44
	6.3	3.3.3	Risk Characterization	45
	6.4 F	Risk Co	onclusions	45
7	FISH	RISK A	SSESSMENT	47
	7.1 H	3ioaccu	ımulative COPCs Risk Assessment	47
	7.1.1		osure Assessment	
	7.1	.1.1	Selenium Exposure Assessment	
	7.1.2	Tiss	ue Based Effects Assessment	
	7.1.3	Risk	Characterization and Uncertainty Analysis	51
	7.1	.3.1	Risk Characterization	51
	7.1	.3.2	Uncertainty Analysis	51

_			
	7.2	Surface Water Direct Contact Risk Assessment	53
	7.3	Risk Conclusions	54
8	RENT	THIC INVERTEBRATE RISK ASSESSMENT	55
U		Sediment Risk Assessment	
	8.1.1		
	8.1.2		
	8.1.3		
	8.1.4		
	8.1.5	·	
	8.1.6		
	8.1.7	-	
	8.	1.7.1 Sensitivity Analysis for Benthic Toxicity	
	8.	1.7.2 Sensitivity Analysis for Optimized Mean Probable Effect Level-Quotient	
	8.	1.7.3 Sensitivity Analysis Benthic Community	. 64
	8.2	Surface Water Risk Assessment	64
	8.2.1	Exposure Assessment	65
	8.2.2	2 Effects Assessment	65
	8.2.3	Surface Water Risk Characterization	66
	8.2.4	Surface Water Uncertainty Analysis	67
	8.3	Risk Conclusions	68
	8.3.1	Sediment	68
	8.3.2	2 Surface Water	69
9	AQU	ATIC PLANTS, REPTILES, AND AMPHIBIANS UNCERTAINTY DISCUSSION.	70
	•	Aquatic Plants	70
	9.2	Reptiles and Amphibians	71
10	0 CON	CLUSIONS AND RISK MANAGEMENT RECOMMENDATIONS	72
I	I KEFE	RENCES	/4
	ist of T		
	able 3-		
1	able 4-	Summary of BERA Dataset	

Table 4-2	Toxic Equivalency Factors for Toxicity Equivalent Calculations
Table 4-3	Summary of Site Surface Sediment Data Evaluated in the BERA Dataset
Table 4-4	Summary of Site Surface Water Data Evaluated in the BERA Dataset
Table 4-5	Summary of Site Tissue Data Evaluated in the BERA Dataset
Table 4-6	Summary of Bioassay/Sediment Toxicity Data Evaluated in the BERA Dataset
Table 4-7	Summary of Benthic Community Data in BERA Dataset
Table 5-1	Receptor Group COPCs
Table 5-2	Complete List of Benthic and Aquatic Invertebrate COPCs
Table 5-3	Complete List of Fish COPC
Table 5-4	Complete List of Aquatic-Dependent Wildlife COPCs
Table 6-1	Summary of Exposure Parameters for Wildlife Dietary Exposure Assessment
Table 6-2	Prey Grouping for Fish and Invertebrate Exposure Point Concentration
	Calculations
Table 6-3	95 UCLs for Fish and Invertebrate Prey Groups
Table 6-4	Surface Weighted Average COPC Concentrations in the Intertidal and
	Supratidal Areas
Table 6-5	Area Use Factor Summary for Wildlife Receptors
Table 6-6	Avian LOAEL TRVs
Table 6-7	Mammalian LOAEL TRVs
Table 6-8	HQs for Wildlife Receptors
Table 6-9	HQ Uncertainty Analysis
Table 6-10	Upper and Lower Bound Area Use Factor Estimates for Wildlife Receptors
Table 6-11	Area Use Factor Uncertainty Analysis for Wildlife Receptors
Table 6-12	PCB Uncertainty Analysis for Wildlife Receptors
Table 7-1	Fish Groupings for Fish Exposure Point Concentration Calculations
Table 7-2	Tissue EPCs for Fish Bioaccumulation Pathways
Table 7-3	Fish LOED TRVs
Table 7-4	HQs for Fish Receptors
Table 7-5	PCB Risk Characterization Uncertainty Assessment for Fish
Table 7-6	Biomagnification Uncertainty Analysis
Table 8-1	Sediment Toxicity Line of Evidence Summary
Table 8-2	Sediment Chemistry Line of Evidence Summary
Table 8-3	Benthic Community Line of Evidence Summary

	Table of Contents
Table 8-4	Weight of Evidence Risk Characterization Summary for Benthic Invertebrates
Table 8-5	Sediment Toxicity Sensitivity Analysis
Table 8-6	Sediment Chemistry Sensitivity Analysis
Table 8-7	Benthic Community Sensitivity Analysis
List of Figu	ires
Figure 1-1	USEPA Eight-Step Process for Ecological Risk Assessment (Adapted from
	USEPA 1997)
П' О 4	D 11D 0 6 101 XV 1 1 M

Figure 2-1 Patrick Bayou Superfund Site Vicinity Map

Baseline Ecological Risk Assessment Conceptual Site Model Figure 3-1

Figure 3-2 Simplified Ecological Conceptual Site Model

Figure 4-1 **BERA Dataset Sediment Sample Locations**

Figure 4-2 BERA Dataset Surface Water Sample Locations

Figure 4-3 BERA Dataset Invertebrate Tissue Sample Locations

Figure 4-4 **BERA Dataset Fish Tissue Sample Locations**

Figure 4-5 BERA Dataset Benthic Toxicity and in situ Community Sample Locations

Figure 6-1 Patrick Bayou Tidal Zones

Figure 8-1 Benthic Invertebrate Risk Characterization for Sediment

Figure 8-2 Relative Contribution of COPC to Optimized Mean PEL-Q by Station

List of Appendices

List of Attachments

Appendix A BERA Dataset

Appendix B Surface Sediment Interpolation and Averaging

Appendix C Wildlife Risk Assessment Supporting Information

Appendix D Fish Risk Assessment Supporting Information

Appendix E Benthic Invertebrate Risk Assessment Supporting Information

Appendix F Comments and Responses Matrix for TCEQ, NOAA, and USEPA – Draft **Ecological Risk Assessment**

Attachment 1 Ecological Fish and Invertebrate Tissue Sampling Data Report

LIST OF ACRONYMS AND ABBREVIATIONS

μg microgram

μg/L micrograms per liter

95 UCL 95 percent upper confidence limit

ACR acute-to-chronic ratio

Ah aryl hydrocarbon ANOVA analysis of variance

AOC Administrative Order on Consent

AQUIRE Aquatic Toxicity Information Retrieval

AUF area use factor

AWQC ambient water quality criteria

BAF bioaccumulation factor
BCF bioconcentration factor

BEHP bis(2-ethylhexyl) phthalate

BERA Baseline Ecological Risk Assessment

BERA Work Plan Final Baseline Ecological Risk Assessment Work Plan

BI benthic index

BMF biomagnification factor

BSAF biota-sediment accumulation factor CCA canonical correspondence analysis

CERCLA Comprehensive Environmental Response, Compensation, and Liability

Act

cm centimeter

COC chemical of concern COI chemical of interest

COPC chemical of potential concern

CSM Conceptual Site Model

DDT dichlorodiphenyltrichloroethane

DL detection limit

DO dissolved oxygen

DOC dissolved organic carbon

DQO data quality objective

DW dry weight

Eco-SSL ecological soil screening level

EPC exposure point concentration

ERA ecological risk assessment

ERED Environmental Residue-Effects Database

ES-BI Engle and Summers Biotic Index

ESBTU Equilibrium Partitioning Sediment Benchmark Toxic Unit

FAV Final Acute Value

FCV Final Chronic value

FCM food chain multiplier

FS Feasibility Study

GOM Gulf of Mexico

GOMA Gulf of Mexico Alliance

GOMA B-IBI Gulf of Mexico Alliance Benthic Index of Biological Integrity

HPAH high molecular weight polycyclic aromatic hydrocarbon

HQ hazard quotient

HSC Houston Ship Channel

JDG Joint Defense Group

kg kilogram km kilometer

km² square kilometer

L/kg liters per kilogram

LOAEL lowest observed adverse effects level

LOE line of evidence

LOED lowest estimated dose

LPAH low molecular weight polycyclic aromatic hydrocarbon

MeHg methylmercury

mg milligram

mg/kg milligrams per kilogram

mg/L milligrams per liter

MHHW mean higher high water

MLLW mean lower low water

NAVD88 North American Vertical Datum 1988

ng/L nanograms per liter

NOAA National Oceanic and Atmospheric Administration

NOAEL no observed adverse effects level

PAH polycyclic aromatic hydrocarbon

PB Patrick Bayou

PBLO Patrick Bayou Lead Organization

PCB polychlorinated biphenyl

PCDD polychlorinated dibenzo-p-dioxin

PCDF polychlorinated dibenzofuran

PEC-Q probable effect concentration quotient

PEL-Q probable effect level quotient

PRG preliminary remediation goal

PRP potentially responsible party

PSCR Preliminary Site Characterization Report

QAPP Quality Assurance Project Plan

RG remediation goal

RI Remedial Investigation

RI/FS Remedial Investigation/Feasibility Study

SAP Sampling and Analysis Plan

SH State Highway

Site Patrick Bayou Superfund Site

SOW Statement of Work

SQG sediment quality guideline

SQG-Q sediment quality guideline quotient

SQT sediment quality triad

SVOC semivolatile organic compound **SWAC** surface weighted average concentration TCDD 2,3,7,8-tetrachlorodibenzo-p-dioxin Texas Commission on Environmental Quality TCEQ TDI total daily intake TEF toxic equivalency factor TEQ toxic equivalent quotient TLtrophic level **TMDL** total maximum daily load **TNRCC** Texas Natural Resource Conservation Commission TOC total organic carbon **TRV** toxicity reference value TU toxic unit **USACE** U.S. Army Corps of Engineers **USEPA** U.S. Environmental Protection Agency VOC volatile organic compound **WOE** weight-of-evidence water quality benchmark **WQB** WQC water quality criteria WW wet weight

waste water treatment plant

WWTP

EXECUTIVE SUMMARY

This Baseline Ecological Risk Assessment (BERA) meets the requirements of Task 4 in the Statement of Work (SOW) of the Administrative Order on Consent (AOC) for the Remedial Investigation/Feasibility Study (RI/FS) of the Patrick Bayou Superfund Site (Site). The approach followed in the BERA provides a characterization of potential Site-related risks to ecological receptors resulting from the presence of hazardous substances in Site media, as presented in the Baseline Ecological Risk Assessment Work Plan (BERA Work Plan; Anchor QEA 2011a).

Problem Formulation

The BERA Conceptual Site Model (CSM) illustrates known and suspected sources of chemical contamination, types of chemicals and affected media, known and potential routes of migration, and known or potential ecological receptors. Complete and potentially significant exposure pathways are identified for the following ecological receptors:

- Benthic invertebrate community
- Fish community
- Sediment-probing birds and omnivorous/herbivorous birds spotted sandpiper
- Carnivorous wading birds composite avian receptor¹
- Piscivorous birds belted kingfisher
- Omnivorous/herbivorous mammals raccoon

The exposure pathways for these receptors include a combination of direct contact with sediment, sediment ingestion, biota ingestion, and contact with porewater and surface water.

The assessment endpoints for the selected receptors are based on protection and maintenance of the communities or populations they represent. Although the goal of the assessment endpoints is based on protection of communities and/or populations, the measurement endpoints for most ecological receptors evaluated in the BERA concern the survival, growth,

¹ A composite avian receptor based on several different species was chosen to represent the range of life histories, physical descriptions (e.g., body weight), and feeding strategies of species within this guild.

and reproduction of the organisms in each receptor group². This practice, which is common in Superfund ecological risk assessment (ERA), requires the extrapolation of individual level effects to assess potential risks to the community or population being considered. As such, thresholds for effects in this BERA are set at a level that may cause limited adverse effects on individuals but, if not exceeded, are not expected to result in adverse effects to the community or population overall.

Estimates of risk for all receptors are performed using deterministic approaches. Single estimates of exposure, such as estimated dietary intake or empirical estimates of concentration of a chemical of potential concern (COPC) in sediment are compared to levels that may result in adverse effects to ecological receptors. The exposure assessment quantifies the magnitude and spatial and temporal patterns of exposure to COPCs for ecological receptors identified during problem formulation. Consistent with the BERA Work Plan, the lowest observed adverse effects levels (LOAELs), or their equivalent, are used to define effects levels for risk estimates.

A hazard quotient (HQ) representing the ratio of estimated exposure to the adverse effects level is calculated for each receptor-COPC pair. Receptor-COPC exposure scenarios that result in HQs less than 1.0 are not expected to result in any adverse effects to either the individual receptor or the overall community or population of receptors represented by the receptor evaluated in the risk assessment. In addition to evaluating quantitative descriptions of ecological risks and threshold concentrations for adverse ecological effects, the risk characterization also presents information on the significance of the identified risks to support risk management decisions.

Several sources of uncertainty are associated with all Superfund risk estimates (USEPA 1997). Uncertainty in this BERA is primarily addressed through sensitivity analysis, whereby different point estimates of parameter uncertainty (e.g., treatment of non-detects) are used to bracket a range of risk estimates around the baseline scenario.

² The exception to this approach is the evaluation of benthic community composition measurement endpoint.

Wildlife Risk Assessment

Dietary exposure of wildlife to Site COPCs is the primary line of evidence (LOE) to assess risk to these receptor groups. Both prey tissue (i.e., fish and shellfish) and incidental sediment ingestion are considered in dietary dose estimates. Several potentially bioaccumulative COPCs, including polychlorinated biphenyls (PCBs), polychlorinated dibenzo-p-dioxin/polychlorinated dibenzofuran (PCDD/PCDFs), polycyclic aromatic hydrocarbons (PAHs), lead, mercury, hexachlorobenzene, hexachlorobutadiene, 1,3-dichlorobenzene, and 1,4-dichlorobenzene, were identified in the BERA Work Plan and are evaluated for wildlife. Only one COPC, PCBs expressed as 2,3,7,8-tetrachlorodibenzo-p-dioxin equivalents (i.e., PCB toxic equivalents [TEQs]), is equal to or exceeds a HQ of 1.0 for two receptors: spotted sandpiper and belted kingfisher. Uncertainty analysis indicates that HQs for these COPC-receptor pairs may be above or below the threshold of concern (HQ of 1.0) depending on the assumptions used to characterize risk; demonstrating that within the ranges of exposure and effects variables evaluated, risks may or may not exceed a threshold of concern for these COPC-receptor pairs.

Fish Risk Assessment

Risks to fish are assessed using a body burden approach, comparing the concentration of COPCs in fish tissue to tissue levels that are expected to have adverse effects on fish. The BERA Work Plan identified PCBs, PCDD/PCDFs, mercury, and selenium as bioaccumulative fish COPCs. Measured or estimated levels of these COPCs in whole body fish are compared to their corresponding effects levels. None of the COPCs exceed their respective effects levels (i.e., all HQs are less than 1.0). As a result, no adverse risk to fish populations from Site COPCs are identified.

Benthic and Aquatic Invertebrate Risk Assessment

Sediment chemistry, Site specific toxicity tests, and data on the condition of the benthic community (e.g., abundance, diversity, etc.) are used to assess risk to benthic invertebrates from exposure to sediment COPCs.

Initially, benthic toxicity models were evaluated to determine if an empirical, Site-specific relationship between surficial sediment chemistry and observed toxicity in laboratory

bioassays using Site sediment could be established. If such a model could be developed, toxicity in Site sediments could then be predicted across the Site using sediment chemistry alone. Several different quotient or toxic unit (TU) models that have been described in the open literature to assess this relationship were evaluated in the BERA Work Plan (Anchor QEA 2011a). The performance of each model was assessed by applying it to Site specific, colocated bulk sediment chemistry data and bioassay (i.e., toxicity) data for the marine amphipod *Leptocheirus plumulosus*. Of the models assessed, a mean quotient model using the Probable Effects Level (Long et al. 2006) was initially selected based on several performance criteria identified in the BERA Work Plan. This model, hereafter referred to as the mean Probable Effects Level-Quotient (PEL-Q), was refined using a series of optimization steps in the BERA Work Plan. This optimized model included four COPCs (total PCBs, total PAHs, lead, and bis(2-ethylhexyl)phthalate [BEHP]) that demonstrated a statistically significant difference in concentration between toxic and non-toxic samples.

This toxicity model initially developed in the BERA Work Plan was refined and reevaluated during the BERA. During the development of the BERA, it became apparent that the toxicity model, as initially conceived, had several limitations that could not be resolved and that led to a significant amount of uncertainty in its utility and relevance as a primary line of evidence. Specifically, and in no particular order of priority, these limitations are: the lack of reference area toxicity tests to account for naturally occurring stressors and confounding factors (e.g., salinity acclimation, porewater ammonia, etc.) that may result in an unknown but potentially significant amount of conservative bias in the toxicity attributed to Site COPCs; the model, when applied to other test species, results in a high percentage of false positives, leading to a potentially significant amount of Site sediments to be falsely predicted as toxic; and the analytical uncertainty in the PCB analysis in the toxicity sample dataset confounds the extrapolation of the sediment chemistry data to more recent samples. Thus, the uncertainty in the model, which is primarily a function of the uncertainty in the data used to develop the model, did not lead to a sufficient level of confidence in the model to define the magnitude and extent of risks to the benthic community.

Therefore, a weight of evidence (WOE) approach was developed in the BERA to assess benthic risk. This WOE approach included the review and analysis of Site-specific bulk sediment chemistry data, sediment bioassay data, and benthic community data as LOEs. The

objective of the WOE approach was to use the apparent correspondence between the values or metrics assigned to the LOEs, and the overall strength of the correspondence, where it existed, to identify areas of the Site where measurable incremental risks to the benthic community due to exposure to Site-related COPCs are deemed probable, indeterminate, or low. The way in which the degree and strength of correspondence among the three LOEs was factored into the consensus-based ranking of different areas of the Site is considered to be conservative (i.e., biased toward identifying areas as probable or indeterminate, rather than low). Twelve different locations within the Site with co-located synoptic bulk sediment chemistry, bioassay data, and benthic community were included in the analysis. Based on the WOE approach, two of these locations were identified as areas where incremental Site-related risks to the benthic community are probable, five locations were identified as indeterminate areas, and five locations were identified as low. PCBs were identified as a primary driver for probable risk locations and are designated as a chemical of concern (COC) for the benthic community.

Finally, the BERA Work Plan identified PCBs as a surface water COPC for benthic and aquatic invertebrate communities. Surface water levels of PCBs did not exceed relevant effect levels for invertebrates. Thus, no risks to invertebrates from PCBs in surface water are identified.

Conclusions and Risk Management Recommendations

The following risk management recommendations are made for the Site based on the outcome of the BERA:

1. PCB TEQ HQs for the sediment-probing and piscivorous bird receptor groups³ are equal to 1.0 and 1.7 for spotted sandpiper and belted kingfisher, respectively. However, uncertainty analyses indicate that HQs for these COPC-receptor pairs may be above or below the threshold of concern (HQ = 1.0) depending on the assumptions used to characterize risk. Thus, within the ranges of exposure and effects variables evaluated, risks may or may not exceed a threshold of concern for individuals exposed to PCBs in Site media.

³ Based on risks to the belted kingfisher.

- 2. Risks to fish populations at the Site are negligible and no risk management for this receptor group is necessary.
- 3. Using a WOE approach, areas of probable benthic risk were identified. Although a quantitative risk characterization for the benthic community could not be performed within the acceptable range of uncertainty, it is apparent that probable risks to the benthic community are likely associated with bulk sediment PCBs. Although no specific risk management recommendations are provided herein, risk management for this receptor group should be considered within the overall context of other risk management considerations (e.g., water quality standards) during the Feasibility Study (FS).
- 4. Ecological and human health risks occur along a continuum and there is not a quantifiable bright line for those risks. Remedial alternatives will be evaluated in the FS that lower the overall Site and sub-area risk for areas that are characterized as indeterminate and probable risks. In concert with USEPA, quantitative tools will be developed to assess the ultimate risk reduction expected from a specific remedial alternative and that risk reduction score will be used as part of the effectiveness assessment for each alternative, along with USEPA's other FS criteria. In addition, the evaluation of surface water impacts as related to sediment concentrations will be a part of the evaluation of remedial alternatives.

1 INTRODUCTION

This document presents the *Baseline Ecological Risk Assessment* (BERA) component of the Patrick Bayou Superfund Site (Site) *Remedial Investigation/Feasibility Study* (RI/FS). It describes the approach, data, and results used to complete the baseline risk assessment for ecological receptors required in Task 4 of the Statement of Work (SOW) of the Administrative Order on Consent (AOC) entered into with the U.S. Environmental Protection Agency (USEPA) for the RI/FS dated on January 31, 2006.

The BERA follows the ecological risk assessment (ERA) approach presented in the *Final Baseline Ecological Risk Assessment Work Plan* (BERA Work Plan; Anchor QEA 2011a). The approach is consistent with *USEPA Guidance for Conducting ERAs* (USEPA 1997, 1998), or as directed by USEPA. In addition, state guidance (Texas Natural Resource Conservation Commission [TNRCC] 2001; Texas Commission on Environmental Quality [TCEQ] 2006) is considered where appropriate.

The ERA process at Superfund sites is described as a series of eight-steps (Figure 1-1; USEPA 1997). The screening-level problem formulation and ecological effects evaluation (Steps 1 and 2 of the USEPA framework) are part of the initial ecological risk screening assessment, which have been completed along with portions of Steps 3 through 6 (problem formulation, study design and data quality objectives (DQOs), field sampling planning, and site investigation) through previous RI/FS documents described in the BERA Work Plan. The BERA Work Plan also provides a refinement of Steps 3 through 6. Step 7 (risk characterization) is performed in this BERA, which also includes summaries of the previous steps, exposure and effects assessments, and an uncertainty analysis. Step 8 (risk management) is the phase where remedial actions are considered and is part of the Feasibility Study (FS).

The general objectives of the BERA are to:

- Identify whether unacceptable ecological risks due to chemicals of potential concern (COPCs) are present at the Site.
- Describe the range and magnitude of those risks to ecological receptors in the absence of any Site remediation.

• Identify, as chemicals of concern (COCs), those chemicals that exceed an acceptable risk threshold for one or more ecological receptors.

The results of the BERA will also support the establishment of risk-based preliminary remediation goals (PRGs) during the FS. Along with other risk management considerations, the risk-based PRGs will be used in the evaluation of remedial options in the FS and, ultimately, identification of final Remediation Goals (RGs) when the remedy is selected.

1.1 Overview of Document

The remainder of this document is organized as follows:

- Section 2 Site Description This section provides general information about the ecological habitat types and species present at the Site.
- Section 3 Problem Formulation This section presents a summary of the ecological COPCs, the conceptual exposure model, and the selected assessment and measurement endpoints.
- Section 4 BERA Data This section presents a summary of the Site data used in the BERA.
- Section 5 Ecological COPC Summary This section summarizes the process used to identify ecological COPCs and presents the COPCs for each ecological receptor group.
- Section 6 Wildlife Risk Assessment This section presents the exposure, effects, and risk characterization evaluation for selected bird and mammal receptors.
- Section 7 Fisk Risk Assessment This section presents the exposure, effects, and risk characterization evaluation for selected fish receptors.
- Section 8 Benthic Invertebrate Risk Assessment This section presents the exposure, effects, and risk characterization evaluation for the benthic invertebrate community.
- Section 9 Aquatic Plants, Reptiles, and Amphibians Uncertainty Discussion This section provides a qualitative discussion of uncertainty regarding risks to these receptors.
- Section 10 Conclusions and Risk Management Recommendations This section presents the overall risk conclusions of the BERA and risk management recommendations.

The following Appendices and Attachments are also included as part of this BERA:

- Appendix A BERA Dataset
- Appendix B Surface Sediment Interpolation and Averaging
- Appendix C Wildlife Risk Assessment Supporting Information
- Appendix D Fish Risk Assessment Supporting Information
- Appendix E Benthic Invertebrate Risk Assessment Supporting Information
- Appendix F Comments and Responses Matrix for TCEQ, NOAA, and USEPA Draft Baseline Ecological Risk Assessment
- Attachment 1 Ecological Fish and Invertebrate Tissue Sampling Data Report

2 SITE DESCRIPTION

Patrick Bayou is a tributary of the Houston Ship Channel (HSC) in Harris County, Texas (Figure 2-1). Patrick Bayou discharges into the south side of the HSC approximately 2.3 miles upstream of its confluence with the San Jacinto River. The Site, its history, and its physical features are described in more detail in the *Preliminary Site Characterization Report* (PSCR; Anchor Environmental 2006), a brief summary is provided in the following paragraphs.

The Site is located north of State Highway (SH) 225 downstream of the City of Deer Park Waste Water Treatment Plant (WWTP) and flows north approximately 10,200 feet (1.9 miles) to the HSC. The drainage upstream of the Site originates in the City of Deer Park and consists of trapezoidal, concrete-lined ditches, which transition into large culverts underneath SH 225. These culverts emerge into a gunite-lined channel at the upstream Site boundary. The gunite-lined portion of the Site extends approximately 1,800 feet north (downstream) and ends near an east-west trending railroad bridge crossing. This gunite-lined portion of the Site has an unconsolidated, earthen bottom. Downstream of the gunite-lined portion of the Site, the channel is composed of natural and armored banks. Nearly the entire Site is tidally influenced. A tributary of Patrick Bayou referred to as the East Fork, joins the Site approximately 6,500 feet upstream of the mouth of Patrick Bayou. The East Fork is a small intermittent stream approximately 5,500 feet long that flows in a northwesterly direction.

To support consistent spatial referencing, the Site is segmented by stations from the mouth of Patrick Bayou at its confluence with the HSC (Station PB-000⁴), up to the Site boundary at the culverts under SH 225 (Station PB-102). These stations provide the approximate linear distance from downstream to upstream in hundreds of feet (e.g., Station PB-102 is approximately 10,200 linear feet from the mouth of Patrick Bayou).

⁴ PB - 'Patrick Bayou'

3 PROBLEM FORMULATION

This section presents the problem formulation for the BERA. The BERA problem formulation is developed in Step 3 of the eight-step risk assessment process and identifies specific factors to be addressed in the ERA (USEPA 1997).

The problem formulation step of the ERA process includes the following:

- Refinement of preliminary COPCs
- Further characterization of the ecological effects of COPCs at the Site
- Review and refinement of information on fate and transport, complete exposure pathways, and ecosystems potentially at risk
- Selection of assessment and measurement endpoints
- Development of a Conceptual Site Model (CSM) and risk questions
- Development of an analysis plan

Detailed information on the BERA problem formulation is presented in the BERA Work Plan and the elements of the problem formulation are summarized here as follows:

- Refined screening process for identifying COPCs (Section 3.1)
- Refined CSM (Section 3.2)
- Refined assessment and measurement endpoints (Section 3.3)
- Analysis plan outlining the methods for conducting the BERA (Section 3.4)

3.1 Identification of Ecological COPCs

As part of the refined screen conducted following the procedures described by USEPA (1997, 2001), data from the Site were screened to identify the BERA COPCs. Refinement of the ecological COPCs was performed in an iterative fashion; refinements were performed as additional relevant data became available during the RI (Anchor Environmental 2006; Anchor 2008a, 2008b). The most recent refinement was conducted as part of the BERA problem formulation presented in the BERA Work Plan. The most recent round of sediment and surface water data (Anchor QEA 2010) was used to refine the final list of COPCs for the BERA. A general summary of the refined screen and ecological COPCs that were identified for each ecological receptor group are presented in a separate section (Section 5).

3.2 Refined Ecological Conceptual Site Model

The ecological CSM for the BERA is one of the four primary products of Step 3 of the eight-step ERA process (USEPA 1997), along with assessment endpoints, exposure pathways, and risk questions. An ecological CSM describes predicted relationships between receptors potentially at risk and the stressors to which they may be exposed.

Consistent with Superfund guidance (USEPA 1997), ecological receptors were identified in the BERA Work Plan to represent the range of organisms exposed to COPCs at the Site. For the BERA, the following species or groups of species (communities) were identified as representative ecological receptors:

- Benthic invertebrate community
- Fish community
- Sediment-probing birds and omnivorous/herbivorous birds spotted sandpiper
- Carnivorous wading birds composite avian receptor⁵
- Piscivorous birds belted kingfisher
- Omnivorous/herbivorous mammals raccoon

Elements of the ecological CSM include information about source media, transport pathways, exposure media, routes of exposure, and receptors. An exposure pathway is considered complete if a chemical of interest (COI) can travel from a source to ecological receptors and is available to the receptors via one or more exposure routes (e.g., ingestion; USEPA 1997). The ecological CSM provides a basis for identifying those exposure pathways and provides a framework for characterizing those risks associated with potentially complete pathways.

Complete pathways can be of varying relevance, so it is important to identify key pathways that reflect the greatest potential exposures to ecological receptors sensitive to the COPCs being considered (USEPA 1997). Exposure pathways for COPCs at the Site were designated in the BERA Work Plan in one the following four ways for each receptor relative to their exposure potential:

⁵ A composite avian receptor based on several different species was chosen to represent the range of life histories, physical descriptions (e.g., body weight), and feeding strategies of species within this guild.

- 1. Complete and significant Complete and significant pathways have a direct link between receptor and COI, and the specific pathway is considered a potentially important contributor to risk. These pathways are quantitatively addressed in the BERA.
- 2. Complete and uncertain Complete and uncertain pathways have a direct link between receptor and COI; however, there are insufficient scientific data available to quantify the significance of the pathway in overall assessment of exposure. These pathways are addressed qualitatively to a level of certainty dependent on available toxicological studies and exposure data.
- 3. Complete and minor Complete and minor pathways have a direct link between receptor and COI; however, the significance of this pathway in terms of overall exposure is not considered significant or is negligible relative to other exposure pathways. These pathways are qualitatively discussed in the risk characterization section for each receptor group.
- 4. Incomplete Incomplete pathways do not have a direct link between receptor and COI. An assessment of these pathways is not performed in the BERA.

The BERA Work Plan identified the following exposure pathways as complete and significant:

- Benthic and aquatic invertebrates:
 - Direct contact and ingestion of sediment and sediment porewater
 - o Biota ingestion
 - Direct contact and ingestion of surface water
- Fish populations
 - o Direct contact and ingestion of sediment and sediment porewater
 - o Ingestion of biota
 - o Direct contact with surface water
- Sediment-probing birds
 - o Ingestion of biota
 - Incidental sediment ingestion
- Carnivorous wading birds
 - Ingestion of biota
 - Incidental sediment ingestion

- Piscivorous birds
 - o Ingestion of biota
- Omnivorous/herbivorous mammals
 - Ingestion of biota
 - o Incidental sediment ingestion

The complete ecological CSM showing source media, transport, exposure media, routes of exposure, and receptors is diagramed in Figure 3-1. A simplified version illustrating exposure media, routes of exposure, and receptors associated with complete and significant pathways is provided in Figure 3-2.

3.3 Assessment Endpoints and Measurement Endpoints

Assessment endpoints are the valued attributes of ecological entities upon which risk management actions are focused (USEPA 1998). The assessment endpoints for all receptors selected in the BERA Work Plan are based on protection and maintenance of the communities or populations they represent. Although the goal of the assessment endpoints is based on protection of communities and/or populations, the assessment endpoints for all ecological receptors evaluated in the BERA concern the survival, growth, and reproduction of the organisms in each receptor group. This practice, which is common in Superfund ERA (primarily for practical reasons), requires the extrapolation of individual level effects to assess potential risks to the community or population being considered. This is considered a generally conservative approach. As such, thresholds for effects in this BERA are set at a level that may cause adverse effects on some individuals, but, if not exceeded, are not expected to result in adverse effects to the community or population overall.

Measurement endpoints are measureable (i.e., quantitative) ecological characteristics that are used to evaluate if the assessment endpoint is protected by addressing a series of risk questions. Measurement endpoints are evaluated with one or more lines of evidence (LOEs).

⁶ An exception to this practice includes the assessment endpoints for threatened or endangered species. In such cases, protection of the individual (in addition to the community or population) is the basis for the assessment and measurement endpoints. As discussed in the BERA Work Plan, no threatened or endangered species were selected as ecological receptors for the BERA due to their absence from the Site.

A LOE is a set of data and associated analyses that can be used to estimate ecological risks. One or more LOE were selected for each measurement endpoint. The assessment endpoints, risk questions, measurement endpoints, and LOE selected for ecological receptors identified in the BERA Work Plan are summarized in Table 3-1.

3.4 Analysis Plan

The analysis plan describes the specific approaches and methods for conducting the risk calculations used to evaluate the risk questions and assessment endpoints (USEPA 1997). Details on the analytical approaches used to evaluate risks to each ecological receptor group are provided in the effects and exposure sections of Sections 6 through 10. A summary of the major components described in the BERA analysis plan is provided below.

Exposure Assessment – All exposure pathways classified as complete and significant in the ecological CSM (Figure 3-2) were evaluated quantitatively. The exposure assessment quantified the magnitude and spatial and temporal patterns of exposure to COPCs for ecological receptors identified during problem formulation. For the BERA, the exposure assessment includes the replacement of many assumptions included in the previous screening level evaluations (Anchor Environmental 2006; Anchor 2008a, 2008b, Anchor QEA 2011a) with Site-specific data to provide a more robust analysis of potential risks.

Effects Assessment – The effects assessment primarily includes an integration of literature derived ecological effects information with any Site-specific data developed during the RI (e.g., toxicity testing). For the wildlife and fish effects assessments, literature derived effects information was used to identify appropriate toxicity reference values (TRVs) and benchmarks. For the benthic invertebrate effects assessment, sediment chemistry, site specific toxicity tests, and data on the condition of the benthic community (e.g., abundance, diversity, etc.) were used to assess effects to benthic invertebrates exposed to sediment COPCs.

Risk Characterization – As part of the risk characterization, information on contaminant exposure and effects were integrated to estimate risks to the receptors. Several different risk estimation methods were used for the various measurement endpoints and LOEs in this

BERA. In addition to presenting quantitative descriptions of ecological risks and threshold concentrations for adverse ecological effects, the risk characterization also presents information on the significance of the identified risks, including: 1) the location and spatial extent of Site contamination exceeding adverse effect thresholds; and 2) the magnitude of or degree to which adverse effect thresholds are exceeded.

Uncertainty Analysis – Several sources of uncertainties are associated with all Superfund risk estimates (USEPA 1997). Although all sources of uncertainty cannot be identified and described within the context of the risk assessment, the major sources of uncertainty in the risk estimates are identified, described, and where possible, quantified. Uncertainty in this BERA is primarily addressed through sensitivity analysis, whereby different point estimates of parameter uncertainty (e.g., treatment of non-detects) are used to bracket a range of risk estimates around the baseline scenario.

4 BERA DATASET

Numerous datasets have been compiled in support of the Patrick Bayou RI/FS over the course of several years. Additional data relevant to the Site were compiled from other sources and screened for inclusion in the RI dataset.

As described in the BERA Work Plan, the BERA uses a subset of the data that comprise the RI dataset (hereafter referred to as the BERA dataset). The BERA utilizes only data collected from within the Site boundaries. The following data are included in the BERA dataset:

- Chemistry data for 52 surface sediment samples collected in 2009 and 2011
- Chemistry data for 26 surface water samples collected in 2009 and 2011
- Chemistry data for 83 whole-body fish and invertebrate tissue samples collected in 2011
- Co-located whole sediment bioassay data and surface sediment chemistry for 51 samples collected between 2001 and 2006

The BERA dataset includes only those media relevant for ecological exposure pathways: surface sediment (0 to 10 centimeter [cm]), surface water, and invertebrate and fish tissue. The BERA dataset, which includes data generated as part of the RI and historical data, is summarized, by medium, in Table 4-1, shown by location in Figures 4-1 through 4-4, and described in detail in the following subsections.

4.1 Data Evaluation

This section summarizes the data quality review process for the BERA dataset and describes the data reduction, summation, and normalization procedures.

4.1.1 Data Quality and Management

Methods for performing data quality reviews for data generated by the Joint Defense Group (JDG) are described in the project Quality Assurance Project Plan (QAPP) (Anchor 2007) and in task specific sampling plans. In addition, detailed review of the quality of the non-JDG datasets were performed prior to acceptance in the project database according to the criteria established for the project (Anchor Environmental 2006). All field data, physical

measurements, and chemical data generated by the JDG or other parties (non-JDG data) were maintained and managed as directed by the project *Data Management Plan* (Anchor 2007). All chemical data were maintained in a relational project database (i.e., EQuiS®) and validated by the Anchor QEA project chemist or qualified third party validator.

4.1.2 Data Treatment, Summation, and Reduction

Rules for the treatment of chemical data in the BERA dataset are described below.

4.1.2.1 General Data Treatment Rules

General data treatment rules for the risk assessment are as follows:

- 1. Sediment data were presented on a dry weight (DW) basis unless stated otherwise.
- 2. Tissue data were presented on a wet weight (WW) basis unless stated otherwise.
- 3. For risk estimation purposes, sediment data generated through duplicate analysis (i.e., field duplicates) were treated as quality control samples and were not included in subsequent calculations or data analysis steps.
- 4. For individual analyte results, a concentration equal to one-half the sample-specific detection limit (DL; as reported by the laboratory) was used for undetected analytes.

4.1.2.2 Summation Rules

Unless stated otherwise, for total values (e.g., total polychlorinated biphenyls [PCBs]) where at least one analyte included in the summation was detected, all detected values were included and non-detect values were included at one-half the DL in the total. For total values where none of the analytes included in the summation were detected, the DL was set to the highest DL for an individual analyte and the total result was labeled as non-detect.

Selected individual analytes included in totals are as follows:

- Total PCBs Total PCBs represent the sum of all reported PCB Congeners (n=209) for all JDG generated data in the BERA dataset. The exception to this general rule applies to the benthic risk assessment. Development of the benthic model (Section 8) utilized non-JDG data that were analyzed for PCB Aroclors.
- Total LPAH Total low molecular weight polycyclic aromatic hydrocarbons [PAHs]

(LPAHs) are the sum of 2-methylnaphthalene, acenaphthene, acenaphthylene, anthracene, fluorine, naphthalene, and phenanthrene.

- Total HPAH Total high molecular weight PAHs (HPAH) are the sum of fluoranthene, pyrene, benzo(a)anthracene, chrysene, benzofluoranthene, benzo(a)pyrene, indeno(1,2,3-c,d)pyrene, dibenzo(a,h)anthracene, and benzo(g,h,i)perylene.
- Total PAH Sum of LPAH and HPAH.

4.1.2.3 Toxic Equivalency Quotients

The 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) toxic equivalency quotient (TEQ) was the primary expression used to assess exposure of wildlife and fish to complex mixtures of polychlorinated dibenzo-p-dioxin/polychlorinated dibenzofuran (PCDDs/PCDFs), and dioxin-like PCBs⁷ (USEPA 2008a). TEQs were calculated using the formula:

$$TEQ = \sum_{n=1}^{k} C_n \times TEF_n$$

where:

 C_n = concentration of dioxin-like chemical n in organism or its diet

 TEF_n = Toxic equivalency factor for dioxin-like chemical n

K = number of dioxin-like chemicals in mixture

TEQs were calculated using fish, bird, and mammalian toxic equivalency factors (TEFs) provided by USEPA (2008). TEFs used to calculate TEQ values are provided in Table 4-2. TEQs were calculated separately for sum PCDD/PCDFs (PCDF/PCDD TEQ) and sum dioxin-like PCBs (PCB TEQ). If at least one congener was detected in a sum, all detected values were included and non-detect values at one-half the DL. If no congeners in a sample were detected, the DL was set to the highest TEF-adjusted congener TEQ DL and the total result was labeled as a non-detect.

⁷ The dioxin-like PCBs include a subset of 12 co-planar PCB Congeners. They are: PCB-77, PCB-81, PCB-105, PCB-114, PCB-118, PCB-123, PCB-126, PCB-156, PCB-157, PCB-167, PCB-169, and PCB-189 (USEPA 2008a).

4.2 Sediment

Surface sediment chemistry from the Site used for the BERA dataset include sediment data collected in October and November of 2009⁸ and August of 2011 (Anchor QEA 2010; 2012). Table 4-3 presents a summary of the sediment samples included in the BERA dataset. These data represent the most recent, comprehensive assessment of conditions at the Site. Previous investigations include data collected between 2000 and 2008. However, the collection methods, field protocols, and quality control protocols differed between sampling events due to the varied purposes (e.g., total maximum daily loading [TMDL] evaluations, sediment transport modeling, etc.), DQOs, and sampling entities (e.g., TCEQ, potentially responsible party [PRP], etc.). The differences represent a potential source of uncertainty in the dataset, which is absent or controlled in the 2009 and 2011 data. Samples from both sampling events were collected from the sediment mixing-zone layer (0 to 10 cm).

The sediment mixing-zone layer is the surface-layer of the sediment bed where active physical mixing occurs due to biological burrowing (bioturbation) and hydrodynamic forces. The thickness of the mixing-zone layer is important because it affects the rate of natural attenuation of surface-layer COPC concentrations at the Site, and it is representative of the depth of the bioactive zone where exposure is likely to occur. Within the mixing-zone layer, bioturbation and hydrodynamic forces tend to homogenize the sediment and chemicals, which produces vertical profiles of chemical concentration that are nearly constant. Results of the Patrick Bayou sediment mixing-zone layer study concluded that the mixing-zone layer depth at the Site, and thus depth of exposure to benthos, is less than 10 cm (Anchor QEA 2009b).

The 2009 data provide a spatially robust characterization of the distribution of COPCs at the Site. Sediment samples were collected in a systematic grid pattern across the Site, and therefore provide a synchronous set of Site-wide data with which to develop surface concentration maps for the risk assessment (Figure 4-1). Forty-six surface sediment samples were collected and analyzed for the list of COPCs identified in the *Ecological COPC Report Addendum* (Anchor QEA 2008b), including metals, semi-volatile organic compounds

⁸ Data collected in October – November 2009 were included in the refinement of COPCs performed in the BERA Work Plan. Data collected in 2011 were subsequent to the BERA Work Plan.

(SVOCs), volatile organic compounds (VOCs), pesticides, PCB Congeners, and PCDDs/PCDFs. The objective of this sampling was to collect a spatially representative sediment dataset for use in the exposure assessment of the BERA. Detailed results of this sampling are provided in the *Sediment and Surface Water COPC Delineation Data Report* (Anchor QEA 2010).

One of the purposes of the 2011 sampling event was to provide additional information on the distribution of PCBs and PAHs in sediments within the upstream portion of the Site (the investigation was conducted from PB-066 to PB-101 and sediment samples were collected between Stations PB-068 and PB-0829; Figure 4-1). Six samples were collected from areas of soft sediment accumulation and analyzed for PAHs and PCB Aroclors. Detailed results of this sampling event are provided in the *Upstream Patrick Bayou Characterization Data Report* (Anchor QEA 2012a).

Surface sediment data were used in the BERA to estimate exposure concentrations for relevant ecological receptors based on direct contact (i.e., benthic invertebrates and fish) and dietary exposure (i.e., fish and wildlife). The compiled results of surface sediment samples from the 2009 and 2011 sampling events included in the BERA dataset are provided in Appendix A-1.

4.3 Surface Water

Surface water chemistry from the Site used for the BERA dataset include surface water data collected in October and November of 2009 and August of 2011. Table 4-4 presents a summary of the surface water samples included in the BERA dataset. These data represent the most recent comprehensive assessment of the conditions at the Site. Sample locations for the 2009 and 2011 sampling events are provided in Figure 4-2.

Four locations within the Site boundary were included in the October-November 2009 sampling effort for a comprehensive list of COPCs, including conventional parameters, metals, SVOCs, VOCs, pesticides, PCBs, and PCDDs/PCDFs. A total of 14 samples were

⁹ Soft-sediments suitable for sampling were not identified upstream of PB-082.

collected from the mid-depth of the water column, as determined in the field, and approximately 6 inches from the bottom at each location. Collection occurred during two tidal conditions: one at approximately slack low tide and the second at approximately midtide (on an outgoing, or ebb tide). Detailed results of this sampling are provided in the Sediment and Surface Water COPC Delineation Report (Anchor QEA 2010).

One of the purposes of the 2011 sampling event was to provide additional surface water data for PCBs in the upstream portion (PB-066 to PB-101) of the Site. Because detectable concentrations of PCBs were reported at Station PB-076, the upstream-most surface water station within Site boundaries during the 2009 sampling event, additional data were required to better characterize the distribution of PCBs in surface water. Detailed surface water sampling results are provided in the *Upstream Patrick Bayou Characterization Report* (Anchor QEA 2012a).

Surface water data are used in the BERA to estimate exposure concentrations for relevant ecological receptors based on direct contact (i.e., benthic invertebrates and fish) and bioaccumulative surface water COPCs. The chemistry results of surface water samples from the 2009 and 2011 sampling events included in the BERA dataset are provided in Appendix A-2.

4.4 Tissue

Tissue chemistry data from the Site included in the BERA dataset were collected during June of 2011 to evaluate risks to selected ecological receptors of concern. Table 4-5 presents a summary of the fish and invertebrate tissue samples included in the BERA dataset. The detailed results of this sampling event have not been reported previously and are therefore attached as a complete data report (Attachment 1) to the BERA.

Invertebrate tissue data included in the BERA dataset include samples for blue crabs (*Callinectes sapidus*), brown shrimp (*Penaeus aztecus*), white shrimp (*Penaeus setiferus*), and oysters (*Crassostrea virginica*). A total of 33 invertebrate samples were collected: 21 blue crab samples, eight brown shrimp samples, three white shrimp samples, and one oyster sample (Table 4-5; Figure 4-3). Invertebrates were divided into two size classes, 2 – 7.5 cm

and 7.5 – 13 cm; representing the prey sizes ingested by the wildlife receptors identified in the BERA Work Plan. Samples were analyzed for wildlife COPCs identified in the BERA Work Plan, including lead, total mercury, PCB Congeners, PCDDs/PCDFs, PAHs, hexachlorobenzene, 1,3-dichlorobenzene, hexachlorobutadiene, and 1,4-dichlorobenzene¹⁰.

Fish tissue data included in the BERA dataset include samples of Gulf killifish (*Fundulus grandis*), Gulf menhaden (*Brevoortia patronus*), pinfish (*Lagodon rhomboides*), sand seatrout (*Cynoscion arenarius*), and striped mullet (*Mugil cephalus*). A total of 50 fish samples were collected: 25 Gulf killifish, ten Gulf menhaden, four pinfish, two sand seatrout, and nine striped mullet (Table 4-5; Figure 4-4). All fish collected for analysis were less than 15 cm (total length) based on the assumed prey size for the wildlife receptors identified in the BERA Work Plan. Fish tissue were analyzed for both fish and wildlife COPCs identified in the BERA Work Plan, including total mercury, hexachlorobenzene, 1,3-dichlorobenzene, PCB Congeners, and PCDDs/PCDFs.

Individual organisms of the same species were composited together to meet analytical tissue volume requirements. If the required amount of tissue volume was not collected from a fishing effort, then samples were either composited over time (i.e., from the same location over the course of several hours or days), or composited spatially from within pre-defined sampling areas. The sampling areas were based on the approximate foraging ranges of the target species. The sampling area for juvenile (2 - 7.5 cm) blue crab, oyster, and killifish was 30 meters. Large juvenile blue crab (7.5 - 13 cm), brown shrimp, white shrimp, Gulf menhaden, pinfish, sand seatrout, and striped mullet had a sampling area of 300 meters. Organisms collected outside of these limits were not composited together.

4.5 Bioassay/Sediment Chemistry Data

Sediment toxicity results and synoptic sediment chemistry analyses from previous studies within the Site boundary provided a foundation for assessing the potential risks to the benthic community resulting from direct exposure to contaminated Site sediments.

¹⁰ Not all samples were analyzed for all COPCs. The target analytes for different size classes were based on expected dietary constituents of wildlife receptors and COPCs identified for wildlife receptors. See Section 6 for more details on wildlife dietary COPCs.

The data included in the BERA benthic toxicity dataset include 51 samples (Table 4-6; Figure 4-5) with co-located whole sediment bioassay data and surface sediment chemistry collected primarily in support of TMDL studies for Patrick Bayou (Parsons et al. 2002, 2004), as well as routine and special studies performed by TCEQ (no date). Of the surface sediment samples, 44 samples were collected within the Site, three samples were collected downstream of the Site in the HSC, and four samples were collected upstream of the Site boundary. Samples were collected between September 2000 and August 2006 and analyzed for metals, PAHs, SVOCs, and PCBs.


4.6 Porewater Data

Porewater data were collected as part of the *Supplemental Work Plan, Patrick Bayou Superfund Site, Deer Park, Texas* (Anchor 2007c). Sixteen porewater samples collected by centrifugation of bulk sediment from the 0-10 cm interval were analyzed for selected SVOCs, metals (total and dissolved), pesticides, PCB Congeners¹¹, ammonia, total sulfide, total organic carbon (TOC), and dissolved organic carbon (DOC). Eight porewater samples collected using passive diffusion sampling were analyzed for VOCs. One-hundred porewater samples collected at ten stations were submitted for mercury speciation (total mercury, methylmercury, and alkylated mercury)¹².

Porewater data were used to identify and refine the list of benthic invertebrate COPCs (Anchor 2008a, 2008b) but was not selected as a LOE to assess risks in this BERA based on the USEPA approved Work Plan (Anchor QEA 2011a). Mercury speciation data were used in this BERA to assess the bioavailability of mercury in sediments consumed by wildlife receptors (Section 6).

4.7 Benthic Community Data

Benthic community data were collected during previous investigations within Patrick Bayou. Site data were collected during third party TMDL investigations performed by the Patrick



¹¹ Only 11 of 16 samples were analyzed for PCB Congeners.

¹² Samples were collected at 2 cm intervals per station to a depth of 20 cm. Every other interval was submitted for analysis.

Bayou Lead Organization (PBLO; Parsons et al. 2002; Parsons et al. 2004)¹³. These samples were collected concomitantly with the sediment samples utilized in the bioassay tests described in Section 4.5.

Sample locations are identified in Figure 4-5. In September 2000, samples were collected from 19 locations. Fifteen stations were located within Patrick Bayou, two stations were located upstream of the Site (i.e., south of SH 225), one station was located in the East Fork Tributary, and one station was located in the HSC just east of the mouth of Patrick Bayou. These locations were sampled again in April 2001. A limited sampling event at four stations within Patrick Bayou was performed in October 2001 and six stations were sampled within Patrick Bayou again in August 2003¹⁵. Records included in the BERA dataset are summarized in Table 4-7.

¹³ Details on sample collection and processing can be found in these documents.

¹⁴ Stations Q, 4A, G, and E.

¹⁵ Stations E, 7, R, 6A, 4A, and 3.

5 ECOLOGICAL COPC SUMMARY

As stated in Section 3.1, ecological COPCs were refined in the BERA Work Plan problem formulation using the Site dataset identified for the BERA (Section 4). COPCs were identified in order to focus the list of chemicals for evaluation of ecological risks. COPCs initially were identified by conducting a screening-level analysis during which the maximum concentrations in various media are compared to conservative thresholds of toxicity. Chemicals with maximum concentrations that did not exceed conservative thresholds of toxicity or were not otherwise identified as COPCs were ruled out as posing unacceptable risk to ecological receptors (Anchor 2008a, 2008b).

In addition, a comprehensive COPC refinement screen was conducted as part of the BERA Work Plan. The initial list of target analytes based on the *COPC Report and Amendment* (Anchor 2008a, 2008b) contained 100 different chemicals.¹⁶ This list was not a likely reflection of the list of chemicals that pose a significant unacceptable risk for adverse effects on ecological receptors. A refinement of the chemicals carried into this BERA was conducted to focus on likely indicator chemicals¹⁷ and to avoid diverting effort in pursuit of chemicals with little likely incremental or measureable contribution to unacceptable risks to ecological receptors.

A total of 44 ecological COPCs were identified for the Site at the conclusion of the BERA Work Plan and are listed by class for each receptor group in Table 5-1. They include eight metals, 19 PAHs (including calculated totals), PCBs¹⁸, PCDD/PCDF¹⁹, three SVOCs, and 12 VOCs. The following sections summarize the selection of COPCs for the BERA for each representative receptor group.

¹⁶ Not including all 209 PCB Congeners, which were treated as one analyte for purposes of the tally.

¹⁷ An indicator chemical or chemical group is one that is the most toxic, persistent, and/or mobile among those substances likely to contribute significantly to the overall risk at the Site (USEPA 1988).

¹⁸ As total congeners and as TEQs using the toxicity equivalent methodology for 2,3,7,8-TCDD and dioxin-like compounds.

¹⁹ As total TEQ using the toxicity equivalent methodology for 2,3,7,8-TCDD and dioxin-like compounds.

5.1 Benthic and Aquatic Invertebrate COPCs

The list of benthic and aquatic invertebrate COPCs was developed through a series of screening and refinement steps prior to the BERA (Anchor 2008a, 2008b, Anchor QEA 2010). A complete description of this screening assessment is provided in the BERA Work Plan.

Based on this refinement, 43 chemicals were identified as sediment COPCs for benthic invertebrates. This included eight metals, 19 individual PAHs or PAH sums, three SVOCs, total PCBs, and 12 VOCs. Only one surface water COPC, total PCBs, was identified for benthic and aquatic invertebrates. Table 5-2 provides a complete list of the benthic and aquatic invertebrate COPCs.

5.2 Fish COPCs

Per the *COPC Report and Amendment* (Anchor 2008a, 2008b), sediment COPCs identified for benthic invertebrates served as a proxy for identifying direct sediment exposure COPCs for fish. For direct surface water exposure, total PCBs were identified as a COPC based on a comparison to screening benchmarks (Anchor QEA 2010). PCDD/PCDF, which lacked a relevant surface water screening value for fish, were retained as a COPC in the BERA and evaluated using an empirical body burden approach (i.e., whole body tissue).

In addition, detected COPCs in surface water and sediment were compared to the list of potentially bioaccumulative chemicals in surface water in TCEQs *Guidance for Conducting Ecological Risk Assessments at Remediation Sites in Texas* update (2006). Chemicals detected in surface water that were included in this list were identified as bioaccumulative COPCs for the BERA. They include mercury, selenium, PCBs, and PCDD/PCDF. For potentially bioaccumulative COPCs in sediment, estimated tissue concentrations of each COPC was calculated using the 95 percent upper confidence limit (95 UCL) of the BERA sediment data multiplied by the geometric mean biota-sediment accumulation factor (BSAF)²⁰. COPCs were compared to their respective TRV²¹ using a hazard quotient (HQ)

²⁰ BSAFs were derived from literature sources and similar sites in the Gulf of Mexico (GOM) region. Details are provided in the COPC Report and Amendment (Anchor 2008a, 2008b).

approach (Anchor 2008b, Anchor QEA 2010). A COPC was retained for the BERA if the Midpoint TRV-based HQ exceeded 1.0. Based on the refined exposure and effects assessment, mercury and total PCBs were identified as bioaccumulative sediment COPCs for fish and were carried forward into the BERA. Table 5-3 provides a complete list of fish COPCs.

5.3 Aquatic-Dependent Wildlife COPCs

The primary route of exposure for aquatic-dependent wildlife is through the ingestion of biota. For some receptors, incidental ingestion of sediment also is considered a complete and significant pathway. During the COPC refinement, TRVs were refined to include recent toxicological data. Specifically, the USEPA's ecological soil screening level (Eco-SSL) documents (USEPA 2005a, 2005b, 2008a, 2007a, 2005c, 2007b, 2007c, and 2007d) were reviewed for arsenic, cadmium, chromium, copper, lead, zinc, dichlorodiphenyltrichloroethane (DDT) and metabolites, and PAHs; respectively. A single TRV based on the arithmetic average of the NOAEL and LOAEL was used (i.e., Midpoint TRV). A COPC was retained for the BERA if the Midpoint TRV based HQ exceeded 1.0, based on the refined exposure assessment.

The Site-wide 95 UCL of the sediment data collected in 2009 was calculated and used as the exposure concentration for the incidental sediment ingestion pathway and was the basis to estimate tissue concentrations using the BSAF approach. Tissue exposure concentrations were estimated using previously calculated geometric mean BSAFs (Anchor 2008a, 2008b) in conjunction with the updated Site-wide 95 UCL. A full explanation of the variables used in the total daily intake (TDI) equation for the ingestion of biota is provided in the BERA Work Plan.

The refined TDI estimates for the aquatic-dependent wildlife receptor (spotted sandpiper, carnivorous wading birds, belted kingfisher, and raccoon) COPCs were compared to their respective Midpoint TRV using a HQ approach. A COPC was retained for the BERA if the Midpoint TRV-based HQ exceeded 1.0. Based on the refined risk characterization, nine

²¹ For the refinement screen of fish bioaccumulative COPCs, a single TRV based on the arithmetic average of the no observed adverse effects level (NOAEL) and lowest observed adverse effects level (LOAEL) was used (i.e., Midpoint TRV).

COPCs: lead, mercury, total PAHs, total PCB Congeners, total PCDD/PCDF, 1,3-dichlorobenzene, 1,4-dichlorobenzene, hexachlorobenzene, and hexachlorobutadiene (Table 5-4), were identified as aquatic-dependent wildlife COPCs for this BERA.

6 WILDLIFE RISK ASSESSMENT

This section presents the risk assessment for wildlife (birds and mammals) at the Site. Dietary exposure is the main pathway by which wildlife receptors are exposed to Site COPCs. To address the different ways wildlife may be exposed to COPCs through their diets, four receptors representing four general feeding guilds were evaluated. They are:

- Sediment-probing birds spotted sandpiper
- Carnivorous birds²²
- Piscivorous birds belted kingfisher
- Omnivorous mammal raccoon

Dietary exposure and potential risks to wildlife receptors were evaluated using TDI as a LOE for all four wildlife receptors. Both prey tissue ingestion and incidental sediment ingestion were accounted for in dietary-dose estimates²³. Several factors had to be accounted for to estimate exposure levels. These included information about feeding rates, foraging areas, prey home ranges, and diets. The sum of exposure from biota and incidental sediment ingestion (i.e., TDIALL) was compared to TRVs expressed as dietary effect thresholds. TRVs were selected from published studies. The TRVs provide the basis for evaluating whether exposure concentrations are at or above a level that may cause an effect on survival, growth, or reproduction of most, but not all, individual organisms in experimentally exposed populations. This follows the conventional practice of using organism-level TRVs defined in this manner to evaluate the potential effects on populations. For the BERA, the LOAEL-based TRVs were used as the effects level.

A HQ was calculated for each COPC and receptor. The HQ was expressed as the ratio of the TDI from all sources (TDIALL) to the LOAEL TRV. A HQLOAEL of 1.0 was considered the point of departure for identifying potentially unacceptable risks to wildlife populations (USEPA 1992, 1997). A HQLOAEL less than 1.0 indicates adverse ecological effects at the population level are unlikely for that receptor group. A HQLOAEL greater than or equal to one indicates

²² Carnivorous birds were evaluated as a composite receptor representing several species (Anchor QEA 2011a).

²³ Ingestion of surface water was not identified as a complete and significant pathway for wildlife receptors in the BERA Work Plan.

that adverse ecological effects may potentially occur. The ecological significance of COPCs with HQ_{LOAEL} equal to or greater than 1.0 are considered in the risk conclusions. The details of this risk assessment for wildlife are presented as follows:

- Section 6.1 presents an overview of the assumptions used to derive exposure concentrations, including how 95 UCLs and area use factors (AUFs) were derived.
- Section 6.2 presents a summary of the effects data. Effects data in this assessment are represented by LOAEL TRVs.
- Section 6.3 presents the risk characterization results and COC-receptor pairs.
- Section 6.4 presents the risk conclusions and uncertainty analyses.

6.1 Exposure Assessment

This section presents the methods and assumptions that were used to estimate wildlife exposures to COPCs²⁴. The measure of exposure selected for wildlife is the dietary ingestion route of exposure. The primary route of exposure for aquatic-dependent wildlife is through the ingestion of biota. For some receptors, incidental ingestion of sediment is considered a complete and significant pathway as well. Thus, the exposure media in the wildlife exposure assessment are prey tissue and surface sediment and the potential risks to each receptor group were evaluated using measured or estimated COPC concentrations in these two media at the Site. Ultimately, the prey tissue and sediment exposure point concentrations (EPCs) were combined (using an assumed fraction of sediment in the diet) to provide a single dietary exposure assessment for each COPC-receptor pair.

The ingested dose received by each representative receptor was estimated by multiplying species-specific food and incidental sediment ingestion rates (normalized to body weight) by the concentrations of COPCs in prey species (fish and invertebrates) and sediment. Modifications based on behavioral factors (i.e., foraging range) were applied where appropriate.

²⁴ COPC-receptor pairs for the BERA were identified in the BERA Work Plan and were summarized in Section 5 (see Table 5-4).

6.1.1 Dietary Exposure Parameters

Ingestion exposure was characterized using a TDI model of the COPCs on a body weight-normalized basis (milligram [mg] of COPC/kilogram [kg] body weight/day). The exposure model representing TDI from environmental media was the sum of incidental sediment TDI and biota TDI:

$$TDI_{all} = TDI_{biota} + TDI_{sediment}$$

The exposure model for biota ingestion was adapted from the USEPA (1993; Equation 4-8) and is represented as:

$$TDI_{biota} = \left[\sum_{k=1}^{m} (C_{k,TS} \times NIR_k)\right] \times AUF$$

where:

TDIbiota = potential average daily dose (mg COPC/kg body weight/day)

 $C_{k,TS}$ = contaminant concentration in k^{th} type of food (mg COPC/kg food)

 NIR_k = normalized ingestion rate of k^{th} type of food on WW basis (kg food/kg

body weight/day)

AUF = area use factor

m = number of contaminated food types

To account for incidental sediment ingestion, the following equation was adopted from USEPA (1993; Equation 4-23):

$$TDI_{sediment} = \left[\left(\sum_{k=1}^{m} \left(C_{k,SD} \times FS \times IR_{total} \right) \right) \middle/ BW \right] \times AUF$$

where:

TDI_{sediment} = total daily intake of COPCs through incidental ingestion of sediment

(mg COPC/kg body weight/day)

 $C_{k,SD}$ = concentration in the k^{th} foraging area sediment on a DW basis (mg/kg)

FS = faction of sediment in diet (as percentage of diet on a DW basis;

unitless)

 IR_{total} = food ingestion rate on a DW basis. For equations using IR_{total} on a WW

basis, conversion to DW ingestion rates would be necessary (mg

food/kg body weight/day)

BW = body weight (kg)

AUF = area use factor

m = number of foraging areas²⁵

A complete discussion of these equations and their input variables is provided in the BERA Work Plan. Parameterization of each variable for representative receptors is also discussed in the BERA Work Plan²⁶. Body weights, food ingestion rates, and sediment ingestion rates vary among bird and mammal receptors. Table 6-1 presents the dietary parameters for bird and mammal receptors. Details and rationale for selected receptor-specific exposure parameters and uncertainties are presented in the BERA Work Plan.

6.1.2 Exposure Point Concentrations

Prey tissue and sediment EPCs were calculated for each COPC-receptor pair. Prey tissue EPCs were determined by calculating the 95 UCL. Estimates of sediment exposure within the foraging area were calculated using surface weighted average concentrations (SWACs). EPCs based on tissue 95 UCLs and sediment SWACs were used to calculate HQs as described below.

²⁵ For the baseline risk characterization, the number of unique foraging areas (m) was set to 1. Receptors were assumed to average their exposure equally across all available habitat.

²⁶ The incidental sediment ingestion rate for the spotted sandpiper is an exception to this statement. A review of the source document for this parameter (Beyer et al. 1994) during preparation of the BERA noted that sediment ingestion by four sandpiper species (i.e., stilt sandpiper, semipalmated sandpiper, least sandpiper, and western sandpiper) ranged from 7 to 30 percent of the dry diet, with an average incidental ingestion rate of 18 percent. The BERA Work Plan listed an incidental sediment ingestion rate of 30 percent for this receptor. However, the average rate of 18 percent was selected for the baseline risk assessment consistent with the selection of more realistic exposure parameters in the BERA and consistent with recent Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) sites (i.e., Portland Harbor). This represents a change from the BERA Work Plan.

6.1.2.1 Fish and Shellfish

Concentrations of COPCs in the diet were estimated using field-collected fish and shellfish tissue data representative of the diets for wildlife receptors. Tissue data were grouped based on the diet composition of the representative receptors; including consideration of species of fish or shellfish and the size class of the organism(s). A discussion of the methods for determining the prey groupings is provided below.

6.1.2.1.1 Prey Groupings

Fish and shellfish tissue data used in the wildlife exposure assessment were collected during a field investigation conducted in 2011 in accordance with the approved *Tissue Sampling and Analysis Plan* (SAP) attached to the BERA Work Plan. The results of this investigation are provided in a data report as an attachment to the BERA (Attachment 1). Prior to estimating tissue EPCs for the wildlife BERA, these results were evaluated using graphical and statistical methods to determine if different prey subgroups existed within the dataset that warranted independent treatment when developing EPCs. Specifically, the data were analyzed to determine whether the different species that were collected and analyzed demonstrated significant differences in the concentration of COPCs in their tissue and/or whether significant spatial trends were apparent in the data. If so, pooling all tissue data prior to calculating the EPC (i.e., 95 UCL) would not be recommended. If multiple populations are present, it is recommended that they be separated and evaluated independently (USEPA 2007).

For each COPC, tissue concentrations in fish and shellfish were evaluated according to the following steps:

- 1. Box plots and histograms of the distribution of COPCs in each species/size class of fish and shellfish collected were constructed.
- 2. Statistical tests (i.e., analysis of variance [ANOVA]) were performed to identify significant differences in COPCs among each species/size class.
- 3. Spatial plots of COPCs results for each species/size class were constructed and evaluated for trends.
- 4. Species were pooled according to the results of steps #1-3 above.
- 5. EPCs were calculated using the pooled groups according to the procedures described in Section 6.1.2.1.2.

Full results of the graphical and statistical analyses are provided as Appendix C-1. COPCs included in the analyses are: PCB TEQ, PCDD/PCDF TEQ, total PCB Congeners, total PAHs, hexachlorobenzene, mercury, and lead. Hexachlorobutadiene, 1,3-dichlorobenzene, and 1,4-dichlorobenzene were not detected in any tissue samples and were not included. Box plots and quantile-quantile plots were constructed on natural logarithms of the tissue data. Statistical tests included ANOVA on natural logarithm transformed data at a significance level (α) of 0.05. Means testing among species were performed using Tukey's HSD.

Analysis by Prey Species

For fish, visual and statistical differences were apparent among species for mercury, total PCB congeners, and PCB TEQ. Gulf menhaden had statistically higher mercury concentrations than other species. Gulf killifish demonstrated significantly higher total PCB Congeners and PCB TEQ concentrations than other fish species and were treated as separate prey groups for the purposes of EPC calculations. Gulf menhaden and striped mullet demonstrated significantly higher PCB TEQ concentrations than pinfish or sand seatrout and were treated as separate groups in EPC calculations as well. For invertebrates, few differences were observed. When both size classes of invertebrates (i.e., 2 to 7.5 cm and 7.5 to 12.5 cm) were pooled, total PCB Congener concentrations in brown shrimp were significantly higher than other species. No other COPCs demonstrated significant differences among invertebrate species. For small size class invertebrates (< 7.5 cm) which constitute the diet of the spotted sandpiper, too few results were available to perform a meaningful comparison among species in this size class. As a result, all species within this size class were pooled for calculating EPCs for spotted sandpiper prey items.

Using the results of this analysis, prey groups were determined for each wildlife receptor in accordance with their expected diets. Prey groups for each receptor are summarized in Table 6-2. The proportion of fish and or shellfish in each receptor's diet was divided evenly among the corresponding prey groups identified for each receptor. For example, a belted kingfisher's diet is assumed to be 89 percent fish (Table 6-1). If two fish prey groups were identified, they were assumed to each be 44.5 percent of the belted kingfisher's diet. Next, an EPC (i.e., 95 UCL) for each prey group was determined as described in the following section.

Spatial Trend Analysis

Plots of tissue results by distance were created to evaluate potential spatial trends. Plots of each COPC by species are located in Appendix C-1. Significant spatial patterns were not apparent for most COPCs and species. Statistically significant linear regressions of concentration by distance were observed for mercury in Gulf killifish, total PCBs and PCB TEQs in Gulf menhaden, and PCDD/PCDF TEQs and PCB TEQs in blue crabs. However, the relationship between concentration and distance was generally weak (e.g., $r^2 < 0.30$ and p-values between 0.05 and 0.01) for most COPC – species pairs. The most significant trends were mercury in Gulf killifish (decreasing downstream to upstream; $r^2 = 0.68$) and total PCBs in Gulf menhaden (increasing downstream to upstream; $r^2 = 0.64$). However, even these trends were not dramatic with less than an order of magnitude difference between downstream and upstream locations. Therefore, for estimating risk to wildlife receptors, no prey subgroups were identified based on spatial differences in COPC concentrations in prey tissue.

6.1.2.1.2 95 UCL

Prey tissue EPCs were calculated using ProUCL Version 4.0 software (USEPA 2007). USEPA's ProUCL software tests the goodness of fit for a given dataset and then computes the appropriate 95 UCL. The ProUCL software used for this analysis allows detected and non-detected values to be considered and creates interpolated values for non-detects based on the perceived distribution of the detected concentrations. Once any necessary interpolation is performed, the software conducts an analysis of the data to determine the most appropriate 95 UCL and makes a recommendation, which was then used as the EPC for the risk calculations. A minimum of six detected concentrations are required to derive a 95 UCL (USEPA 2007). In the case where an insufficient number of detected values were available, the maximum concentration was used to represent the EPC. Table 6-3 presents the EPCs for each receptor and their respective prey groups. All ProUCL outputs for the 95 UCL computations for each prey subgroup are provided in Appendix C-2.

Based on the foraging ranges of the receptors, their prey, and the distribution of habitat types, it was assumed that most aquatic and mammalian receptors would generally average their exposure to fish, invertebrates, and other prey items across the Site. Therefore, the

95 UCLs were calculated based on Site-wide exposure and sub-areas were not defined for the baseline scenario.

6.1.2.2 Sediment

It was assumed that receptors would only have access to the supratidal and intertidal areas for wading based on the size (due to constraints with water depth), foraging habits, and life history of the receptors and the guilds they represent. Therefore, the Site was divided into three areas: 1) subtidal (< 0.5 feet elevation²⁷); 2) intertidal (> 0.5 feet to < 1.5 feet elevation); and 3) supratidal (> 1.5 feet elevation) (Figure 6-1). These elevations represent the mean lower low water (MLLW) (0.5 feet) and mean higher high water (MHHW) (1.5 feet) for the Site.

6.1.2.2.1 Sediment Surface Weighted Average Concentration

Receptors were assumed to average their exposure across the supratidal and intertidal zones. Thus, SWACs were calculated for each COPC for the combined supratidal and intertidal zones as defined above. This SWAC was used as the sediment EPC for incidental sediment ingestion calculations. Table 6-4 presents the results of the SWAC calculations.

Methylmercury Estimates in Sediment

Mercury in sediments occurs in two general forms, organic and inorganic. Organic mercury is primarily present as methylmercury (CH₃Hg⁺) (MeHg). MeHg occurs as an aqueous species and can be adsorbed on to sediment organic carbon through equilibrium partitioning processes. It is the most prevalent and most biochemically active form of organic mercury in sediments. Relative to inorganic forms of mercury, MeHg is significantly more toxic (Wolfe et al. 1998).

During the selection of COPCs for the BERA (Anchor 2008a, 2008b), measures of total mercury in bulk sediment were conservatively estimated to be in the methylated form. As

²⁷ In North American Vertical Datum 1988 (NAVD88). These values were derived from the gauge datum's for the NOAA/TCOON station located at Battleship Texas State Park (Station ID 8770743). Station information can be located at:

 $http://tides and currents.noaa.gov/data_menu.shtml?stn=8770743\%20 Battleship\%20 Texas\%20 State\%20 Park,\%20 TX\&type=Datums$

such, estimates of incidentally ingested mercury in sediment were compared to a MeHg TRV. This was a highly conservative assumption as MeHg is typically present as only a small proportion of total mercury in sediment (USEPA 1997). Based on previous investigations for the Site (Anchor 2008a, 2008b), methylation in Site sediments is limited, due to high concentrations of sulfur present in surface sediments. Because there was a concern that mercury methylation could occur at centimeter scale intervals, a detailed evaluation of mercury in pore water was conducted over 2 cm intervals in the top 20 cm of surface sediments throughout the Site (Anchor 2007c). These measurements were taken in lieu of determining the concentration of MeHg in bulk sediment to determine if there were any vertical zones of higher potential MeHg production, and to determine the speciation of mercury in porewater and sediment including MeHg, mercury chloride, divalent mercury, and mercury sulfide. Because MeHg occurs as an aqueous species, actual pore water measurements versus bulk sediment measurements provide more accurate and conservative estimates of the methylation potential, location, and bioavailability. Equilibrium partitioning theory allows the calculation of MeHg measured in porewater to MeHg in bulk sediment if DOC and TOC are known.

To limit the uncertainty in the exposure assessment for mercury in sediments, available Site data were evaluated to estimate the proportion of MeHg to total mercury in Site sediments on a bulk sediment basis using the existing sediment and porewater data. These proportions were used to estimate bulk sediment concentrations of MeHg and inorganic mercury for the incidental sediment ingestion pathway exposure assessment. TRVs for MeHg and inorganic mercury were developed in Section 6.3 for each form of mercury ingested by wildlife receptors.

Estimates of MeHg in sediments were developed using simple equilibrium partitioning models and Site-specific measurements of porewater MeHg, DOC, and bulk sediment TOC. Results of MeHg porewater analysis (0-10 cm interval) at nine locations within the Site were evaluated. Estimates of bulk sediment MeHg results were then compared to observed bulk sediment total mercury results in co-located or nearby surface sediment samples. Results of this analysis are provided in Appendix C-3. This analysis concluded that rates of methylation in Site surface sediments are low; estimated MeHg did not exceed 0.2 percent of total mercury in samples evaluated. Uncertainty analysis using the most conservative observed

Site values for DOC, TOC, and total mercury²⁸ did not result in any estimates of MeHg exceeding 2.3 percent of total mercury.

For the exposure assessment, the total mercury SWAC was divided into MeHg and inorganic forms for estimates of total daily intake. One percent of the total mercury SWAC was assumed to be MeHg and the remainder (99 percent) inorganic mercury. Thus, the TDIsediment for total mercury included MeHg and inorganic mercury. In the risk characterization step, the MeHg TDIsediment was compared to the MeHg TRV²⁹. Similarly, the inorganic mercury TDIsediment was compared to an inorganic mercury TRV. Risk estimates for total mercury were then calculated as the sum of the HQ for MeHg and inorganic mercury in Section 6.4.

6.1.2.3 Terrestrial Animals and Plant Matter

Carnivorous birds and raccoons are the only receptors that may be exposed to possible Site COPCs through ingestion of terrestrial animals and plant matter at the Site. Terrestrial animal components of carnivorous bird and raccoon diets are expected to include primarily insects and occasional small mammals, reptiles, and amphibians. Empirical data to estimate concentrations in terrestrial animals at the Site are not available. To account for carnivorous bird and raccoon COPC exposure due to ingestion of contaminated terrestrial animal matter, COPC concentrations in a hypothetical omnivorous mammal, the marsh rice rat, were estimated using Equation 5-12 from the *Screening Level Ecological Risk Assessment Protocol for Hazardous Waste Combustion Facilities* (USEPA 1999). The marsh rice rat is a small, semi-aquatic omnivore that inhabits marshy areas, is common in eastern Texas and is often used as target receptors or prey items in risk assessment (Anchor QEA 2011a). It was assumed that half of the marsh rice rats consumed by carnivorous birds and raccoons were exposed to the Site (i.e., FR_{animal} =0.5).

USEPA (1999) Equation 5-12 utilizes bioconcentration factors (BCFs) and prey-item trophic level information to estimate marsh rice rat COPC tissue concentrations from consumption of contaminated prey items, plants, sediment and water. Site surface water is brackish and is generally unsuitable for drinking (Anchor QEA 2011a). Because of this, the contribution of

²⁸ Within the dataset used for the evaluation of mercury partitioning in Appendix C-3.

²⁹ The MeHg TDI_{sediment} is in addition to the MeHg TDI_{biota}. All mercury ingested by a receptor through ingestion of food (i.e., biota) was assumed to be MeHg.

Site water ingestion to marsh rice rat tissue COPC concentrations was not quantified. Equation input variables included BCFs and default factors included in USEPA (1999)³⁰, Sitespecific information (e.g., sediment and fish COPC concentrations), or were estimated using best professional judgment. Equation 5-12 accounts for the proportion that each contaminated animal or plant item contributes to the marsh rice rat diet. These omnivores typically consume equal parts vegetation and animal matter (Anchor QEA 2011a). The marsh rice rat diet was therefore assumed to be composed of 50 percent vegetation and 50 percent animal matter (25 percent invertebrates and 25 percent fish). The marsh rice rat PCB Congener TEQ tissue concentration was preliminarily calculated using either the default Aroclor 1254 BCFs or the 2,3,7,8-TCDD BCFs (USEPA 1999). The 2,3,7,8-TCDD BCFs resulted in a more conservative estimate of PCB Congener TEQ tissue concentration. For this reason, the 2,3,7,8-TCDD BCFs were used for estimating both marsh rice rat and plant tissue concentrations. The variables used for calculating marsh rice rat mercury and PCB Congener TEQ tissue concentrations are included in Appendix C-4 (Table C-4.2).

For plant matter, carnivorous birds and raccoons were assumed to ingest rooted aquatic plants. Mercury and PCBs were identified as COPCs for raccoon and carnivorous birds (Section 6.1).

Mercury and PCB Congener TEQ uptake by plants was assessed using a simple bioaccumulation factor (BAF) approach. Soil-to-plant BAFs for inorganic mercury (as mercuric chloride) and PCB congener TEQ (as 2,3,7,8-TCDD) were selected from USEPA (1999). The recommended BAF is 0.0375 kg/kg for mercuric chloride and 0.0056 kg/kg for 2,3,7,8-TCDD. The BAFs for mercury and PCB Congener TEQ were used to estimate exposure through ingestion of plant matter according to the following equation:

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

³⁰ Some equation variables were based on the knowledge of a COPC's log octanol-water partition coefficient (Log K_{ow}) which was included in USEPA (1999). No methylmercury Log K_{ow} was available in the USEPA (1999). A methylmercury Log K_{ow} was alternatively adopted from the *Canadian Tissue Residue Guidelines for the Protection of Wildlife Consumers of Aquatic Biota: Methylmercury* (Environment Canada 2002). The highest Log K_{ow} in the range provided Table 2 (Environment Canada 2002) was conservatively used.

where:

CPLANT = Concentration of mercury in plant tissue

CSED = Concentration of mercury in sediment

BAFPLANT = Bioaccumulation factor for mercury (kgsed / kgplant)

C_{PLANT} was converted to a WW basis prior to estimating total daily intake for plant matter assuming an average water content of 63 percent³¹ (USEPA 1993) according to the following equation:

$$C_{PLANT,WW} = C_{PLANT,DW} \times \left(\frac{(100 - \% moisture)}{100}\right)$$

where:

CPLANT, www = Concentration in plant tissue as WW

CPLANT,DW = Concentration in plant tissue as DW

% moisture = Moisture content of plant tissue

The sediment EPC based on the SWAC for total mercury and PCB Congener TEQs (Section 6.2.2.2) were used as the CSED term to estimate CPLANT. Estimates of CPLANT for mercury and PCB Congener TEQs are provided in Appendix C-4 for raccoon and carnivorous birds.

6.1.3 Area Use Factors

An AUF represents a ratio of the total area available to a receptor to forage, roost, breed, etc. within a site relative to its entire home range. In some cases, it may also represent the proportion of time a receptor may spend at a site due to migratory considerations and other factors. In this analysis, all receptors are assumed to be non-migratory and non-habitat selective due to breeding status (e.g., breeding, rearing, dispersal, etc.). Only differences in home range relative to habitat available at the Site were considered³².

³¹ Mid-point of emergent aquatic vegetation values from Table 4-2 of USEPA (1993).

³² Temporal use was not considered because: 1) within each guild there are some species that are non-migratory and; 2) quantitative assessment of temporal use due to breeding status is highly uncertain based on available data.

In the BERA Work Plan, an extensive review of each receptor's life history and habitat requirements was performed and compared to the available habitat within the Site. In general, the Site provides medium to poor quality habitat for most wildlife receptors due to the lack of natural conditions within the riparian and upland habitats which provide suboptimal conditions for denning, nesting, roosting, foraging, and cover. Sufficient intertidal zone habitat for wading, gleaning, and probing is generally limited as well.

Table 6-5 provides an estimate of foraging ranges for ecological receptors based on available literature. This analysis is in addition to what was performed in the BERA Work Plan and was used to develop AUFs based on relative habitat available for each receptor. Because the Site itself provides relatively poor quality habitat to the receptor groups identified, the upper quartile (i.e., 75th percentile) of the home ranges identified in Table 6-5 was used for the baseline exposure assessment³³. In general, higher quality habitat results in smaller home ranges; particularly for territorial species. Conversely, poor quality habitat requires larger home ranges to meet the minimum requirements (e.g., food, water) for most species; meaning that species that utilize the Site are likely to be near the upper bound limit of their home ranges. The home ranges from the reported studies were compared to the available habitat in terms of tidal stream distance in kilometers (km) or foraging area in square kilometers (km²) consistent with the home ranges reported in the literature. Given the previously described habitat quality at the Site, the AUFs presented in Table 6-5 are considered reasonable estimates of the proportion of habitat that is available to meet the needs of the ecological receptors evaluated in this BERA.

6.2 Effects Assessment

This section presents the selected TRVs used to characterize potential effects for wildlife COPC-receptor pairs. The evaluation used dietary-based TRVs, which were based on LOAELs derived from toxicological literature and approved by USEPA in the BERA Work Plan. The effects data presented in this section are used in combination with the exposure data presented in Section 6.1 to complete the risk characterization step (Section 6.3).

 $^{^{33}}$ The exception is the raccoon which had too few relevant home range studies (n=2) to estimate a percentile. The Midpoint (e.g., media) value was used for this receptor.

The identification of threshold effect concentrations focused on ecologically relevant effects such as survival, reproduction, and growth. Derivation of the TRVs is provided in the *COPCs Report and Amendment* (Anchor 2008a, 2008b) and the BERA Work Plan. All TRVs were approved by USEPA in the BERA Work Plan. The TRVs for birds and mammals are presented in Tables 6-6 and 6-7, respectively.

6.2.1 Development of Inorganic Mercury TRV

As discussed in Section 6.1.2.2.1, incidental ingestion of mercury in sediment was evaluated for both methylmercury and inorganic mercury. The mammalian and avian TRV in the BERA Work Plan were based on studies performed using methylmercury (e.g., methylmercury chloride). To address the uncertainty associated with applying a methylmercury TRV to an ingested dose of inorganic mercury, the literature was reviewed to identify an appropriate inorganic mercury TRV.

Organic mercury and inorganic forms of ingested mercury have distinctly different toxicokinetics. MeHg is readily absorbed in the gut (90-95 percent for mammals and nearly 100 percent in birds) while inorganic mercury is less readily absorbed (7-15 percent for mammals and only a few percent for birds) and tends to pass through the digestive tract and is excreted in feces (Wolfe et al. 1998). Once absorbed, MeHg acts primarily as a neurotoxicant and can readily cross the blood-brain barrier; inorganic mercury cannot. Both forms can exert immunotoxic and genotoxic effects on mammals with MeHg being a much more potent form (Wolfe et al. 1998). In birds, MeHg acts primarily as a potent neurological and embryo toxicant while inorganic salts tend to affect the kidneys.

Hill et al. (1981) observed that in studies with Japanese quail (Coturnix), MeHg was always more toxic than inorganic mercury (mercury chloride). Similarly, mammals are more sensitive to MeHg than inorganic forms of mercury in all studied cases (Wolfe et al. 1998; Eisler 1987).

Hill and Shaffner (1976) exposed Japanese quail to mercury chloride in the diet for one year at five dose levels (2 – 32 milligrams per kilogram [mg/kg]) and measured reproductive success. Adverse effects to fertility and hatchability were observed at the 8 mg/kg dose but were not apparent at 4 mg/kg. Thus, the 8 mg/kg dose was selected as a chronic LOAEL.

Food consumption rates and body weights were not specified in the study. A body weight of 0.15 kg and food ingestion rate of 0.0169 kg/day were assumed per Sample et al. (1996). Using these assumptions, a chronic LOAEL can be expressed as 0.9 mg/kg/day. This LOAEL was selected as the avian TRV for assessing risks to birds from incidental sediment ingestion of inorganic mercury. This TRV is listed in Table 6-6.

Auerlich et al. (1974) conducted a study with pregnant mink exposed to mercury chloride through the diet at a daily dosage of 7.39 mg/kg food for six months. They observed significant reductions in kit weight but no reductions in fertility or kit survival. Based on a food ingestion rate of 0.137 kg/d (Bleavins and Aulerich 1981) and an assumed body weight of 1.0 kg (USEPA 1993), a NOAEL based on a reproductive endpoint of 1.01 mg/kg body weight/day can be derived. Using the allometric scaling equations of Sample et al. (1996)³⁴ and an assumed raccoon body weight of 3.99 kg, a body-weight adjusted NOAEL TRV for raccoon is 0.715 mg inorganic mercury/kg body weight/day. Converting this NOAEL to a LOAEL based value using a factor of five results in a LOAEL TRV of 3.58 mg/kg body weight /day³⁵. This TRV was selected to assess risks to raccoon from incidental ingestion of inorganic mercury in sediment. This TRV is listed in Table 6-7.

6.3 Risk Characterization and Uncertainty Analysis

This section presents the risk characterization process and results for aquatic-dependent wildlife receptors based on the dietary-dose exposure. A deterministic risk characterization was completed to characterize risks to wildlife receptors. Specifically, a risk characterization of wildlife COPC-receptor pairs based on a dietary-dose exposure compared to a dietary-based TRV was conducted using a HQ approach that integrated exposure and effects data. All baseline risk characterization calculations are provided in Appendix C-4.

6.3.1 Characterizing PCB Risks

In the exposure and effects assessments, PCBs were considered both as a sum of total congeners, and as TEQ based on toxicity relative to 2,3,7,8-TCDD. For the baseline risk characterization, PCB risks are estimated using TEQ for dioxin-like PCB Congeners only.

³⁴ Scaling was performed for all mammalian TRVs in the approved BERA Work Plan.

³⁵ A NOAEL to LOAEL conversion factor of five was used in the approved BERA Work Plan TRV derivations.

USEPA (2008) notes that "large uncertainty may be introduced in assessing exposure and effects by assuming that congener profiles present in commercial mixtures used in toxicity tests (e.g., Aroclors) are representative of PCB profiles in weathered environmental samples (either exposure media or biota)." Unfortunately, effects information in the literature is primarily defined in terms of commercial mixtures. In addition, USEPA (2003) observed that while data currently available provide evidence to indicate that wildlife species are susceptible to effects from non-dioxin like PCBs, the existing data are inadequate for use in quantitative risk assessment. They conclude that the lack of data regarding dose-response relationships for non-dioxin like PCBs preclude performing quantitative ERAs for these congeners, either individually or cumulatively.

Based on this information, risk characterization for birds and mammals was not performed on a total PCB basis but rather was assessed for the dioxin-like PCBs using the TEQ approach. The HQs associated with the TEQ of dioxin-like PCBs were used to make conclusions regarding PCB risk and to identify COC. The uncertainty associated with this approach was evaluated as part of the uncertainty analysis by estimating HQs using the exposure and effects data for total PCBs that were developed in Sections 6.2 and 6.3.

6.3.2 Risk Estimates

6.3.2.1 Belted Kingfisher

Five COPCs were identified for the belted kingfisher in the BERA Work Plan: mercury, total PCB Congeners, hexachlorobenzene, 1,3-dichlorobenzene, and PCB Congener TEQ. The kingfisher diet included fish and invertebrates. Eighty-nine percent of the diet was assumed to be made up of fish, and 11 percent was assumed to be made up of invertebrates. No incidental sediment ingestion was assumed for this receptor. Only PCB Congener TEQ has a HQ that exceeds 1.0 (Table 6-8).

6.3.2.2 Carnivorous Birds

Three COPCs were identified for the carnivorous bird receptor group in the BERA Work Plan: mercury, total PCB Congeners, and PCB Congener TEQ. The carnivorous bird diet included fish, invertebrates, and terrestrial mammals. Thirty-three percent of the diet was assumed to be made up of fish, 33 percent was assumed to be made up of invertebrates, and 34 percent made up of mammals. Incidental sediment ingestion was included as 1 percent of

the total daily intake of food. None of the COPCs were retained as COCs for carnivorous birds based on the HQ calculations (Table 6-8).

6.3.2.3 Raccoon

Three COPCs were identified for the raccoon in the BERA Work Plan, mercury, total PCB Congeners, and PCB Congener TEQ. The raccoon diet included fish, invertebrates, terrestrial mammals, and plants. The diet was assumed to be 23 percent fish, 23 percent invertebrates, 30 percent plants, and 24 percent mammals. Incidental sediment ingestion was included as 9.4 percent of the total daily intake of food. None of the COPCs were retained as COCs for raccoons based on the HQ calculations (Table 6-8).

6.3.2.4 Spotted Sandpiper

Ten COPCs were identified for the spotted sandpiper in the BERA Work Plan: lead, mercury, total PAH, total PCB Congeners, hexachlorobenzene, hexachlorobutadiene, 1,3-dichlorobenzene, 1,4-dichlorobenzene, PCDD/PCDF TEQ, and PCB Congener TEQ. One-hundred percent of the spotted sandpiper diet was assumed to be made up of invertebrates. Incidental sediment ingestion was included at 18 percent of the total daily intake of food. PCB Congener TEQ has a HQ of 1.0; all other COPCs HQ were less than 1.0. (Table 6-8).

6.3.3 Uncertainty Analysis

As described in the BERA Work Plan, the emphasis of the uncertainty assessment is to focus on a quantitative assessment of uncertainty in the exposure and effects assessment for each receptor group. Sources of uncertainty in the wildlife risk assessment include uncertainty in the various exposure factors selected to describe the exposure of the various receptors (e.g., body weight), potential error in the grouping of prey items that constitute the diet, estimates of Site use, TRV selection, and risk characterization for PCBs. Specific sources of uncertainty are discussed in the following subsections.

6.3.3.1 Exposure Assessment

6.3.3.1.1 Prey Item Grouping

Prey items were grouped within the diet of the receptors based on apparent and statistical differences in concentrations among the different species. The groupings of these species may affect statistical estimates of the population mean (i.e., 95 UCL). In addition, it was assumed that these species constitute equal proportions of a receptor's diet of that particular group (e.g., fish). To evaluate this uncertainty, several different scenarios were examined with regard to prey item grouping and proportion of species within the diet. The following scenarios were examined:

- 1. The proportion of the species groupings in the baseline scenario were adjusted based on the relative biomass of the species collected during the 2011 field investigation³⁶ (Appendix C-5.1).
- 2. All species were pooled within their respective prey types (fish, shellfish) and a 95 UCL was estimated for each group (Appendix C-5.2).
- 3. All species were evaluated independently and a 95 UCL (or maximum value for species represented by less than six samples) was calculated (Appendix C-5.3).

The EPCs and adjusted prey proportions, along with resulting HQs, are summarized in Table 6-9. Overall, the uncertainty associated with the prey item grouping and resultant EPC values is low with regard to changes in the risk characterization (i.e., HQ). No COC would be dropped or added from the baseline scenario in Section 6.4 for any receptor.

6.3.3.1.2 Surface Weighted Averages

Surface weighted averages of COPCs were estimated and used to develop sediment EPCs for the BERA. Surface weighted averaging methods are described in Appendix B. Several different methods were evaluated and compared prior to final interpolation, resulting in a high degree of confidence in the interpolations and resulting surface weighted averages. Definition of the foraging zones within the Site were based on empirical analysis of bathymetric data and established MHHW and MLLW datum from nearby National Oceanic and Atmospheric Administration (NOAA) gauges.

 $^{^{36}}$ Biomass proportions for prey items are summarized in Appendix C-5 (Table C-5.1) and in Attachment 1.

The supratidal and intertidal habitat were included in the baseline SWAC calculations. To evaluate the uncertainty in the identification of these tidal zones, sensitivity analysis was performed by estimating the SWAC across all tidal zones (supratidal, intertidal, and subtidal) as part of the uncertainty analysis. Results of this analysis are provided in Appendix C-5.4. The HQs for SWACs calculated using all tidal zones are also provided in Table 6.9. The inclusion of the subtidal zone in the SWAC calculations has no apparent effect on HQs for wildlife.

The gunite channel, which is defined as supratidal and intertidal habitat, was included in the baseline SWAC calculations. However, there is a lower probability that wildlife will forage in this area due to the industrial nature of the surrounding area. To assess the uncertainty in foraging within this area, SWAC were calculated excluding this portion of the Site. Results of this analysis are provided in Appendix C-5.8 and HQ based on SWAC excluding the gunite channel are presented in Table 6-9. Including or excluding the gunite channel has no apparent effect on HQs for wildlife, most likely due to the smaller area relative to the entire Site.

An analysis was also conducted to evaluate whether areas of elevated COPC sediment concentrations would lead to higher EPCs than indicated by the Site-wide SWAC. This evaluation was conducted for the spotted sandpiper and carnivorous bird receptors³⁷ which have complete and significant incidental sediment ingestion pathways. For each COPC, sediment sample locations with a COPC concentration equal to or higher than the 90th percentile of the Site-wide concentrations were selected. Using GIS, a buffer was created centering the minimum reported home-range distance for a given receptor (Table 6-5) on these sample locations. A SWAC was calculated for each of the areas³⁸. The highest SWAC was then substituted into the risk equations as the sediment EPC to evaluate how it would affect the HQs. Use of the elevated concentration subarea SWACs as the sediment EPCs resulted in small increases (<0.1) in the HQs. A comparison of the Site-wide and elevated subarea HQs are summarized in Table 6-9. Figures depicting the hot spot areas for each COPC are included in Appendix C-5.9.

³⁷ The raccoon's minimum reported home-range was larger than the Site. For this reason, SWACs were not calculated for the raccoon.

³⁸ The hot spot SWAC was calculated for the intertidal and supratidal tidal zones.

Based on these conditions and analyses, the uncertainty in the assessment of exposure of receptors to sediment COPCs is considered low.

6.3.3.1.3 Terrestrial Animals and Plant Matter

Terrestrial animal prey (marsh rice rat) and plant matter mercury and PCB Congener TEQ concentrations were estimated using USEPA (1999) methods and included as appropriate in the carnivorous bird and raccoon risk estimation calculations. These dietary components contributed approximately 12 percent (mercury) and 3 percent (PCB Congener TEQ (Mammal)) to the raccoon HQs, and the marsh rice rat contributed approximately 18 percent (mercury) and 64 percent (PCB Congener TEQ (Avian)) to the carnivorous bird HQs.

The exposure assessment for these diet components were based on simple bioaccumulation models using USEPA derived BCFs, BAFs, and FCMs. Use of modeled COPC concentrations based on literature values in the diet of a receptor generally has a greater degree of uncertainty relative to empirical data (e.g., tissue data). However, it is expected that the uncertainty in the contribution of plant and terrestrial animal matter to the exposure of raccoons and carnivorous birds to Site COPC is not great enough to affect the risk conclusions for these receptors.

6.3.3.1.4 Area Use Factor

Predicted home ranges for representative receptors were evaluated based on literature data, empirical Site data (e.g., intertidal area, stream length), and qualitative habitat information. The 75th percentile of reported home ranges for representative receptors were compared to empirical Site data to develop AUFs for wildlife receptors. Given the poor quality habitat at the Site, this is considered a reasonable assumption.

Other exposure modifying factors such as migration, and life-cycle variables (e.g., breeding) were not considered in the baseline exposure assessment due to the uncertainty associated with quantifying such variables. However, many of the species within the avian guilds represented in the exposure assessment are migratory; particularly shorebirds and herons. In addition, several species (i.e., belted kingfisher) do not have suitable nesting sites within or adjacent to the Site itself. The result being that even resident species would likely move

off-Site for a significant portion of their life cycle. The aggregate effect of these factors would be to reduce exposure to the Site for most avian receptor groups evaluated in this BERA. Although this reduction in exposure due to exposure modifying factors has not been quantified, it would indicate that the exposure assessment is conservative with respect to these factors.

To evaluate the potential uncertainty associated with the AUF selected for the BERA, the AUF was varied for each receptor using the upper bound and lower bound home range estimates presented in Table 6-10. The lowest and highest average home range for each receptor was selected and compared to the available Site habitat to determine a range of AUFs (Table 6-10). HQs for each scenario were then determined (Table 6-11; Appendices C-5.5 and C-5.6, respectively).

Differences in HQ based on upper- and lower-bound estimates of the AUF are small. Three COPC-receptor pairs have an HQ > 1.0 based on the upper-bound AUF (smallest home range) but an HQ < 1.0 based on a lower-bound estimate of its AUF (largest home range). They include spotted sandpiper – PCB TEQ (HQ = 3.1-0.8), belted kingfisher – PCB TEQ (HQ = 3.0-1.7), and belted kingfisher – mercury (HQ = 1.0-0.58). Thus, the uncertainty associated with the range of AUF values used in the baseline risk characterization is considered low for most COPC-receptor pairs. While the AUFs for spotted sandpiper and belted kingfisher are representative of baseline conditions, the selection of the AUF term does have a significant effect on whether the baseline HQ for these COPC-receptor pairs fall above or below a HQ of 1.0.

6.3.3.2 Effects Assessment

By their nature, ERAs rely on limited toxicological information to evaluate the effects of contaminants in the environment. Testing is often done using laboratory or domesticated animals in a highly controlled environment which can create significant uncertainty in extrapolating those effects to wild species and populations. To address this uncertainty, effects data are chosen using conservative approaches in ERAs. However, the lack of robust datasets that allow comparisons among species (both wild and laboratory) means that assessing the degree of conservatism is a highly uncertain undertaking.

6.3.3.3 Risk Characterization

Risk characterization was performed using deterministic methods which do not provide any quantitative assessment of the probability that a risk threshold has been exceeded. Therefore, parameters in the risk assessment are set at conservative thresholds to allow conclusions regarding negligible risk to be made with confidence. As a result, conclusions regarding risk at or near a threshold of concern should be made considering the conservative bias of the assessment.

6.3.3.3.1 PCB Risk Characterization

As discussed in Section 6.3.1, risks from PCBs to wildlife receptors were characterized using exposure and effects assessments based on PCB TEQs rather than total PCBs due to the uncertainty in assessing risks based on environmental mixtures of dioxin like and non-dioxin like PCBs. To characterize the potential uncertainty associated with this approach, HQs were determined using exposure and effects data based on total PCBs. Exposure estimates (i.e., TDI) were estimated using total PCB Congeners (Section 6.2) and compared to effects values for total PCBs in the literature primarily based on commercial mixtures of PCB Aroclors. TRVs for birds and mammals were derived on this basis in Section 6.3. Risk estimates (HQs) were then determined for each receptor on a total PCB basis and compared to risk estimates based on PCB TEQs (Table 6-12; Appendix C-5.7).

HQs based on total PCB Aroclors are lower than those based on PCB TEQ. Reasons for this difference are not clear but may be due to a lower proportion of dioxin-like PCBs in the Aroclor mixture used to develop the total PCB TRVs relative to what is in Site media. Regardless, risk characterization based on PCB TEQs is more conservative than for total PCBs and is likely to be less uncertain (for the reasons described in Section 6.3.1). Therefore, conclusions regarding risk from PCBs to wildlife receptors have a higher degree of confidence than would be the case using total PCBs.

6.4 Risk Conclusions

Risk to avian and mammalian representative ecological receptors due to ingestion of COPCs from the Site has been evaluated and characterized in the previous sections. Results of this analysis indicate that for all COPCs, except PCBs on a TEQ basis, risks do not exceed a dietary threshold associated with adverse effects to the survival, growth, or reproduction of

wildlife communities or populations. Risks due to PCB TEQs exceed this threshold for spotted sandpiper and belted kingfisher (HQ of 1.0 and 1.7, respectively).

7 FISH RISK ASSESSMENT

This section presents the risk assessment for fish at the Site. Fish may be exposed to contaminants through direct contact with Site media (i.e., sediment and surface water) and/or ingestion of contaminated biota. Bioaccumulation based exposure is likely the most important pathway by which fish receptors are exposed to Site COPCs.

For bioaccumulative COPCs, empirical or estimated whole body tissue concentrations in fish were compared to literature-derived fish TRVs using a HQ approach. Surface water risk was evaluated using surface water benchmarks compared to Site surface water data using a HQ approach. The direct contact with sediment pathway was evaluated per the *COPC Report and Amendment* (Anchor 2008a, 2008b) using the results of the risk assessment for benthic invertebrates. Sediment COPCs identified for benthic invertebrates serve as a proxy for identifying direct sediment exposure COPCs for fish.

The details of the risk assessment for fish are presented as follows:

- Section 7.1 presents the assessment for bioaccumulation-based exposure, including:
 - An overview of the assumptions used to derive exposure concentrations, including how 95 UCLs were derived
 - A summary of the effects data. Effects data in this assessment are represented by tissue based lowest estimated dose (LOED) TRVs
 - The risk characterization results and uncertainty analysis
- Section 7.2 presents the assessment for surface water based exposure, including:
 - o An overview of the assumptions used to derive exposure concentrations
 - A summary of the effects data for surface water exposure, including an alternate water quality benchmark (WQB)
 - o The risk characterization results
- Section 7.3 presents the risk conclusions

7.1 Bioaccumulative COPCs Risk Assessment

Based on the screening performed by Anchor (2008a, 2008b, Anchor QEA 2011a), the potentially bioaccumulative COPCs included in the BERA for fish are:

- Mercury³⁹
- PCBs
- Selenium
- PCDD/PCDFs

Mercury and PCBs were identified as potentially bioaccumulative COPCs for both sediment and surface water. Selenium was identified as a potentially bioaccumulative COPC for fish based on its presence in surface water but was eliminated as a bioaccumulative COPC in sediment (Anchor QEA 2011a). PCDD/PCDFs were identified as a fish COPC based on its presence in surface water. Mercury, PCBs, and PCDD/PCDFs were evaluated using empirical tissue data collected from the Site in 2011, as described in Section 4.4. Similar to the wildlife risk assessment, exposure and effects assessment for PCBs was performed using both PCB TEQ and total PCBs, but risk was characterized based on PCB TEQ only. PCDD/PCDFs were evaluated on a TEQ basis. Selenium, which was not included in the tissue sampling program (Anchor QEA 2011a), was evaluated using estimated tissue concentrations based on a simple bioaccumulation modeling approach. Methods and results of these risk analyses are described in the following subsections.

7.1.1 Exposure Assessment

Fish tissue samples used in the fish exposure assessment were collected during a field investigation conducted in 2011 in accordance with the approved SAP attached to the BERA Work Plan. The results of this investigation are provided in a data report as an attachment to the BERA (Attachment 1). Prior to estimating tissue EPCs for the BERA, these results were evaluated using graphical and statistical methods to determine if different subgroups existed within the dataset that warranted independent treatment when developing EPCs (please review Section 6.1.2.1.1 for the detail on the prey grouping evaluation for the wildlife risk assessment). Similar groupings are used for the fish exposure estimate. Table 7-1 presents the fish grouping for fish tissue EPC calculations. For PCB TEQ and total PCBs, exposure

³⁹ Mercury in tissue was analyzed as total mercury (i.e., USEPA Method 7471A). It has been shown that the ratio of methylmercury to total mercury in estuarine fish can range between 71 – 97 percent (USEPA 2010). For purposes of the exposure assessment for fish, mercury in tissue was assumed to be 100 percent methylmercury.

estimates were evaluated for 1) Gulf killifish⁴⁰ and 2) all fish without Gulf killifish. For PCDD/PCDF TEQ, exposure estimates were determined for 1) Gulf menhaden; 2) pinfish, striped mullet, and Gulf killifish; and 3) sand seatrout (Appendix D-1.1). For mercury, exposure estimates were evaluated for 1) Gulf menhaden and 2) all fish without Gulf menhaden. In addition, exposure estimates were estimated by pooling all five species for each COPC. Finally, selenium, which was evaluated using estimated fish tissue concentrations, was not evaluated on a species specific basis (see Section 7.1.1.1).

For fish tissue EPCs, 95 UCLs were calculated using ProUCL Version 4.0 software (Appendix D-1.2). As with the wildlife exposure assessment, ProUCL software used for this analysis allowed detected and non-detected values to be considered and creates interpolated values for non-detects based on the perceived distribution of the detected concentrations. The ProUCL software analyzed the data to determine the most appropriate UCL and made a recommendation, which was then used as the EPC for the fish tissue risk calculations. Table 7-2 presents the fish tissue EPCs for mercury, total PCBs, PCB TEQs, and PCDD/PCDF TEQs.

7.1.1.1 Selenium Exposure Assessment

For selenium, the fish tissue EPC was estimated using a simple bioaccumulation model due to the lack of empirical selenium tissue data. The fish tissue EPC for selenium is based on the 95 UCL EPC for dissolved selenium in surface water at the Site and a literature based BCF. The dissolved concentration of selenium was selected as it best reflects the bioavailable fraction for fish (USEPA 2004). The literature based BCF selected is 129 liters per kilogram (L/kg) as WW tissue which was derived from the USEPA report, *Screening Level Ecological Risk Assessment Protocol for Hazardous Waste Combustion Facilities - Volume One* (USEPA 1999). USEPA recommends the use of this BCF, which is based on the geometric mean of the 12 BCFs listed in the report. This BCF was also adopted by TCEQ during their evaluation to identify potentially bioaccumulative COPCs in surface water (Reat 2008).

⁴⁰ PCBs had statistically significant higher concentrations in Gulf killifish than other species sampled. Similarly, Gulf menhaden had significantly higher mercury concentrations than other species.

The UCL 95 for dissolved selenium in Site surface water is 12 micrograms per liter (μ g/L) (Appendix D-1.2). The tissue concentration is calculated as follows:

$$Tissue\ Residue_{Fish} \frac{mg}{kg} = 95\ UCL_{surface\ water} \times BCF_{fish} \times CF$$

where:

95 UCL-surface water = Dissolved selenium 95 UCL at the Site, μ g/L

BCF-fish = USEPA recommended value, L/kg (WW tissue)

CF = Conversion factor is 0.001 (mg/1000 micrograms $[\mu g]$)

The estimated tissue EPC for selenium is 1.55 mg/kg WW. This EPC was used to estimate risks to fish using a HQ approach.

7.1.2 Tissue Based Effects Assessment

For fish, the measure of effect for this endpoint is a tissue-based threshold effect concentration. The primary source of tissue-based effects concentrations is the Environmental Residue-Effects Database (ERED) (http://el.erdc.usace.army.mil/ered/). The ERED is a compilation of literature effects data from over 2,000 studies published between 1964 and 2007 where tissue concentrations were measured in fish for the targeted effect. A copy of the most current database (last downloaded September 2011) was obtained from the U.S. Army Corps of Engineers (USACE). At present, the database contains only those results where an observed effect is reported as being directly related to a chemical in the tissue. The ERED includes the effects data presented in Jarvinen and Ankley (1999), as well as multiple other sources.

To identify an appropriate tissue-based threshold effect concentration, the geometric mean of reported LOED-equivalent fish tissue residues found in the ERED was used. The ERED dataset used for the BERA was evaluated using the following data requirements defined for the purposes of identifying an appropriate effects concentration for whole body fish tissue:

- Exposure route of 'ingestion' ('water', 'food', 'combined') or 'absorption'
- 'Whole-body' or 'carcass' tissue concentrations
- Endpoints of 'reproduction', 'growth', 'mortality', or 'survival'
- Exposure medium of 'sediment', 'water', 'water/food', 'food', or 'combined'

LOED for low effect results

The geometric mean of the derived or measured LOED values from the final list of studies was used as the TRV for the effects endpoint. This database was the primary source for selection of effects thresholds for the BERA. A review of the ERED database sources, as well as other literature sources, was performed during the BERA to identify any additional studies that could add to the body of information currently available for selecting measures of effect. Table 7-3 presents the Fish LOED TRVs. Effects on fish due to 2,3,7,8-TCDD exposure were used as the surrogate for effects due to exposure to PCB TEQ and PCDD/PCDF TEQ. Details on the data used to derive the mercury, PCB and PCDD/PCDF TEQ, total PCB, and selenium TRVs are presented in Appendices D-2 through D-5.

7.1.3 Risk Characterization and Uncertainty Analysis

This section presents the risk characterization and uncertainty analysis for fish receptors based on a bioaccumulation exposure. A deterministic risk characterization was conducted to characterize risks to fish receptors. A COPC with a HQ greater than 1.0 was retained as a COC.

7.1.3.1 Risk Characterization

All HQs for mercury, PCB TEQ, and PCDD/PCDF TEQ were less than 1.0 for each fish receptor grouping. For selenium, the HQ based on the estimated tissue EPC is 0.81. Therefore, none of the COPCs were retained as COCs for any of fish receptor groups based on the HQ calculations (Table 7-4).

7.1.3.2 Uncertainty Analysis

7.1.3.2.1 PCB Risk Characterization

As discussed in Section 6.3.1, there is significant uncertainty in assessing risks to fish and wildlife using total PCBs effects data and environmental mixtures of PCBs. In the previous section, risks to fish were characterized using PCB TEQ exposure and effects data. To assess the potential uncertainty associated with risks due to non dioxin-like PCBs, HQs were calculated using total PCB exposure and effects data. The HQs based on total PCBs for the different prey groupings defined in the exposure assessment were all less than 1.0

(Table 7-5). Thus, the uncertainty associated with concluding there is negligible risk to fish from PCBs is low.

7.1.3.2.2 Risk to Larger Size Class Fish

Risk characterization for bioaccumulative COPC is based on empirical tissue data for fish less than 15 cm in total length. There is some concern that risks to larger, higher trophic level fish may be higher due to biomagnification of COPC in prey items consumed by these higher trophic levels. Biomagnification through the food chain to higher trophic levels can be expressed using a biomagnification factor (BMF; USEPA 1999). Estimation of a BMF for higher trophic level organisms can be performed using the ratio of food chain multipliers⁴¹ (FCM) for the receptor and prey using the following formula (USEPA 1999):

$$BMF = \frac{FCM_{receptor}}{FCM_{prey}}$$

where:

BMF

= biomagnification factor between prey and receptor

 $FCM_{receptor}$

= food chain multiplier to account for dietary uptake of a COPC by

the receptor of interest

FCM_{prey}

= food chain multiplier to account for dietary uptake of a COPC by a

receptor's prey

USEPA (1999) provides FCM values for a range of trophic levels and K_{ow} values. To evaluate the uncertainty associated with risks to higher trophic level fish, BMFs were estimated using the FCM values in USEPA (1999) and used to estimate potential whole body COPC concentrations. It was assumed that the fish collected for the BERA were trophic level (TL) 2 and that larger size class fish would be TL 4 fish. The estimated BMFs were then applied to

⁴¹ Food chain multipliers are used to account for dietary uptake of a COPC for an ecological receptor based on its trophic level. The FCM is generally used in conjunction with a BCF and media specific COPC concentration (i.e., surface water) to estimate a receptor's or its prey's whole body tissue concentration.

the fish tissue EPC⁴² in Table 7-2 to estimate potential exposure of larger, higher trophic level fish using the following equation:

$$C_{TL4} = C_{TL2} \times BMF$$

where:

CTL4 = Estimated tissue concentration in larger TL 4 fish

 C_{TL2} = Empirically measured or estimated⁴³ tissue EPC for small size class (TL

2) fish

BMF = Biomagnification factor

Finally, the estimated COPC concentrations in TL 4 fish were compared to their respective whole body fish tissue TRVs in Table 7-3 to estimate an HQ for each COPC. Results are provided in Table 7-6. No HQ based on estimated whole body tissue concentrations for larger, higher trophic level fish exceed 1.0. Thus, the uncertainty that fish risk characterization based on empirically measured tissue in smaller size class fish are not protective of larger size class fish is considered low.

7.2 Surface Water Direct Contact Risk Assessment

Total PCBs were identified as the only direct contact COPCs for fish based on a comparison of Site surface water concentrations of total PCBs to the USEPA saltwater ambient water quality criteria (AWQC) (USEPA 1980). However, this criterion is for the protection of a fish eating mammals that use aquatic systems (USEPA 1985). This criterion has little ecological relevance for protection of fish populations at the Site.

Rather, the most appropriate approach to assess risks to fish populations from PCBs in surface water is through the body burden approach described in Section 7.1. Unlike some organic chemicals (e.g., PAHs), PCBs are not readily metabolized by fish and, therefore, can provide useful information on both exposure and toxicity. Field (2008) noted that tissue residues of PCBs play an integral role in assessing exposure via all pathways (i.e., sediment, surface water, and diet) and provides a time- and location-integrated assessment of exposure, which

⁴² The pooled tissue EPC was selected for this analysis. It was assumed that larger size class fish would feed opportunistically across all species of smaller size class fish.

⁴³ In the case of selenium

abiotic samples generally do not. Finally, early life-stage effects based on larval fish exposed to PCBs through maternal transfer of PCBs to eggs is one of the most sensitive toxicity endpoints for PCB effects on fish. These particular effects are related to the tissue body burden of fish rather than direct contact with PCBs in surface water. Thus, use of a tissue residue approach to assess risks to fish from all potential sources, including surface water, is a more robust approach to assessing risk. As a result, the bioaccumulative COPCs risk assessment in Section 7.1, which concluded that risks to fish from PCBs is negligible, demonstrates that risk to fish from PCBs in surface water is, therefore, negligible.

7.3 Risk Conclusions

Based on the assessment of bioaccumulation-based and surface water exposures for fish, there are no COCs for fish.

8 BENTHIC INVERTEBRATE RISK ASSESSMENT

This section presents the baseline risk assessment for benthic invertebrates at the Site. The CSM identified direct contact with sediment and surface water as the primary exposure pathways for benthic invertebrates. Risks to benthic invertebrates from sediment and surface water are evaluated in the following sections.

8.1 Sediment Risk Assessment

As described in the BERA Work Plan, the primary approach to evaluate potential risks to the benthic community included the development of a site-specific predictive benthic toxicity model for Patrick Bayou. Both sediment quality guidelines (SQGs) and benthic toxicity predictive tools have been developed from paired toxicity and chemistry data for national and site-specific programs to allow for the estimation of toxicological impacts. Several predictive toxicity models were evaluated in the development of the BERA in an attempt to define the relationship between site surface sediment chemistry and toxic responses in sediment bioassays.

The models evaluated generally fall into two groups: 1) generic sediment quality guideline quotient (SQG-Q) models (Long et al. 1998; Ingersoll et al. 2001; MacDonald et al. 1996); and 2) mechanistic models (USEPA 2003b, 2005, 2008). Each of these approaches are considered 'generalized' models in the sense that they rely on data from a compilation of sites and/or studies performed under a wide range of locations and conditions. Thus, they are intended to be representative of a broad range of conditions but may not account for potentially unique or confounding conditions that may be present at a site to which the model is extrapolated. To overcome this potential shortcoming, site-specific toxicity and chemical information was integrated into a generalized model framework to evaluate the applicability of a site-specific benthic toxicity predictive model for the Site.

Specifically, the performance of the generalized models was evaluated in the BERA Work Plan using 51 samples with co-located sediment chemistry and toxicity test data from the Site. In the BERA Work Plan, predictive model development focused on survival of *Leptocheirus plumulosus* in 10-day whole sediment bioassay tests as the effect endpoint for the model.

The predictive abilities of SQG-Q and Equilibrium Partitioning Sediment Benchmark Toxic Unit (ESBTU) models were evaluated by comparing the observed toxicity in the bioassay tests and the predicted toxicity based on chemistry data for each sample. Models that failed to meet the acceptability criteria identified in the Work Plan were dropped from further consideration. Only two models, the mean probable effect level-quotient (PEL-Q) and mean probable effect concentration-quotient (PEC-Q), met the acceptability criteria set forth in the Work Plan for one species, *Leptocheirus plumulosus*. Additional optimization steps were performed to improve the model performance. Ultimately, four COPCs were included in the optimized model based on *Leptocheirus plumulosus*: total PAHs, total PCBs, lead, and bis(2-ethylhexyl) phthalate (BEHP). Appendix E-1 summarizes the benthic toxicity modeling effort performed in the BERA Work Plan.

As noted above, this benthic toxicity model was identified as the the primary LOE to characterize the risk to the benthic community in the BERA Work Plan. However, during preparation of the BERA it became apparent that the toxicity model had several limitations that could not be resolved as initially conceived in the BERA Work Plan; leading to a significant amount of uncertainty in its utility and relevance as a primary line of evidence. Specifically, and in no particular order of priority, these limitations are: 1) the lack of reference area toxicity tests to account for naturally occurring stressors and the presence of confounding factors (e.g., salinity acclimation, porewater ammonia, etc.) that may result in an unknown but potentially significant amount of conservative bias in the toxicity attributed to Site COPCs; 2) the model, when applied to other test species, results in a high percentage of false positives, leading to a potentially significant amount of Site sediments to be falsely predicted as toxic; and 3) the analytical uncertainty in the PCB analysis in the toxicity sample dataset confounds the extrapolation of the sediment chemistry data to more recent samples. In this case, the uncertainty in the model, which is primarily a function of the uncertainty in the data used to develop the model, did not lead to a sufficient level of confidence in the model to define the magnitude and extent of risks to the benthic community using this LOE alone. Additional model evaluations performed as part of the BERA are described and presented in Appendix E-1 and reflects additional evaluations requested by USEPA.

As a result of these uncertainties in the predictive ability of the model identified during the development of the BERA, a more comprehensive alternative approach including several

LOE within a weight of evidence (WOE) framework was considered more appropriate for this Site. This WOE approach is described in detail in the following section.

8.1.1 Overview of the Benthic Invertebrate WOE Approach

Assessments of contaminant-related impacts to the benthic community in estuarine and marine environments may include chemical, toxicological, and biological evaluations to determine whether: 1) contaminants are present at the Site at concentrations that may be associated with adverse effects; 2) the sediment is toxic to benthic organisms; and 3) the benthic community structure is adversely affected. These multiple LOEs provide a WOE approach that can be used to estimate the likelihood of contaminant-related impacts. This type of approach was first proposed by Long and Chapman (1985) to define sediment quality and is often termed a sediment quality triad (SQT) approach. Chemical, toxicological, and biological evaluations are independent measures; therefore, any causal relationships can only be implied by correlations between the chemical, toxicological, and biological endpoints. For example, the dynamic nature of estuarine environments, where non-chemical related factors (e.g., salinity, temperature, dissolved oxygen) often drive the structure and health of the benthic community, can lead to difficulties in detecting chemical-related impacts in this system.

To evaluate benthic risk at the Site, three LOEs were considered: toxicity, chemistry, and benthic community data within a SQT approach. Specifically, risk associated with each independent LOE was determined using categorical criteria – low, indeterminate, or high – based on the relative response (i.e., toxicity, benthic community) or magnitude (i.e., sediment chemistry) of each LOE. Next, all three LOE were considered together for each sampling location using a WOE framework. It should be recognized that ecological risks occur along a continuum and there is not a quantifiable bright line for those risks. In cases where there may be a high degree of uncertainty in risk estimates, the term "indeterminate" has been used by USEPA Region 6 at other aquatic based sites (e.g., Calcasieau Estuary Remedial Investigation). In the FS, remedial alternatives will be evaluated that lower the overall site and sub-area risk for areas that are characterized as indeterminate or probable risk. The WOE approach is based on the following simple premises:

Consensus among the three LOE indicating adverse effects is a strong indication that
 COPC in sediments potentially pose a significant incremental risk to the benthic

- community. Likewise, consensus among the three LOE that adverse risks are not apparent is a strong indication that COPC in sediments do not pose an incremental risk to the benthic community.
- Lack of consensus among the three LOE indicate that COPC in sediments may or may
 not pose a significant incremental risk to the benthic community but there are
 insufficient data to conclusively identify the source of apparent adverse effects
 (environmental vs. anthropogenic). Thus, risks at these locations are indeterminate
 with respect to CERCLA related releases.

The following sections describe the dataset used to evaluate benthic risk, the evaluation and characterization of risk for each LOE, and the consensus based risk characterization using the WOE framework for each location within the Site.

8.1.2 Data Used for the WOE Approach

The benthic invertebrate risk assessment is based on co-located sediment toxicity, chemistry, and benthic community samples collected between 2000 and 2006 as part of historical investigations (Parsons Engineering et al. 2002, 2004; TCEQ n.d.).

The dataset includes the following:

- 12 stations (gunite and upstream samples excluded; see Figure 4-5)
- 30 discrete bulk sediment samples
- 30 sediment chemistry analyses⁴⁴
- 75 toxicity test results⁴⁵
- 23 benthic community evaluations

Stations located outside the site boundaries (2, 8, 9, 10) and those within the gunite channel portion of the Site were excluded from this risk assessment. The gunite stations (7, Y, R) were not included in the evaluation, as this section of the Site does not provide appropriate habitat for a benthic community.

⁴⁴ Not all chemical analytes were analyzed in all samples.

⁴⁵ Multiple species were frequently tested for each discrete sediment sample.

8.1.3 Benthic Toxicity Line of Evidence

To evaluate benthic toxicity as a LOE, test species survival in the 75 toxicity test results in the dataset listed above was used as the endpoint of concern. Data used in the benthic toxicity LOE are provided in Appendix E-2. A test result was considered to be toxic when the control-adjusted survival was less than 60 percent. The selection of a 60 percent control-adjusted survival threshold to define a toxic response was intended to account for the lack of reference area results, which would likely indicate survival levels significantly less than control tests, and to account for the fact that small differences between reference area test results and Site results are not likely to be ecologically significant and should not solely drive remedial decision-making. The control-adjusted 60 percent survival threshold should account for these adjustments. Species tested included: the amphipod, *Leptocheirus plumulosus*, the polychaete, *Neanthes arenaceodentanta*; the amphipod, *Ampelisca abdita*; and the mysid, *Americamysis bahia*⁴⁶.

To characterize toxicity for a specific location, the proportion of toxic samples was determined for each location. The proportion toxic was calculated as follows:

Proportion toxic = Toxic results / Total results

Proportion toxicity estimates for each sample are presented in Appendix E-2. Next, overall incremental risk to the benthic community due to sediment toxicity was categorized for each location using the proportion of toxic samples according to the following criteria:

- Probable Risk the proportion of toxic samples was equal to or exceeded fifty percent
- Indeterminate Risk the proportion of toxic samples was between 25 and 50 percent (exclusive)
- Low Risk the proportion of toxic samples was less than or equal to 25 percent

The results of this evaluation are presented in Table 8-1. For two stations, 4A and 6A, there is probable risk to the benthic community as the proportion toxic exceeded 50 percent. For station 3, the risk is indeterminate as the proportion toxic was less than 50 percent but greater than or equal to 25 percent. All other stations were characterizes as low benthic

⁴⁶ Test conditions (e.g., duration, etc.) are described in the BERA Work Plan.

toxicity risk (less than 25 percent toxic). The degree to which risks to the benthic community at locations identified using this LOE are caused by exposure to Site-related COPCs, and the degree to which exposure to COPCs or other stressors is causing measureable adverse responses in the benthic community, and are evaluated through the two other LOEs, as described below.

8.1.4 Sediment Chemistry Line of Evidence

The mean PEL-Q model developed in the BERA Work Plan was considered an appropriate tool to characterize the relative risk to the benthic community based on the 30 sediment chemistry analyses in the WOE dataset for the four COPC included in the model (PAHs, PCBs, lead, BEHP). To evaluate the mean PEL-Q as a LOE, a comparison of relative differences in optimized mean PEL-Q by station was conducted within the Site. For each sample and station, the relative percent difference from the median of the optimized mean PEL-Q was calculated:

$$PEL-Q \% \Delta = ([PEL-Q]-[PEL-Q] \text{ median}) / [PEL-Q] \text{ median} * 100$$

The relative percent difference from the median was used in lieu of an absolute value for the mean PEL-Q at a given station because of the uncertainty associated with attributing a specific mean PEL-Q value with a specific adverse response due to limitations in the benthic toxicity model discussed above (e.g., lack of a reference area, etc.). The use of an absolute value would imply a level of precision that does not exist. Sediment chemistry data and calculations of percent difference from median of the optimized mean PEL-Q estimates are presented in Appendix E-2.

The overall incremental risk to the benthic community due to exposure to sediment-based COPCs was categorized for each location according to the following criteria:

- Probable Risk relative optimized mean PEL-Q exceeded the 85th percentile (highest 15 percent) of locations within the Site
- Indeterminate Risk relative optimized mean PEL-Q was less than the 85th percentile but was equal to or greater than the Site-wide average
- Low Risk relative optimized mean PEL-Q was less than the Site-wide average

The results of this evaluation are presented in Table 8-2. Two stations, 4A and 6A, are characterized as probable risk based on sediment chemistry LOE. All other stations are characterized as low risk⁴⁷.

8.1.5 Benthic Community Line of Evidence

To evaluate benthic community as a LOE, relative Engle and Summers Biotic Index (ES-BI) scores were calculated for each of the 23 benthic community evaluations in the WOE dataset. Samples included in this LOE were collected from Patrick Bayou in 2000 and 2001. This difference in collection period may create an uncertainty in the interpretation of the results as salinity, dissolved oxygen and flushing flows may have been different before each monitoring event. As summarized in Appendix E-3, these physical factors have an important influence on benthic community condition. Nevertheless, in order to maximize the use of the available data, and not bias the data seasonally, and because the Patrick Bayou data were not being compared to off-Site reference areas, the average ES-BI score was then calculated for each of the twelve stations included in the analysis. Details on how the ES-BI scores are calculated and averaged is described in Appendix E-4. Next, overall incremental risk to the benthic community risk was categorized for each location according to the following criteria:

- Probable Risk the average relative ES-BI score was less than the 15th percentile (lowest 15 percent) of locations within the Site
- Indeterminate Risk the average relative ES-BI score was greater than the 15th percentile but was equal to or less than the Site-wide average
- Low Risk relative ES-BI score was greater than the Site-wide average

The results of this evaluation are presented in Table 8-3. Two stations, 6A and Q, are classified as representing a probable risk to the benthic community as the ES-BI scores were less than the 15th percentile of all stations. For four other stations, there is indeterminate risk. Finally, for all other stations there is low risk.

⁴⁷ Due to the relatively high values at Station 4A and 6A, no stations exceeded the average but were less than the 85th percentile (i.e., indeterminate risk).

8.1.6 Weight of Evidence Risk Characterization

To evaluate the overall risk to the benthic community the three LOEs were evaluated together using a consensus based approach. The overall by station-specific benthic invertebrate risk characterization is based on these LOE categorizations:

- Low Risk If all three LOE were characterized as low risk, then that location was characterized as low risk.
- Indeterminate Risk If none of the LOE was categorized as probable risk but were categorized as indeterminate or a mix of indeterminate and low risk, then that location was characterized as indeterminate risk.
- Probable Risk If any one of the three LOE was categorized as probable risk for a location, then that location was characterized as representing a probable risk to the benthic community.

The results of the assessment for all three LOE are presented in Table 8-4 and Figure 8-1. Five of 12 stations are categorized as low risk based on toxicity, chemistry, and benthic community condition. Four of 12 stations are categorized as indeterminate risk based on these LOE. Three of 12 locations (4A, 6A, and Q) demonstrate probable risk based on these LOE.

As noted in Section 8.1.1, ecological and human health risks occur along a continuum and there is not a quantifiable bright line for those risks. Remedial alternatives will be evaluated in the FS that lower the overall site and sub-area risk for areas that are characterized as indeterminate and probable risks. In concert with USEPA, quantitative tools will be developed to assess the ultimate risk reduction expected from a specific remedial alternative and that risk reduction score will be used as part of the effectiveness assessment for each alternative, along with USEPA's other FS criteria. In addition, the evaluation of surface water impacts as related to sediment concentrations will be a part of the evaluation of remedial alternatives.

8.1.7 Benthic Invertebrate Risk Assessment Uncertainty Analysis

To evaluate the potential uncertainty in the benthic risk characterization, sensitivity analysis was conducted on all three LOE. The analysis was performed as follows:

• The benthic toxicity LOE was evaluated using a higher (80 percent) survival

- threshold for identifying toxic and non-toxic samples
- The probable risk based on the mean PEL-Q was categorized using a more conservative threshold (75th percentile)⁴⁸
- The benthic community LOE and used more conservative threshold (25th percentile) to categorize a location as probable risk. In addition, this LOE considered the draft TCEQ Index (L. Broach, 2012) as a LOE.

8.1.7.1 Sensitivity Analysis for Benthic Toxicity

For the benthic toxicity sensitivity analysis, a higher threshold was used to evaluate benthic toxicity. In the baseline assessment, the threshold for identifying toxic and non-toxic samples was 60 percent control-adjusted survival. For the sensitivity analysis, an 80 percent control-adjusted survival threshold was used for identifying toxic and non-toxic samples. For comparison purposes, two other thresholds were also evaluated, 50 percent and 70 percent.

The results of the benthic toxicity sensitivity analyses are presented in Table 8-5. Selection of a higher survival threshold to categorize a test as non-toxic does not dramatically affect outcome. Only one station (T) would be re-categorized from low to indeterminate risk based on the WOE framework in Section 8.2.

8.1.7.2 Sensitivity Analysis for Optimized Mean Probable Effect Level-Quotient

For the baseline relative optimized mean PEL-Q risk characterization, the threshold for probable risk was the 15th percentile of the average station mean PEL-Q values. For the sensitivity analysis, the threshold was set to a more conservative value, the 25th percentile, to identify probable risk. The results of the sensitivity analysis are presented in Table 8-6. Selection of a higher threshold to identify probable risk has minimal effect on outcome for the relative optimized mean PEL-Q. One station (3) would be re-categorized from indeterminate to probable risk.

⁴⁸ The upper 25th percentile.

8.1.7.3 Sensitivity Analysis Benthic Community

A second benthic invertebrate community index was selected for a sensitivity analysis to evaluate Site benthic invertebrate community data. A BI currently under development by TCEQ (L. Broach 2012) for tidal streams in Texas was used in this evaluation. The TCEQ BI is in draft form and has not yet been peer-reviewed. However, this index was used here as requested by USEPA. This index has six different metrics that are described in Appendix E-4. The benthic community data were scored using the approach described by TCEQ (see Appendix E-4). However, the evaluation of a score for a station was not based on a comparison to a reference condition (e.g., stressed or not stressed) but in the context of the other scores at the Site (a relative analysis).

In addition, for the sensitivity analysis of the benthic community, the ES-BI and TCEQ relative benthic indices were also evaluated using a more conservative threshold, the lower 25th percentile, to identify probable risk. The results of the sensitivity analyses are presented in Table 8-7. The selection of a higher threshold, 25 percent, for the ES-BI affected one station. Station U would be re-categorized from indeterminate to probable risk.

The addition of TCEQ BI adds some uncertainty to the assessment. Two stations (U and 5) would be re-categorized from indeterminate to probable risk. One station (V) would be re-categorized from indeterminate to low risk. One station (Q) would be re-categorized from probable to indeterminate risk. In addition, there is no difference between 15 percent and 25 percent threshold for TCEQ BI. The differences between the ES-BI and TCEQ BI scores may be due to the different factors that have been demonstrated to affect benthic community structure and the metrics used to evaluate that structure. Direct causal relationships are typically not evident (Long and Chapman 1985; Gaston and Edds 1994). However, the influence of grain size and salinity on the distribution of benthic infauna in soft-sediments has been well documented and may have an influence on the metrics used with the two BI's (see Appendix E-3).

8.2 Surface Water Risk Assessment

Benthic invertebrates may be directly exposed to surface water at the sediment-water interface or, for some burrowing organisms, through irrigation of their burrows. Risks to

benthic invertebrates exposed to surface water were assessed by comparing the concentration of Site surface water COPCs to effects values using a HQ approach.

Surface water data collected in 2009 were evaluated in the BERA Work Plan to identify COPCs. Total PCBs was the only surface water COPC identified for benthic invertebrates based on the initial and refined risk-based screening.

8.2.1 Exposure Assessment

The exposure assessment for surface water COPCs includes surface water sampling conducted in 2009 and 2011. Samples were collected at eight stations (see Figure 4-2). Discrete samples were collected at mid-depth and within six inches of the bottom. Stations sampled in 2009 (n=4) included two sampling events. One round of sampling was conducted during low-tide, while the second was collected on an outgoing tide to capture any potential upstream flow and to avoid potential flow-reversal conditions from the HSC. Stations sampled in 2011 (n=4) were sampled at mid-depth during a single event. The total fraction (i.e., unfiltered) samples were analyzed for PCB Congeners (n=209). A more detailed discussion of the surface water dataset is provided in Section 4.4.2.

The EPC for total PCBs was calculated as the 95 UCL of the surface water dataset. Total PCB Congeners were calculated using all 209 congeners with non-detect results set at one-half the DL. Using USEPA's ProUCL software (Version 4.0; (USEPA 2002), the recommended total PCB UCL 95 was calculated to be 193 nanograms/liter (ng/L). Calculations are summarized in Appendix E-5.

8.2.2 Effects Assessment

The national recommended AWQC saltwater criteria continuous concentration (USEPA 2012) of 30 ng/L⁴⁹ was initially selected as the effects value for total PCBs in surface water in the BERA Work Plan. The narrative intent of the national recommended AWQC is to be protective of 95 percent of all aquatic receptors, including benthic invertebrates. However, the PCB AWQC for aquatic receptors (USEPA 1980) was derived using a tissue-based final

⁴⁹ The Texas Surface Water Quality Standard for PCBs (TCEQ 2012) has adopted USEPA's recommended AWQC CCC.

residue value approach intended to be protective of mink exposed to PCB through ingestion of aquatic biota (e.g., fish and invertebrates). Thus, risk estimates using this effects value are associated with substantial uncertainty for direct effects on benthic invertebrates.

Fuchsman, et al. (2006) derived an alternative WQB specifically for aquatic invertebrates based on published data obtained from USEPA's online Aquatic Toxicity Information Retrieval (AQUIRE) database. Although derivation of WQBs and criteria typically incorporates toxicity data for both fish and invertebrates, the authors limited their analysis to invertebrates due to concerns over differences in the mechanism(s) of toxicity exerted by PCBs in fish and invertebrates. While both groups are susceptible to PCBs via narcosis, invertebrates lacking the Ah receptor are not as susceptible to co-planar PCBs, which can act as ligands for the Ah receptor found in most vertebrates, resulting in production of toxic metabolites or alteration of gene transcription. The authors applied the USEPA guidelines (Stephan, et al. 1985) to identify appropriate data and to develop a final acute value (FAV). Aroclor 1254 was selected as the commercial PCB mixture for their analysis due to: 1) the availability of data meeting USEPA guidelines; and 2) generally greater toxicity relative to less or more-chlorinated Aroclor mixtures. They derived an FAV of 6,960 ng/L for Aroclor 1254. Again, using the USEPA guidelines, an acute-to-chronic ratio (ACR) of 13 was derived using paired acute and chronic tests. Finally, they calculated a final chronic value (FCV) of 540 ng/L by dividing the FAV by the ACR. This FCV is nearly equal to the lowest chronic effect concentration reported for mortality and is lower than most chronic values identified for reproduction, growth, or development. Therefore, the authors concluded, "the FCV of [540 ng/L] is a reasonable estimate of a chronic toxicity threshold for sensitive invertebrate taxa."

This FCV, based on aquatic invertebrate toxicity data, represents a more ecologically relevant effects threshold for benthic invertebrates compared to the USEPA WQC. As such, this alternative TRV was used to derive surface water HQs and identify COC for benthic invertebrates.

8.2.3 Surface Water Risk Characterization

The surface water exposure risk characterization for benthic invertebrates was conducted by comparing the 95 UCL to the alternative TRV using a HQ approach. Using the alternative

TRV specific for benthic invertebrates the HQ is 0.4. In fact, none of the surface water samples from the Site exceeded the alternative effects value for benthic invertebrates.

8.2.4 Surface Water Uncertainty Analysis

Uncertainty in the benthic invertebrate risk assessment for exposure to surface water includes uncertainty in the exposure and effects assessment. Sampling for the exposure assessment consisted of two different sampling events. The first event involved multiple locations, depths, and tidal cycles and is considered representative of Site conditions. The second event was based on a biased design to target areas with potentially elevated concentrations based on the results of sediment testing. Thus, the data are considered representative but biased conservatively. Therefore, the uncertainty associated with the effects assessment is considered low.

During the initial screen of the surface water data in the BERA Work Plan, conservative screening values were selected to identify COPCs. The only COPC to exceed its screening benchmark was PCBs. Thus, the selection was conservative and the uncertainty associated with the screening is low. The TRV selected in the effects assessment is based on a comprehensive review of available studies on the effects of PCBs to benthic invertebrates. The derivation of the TRV was based on methods and principles similar to the derivation of water quality criteria (WQC). Thus, the derivation of the TRV is considered robust and conservative. As a result, the uncertainty associated with the selection of the benthic invertebrate surface water TRV for PCBs is considered low as well.

There is some potential uncertainty associated with the comparison of a PCB Aroclor based TRV to an EPC based on total PCB congeners due to differences in the analytical basis for the two values. The FAV and ACR values identified by Fuchsman et al. (2006) to calculate the TRV are based on Aroclor 1254 and Aroclor 1242, 1248, and 1254; respectively. These Aroclors, particularly Aroclor 1254 and 1248 are typically dominated by tri-chlorinated and tetra-chlorinated PCB congeners (Frame, et al. 1996). Surface water data for the Site used in the exposure assessment (Appendix A) indicates that tri-chlorinated and tetra-chlorinated PCB congeners predominate. In addition, surface sediments and historical surface water data analyzed for PCB Aroclors frequently had only detectable levels of Aroclor-1248 (and to a lesser extent, Aroclor-1254). This suggests that the congener profiles of the effects data used

to derive the TRV and the exposure assessment for surface water are comparable and would not be expected to contribute a significant degree of uncertainty in the risk characterization for PCBs in surface water.

Overall, the uncertainty associated with the surface water risk assessment for benthic invertebrates is considered low and decisions regarding risk for this receptor-pathway pair can be made with an acceptable degree of confidence.

8.3 Risk Conclusions

This section presents overall conclusions regarding risks to benthic invertebrates from direct contact with sediment using a WOE approach and direct contact with surface water using a deterministic comparison to TRVs.

8.3.1 Sediment

The condition of the benthic community at the Site reflects primarily the temporal and spatial gradients and dynamics of non-chemical stressors that are present in a tidal stream located in a highly developed, industrialized watershed, as well as CERCLA related releases. The Site has been significantly modified from its natural state (e.g., channelized, armored), which has likely increased the variability of many non-chemical stressors (e.g., salinity, dissolved oxygen, temperature). Upstream urbanization contributes to the benthic conditions at the Site through organic enrichment and increased flow during storm events. Thus, interpretation of the LOE for risks to the benthic community are problematic with respect to environmental conditions vis-à-vis CERCLA related releases.

Nonetheless, areas of probable risk to the benthic community have been identified using a WOE approach. Higher mean PEL-Q values are associated with areas of probable risk. Of the four COPC (PCBs, PAHs, lead, BEHP) represented by the mean PEL-Q, the contribution of PCBs generally exceeds those of other COPC and, in the case of stations identified as probable risk, accounts for a significant proportion of the mean PEL-Q value (Figure 8-2). This suggests that PCBs are the primary risk driver for sediment benthic risk. As such, PCBs are identified as sediment COC for the benthic community.

8.3.2 Surface Water

Risk to benthic invertebrates from PCBs in surface water is determined to be negligible and no surface water COCs are identified.

9 AQUATIC PLANTS, REPTILES, AND AMPHIBIANS UNCERTAINTY DISCUSSION

Risks to aquatic plants, reptiles, and amphibians were not addressed quantitatively per the problem formulation and ecological CSM (Anchor QEA 2011a). In general, relevant exposure and effects data for these receptors groups are not available to quantify risk. Rather, a qualitative assessment of the uncertainty associated with potential risks was performed for these groups.

9.1 Aquatic Plants

Evaluating risks to aquatic plants is uncertain primarily due to: 1) the lack of standardization in test endpoints available for aquatic plants (e.g., test duration, effect type, etc.); and 2) unknown sensitivity of test species to a chemical relative to other aquatic plant species (USEPA 2010). Relationships between sublethal biochemical effects observed in many plant studies (e.g., carbon fixation, lipid peroxidation, etc.) and typical assessment endpoints regarding survival, growth, or reproduction are poorly defined. Thus, the pool of ecologically relevant studies is quite limited (USEPA 2010). In addition, many of the available toxicity studies with aquatic plants are limited to macroalgae or Lemna sp. (i.e., duckweed; a free-floating aquatic plant). In addition, very few tests have been performed with marine or brackish water species most relevant to the Site. A review performed in 2008 of the nearly 23,000 aquatic plant entries in the ECOTOX database (USEPA 2010) indicated that less than 10 percent were performed using submerged grasses or vascular plants (other than Lemna sp.). Similarly, only 14 percent were performed using marine species (primarily marine algae). Finally, there are no widely accepted approaches to developing or validating extrapolation factors that may be used to address interspecies sensitivity or differences in test endpoints.

Thus, while there are effects thresholds for aquatic plants reported in the literature, the uncertainty associated with applying those values to quantitatively assess risks to aquatic plant communities at the Site is high enough to preclude a meaningful characterization of risk. However, many of the WQB (i.e., state and federal WQC) used to identify COPCs for other receptor groups (e.g., fish, benthic invertebrates) are based on criteria considered protective of aquatic plants. Thus, risk conclusions and risk management decisions based on other receptor groups are likely to be protective of aquatic plants.

9.2 Reptiles and Amphibians

Reptiles and amphibians were not selected as receptors of concern due to the insufficient toxicological data for detailed quantitative evaluation of potential adverse effects on this group. In addition, amphibians are not an ecologically relevant receptor group given the brackish nature of the Site; few amphibians that are potentially present in this area are tolerant of brackish waters. Thus, exposure to Site COPCs is considered negligible, risks are therefore negligible as well, and the uncertainty associated with this conclusion is considered low.

As ectothermic vertebrates, reptiles have different anatomies, physiologies, metabolisms, behaviors, exposures, and responses to environmental contaminants compared to birds and (even more distantly related) mammals that would, by necessity, be used as surrogates in the BERA due to the lack of availability of toxicity reference values for reptiles. This is recognized as a potentially important uncertainty, as the relatively sedentary behavior of many reptiles may increase their vulnerability to exposures (compared to birds and mammals). However, because reptile metabolic demands are relatively lower than birds or mammals, exposure through ingestion would be expected to be less than similar sized birds or mammals. Finally, dermal exposure may be a relatively more important exposure route for reptiles than birds or mammals. Weir et al. (2010) observed that reptiles may receive a higher daily dose of lipid soluble contaminants due to the lack of feathers and fur and the higher lipid content of reptile skin. Unfortunately, there are no means to evaluate this statement quantitatively with regards to assessing exposure to Site COPCs.

In conclusion, risks associated with most wildlife COPCs are considered negligible for reptiles given the negligible risks for birds and mammals. Because some avian receptors have estimated exposures to PCBs at the Site in excess of potentially adverse effects levels (i.e., LOAEL TRVs), it may also be expected that some reptiles, assuming similar exposures, would have similar risks. However, there are no means to quantify potential risks to reptiles, given the lack of relevant exposure and effects information for PCBs.

10 CONCLUSIONS AND RISK MANAGEMENT RECOMMENDATIONS

This section summarizes the results of the risk characterization and conclusions to identify Site COCs that might cause potentially unacceptable ecological risk and to provide risk management recommendations based on magnitude and extent of risks and their associated uncertainties.

Potential unacceptable ecological risks are associated with protection and maintenance of piscivorous and shorebird populations 50 . Baseline PCB TEQ HQs for these receptor groups are equal to 1.0 and 1.7 for spotted sandpiper and belted kingfisher, respectively. However, uncertainty analyses indicate that HQs for these COPC-receptor pairs may be above or below the threshold of concern (HQ = 1.0) depending on the assumptions used to characterize risk. Thus, within the ranges of exposure and effects variables evaluated, risks may or may not exceed a threshold of concern for individuals exposed to PCBs in Site media.

Using a WOE approach that included three LOE (sediment chemistry, sediment toxicity tests, and benthic community data), PCBs were identified as COC for the benthic community. Areas of probable risk were identified based on a consensus of the three LOE evaluated. However, the available data did not support a quantitative estimate of the magnitude of risk within these areas.

In summary incremental risk to benthos and wildlife are driven primarily by exposure to PCBs. Although no specific risk management recommendations are provided herein, risk management based on risks to ecological receptors should be considered within the overall context of other risk management considerations (e.g., water quality standards) during the FS.

⁵⁰ No potentially unacceptable risks from Site COPCs were identified for carnivorous birds, raccoons, fish populations, or benthic invertebrate populations.

11 REFERENCES

- Anchor Environmental, LLC (Anchor), 2006. *Preliminary Site Characterization Report.*Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. May 2006.
- Anchor, 2007. *Quality Assurance Project Plan.* Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group.
- Anchor, 2007a. Remedial Investigation Work Plan Remedial Investigation/Feasibility Study.

 Patrick Bayou Superfund Site, Deer Park, Texas. Prepared for USEPA and Patrick

 Bayou Joint Defense Group. January 2007.
- Anchor, 2007b. Work Package 2 Vertical Profiling Hydrodynamic Field Data Collection and Contaminant Source Evaluation. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. April 2007.
- Anchor, 2007c. Draft Supplemental Work Plan. Additional Hydrodynamic Data Collection. Geotechnical Data Collection. Sediment Accretion Evaluations. Porewater Sampling and Estimation of Distribution Coefficients. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. May 2007.
- Anchor, 2008a. Selection of Contaminants of Potential Concern for Ecological Risk

 Assessment Technical Memorandum. Patrick Bayou Superfund Site. Deer Park,

 Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. May 2008.
- Anchor, 2008b. Amendment to the Selection of Contaminants of Potential Concern for Ecological Risk Assessment Technical Memorandum. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. October 2008.
- Anchor, 2008c. *Draft Mixing Zone Evaluation Work Plan.* Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. October 2008.
- Anchor QEA, 2009a. Surface Sediment Contaminant of Potential Concern Delineation and Surface Water Sampling and Analysis Plan. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. September 2009.

- Anchor QEA, 2009b. *Patrick Bayou Sediment Mixing-Zone Layer Study Memorandum*. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. June, 2009.
- Anchor QEA, 2010. Sediment and Surface Water Contaminant of Potential Concern

 Delineation Report. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for

 USEPA and Patrick Bayou Joint Defense Group. May 2010.
- Anchor QEA, 2011a. Final Baseline Ecological Risk Assessment Work Plan. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. May 2011.
- Anchor QEA, 2011b. Final Upstream Patrick Bayou Characterization Sampling and Analysis Plan. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. June 2011.
- Anchor QEA, 2011c. *Sediment Transport Modeling Report, Patrick Bayou.* Prepared for Patrick Bayou Joint Defense Group. September 2011.
- Anchor QEA, 2012. *Upstream Patrick Bayou Characterization Data Report.* Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. February 2012.
- Anchor QEA, 2013 (in progress). *Draft Patrick Bayou Remedial Investigation Report*, Patrick Bayou Superfund Site, Deer Park, Texas.
- Aulerich, R. J., R.K. Ringer, and S. Iwamoto, 1974. Effects of dietary mercury on mink. *Arch. Environ. Contam. Toxicol.* 2: 43-51.
- Beyer, N., E. Connor, and S. Gerould, 1994. Estimates of Soil Ingestion by Wildlife. *Journal of Wildlife Management* 58:375-382.
- Bleavins, M.R., and R.J. Aulerich, 1981. Feed Consumption and Food Passage Time in Mink (Mustela vison) and European Ferrets (Mustela putorius furo). *Lab. Anim. Sci.* 31:268–269.
- Brooks R.P., and W.J. Davis, 1987. Habitat Selection by Breeding Belted Kingfishers (Ceryle alcyon). *American Midland Naturalist* 117:63–70.

- Butler R.W., P.C.F. Shepherd, and M.J.F. Lemon, 2002. Site fidelity and local movements of migrating western sandpipers on the Fraser River estuary. *Wilson Bulletin* 114:485–490.
- Colwell, M.A., R.L. Mathis, L.W. Leeman, and T.S. Leeman, 2002. Space use and diet of territorial long-billed curlew (Numenius americanus) during the non-breeding season. *Northeastern Naturalist* 83:47–56.
- Cornwell G.W., 1963. Observations on the Breeding Biology and Behavior of a Nesting Population of Belted Kingfishers. *The Condor* 65:426–431.
- Custer, C.M., and J. Galli, 2002. Feeding habitat selection by Great Blue Herons and Great Egrets nesting in east central Minnesota. *Waterbirds* 25: 115-124.
- Dauer, D.M., 2003. Biological criteria, environmental health and estuarine macrobenthic community structure. *Marine Pollution Bulletin*, 26:5, pp. 249-257.
- Dobberstine, J., J. Horne, L. Brzuzy, and C. Howard, 2007. Identifying suitable reference sites for impacted sites along the Houston Ship Channel. Proceedings of the Eighth Biennial State of the Bay Symposium, Galveston Bay Estuary Program.
- Dumas J.V., 2000. Roseate Spoonbill (Platalea ajaja). A. Poole, F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/490/articles/introduction.
- Eisler, R., 1987. *Mercury Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review.*Biological Report 85(1.10). Laurel, MD: U.S. Fish and Wildlife Service.
- Engle, V.D., and J.K. Summers, 1999. Refinement, Validation, and Application of a Benthic Condition Index for Northern Gulf of Mexico Estuaries. *Estuaries* 22: 624-635.
- Environment Canada, 2002. Canadian Tissue Residue Guidelines for the Protection of Consumers of Aquatic Life: Methylmercury. Scientific Supporting Document. Ecosystem Health: Science-based Solutions Report No. 1-4. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada. Ottowa.
- Field, J., 2008. *Use of Tissue Residue Data in Exposure and Effects Assessment for Aquatic Organisms*. In: National sediment bioaccumulation conference proceedings. Bethesda, MD: USEPA.

- Fisher, C.E., 2007. Ecology of raccoons in Central Alabama: A study of survival, space use, and habitat selection.
- Frame, G.M., J.W. Cochran, and S.S. Boewadt, 1996. Complete PCB congener distributions for 17 Aroclor mixtures determined by 3 HRGC systems optimized for comprehensive, quantitative, congener-specific analysis. *J. High Resol. Chromatogr.*, 19:657-668.
- Frederick P.C., 1997. Tricolored Heron (Egretta tricolor). A. Poole, F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/306/articles/introduction
- Fuchsman, P.C., T. R. Barber, J.C. Lawton, K. B. Leigh, 2006. An evaluation of cause-effect relationships between polychlorinated biphenyl concentrations and sediment toxicity to benthic invertebrates. *Environmental Toxicology and Chemistry* 25:2601–2612.
- Gaston, G.R., and K.A. Edds, 1994. Long-term study of benthic communities on the continental shelf off Cameron, Louisiana: A review of brine effects and hypoxia. Gulf Research Reports 9:57-64.
- Gibbs, J.P., S. Woodward, and M.L. Hunter, 1987. Determinants of great blue heron colony distribution in coastal Maine. *Auk* 104:38–47.
- Gibbs, J.P., 1991. Spatial relationships between nesting colonies and foraging areas of great blue herons. *Auk* 108:764–770.
- Haig S.M., L.W. Oring, P.M. Sanzenbacher, and O.W. Taft, 2002. Space Use, Migratory Connectivity, and Population Segregation among Willets Breeding in the Western Great Basin. *The Condor* 104:620–630.
- Heath, J.A., P. Frederick, J.A. Kushlan, and K.L. Bildstein, 2009. White Ibis (Eudocimus albus), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Laboratory of Ornithology. Available form: http://bna.birds.cornell.edu/bna/species/009.
- Heinz, G.H., 1979, "Methylmercury: Reproductive and Behavioral Effects on Three Generations of Mallard Ducks," *J. Wildl. Manage*. 43(2):394-401.
- Hill, E.F., and C.S. Shaffner, 1976. Sexual Maturation and Productivity of Japanese Quail Fed Graded Concentrations of Mercuric Chloride. *Poult. Sci.* 55:1449–1459.

- Hill, E.F., 1981. Inorganic and Organic Mercury Chloride Toxicity to *Coturnix*: Sensitivity Related to Age and Quantal Assessment of Physiological Responses, Ph.D. Thesis, University of Maryland, College Park. 221 p.
- Ingersoll, C.G., D.D. MacDonald, N. Wang, 2001. Predictions of sediment toxicity using consensus-based freshwater sediment quality guidelines. *Archives of Environmental Contamination and Toxicology* 41:8–21.
- Jarvinen, A.W., and G.T. Ankley, 1999. Linkage of Effects to Tissue Residues: Development of a Comprehensive Database for Aquatic Organisms Exposed to Inorganic and Organic Chemicals. SETAC Press: Pensacola, Florida, 364 pp.
- Klima J., and J.R. Jehl Jr., 1998. Stilt sandpiper (Calidris himantopus) [online]. In: A. Poole, editor. The birds of North America online. Ithaca, NY: Cornell Laboratory of Ornithology.
- Lanctot R.B., and C.D. Laredo, 1994. Buff-breasted Sandpiper (Tryngites subruficollis). A. Poole and F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/091/articles/introduction
- Long, E.R., and P.M. Chapman, 1985. A sediment quality triad: Measures of sediment contamination, toxicity and infaunal community composition in Puget Sound. *Marine Bulletin* 16:405-415.
- Long, E.R., L.J. Field, D.D. MacDonald, 1998. Predicting toxicity in marine sediments with numerical sediment quality guidelines. *Environmental Toxicology and Chemistry* 17:714–727.
- Long, E.R., C.G. Ingersoll, and D.D. MacDonald, 2006. Calculation and Uses of Mean Sediment Quality Guideline Quotients: A Critical Review. Environmental Science & Technology, 40:1726-1736.
- MacDonald, D.D., R.S. Carr, F.D. Calder, E.R. Long, and C.G. Ingersoll, 1996. Development and Evaluation of Sediment Quality Guidelines for Florida Coastal Waters. *Ecotoxicology* 5:253–278.
- MacDonald, T.A., B.J. Burd, V.I. MacDonald, and A. van Roodselaar, 2010. Taxonomic and feeding guild classification for the marine benthic macroinvertebrates of the Strait of Georgia, British Columbia. *Canadian Technical Report of Fisheries and Aquatic Sciences.* 2874: iv + 63 p.

- Macwhirter B., P. Austin-Smith, Jr., D. Kroodsma, 2002. Sanderling (Calidris alba). A. Poole F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/653/articles/introduction.
- Mathisen, J., and A. Richards, 1978. Status of great blue herons on the Chippewa National Forest. *Loon* 50:104–106.
- Mazeika, S., P. Sullivan, M.C. Watzin, and W.C. Hession, 2006. Differences in the reproductive ecology of belted kingfishers (Ceryle alcyon) across streams with varying geomorphology and habitat quality. Waterbirds: *The International Journal of Waterbird Biology*, 29:258–270.
- Mccrimmon, Jr. D.A., J.C. Ogden, and G.T. Bancroft, 2001. Great Egret (Ardea alba). A. Poole, F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/570/articles/introduction.
- Parsons K.C., and T.L. Master, 2000. Snowy Egret (Egretta thula). A. Poole, F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/489/articles/introduction.
- Parsons et al., 2002. Assessment of Sediment Toxicity and Quality in Patrick Bayou, Segment 1006, Harris County, Texas. Prepared for Patrick Bayou TMDL Lead Organization.
- Parsons et al., 2004. Assessment of Sediment Toxicity and Quality in Patrick Bayou, Segment 1006, Harris County, Texas. Prepared for Patrick Bayou TMDL Lead Organization.
- Patton, J.F., and M.P. Dieter, 1980 "Effects of Petroleum Hydrocarbons on Hepatic Function in the Duck," *Comp. Biochem. Physiol. Part C: Comp. Pharmac.* 65(1):33-36.
- Peifer, R.W., 1979. Great blue herons foraging for small mammals. *Wilson Bulletin* 91:630–631.
- Prose, B.L., 1985. Habitat suitability index models: belted kingfisher. *U.S. Fish and Wildlife Service Biological Report 82*.
- Reat, Vickie, 2008. Re: Document availability for evaluation of bioaccumulative COC with water quality standards. Texas Commission on Environmental Quality.
- Rodgers, Jr. J.A., and H.T. Smith, 1995. Little Blue Heron (Egretta caerulea). A. Poole, F. Gill, editors. The Birds of North America Online [Internet]. Available from: http://bna.birds.cornell.edu/bna/species/145/articles/introduction.

- Sample, B.E., D.M. Opresko, and G.W. Suter II, 1996. *Toxicological Benchmarks for Wildlife: 1996 Revision*. Oak Ridge, TN: Oak Ridge National Laboratory.
- Sanzenbacher P.M., and S.M. Haig, 2002. Residency and Movement Patterns of Wintering Dunlin in the Willamette Valley of Oregon. *The Condor* 104:271–280.
- Stephan et al. 1985. Guidelines for Deriving Numerical National Water Quality Criteria For The Protection of Aquatic Organisms and Their Uses. Office of Research And Development, Environmental Research Laboratories, U.S. Environmental Protection Agency. Duluth, Minnesota. PB85-227049
- Stuewer, F.W., 1943. Raccoons: Their habits and management in Michigan. *Ecology Monographs* 13:203–257.
- Texas Commission on Environmental Quality (TCEQ), 2003. An evaluation of macrobenthic community structure in the Galveston Bay System from 1997 to 2000. Texas Commission on Environmental Quality, Field Operations Division. AS-191. May 2003.
- TCEQ, 2006. *Update to Guidance for Conducting Ecological Risk Assessment at Remediation Sites in Texas RG-263 (Revised)*. Austin, TX: Texas Commission on Environmental Quality Available from:

 http://www.tceq.state.tx.us/assets/public/remediation/eco/0106eragupdate.pdf.
- TCEQ, 2012. Aquatic Life Surface Water Risk-Based Exposure Limits (SWRBELs). TRRP Protective Concentration Levels, Texas Commission on Environmental Quality. http://www.tceq.state.tx.us/remediation/trrp/trrppcls.html.
- TCEQ, 2012. Draft, Development of a Tidal Stream Benthic Index in Galveston Bay, Texas. Linda Broach, May 2012.
- TNRCC, 2001. Guidance for Conducting Ecological Risk Assessments at Remediation Sites in Texas. Austin, TX: Texas Natural Resource Conservation Commission.
- USACE (United States Army Corps of Engineers), 2011. Environmental Effects Residue Database. U.S. Army Corps of Engineers, Dredging Operations Technical Support Program. http://el.erdc.usace.army.mil/ered/. Accessed: September 2011 (Last USACE update October 2011).
- USEPA (U.S. Environmental Protection Agency), 1980. Ambient Water Quality Criteria For Polychlorinated Biphenyls. Prepared By, U.S. Environmental Protection Agency,

- Office of Water Regulations and Standards, Criteria and Standards Division. Washington, D.C., EPA 440/5-80-068.
- USEPA, 1985. Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses. Office of Research and Development. PB85-227049. January 1985.
- USEPA, 1988. Guidance for Conducting Remedial Investigations and Feasibility Studies under CERCLA. Interim Final. EPA/540/G-89/004. Washington, DC: Office of Emergency and Remedial Response.
- USEPA, 1992. *Framework for Ecological Risk Assessment*. EPA/630/R-92/001. Risk Assessment Forum, USEPA, Washington, DC.
- USEPA, 1993. *Wildlife Exposure Factors Handbook.* EPA/600/R-93/187. Washington, DC: Office of Research and Development.
- USEPA, 1997. Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments. Solid Waste and Emergency Response. OSWER Directive 9285.7-25. USEPA 540-R-97-006.
- USEPA, 1998. *Guidelines for Ecological Risk Assessment*. Risk Assessment Forum. USEPA/630/R-95/002F. Available from: http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=12460.
- USEPA, 1999. Screening Level Ecological Risk Assessment Protocol for Hazardous Waste Combustion Facilities. Peer Review Draft. Washington, DC: Office of Solid Waste.
- USEPA, 2001. The Role of Screening-Level Risk Assessments and Refining Contaminants of Concern in Baseline Ecological Risk Assessments. USEPA540/F-01/014. Office of Solid Waste. June 2001.
- USEPA, 2002. *Calculating Upper Confidence Limits for Exposure Point Concentrations at Hazardous Waste Sites.* Office of Emergency and Remedial Response, USEPA, Washington, DC. OSWER 9285.6-10.
- USEPA, 2003a. Generic Ecological Assessment Endpoints (GEAEs) for Ecological Risk Assessment. Risk Assessment Forum, USEPA. Washington, DC. USEPA/630/P-02/004F. October 2003.

- USEPA, 2003b. Procedures for the derivation of equilibrium partitioning sediment benchmarks (ESBs) for the protection of benthic organisms: PAH mixtures. Washington, DC: Office of Research and Development.
- USEPA, 2004. Draft Aquatic Life Water Quality Criteria for Selenium. U.S. Environmental Protection Agency Office of Water Office of Science and Technology. Washington, D.C. EPA-822-D-04-001. November 2004.
- USEPA, 2005. Procedures for derivation of equilibrium partitioning sediment benchmarks (ESBs) for the protection of benthic organisms: Metal mixtures (cadmium, copper, lead, nickel, silver, and zinc). Washington, DC: Office of Research and Development.
- USEPA, 2005a. *Ecological Soil Screening Levels for Arsenic*. OSWER Directive 9285.7-62. Office of Solid Waste and Emergency Response, Washington, DC. March 2005.
- USEPA, 2005b. *Ecological Soil Screening Levels for Cadmium*. OSWER Directive 9285.7-65. Office of Solid Waste and Emergency Response, Washington, DC. March 2005.
- USEPA, 2005c. *Ecological Soil Screening Levels for Lead.* OSWER Directive 9285.7-70. Office of Solid Waste and Emergency Response, Washington, DC. March 2005.
- USEPA, 2007. ProUCL Version 4.0 Technical Guide. EPA/600/R-07/041. Washington, DC: Office of Research and Development.
- USEPA, 2007a. *Ecological Soil Screening Levels for Copper*. OSWER Directive 9285.7-68. Office of Solid Waste and Emergency Response, Washington, DC. February 2007.
- USEPA, 2007b. *Ecological Soil Screening Levels for Zinc*. OSWER Directive 9285.7-73. Office of Solid Waste and Emergency Response, Washington, DC. June 2007.
- USEPA, 2007c. *Ecological Soil Screening Levels for DDT and Metabolites*. OSWER Directive 9285.7-57. Office of Solid Waste and Emergency Response, Washington, DC. April 2007.
- USEPA, 2007d. *Ecological Soil Screening Levels for Polycyclic Aromatic Hydrocarbons* (*PAHs*). OSWER Directive 9285.7-78. Office of Solid Waste and Emergency Response, Washington, DC. June 2007.
- USEPA, 2008. *Ecological Soil Screening Levels for Chromium*. OSWER Directive 9285.7-66. Office of Solid Waste and Emergency Response, Washington, DC. June 2008.

- USEPA, 2008a. Framework for application of the toxicity equivalence methodology for polychlorinated dioxins, furans, and biphenyls in ecological risk assessment.

 Washington, DC: Office of the Science Advisor.
- USEPA, 2008b. Procedures for the derivation of equilibrium partitioning sediment benchmarks (ESBs) for the protection of benthic organisms: compendium of Tier 2 values for nonionic organics. Washington, DC: Office of Research and Development.
- USEPA, 2008c. EPA's 2008 Report on the Environment. National Center for Environmental Assessment, Washington, DC; EPA/600/R-07/045F. Available from the National Technical Information Service, Springfield, VA, and online at http://www.epa.gov/roe.
- USEPA, 2010. ProUCL Version 4.1.00 Technical Guide. EPA/600/R-07/041. Washington, DC: Office of Research and Development. May 2010.
- USEPA, 2012. National Recommended Water Quality Criteria. Available from: http://www.epa.gov/ost/criteria/wqctable/. Last accessed March 2012.
- Van den Berg, M., L. Birnbaum, A.T. Bosveld, B. Brunstrom, P. Cook, M. Feeley, J.P. Giesy, A. Hanberg, R. Hasegawa, and S.W. Kennedy, 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environ. Health Perspect.* 106, 775–792.
- Van den Berg, M., L.S. Birnbaum, M. Denison, M. De Vito, W. Farland, M. Feeley, H. Fiedler, H. Hakansson, A. Hanberg, L. Haws, M. Rose, S. Safe, D. Schrenk, C. Tohyama, A. Tritscher, J. Tuomisto, M. Tysklind, N. Walker, and R.E. Peterson, 2006. The 2005 World Health Organization Reevaluation of Human and Mammalian Toxic Equivalency Factors for Dioxins and Dioxin-Like Compounds. Toxicological Sciences, 93(2), 223–241.
- Vennesland, R., and R.W. Butler, 2011. Great Blue Heron (Ardea herodias), The Birds of North America Online (A. Poole, Ed.). Cornell Lab of Ornithology, Ithaca, New York.
- Vos, J.G., H.L. Van Der Maas, A. Musch, and E. Ram, 1971. Toxicity of hexachlorobenzene in Japanese quail with special reference to porphyria, liver damage, reproduction, and tissue residues. *Toxicol. Appl. Pharmacol.* 18:944-957.

- Warnock S.E., and J.Y. Takekawa, 1995. Habitat Preferences of Wintering Shorebirds in a Temporally Changing Environment: Western Sandpipers in the San Francisco Bay Estuary. *The Auk* 112:920–930. http://bna.birds.cornell.edu/bna/species/161/articles/introduction.
- Weir, S.M., J.G. Suski, and C.J. Salice, 2010. Ecological risk of anthropogenic pollutants to reptiles: Evaluating assumptions of sensitivity and exposure. Environmental Pollution 158:3596-3606.
- Wolfe, M.F., S. Schwarzbach, and R.A. Sulaiman, 1998. Effects of Mercury on Wildlife: A Comprehensive Review. *Environ. Toxicol.* Chem. 17:146–160.

C

TABLES

C

C



Assessment Endpoints, Measurement Endpoints, and Lines of Evidence

Assessment Endpoint	Ecological Receptor	Risk Questions	Measurement Endpoint	Lines of Evidence		
		Is the survival of benthic invertebrates observed or predicted from sediment COPCs below acceptable thresholds?	Bulk surface sediment concentrations	Predicted toxicity of Site sediments based on Site-specific empirical sediment chemistry and bioassay data (i.e. Site-specific predictive toxicity model).		
Survival, growth, and reproduction of the benthic and aquatic invertebrate community	Invertebrates	Is the condition of the benthic community below acceptable thresholds due to exposure to sediment COPC?	Bulk surface sediment concentrations	Benthic community indices describing the condition of the observed benthic community relative to regional and local reference conditions.		
		Do COPC concentrations in surface water exceed benchmarks protective of the benthic invertebrate community?	Surface water concentrations	Concentrations of surface water COPC compared to effects- based values that are protective of benthic invertebrate communities.		
Survival, growth, and reproduction of the fish community		Do COPC concentrations in surface water exceed benchmarks protective of the fish community?	Surface water concentrations	Comparison of surface water COPC concentrations compared to effects-based benchmarks protective of the survival, growth, and reproduction of fish communities.		
	Fish	Do COPC concentrations in whole body tissues of fish exceed benchmarks protective of the fish community?	Whole body fish	Comparison of whole body fish tissue to appropriate tissue residue levels protective of the survival, growth, and reproduction of fish communities.		
		Do COPC concentrations in sediment exceed benchmarks protective of the fish community?	Bulk surface sediment concentrations	Comparison of sediment COPC concentrations compared to effect levels predicted to reduce survival of fish communities ^a .		
Survival growth and reproduction of sediment probing and omnivorous/herbivorous bird communities	Spotted Sandpiper	Does the dietary dose of COPCs received due to consumption of prey species captured at the Site, as well as incidental ingestion of Site sediment, exceed TRVs for survival, growth, and reproduction?	Species-specific dietary dose of COPC in prey and incidentally ingested sediment	Comparison of estimated dietary dose to dietary TRVs protective of sediment probing and omnivorous/herbivorous bird communities.		



Assessment Endpoints, Measurement Endpoints, and Lines of Evidence

Assessment Endpoint	Ecological Receptor	Risk Questions	Measurement Endpoint	Lines of Evidence				
Survival growth and reproduction of carnivorous wading bird communities	Composite carnivorous wading bird ^b	Does the dietary dose of COPCs received due to consumption of prey species captured at the Site, as well as incidental ingestion of Site sediment, exceed TRVs for survival, growth, and reproduction?	Species-specific dietary dose of COPC in prey and incidentally ingested sediment	Comparison of estimated dietary dose to dietary TRVs protective of carnivorous wading bird communities.				
Survival growth and reproduction of piscivorous bird communities	Belted Kingfisher	Does the dietary dose of COPCs received due to consumption of prey species captured at the Site, as well as incidental ingestion of Site sediment, exceed TRVs for survival, growth, and reproduction?	Species-specific dietary dose of COPC in prey	Comparison of estimated dietary dose to dietary TRVs protective of piscivorous bird communities.				
Survival growth and reproduction of aquaticassociated omnivorous mammal communities	ction of aquatic- ted omnivorous Raccoon Raccoon Site sediment, exce		1	Comparison of estimated dietary dose to dietary TRVs protective of omnivorous mammals.				

Notes:

^a - Benthic invertebrate LOE for sediment used as proxy for fish. See text for explanation.

^b - A composite receptor representing the average or mid-point exposure factors (e.g., body weight) of the species in this guild were used (see Anchor QEA 2009).

Table 4-1
Summary of BERA Dataset

Medium	Data Type	Number of Samples		
Surface Sediment	Chemical concentrations in all surface sediment collected within the 0 - 10 cm depth interval in the Site.			
Surface Water	Chemical concentrations in surface water collected using a peristaltic pump and horizontal van Dorn bottle from within the Site.	26		
Invertebrate, Whole Body Fish Tissue	Chemical concentrations in oyster, blue crab, brown shrimp, white shrimp, Gulf killifish, Gulf menhaden, pinfish, sand seatrout, and striped mullet collected from within the Site.	83		
Sediment Toxicity	Toxicity response endpoints for surface sediment samples tested with four invertebrate species.	51		
Benthic Community	Invertebrates from each sample were identified to the lowest practical taxon and enumerated. The data were used to calculate	48 ^a		
benuite Community	various indices including richness, diversity, and indices of relative abundance (e.g., relative abundance of Bivalvia).	26 ^b		

Notes:

- a Patrick Bayou Lead Organization Site samples (Parsons et al. 2002, 2004)
- b Comparison bayou samples (Dobberstine 2007)

Table 4-2
Toxic Eqivalency Factors for Toxicity Equivalent Calculations

	TEF									
Congener	Mammals ¹	Birds ²	Fish ²							
Dioxins										
2,3,7,8-TCDD	1	1	1							
1,2,3,7,8-PeCDD	1	1	1							
1,2,3,4,7,8-HxCDD	0.1	0.05	0.5							
1,2,3,6,7,8-HxCDD	0.1	0.01	0.01							
1,2,3,7,8,9-HxCDD	0.1	0.1	0.01							
1,2,3,4,6,7,8-HpCDD	0.01	0.001	0.001							
OCDD	0.0003	0.0001	0.0001							
Furans										
2,3,7,8-TCDF	0.1	1	0.05							
1,2,3,7,8-PeCDF	0.03	0.1	0.05							
2,3,4,7,8-PeCDF	0.3	1	0.5							
1,2,3,4,7,8-HxCDF	0.1	0.1	0.1							
1,2,3,6,7,8-HxCDF	0.1	0.1	0.1							
1,2,3,7,8,9-HxCDF	0.1	0.1	0.1							
2,3,4,6,7,8-HxCDF	0.1	0.1	0.1							
1,2,3,4,6,7,8-HpCDF	0.01	0.01	0.01							
1,2,3,4,7,8,9-HpCDF	0.01	0.01	0.01							
OCDF	0.0003	0.0001	0.0001							
Non-ortho PCBs										
3,3',4,4'-TCB (77)	0.0001	0.05	0.0001							
3,4,4',5-TCB (81)	0.0003	0.1	0.0005							
3,3',4,4',5-PeCB (126)	0.1	0.1	0.005							
3,3',4,4',5,5'-HxCB (169)	0.03	0.001	0.00005							
Mono-ortho PCBs	-									
2,3,3',4,4'-PeCB (105)	0.00003	0.0001	0.000005							
2,3,4,4',5-PeCB (114)	0.00003	0.0001	0.000005							
2,3',4,4',5-PeCB (118)	0.00003	0.00001	0.000005							
2',3,4,4',5-PeCB (123)	0.00003	0.00001	0.000005							
2,3,3',4,4',5-HxCB (156)	0.00003	0.0001	0.000005							
2,3,3',4,4',5'-HxCB (157)	0.00003	0.0001	0.000005							
2,3',4,4',5,5'-HxCB (167)	0.00003	0.00001	0.000005							
2,3,3',4,4',5,5'-HeCB (189)	0.00003	0.00001	0.000005							

Sources:

- 1 Van den Berg et al., 2006
- 2 Van den Berg et al., 1998

Table 4-3
Summary of Site Surface Sediment Data Evaluated in the BERA Dataset

Summary of Site Surface Sediment Data Evaluated in the BERA Dataset								_						
Sample ID	Mercury	Total Metals ^a	Dioxins/Furans	PCB Congeners	Pesticides	svocs	SOOA	Ammonia	T0C	Grain Size	Specific Gravity	AVS/SEM	PCB Aroclors	PAHs
PB001.1-1SS010-091027-N	Х	Х	Х	Х	Х	Х	Х	Χ	X	Х		Χ		
PB001.2-1SS010-091027-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB001.3-1SS010-091027-N	Х	Х	Х	Х	х	Х	Х	Х	Х	Х	Х	Х		
PB002-1SS010-091027-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB004-1SS010-091027-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB005-1SS010-091027-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB007.1-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB007.2-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Х	Χ	Х					
PB009-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Χ	Χ	Х	Х		Х		
PB011-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Х	Χ	Х					
PB013.1-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB013.2-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB015-1SS010-091028-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB016-1SS010-091101-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB018-1SS010-091101-N	Х	Х	Х	Х	Х	Х	Х	Χ	Х	Х		Х		
PB019-1SS010-091101-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB022-1SS010-091101-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB023-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					*
PB024-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		
PB026-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB028-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Χ	Х	Х		Х		
PB032-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB034-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB037-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB041-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB043-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB044-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB047.1-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Χ	Х	Х					
PB047.2-1SS010-091030-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB048-1SS010-091031-N	Х	Х	Х	Х	Х	Х	Х	Χ	X					
PB049-1SS010-091031-N	Х	Х	Х	Х	Х	Х	Х	Х	X	Х	Х	Х		
PB053-1SS010-091031-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB056-1SS010-091031-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB057-1SS010-091031-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB059.1-1SS010-091102-N	Х	Х	Х	Χ	Х	Х	Х	Х	Х	Х		Х		
PB059.2-1SS010-091031-N	Х	Х	Х	Х	Х	Х	Х	Х	Х			,		
PB063.1-1SS010-091031-N	Х	Х	Х	Χ	Х	Х	Х	Х	Х	Х		Х		
PB063.2-1SS010-091102-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB064-1SS010-091102-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB066-1SS010-091101-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
	Sample ID PB001.1-1SS010-091027-N PB001.2-1SS010-091027-N PB001.3-1SS010-091027-N PB002-1SS010-091027-N PB004-1SS010-091027-N PB005-1SS010-091027-N PB007.1-1SS010-091028-N PB007.2-1SS010-091028-N PB009-1SS010-091028-N PB011-1SS010-091028-N PB013.1-1SS010-091028-N PB013.2-1SS010-091028-N PB015-1SS010-091028-N PB016-1SS010-091028-N PB016-1SS010-091101-N PB019-1SS010-091101-N PB022-1SS010-091101-N PB022-1SS010-091030-N PB024-1SS010-091030-N PB024-1SS010-091030-N PB028-1SS010-091030-N PB032-1SS010-091030-N PB034-1SS010-091030-N PB041-1SS010-091030-N PB041-1SS010-091030-N PB047.1-1SS010-091030-N PB047.1-1SS010-091030-N PB047.1-1SS010-091031-N PB048-1SS010-091031-N PB049-1SS010-091031-N PB059-1-1SS010-091031-N PB059-1-1SS010-091031-N PB059.2-1SS010-091031-N PB059.2-1SS010-091031-N PB063.2-1SS010-091031-N PB063.1-1SS010-091031-N PB063.1-1SS010-091031-N PB063.1-1SS010-091031-N	Sample ID PB001.1-15S010-091027-N X PB001.2-15S010-091027-N X PB001.3-15S010-091027-N X PB002-15S010-091027-N X PB004-15S010-091027-N X PB005-15S010-091027-N X PB007.1-15S010-091028-N X PB007.2-15S010-091028-N X PB009-15S010-091028-N X PB011-15S010-091028-N X PB013.1-15S010-091028-N X PB013.2-15S010-091028-N X PB015-15S010-091028-N X PB015-15S010-091028-N X PB016-15S010-091028-N X PB018-15S010-091010-N X PB022-15S010-091101-N X PB022-15S010-091101-N X PB023-15S010-091030-N X PB024-15S010-091030-N X PB028-15S010-091030-N X PB034-15S010-091030-N X PB034-15S010-091030-N X PB037-15S010-091030-N X PB047-15S010-091030-N X PB047-15S010-091030-N X PB047-15S010-091030-N X PB047-15S010-091030-N X PB047-15S010-091030-N X PB047-15S010-091030-N X PB047-15S010-091031-N X PB048-15S010-091031-N X PB049-15S010-091031-N X PB059-1-15S010-091031-N X PB059-1-15S010-091031-N X PB059-2-15S010-091031-N X PB063.2-15S010-091031-N X PB063.2-15S010-091031-N X PB064-15S010-091031-N X PB063.2-15S010-091031-N X PB064-15S010-091031-N X PB059-2-15S010-091031-N X PB059-2-15S010-091031-N X PB063.2-15S010-091031-N X PB064-15S010-091031-N X PB064-15S010-091031-N X	Sample ID PB001.1-15S010-091027-N	Sample ID	Sample ID <	Sample ID	Sample ID	Sample ID	Sample ID	Sample ID				

Table 4-3
Summary of Site Surface Sediment Data Evaluated in the BERA Dataset

Station ID	Sample ID		Total Metals ^a	Dioxins/Furans	PCB Congeners	Pesticides	svocs	vocs	Ammonia	тос	Grain Size	Specific Gravity	AVS/SEM	PCB Aroclors	PAHs
PB069	PB069-1SS010-091101-N	X	X	X	Х	Х	Х	Х	Х	Х	Х	Х	Х		
PB074	PB074-1SS010-091101-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB081	PB081-1SS010-091103-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Х		
PB093	PB093-1SS010-091103-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х		Χ		
PB097	PB097-1SS010-091103-N	Х	Х	Х	Х	Х	Х	Х	Х	Х					
PB101	PB101-1SS010-091103-N	Х	Х	Х	Х	Х	X	Х	Х	Х	Х		X		
PB076.1	PB076.1-1SS010-20110802-N													Х	Х
PB070A	PB070A-1SS010-20110802-N													Х	Х
PB081.1	PB081.1-1SS010-20110803-N													Х	Х
PB082	PB082-1SS010-20110803-N													Χ	Х
PB069.1	PB069.1-1SS008-20110804-N													Χ	· X
PB068	PB068-1SS010-20110804-N													Х	Х

PCB - polychlorinated biphenyl

SVOC - semi-volatile organic compound

VOC - volatile organic compound

TOC - total organic carbon

AVS - acid volatile sulfide

SEM - simultaneously extracted metals

PAH - polycyclic aromatic hydrocarbon

a - Metals include arsenic, cadmium, chromium, copper, lead, mercury, nickel, selenium, and zinc



Summary of Site Surface Water Data Evaluated in the BERA Dataset

Station ID	Sample ID		Dioxins/Furans	PCB Congeners	Pesticides	SNOCs	VOCs	TSS SST	TDS	T0C	рос	Phosphorous	TKN	Total Cyanide	Alkalinity	Chloride, Fluoride, Sulfate, Nitrate	Selenium
	PB006-1SWMID-091104-N	Χ	Χ	Х	Х	Χ	Х	Х	Χ	Χ	Х	Х	Χ	Χ	Х	Х	Х
PB006	PB006-1SWMID-091105-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
PBOOG	PB006-1SWNBT-091104-N	Х	Х	Х	Х	Х	Х	Χ	Х	Х	Х	Х	Х	Х	Х	Х	Х
	PB006-1SWNBT-091105-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
	PB031-1SWMID-091104-N	Χ	Х	Χ	Х	Х	Х	Х	Х	X	Χ	Х	Х	Х	Х	Х	Х
PB031	PB031-1SWMID-091105-N	Χ	Х	Χ	Х	Х	Х	Х	Х	Х	Χ	Х	Х	Х	Х	Χ	Х
PB031	PB031-1SWNBT-091104-N	Χ	Х	Χ	Х	Х	Х	Х	Х	Χ	Χ	Х	Х	Х	Х	Х	Х
	PB031-1SWNBT-091105-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Χ	Х	Х	Х	Х	Х	Х
PB059	PB059-1SWMID-091104-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
120033	PB059-1SWMID-091105-N	Χ	Χ	Х	Х	Х	Х	Х	Χ	Х	X	Х	Х	Х	Х	Χ	Х
	PB076-1SWMID-091104-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
PB076	PB076-1SWMID-091105-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
PBU/6	PB076-1SWNBT-091104-N	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
	PB076-1SWNBT-091105-N	Х	Х	Х	Х	Х	Х	Х	X.	Х	Х	Х	Х	Х	Х	Х	Х
PB066B	PB066B-1SWMID-20110808-N	,		Х				Х		Х							
PB101C	PB101C-1SWMID-20110808-N			Х				Х		Х							
PB082.1	PB082.1-1SWMID-20110808-N			Х				Х		Х							
PB080	PB080-1SWMID-20110808-N			Х				Х		Х							

Notes:

PCB - polychlorinated biphenyl

SVOC - semi-volatile organic compound

VOC - volatile organic compound

TSS - total suspended solids

TDS - total dissolved solids

TOC - total organic carbon

DOC - dissolved organic carbon

TKN - total kjeldahl nitrogen

Table 4-5
Summary of Site Tissue Data Evaluated in the BERA Dataset

Species	Size Class	Number of Samples	Mercury	Lead	1,3-Dichlorobenzene	1,4-Dichlorobenzene	Hexachlorobenzene	Hexachlorbutadiene	PAHs	Dioxins/Furans	PCB Congeners	Lipid	Percent Moisture
Fish		·-							r				
Gulf Killifish	A (<15 cm)	25	Х		Х		X			X	Х	X	Х
Gulf Menhaden	A (<15 cm)	10	Х		Х		Х			Х	Х	X	Х
Pinfish	A (<15 cm)	4	Х		Х		Х			Х	Х	Х	Х
Sand Seatrout	A (<15 cm)	2	Х		Х		X			Х	Х	Х	Х
Striped Mullet	A (<15 cm)	9	Х		Х		Х			Х	Х	Х	Х
Invertebrates		-											
Blue Crab	A (2-7.5 cm)	18	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
Blue Crab	B (7.5-13 cm)	3	Х		Х		Х			Х	Х	Х	Х
Brown Shrimp	B (7.5-13 cm)	8	Х		Х		Х				Х	Х	Х
Oyster	A (2-7.5 cm)	1	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
White Shrimp	A (2-7.5 cm)	1	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
White Shrimp	B (7.5-13 cm)	2	Х		Χ		Χ				Х	Х	Х

PCB - polychlorinated biphenyl

PAH - polycyclic aromatic hydrocarbon



Summary of Bioassay/Sediment Toxicity Data Evaluated in the BERA Dataset

Agency/Entity	Activity	Station ID	Sample ID	Matrix	Sample Date	Lephtocheirus plumulosus	Neanthes arenaceodentata	Americamysis bahia	Ampelisca abdita
TCEQ/USEPA	SS	3	3-8/12/2003-EPA/TCEQ	Whole Sediment	8/12/2003			Х	Х
TCEQ/USEPA	SS	3	3-8/31/2006-TCEQ	Whole Sediment	8/31/2006			Х	Х
TCEQ/USEPA	SS	4A	4A-8/12/2003-EPA/TCEQ	Whole Sediment	8/12/2003			Х	Х
TCEQ/USEPA	SS	4A	4A-8/31/2006-TCEQ	Whole Sediment	8/12/2003			Х	Х
TCEQ/USEPA	SS	6A	6A-8/12/2003-EPA/TCEQ	Whole Sediment	8/31/2006			X	Х
TCEQ/USEPA	SS	6A	6A-8/31/2006-TCEQ	Whole Sediment	8/12/2003			Х	Х
TCEQ/USEPA	SS	7	7-8/12/2003-EPA/TCEQ	Whole Sediment	8/31/2006			Х	Х
TCEQ/USEPA	SS	9	9-8/12/2003-TCEQ	Whole Sediment	8/12/2003			х	Х
PBLO	TMDL	2	2-4/23/2001	Whole Sediment	4/23/2001	Х	Х		
PBLO	TMDL	2	2-9/5/2000	Whole Sediment	9/5/2001	Х	Х		
PBLO	TMDL	2.5	2.5-4/23/2001	Whole Sediment	4/23/2001	Х	Х		
PBLO	TMDL	2.5	2.5-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	3	3-4/24/2001	Whole Sediment	4/24/2001	X	Х		
PBLO	TMDL	3	3-8/12/2003	Whole Sediment	8/12/2003	Х		Х	Х
PBLO	TMDL	3	3-9/6/2000	Whole Sediment	9/6/2000	Х	Х		
PBLO	TMDL	5	5-4/24/2001	Whole Sediment	4/24/2001	Х	Х		
PBLO	TMDL	5	5-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	7	7-4/26/2001	Whole Sediment	4/26/2001	Х	Х		
PBLO	TMDL	7	7-8/12/2003	Whole Sediment	8/12/2003	Х		Х	Х
PBLO	TMDL	7	7-9/6/2000	Whole Sediment	9/6/2000	Х	Х		
PBLO	TMDL	8	8-4/25/2001	Whole Sediment	4/25/2001	Х	Х		
PBLO	TMDL	8	8-9/6/2000	Whole Sediment	9/6/2000	Х	Х		
PBLO	TMDL	9	9-8/12/2003	Whole Sediment	8/12/2003	Х		Х	Х
PBLO	TMDL	9	9-9/6/2000	Whole Sediment	9/6/2000	Х	Х		
PBLO	TMDL	10	10-4/25/2001	Whole Sediment	4/25/2001	Х	Х		
PBLO	TMDL	10	10-9/5/2000	Whole Sediment	9/5/2001	Х	Х		
PBLO	TMDL	4A	4A-4/24/2001	Whole Sediment	4/24/2001	Х	Х		
PBLO	TMDL	4A	4A-8/12/2003	Whole Sediment	8/12/2003	Х		Х	Х
PBLO	TMDL	4A	4a-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	6A	6A-4/25/2001	Whole Sediment	4/25/2001	Х	Х		
PBLO	TMDL	6A	6A-8/12/2003	Whole Sediment	8/12/2003	Х		Х	Х
PBLO	TMDL	6A	6a-9/5/2000	Whole Sediment	9/5/2000	Х	X		
PBLO	TMDL	E	E-4/23/2001	Whole Sediment	4/23/2001	Х	Х		
PBLO	TMDL	E	E-9/6/2000	Whole Sediment	9/6/2000	X	Х		
PBLO	TMDL	G	G-4/24/2001	Whole Sediment	4/24/2001	Х	Х		
PBLO	TMDL	G	G-9/6/2000	Whole Sediment	9/6/2000	X	X		



Summary of Bioassay/Sediment Toxicity Data Evaluated in the BERA Dataset

Agency/Entity	Activity	Station ID	Sample ID	Matrix	Sample Date	Lephtocheirus plumulosus	Neanthes arenaceodentata	Americamysis bahia	Ampelisca abdita
PBLO	TMDL	Q	Q-4/25/2001	Whole Sediment	4/25/2001	X	Х		
PBLO	TMDL	Q	Q-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	R	R-4/26/2001	Whole Sediment	4/26/2001	Х	Х		
PBLO	TMDL	R	R-8/12/2003	Whole Sediment	8/12/2003	Х		Х	Х
PBLO	TMDL	R	R-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	S	S-4/24/2001	Whole Sediment	4/24/2001	Х	Х		
PBLO	TMDL	S	S-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	Т	T-4/25/2001	Whole Sediment	4/25/2001	Х	Х		
PBLO	TMDL	Т	T-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	U	U-4/24/2001	Whole Sediment	4/24/2001	Х	X		
PBLO	TMDL	U	U-9/6/2000	Whole Sediment	9/6/2000	Х	Х		
PBLO	TMDL	V	V-4/23/2001	Whole Sediment	4/23/2001	Х	Х		
PBLO	TMDL	V	V-9/5/2000	Whole Sediment	9/5/2000	Х	Х		
PBLO	TMDL	Υ	Y-4/26/2001	Whole Sediment	4/26/2001	Х	Х		
PBLO	TMDL	Υ	Y-9/6/2000	Whole Sediment	9/6/2000	Х	Х		

Notes:

-- No Testing Performed

TMDL - Total Maximum Daily Load

PBLO - Patrick Bayou Lead Organization

TCEQ - Texas Commission on Environmental Quality

USEPA - U.S. Environmental Protection Agency

SS - Special Study



Summary of Benthic Community Data in BERA Dataset

Agency/Entity	Location	Station ID	Sample ID	Matrix	Sample Date	Benthic Community Data
PBLO	Patrick Bayou	2	2-4/23/2001	Whole Sediment	4/23/2001	X
PBLO	Patrick Bayou	2	2-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	2.5	2.5-4/23/2001	Whole Sediment	4/23/2001	X
PBLO	Patrick Bayou	2.5	2.5-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	3	3-4/24/2001	Whole Sediment	4/24/2001	Х
PBLO	Patrick Bayou	3	3-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	3	3-9/6/2000	Whole Sediment	9/6/2000	Х
PBLO	Patrick Bayou	5	5-4/24/2001	Whole Sediment	4/24/2001	X
PBLO	Patrick Bayou	5	5-9/5/2000	Whole Sediment	9/5/2000	X
PBLO	Patrick Bayou	7	7-4/26/2001	Whole Sediment	4/26/2001	Х
PBLO	Patrick Bayou	7	7-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	. 7	7-9/6/2000	Whole Sediment	9/6/2000	X
PBLO	Patrick Bayou	8	8-4/25/2001	Whole Sediment	4/25/2001	X
PBLO	Patrick Bayou	8	8-9/6/2000	Whole Sediment	9/6/2000	X
PBLO	Patrick Bayou	9	9-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	9	9-9/6/2000	Whole Sediment	9/6/2000	X
PBLO	Patrick Bayou	10	10-4/25/2001	Whole Sediment	4/25/2001	X
PBLO	Patrick Bayou	10	10-9/5/2000	Whole Sediment	9/5/2000	X
PBLO	Patrick Bayou	4A	4A-4/24/2001	Whole Sediment	4/24/2001	Х
PBLO	Patrick Bayou	4A	4A-10/1/2001	Whole Sediment	10/1/2001 ^a	Х
PBLO	Patrick Bayou	4A	4A-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	4A	4a-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	6A	6A-4/25/2001	Whole Sediment	4/25/2001	Х
PBLO	Patrick Bayou	6A	6A-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	6A	6a-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	E	E-4/23/2001	Whole Sediment	4/23/2001	Х
PBLO	Patrick Bayou	E	E-10/1/2001	Whole Sediment	10/1/2001	Х
PBLO	Patrick Bayou	E	E-9/6/2000	Whole Sediment	9/6/2000	Х
PBLO	Patrick Bayou	E	E-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	G	G-4/24/2001	Whole Sediment	4/24/2001	Х
PBLO	Patrick Bayou	G	G-10/1/2001	Whole Sediment	10/1/2001	Х
PBLO	Patrick Bayou	G	G-9/6/2000	Whole Sediment	9/6/2000	Х



Summary of Benthic Community Data in BERA Dataset

Agency/Entity	Location	Station ID	Sample ID	Matrix	Sample Date	Benthic Community Data
PBLO	Patrick Bayou	Q	Q-4/25/2001	Whole Sediment	4/25/2001	Х
PBLO	Patrick Bayou	Q	Q-10/1/2001	Whole Sediment	10/1/2001	Х
PBLO	Patrick Bayou	Q	Q-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	R	R-4/26/2001	Whole Sediment	4/26/2001	Х
PBLO	Patrick Bayou	R	R-8/12/2003	Whole Sediment	8/12/2003	Х
PBLO	Patrick Bayou	R	R-9/5/2000	Whole Sediment	9/5/2000	X
PBLO	Patrick Bayou	S	S-4/24/2001	Whole Sediment	4/24/2001	Х
PBLO	Patrick Bayou	S	S-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	Т	T-4/25/2001	Whole Sediment	4/25/2001	X
PBLO	Patrick Bayou	Т	T-9/5/2000	Whole Sediment	9/5/2000	Χ -
PBLO	Patrick Bayou	U	U-4/24/2001	Whole Sediment	4/24/2001	Х
PBLO	Patrick Bayou	U	U-9/6/2000	Whole Sediment	9/6/2000	X
PBLO	Patrick Bayou	V	V-4/23/2001	Whole Sediment	4/23/2001	Х
PBLO	Patrick Bayou	V	V-9/5/2000	Whole Sediment	9/5/2000	Х
PBLO	Patrick Bayou	Y	Y-4/26/2001	Whole Sediment	4/26/2001	X
PBLO	Patrick Bayou	Υ	Y-9/6/2000	Whole Sediment	9/6/2000	Х

Notes:

PBLO - Patrick Bayou Lead Organization

^a - Exact date is not known for the PBLO samples collected in October 2001.

Table 5-1
Receptor Group COPCs

		Chemicals of Potential	Concern
Receptor Group	Surface Water	Sediment	Bioaccumulative
Benthic Invertebrates	PCBs	Metals, PAHs, PCBs, and SVOCs ^a	
Fish	PCBs	Metals, PAHs, PCBs, and SVOCs ^a	Mercury, Selenium, PCBs, Dioxin/Furans
Wildlife (birds and mammals)		Mercury, Metals, PAHs, PCBs, Dioxin/Furans, and SVOCs ^b	Mercury, Metals, PAHs, PCBs, Dioxin/Furans, and SVOCs ^b

^a - See Table 5-3 for complete list.

PAH - polycyclic aromatic hydrocarbon

PCBs - polychlorinated biphenyl

SVOC - semi-volatile organic compound

-- no COPCs

^b - See Table 5-4 for complete list.

Table 5-2
Complete List of Benthic and Aquatic Invertebrate COPCs

COPC	Sediment	Surface Water
Metals		
Barium	Х	:
Cadmium	Х	
Chromium	Х	
Lead	х	
Mercury	х	
Nickel	Х	
Silver	х	
Zinc	х	
PAHs		
2-Methylnaphthalene	Х	
Acenaphthene	х	
Acenaphthylene	х	
Anthracene	X	
Benzo(a)anthracene	×	
Benzo(a)pyrene	Х	
Benzo(e)pyrene	X	
Chrysene	X	
Dibenzo(a,h)anthracene	X	
Fluoranthene	X	
Fluorene	X	· · · · · · · · · · · · · · · · · · ·
Indeno(1,2,3-cd)pyrene	X	
Naphthalene	X	
Perylene	$\frac{\hat{x}}{x}$	
Phenanthrene	X	
Pyrene	$\frac{\hat{x}}{x}$	
Total HMW PAH	$\frac{\hat{x}}{x}$	
Total LMW PAH	X	
Total PAH	$\frac{\hat{x}}{x}$	
PCBs		
PCB Congeners	Тх	X
SVOC / VOC	^	
Benzene	Х	
Benzidine	 	
		
Bis(2-Chloroisopropyl)ether	X	
Bis(2-Ethylhexyl)phthalate	X	
Chlorobenzene	X	
Ethylbenzene	X	
Hexachlorobenzene	X	-
Hexachlorobutadiene	X	
Isopropylbenzene	X	+
n-Butylbenzene		-
n-Propylbenzene	X	
Sec-Butylbenzene	X	+
Tert-Butylbenzene	X	+
Toluene Total xylene	X	

^a List of COPC taken from Table 7 of BERA Work Plan



Complete List of Fish COPC

Sediment	Surface Water	Bioaccumulative
Same as Table 5-2	Total PCBs	Total PCBs
		PCB TEQ
		PCDD/PCDF TEQ ¹
		Mercury
		Selenium

Notes:

PCB - polychlorinated biphenyl

TEQ - 2,3,7,8-TCDD toxic equivalency quotient

¹Based on detected levels in surface water



Complete List of Aquatic-Dependent Wildlife COPCs

COPC	Belted Kingfisher	Sandpiper	Carnivorous Wading Bird	Raccoon
Lead		Х		-
Mercury	Х	Х	X	Х
Total PAHs		Х		
Total PCB Congeners	Х	Х	Х	Х
PCB Congeners TEQ (Avian)	Х	Х	Х	
PCB Congeners TEQ (Mammal)				Х
Hexachlorobenzene	X	Х		
Hexachlorobutadiene		Х		
1,3-Dichlorobenzene	X	Х	•	
1,4-Dichlorobenzene		х		
Dioxin/Furan TEQ (Avian)		Х		

Notes:

COPC - chemical of potential concern

PAH - polycyclic aromatic hydrocarbon

PCB - polychlorinated biphenyl



Summary of Exposure Parameters for Wildlife Dietary Exposure Assessment

B			IR (mg food/kg	NIR (kg food/kg		Proportio	on in Diet		Tissue	Sediment EPC	
Receptor ^a	BW (kg)	FS	BW/day)	BW/day)	Fish	Fish Invertebrate		Terrestrial Animal	EPC (C _{TS} ; mg/kg)	(C _{so} ; mg/kg)	AUF
Belted Kingfisher	0.136	0	0.024	0.682	0.89	0.11	0	0			
Carnivorous Bird	0.87	0.01	0.074	0.327	0.33	0.33	0	0.34	See Table	See Table	See Table
Spotted Sandpiper	0.043	0.18	0.012	0.992	0	1	0	0	6-3	6-4	6-5
Raccoon	3.99	0.094	0.094	0.09	0.232	0.234	0.3	0.234			

^a - Sources for all exposure parameters and exposure parameter calculations are provided in the BERA Work Plan.

Notes:

kg - kilogram

mg - milligram

BW - body weight

FS - fraction of sediment in the diet

IR - food ingestion rate

FR - fraction of food intake from the Site

NIR - normalized ingestion rate

EPC - exposure point concentration

AUF - area use factor



Prey Grouping for Fish and Invertebrate Exposure Point Concentration Calculations

		Fish		-	Invertebra	ates
Receptor	СОРС	Group 1	Group 2	Group 3	Group 1	Group 2
	Mercury	GM	GKF, STM, PNF, SAS		BCR, BRS, WHS, OYS	
	Total PCB Congeners	GKF	GM, STM, PNF, SAS		BRS	BCR,WHS, OYS
Belted Kingfisher	Hexachlorobenzene	GKF, GM, STM, PNF, SAS			BCR, BRS, WHS, OYS	
	1,3-Dichlorobenzene	GKF, GM, STM, PNF, SAS			BCR, BRS, WHS, OYS	
	PCB TEQ	GKF	GM, STM	PNF, SAS	BCR, BRS, WHS, OYS	
	Mercury	GM	GKF, STM, PNF, SAS		BCR, BRS, WHS, OYS	
Carnivorous Bird	Total PCB Congeners	GKF	GM, STM, PNF, SAS		BRS	BCR,WHS, OYS
	PCB TEQ	GKF	GM, STM	PNF, SAS	BCR, BRS, WHS, OYS	
	Mercury	GM	GKF, STM, PNF, SAS		BCR, BRS, WHS, OYS	
Raccoon	Total PCB Congeners	GKF	GM, STM, PNF, SAS		BRS	BCR,WHS, OYS
,	PCB TEQ	GKF	GM, STM, PNF, SAS		BCR, BRS, WHS, OYS	
	Lead				BCR, OYS, WHS ^a	
	Mercury				BCR, OYS, WHS ^a	
	Total PAH				BCR, OYS, WHS ^a	<u></u>
	Total PCB Congeners				BCR, OYS, WHS ^a	
Spotted	Dioxin/furan TEQ				BCR, OYS, WHS ^a	
Sandpiper	PCB TEQ				BCR, OYS, WHS ^a	
	Hexachlorobenzene		-		BCR, OYS, WHS ^a	
	Hexachlorobutadiene				BCR, OYS, WHS ^a	
	1,3-Dichlorobenzene				BCR, OYS, WHS ^a	
	1,4-Dichlorobenzene				BCR, OYS, WHS ^a	

^a Individuals 2 to 7.5 cm total length only.

Notes:

COPC - chemical of potential concern

PCB - polychlorinated biphenyl

TEQ - toxic equivalent

PAH - polycyclic aromatic hydrocarbon

BCR - Blue crab

OYS - Oyster

WHS - White shrimp

BRS - Brown shrimp

GKF - Gulf killifish

GM - Gulf menhaden

STM - Striped mullet

PNF - Pinfish

SAS - Sand seatrout

Table 6-3
95 UCLs for Fish and Invertebrate Prey Groups

		Chemical of Potential Concern ^{a,b}									
Prey Grouping	Lead	Mercury	Total PCB Congener	Hexachlorobenzene	1,3-Dichlorobenzene	PCB Congener TEQ (Avian)	PCB Congener TEQ (Mammal)	Total PAH	Dioxin/Furan TEQ	Hexachlorobutadiene	1,4-Dichlorobenzene
GM		0.145	_ <u>-</u> -								
GKF			5.75			0.000868	0.0000870				
GKF, GM, STM, PNF, SAS				0.646	0.769 ^d						
GKF, STM, PNF, SAS		0.0580									
GM, STM, PNF, SAS			2.88				0.0000319				
GM, STM						0.000482					
PNF, SAS						0.000160					
BCR, BRS, WHS, OYS		0.0515		0.976	0.758 ^d	0.000452	0.0000279				
BRS			2.01								
BCR,WHS, OYS			0.813								
BCR, OYS, WHS ^c	0.292	0.0533	0.821	1.13	0.373 ^d	0.000430		0.140	0.0000223	0.373 ^d	0.373 ^d

^a - All units are mg/kg.

TEQ - toxic equivalent

PCB - polychlorinated biphenyl

PAH - polycyclic aromatic hydrocarbon

BCR - Blue crab

OYS - Oyster

WHS - White shrimp

BRS - Brown shrimp

GKF - Gulf killifish

GM - Gulf menhaden

STM - Striped mullet

PNF - Pinfish

SAS - Sand seatrout

^b - All totals and TEQs calculated using non-detects at 1/2 the detection limit.

^c - Individuals 2 to 7.5 cm total length only.

^d - Maximum non-detect value used

Table 6-4
Surface Weighted Average COPC Concentrations in the Intertidal and Supratidal Areas

СОРС	COPC Units Intertidal SWAC Supratidal		Supratidal SWAC	Intertidal and Supratidal Weighted Average ^a
Lead	mg/kg	45.0	53.5	48.9
Mercury (Total)	mg/kg	6.01	3.38	4.79
Total PAH	μg/kg	6660	8040	7310
1,3-Dichlorobenzene	μg/kg	78.6	67.9	73.6
1,4-Dichlorobenzene	μg/kg	76.2	72.5	74.4
Hexachlorobenzene	μg/kg	2140	2130	2140
Hexachlorobutadiene	μg/kg	3570	4130	3830
Total PCB Congener	μg/kg	9240	6300	7870
PCB Congener TEQ (Avian)	ng/kg	2530	1570	2080
PCB Congener TEQ (Mammal)	ng/kg	137	84.0	112
Dioxin/Furan TEQ (Avian)	ng/kg	122	91.1	108

^a - Intertidal and Supratidal Weighted Average is the sediment exposure point concentration.

COPC - chemical of potential concern

SWAC - surface weighted average concentration

PAH - polycyclic aromatic hydrocarbon

PCB - polychlorinated biphenyl

TEQ - toxic equivalent

mg/kg - milligram per kilogram

μg/kg - microgram per kilogram





Receptor	Surrogate Species	Home Range ^d	Units	Source	Comments
	Buff-breasted Sandpiper	3.4	km	Lanctot and Laredo (1994)	Average breeding home range
	Killdeer	4.6	km	Sanzenbacher 2002	Average linear distance traveled from capture site
	Stilt Sandpiper	8	km	Klima and Jehl (1998)	Maximum flight distance from nest
	Piping Plover	3.3	km	Drake (2001)	Average of all seasons and locations
	Western Sandpiper	4.9	km	Butler et al. (2002), Warnock and Takekawa (1995)	
	Sanderling	1.5	km	MacWhirter et al. (2002)	Average non-breeding range
Spotted Sandpiper	Willet	9.6	km	Haig et a.l (2002)	Average non-breeding home range
	Long-billed Curlew	1.95	km	Colwell et al. (2002)	Midpoint of reported home range for observed individuals
	Average home range:	4.7	km		
	Upper 75th Percentile ^b	7.3	km		
	Tidal stream within Site	2.5	km		
	AUF _{mean} :	0.54			
	AUF _{75th} :	0.34			
	Belted Kingfisher	2.2	km	Brooks and Davis (1987)	Breeding range
	Belted Kingfisher	1.0	km	Brooks and Davis (1987)	Breeding range
	Belted Kingfisher	1.0	km	Brooks and Davis (1987)	Non-breeding range
	Belted Kingfisher	0.4	km	Brooks and Davis (1987)	Non-breeding range
	Belted Kingfisher	3.6	km	Prose (1985)	Reported as range (2.4 - 4.8 km). Midpoint of range taken as average
	Belted Kingfisher	1.6	km	Mazeika et al. 2006	
Belted Kingfisher	Belted Kingfisher	4.4	km	Cromwell (1963)	Distance observed from nest site; midpoint of reported range (0.81 - 8.0 km)
	Average home range:	2.0	km		
	Upper 75th Percentile ^b	3.6	km		
	Tidal stream within Site ^a	2.5	km		
	AUF _{mean} :	1.3		1	
	AUF _{75th} :	0.69		1	
	Great Blue Heron	2.7	km	Custer and Galli (2002)	Median flight distance for Minnesota population
	Great Blue Heron	1,8	km	Mathisen and Richards (1978)	Taken from USEPA (1993)
	Great Blue Heron	7	km	Parnell and Soots, 1978, cited in Short and Cooper, 1985	Taken from USEPA (1993)
	Great Blue Heron	15	km	Gibbs et al., 1987; Gibbs, 1991; Peifer, 1979	Taken from USEPA (1993)
	Yellow-crowned Night Heron	1.4	km	Watts 2011	Breeding population
Carnivorous Birds	Tri-colored Heron	5.6	km	Frederick 1997	Breeding population
	Little Blue Heron	6.55	km	Rodgers and Smith 1995	Breeding population; midpoint of reported average home ranges (2.9 - 10.2 km)
	Snowy Egret	2.8	km	Parsons and Master 2000	Breeding population
	Great Egret	6.3	km	Mccrimmon et al. 2011	Colonial population
	Great Blue Heron	4.4	km	Vennesland and Butler 2011	Midpoint of reported average home ranges (2.3 - 6.5 km)
	Roseate Spoonbill	12	km .	Dumas 2000	Breeding population
	White Ibis	2.7	km	Heath et al. 2009	



Area Use Factor Summary for Wildlife Receptors

Receptor	Surrogate Species	Home Range ^d	Units	Source	Comments
	Average home range:	5.7	km		
	Upper 75th Percentile ^b	6.9	km		
Carnivorous Birds	Tidal stream within Site ^a	2.5	km		,
	AUF _{mean} :	0.44			
	AUF _{75th} :	0.36			
	Raccoon	1.56	km²	Stuewer (1943)	Average riparian habitat range for males and females; taken from USEPA (1993)
	Raccoon	1.33	km²	Fisher (2007)	Weighted average of both sexes in riverine habitat during breeding, rearing, and dispersal
Raccoon	Average home range ^c :	1.45	km²		
	Area of intertidal/supratidal		, 2		
•	zone within Site ^a :	0.03	km²		
	AUF _{mean} :	0.021	<u> </u>		

^a Gunite channel at south end of site not included in available habitat calculations. Represents approximately 0.6 km of tidal stream and 0.006 km² of intertidal habitat.

Notes:

km - kilometer

AUF - area use factor

USEPA - U.S. Environmental Protection Agency

km² - square kilometers

 $^{^{\}rm b}$ Upper 75th percentile based on untransformed data. See Appendix C for derivation.

^c Insufficient data to derive percentiles.

^d Home Range Distribution Analysis provided in Appendix C-4, Exhibit C-4.2

Table 6-6
Avian LOAEL TRVs

СОРС	LOAEL (mg/kg BW/day)	Source
Lead	3.26	Eco-SSL (USEPA 2005d)
Mercury (Methyl)	0.064	Heinz (1979)
Mercury (Inorganic)	0.9	Hill and Schaffner (1976)
Total PAH 325		Patton and Dieter (1980); Petroleum Hydrocarbon Mixtures
Total PCB Congener	2.05	Sample et al. (1996) as Aroclor 1242
Dioxin/furan Congener TEQ	0.00014	Sample et al. (1996); 2,3,7,8-TCDD
Hexachlorobenzene	3.35	Vos et al. (1971)
Hexachlorobutadiene	20	ECOTOX (ref 35430) (USEPA 2007b)
,3-Dichlorobenzene 160.8		1,4-Dichlorobenzene used as surrogate
1,4-Dichlorobenzene	160.8	ECOTOX (ref 344) (USEPA 2007b)
PCB Congener TEQ 0.00014		Sample et al. (1996); 2,3,7,8-TCDD

LOAEL - lowest observed apparent effects level

TRV - toxicity reference value

mg/kg - milligram per kilogram

BW - body weight

COPC - chemical of potential concern

PAH - polycyclic aromatic hydrocarbon

PCB - polychlorinated biphenyl

Table 6-7
Mammalian LOAEL TRVs

СОРС	LOAEL (mg/kg BW/day)	Source
Mercury (Methyl)	0.0177	Sample et al. (1996)
Mercury (Inorganic)	3.54	Aulerich et al. (1974)
Total PCB Congener	0.106	Sample et al. (1996); as Aroclor 1248
PCB Congener TEQ	5.4E-07	Sample et al. (1996); 2,3,7,8- TCDD

LOAEL - lowest observed apparent effects level

TRV - toxicity reference value

COPC - chemical of potential concern

mg/kg - milligram per kilogram

BW - body weight

PCB - polychlorinated biphenyl



HQs For Wildife Receptors

СОРС	Spotted Sandpiper	Carnivorous Birds	Belted Kingfisher	Raccoon
Lead	0.04			
Mercury (Methyl)	0.28	0.11	0.71	0.0049
Mercury (Inorganic)	0.004	0.0014	NA	0.00026
Total Mercury (Methyl & Inorganic) ^a	0.28	0.11	0.71	0.0052
Total PAH	0.00016			
Dioxin/Furan Congéner TEQ	0.05			
Hexachlorobenzene	0.11		0.10	
Hexachlorobutadiene	0.006			
1,3-Dichlorobenzene	0.0008		0.0022	
1,4-Dichlorobenzene	0.0008			
PCB Congener TEQ	1.0	0.51	1.7	0.16

^a Sum of methylmercury and inorganic mercury HQs.

Notes:

Highlighted cells have HQ greater than or equal to 1.0

-- Not assessed; Belted Kingfisher are not assumed to incidentally ingest sediment from the Site

HQ - hazard quotient

COPC - chemical of potential concern

PAH - polycyclic aromatic hydrocarbon



HQ Uncertainty Analysis

	l	Uncertainty	Uncertainty	Uncertainty	Uncertainty	Uncertainty	Uncertainty		
СОРС	Baseline	Analysis Scenario	Analysis	Analysis	Analysis	Analysis Scenario	l -		
		1	Scenario 2	Scenario 3	Scenario 4	5	6		
Raccoon					_				
Mercury (Methyl)	0.0044	0.0044	0.0045	0.0040	0.0054	0.005			
Mercury (Inorganic)	0.00026	0.00026	0.00026	0.00026	0.00054	0.00031			
PCB Congener TEQ (Mammal)	0.16	0.140	0.17	0.16	0.16	0.16			
Kingfisher	•								
Mercury (Methyl)	0.7	0.60	0.62	0.49	0.71	0.71			
Mercury (Inorganic)	NA	NA NA	NA	NA	NA	NA			
Hexachlorobenzene	0.10	0.13	0.09	0.10	0.10	0.10			
1,3-Dichlorobenzene	0.0022	0.0020	0.0022	0.0022	0.0022	0.0022			
PCB Congener TEQ (Avian)	1.7	1.5	2.4	2.0	1.7	1.7			
Carnivorous Bird									
Mercury (Methyl)	0.11	0.080	0.410	0.080	0.11	0.11	0.11		
Mercury (Inorganic)	0.0014	0.0014	0.0014	0.0014	0.0030	0.0017	0.0017		
PCB congener TEQ (Avian)	0.51	0.48	0.59	0.55	0.51	0.51	0.76		
Spotted Sandpiper						•	·		
Lead	0.04	0.04	0.05	0.04	0.04	0.04	0.04		
Mercury (Methyl)	0.28	0.26	0.28	0.28	0.28	0.28	0.28		
Mercury (Inorganic)	0.0040	0.006	0.006	0.0040	0.008	0.005	0.005		
Total PAH	0.00016	0.00030	0.00017	0.00016	0.00017	0.00016	0.00016		
Dioxin/Furan Congener TEQ (Avian)	0.05	0.06	0.06	0.05	0.06	0.05	0.05		
Hexachlorobenzene	0.11	0.08	0.11	0.11	0.11	0.11	0.11		
Hexachlorobutadiene	0.006	0.010	0.010	0.010	0.006	0.006	0.006		
1,3-Dichlorobenzene	0.0008	0.0015	0.008	0.0008	0.0008	0.0008	0.0008		
1,4-Dichlorobenzene	0.0008	0.0015	0.008	0.0008	0.0008	0.0008	0.0008		
PCB Congener TEQ (Avian)	1.0	0.91	1.1	1.1	1.1	1.0	1.0		



Baseline Analysis: Original fish and invertebrate species groupings; sediment concentration is weighted average of intertidal and supratidal SWACs. See Appendix C-4.

Uncertainty Analysis 1: Each prey species entered into risk sheets individually; sediment concentration is weighted average of intertidal and supratidal SWACs. See Appendix C-5.3

Uncertainty Analysis 2: All fish species grouped and all invert species grouped; sediment concentration is weighted average of intertidal and supratidal SWACs. See Appendix C-5.2

Uncertainty Analysis 3: Original fish and invertebrate species groupings but proportions entered dependent upon catch biomass ratios; sediment concentration is weighted average of intertidal and supratidal SWACs. See Appendix C-5.1

Uncertainty Analysis 4: Original fish and invertebrate species groupings; sediment concentration is weighted average of subtidal, intertidal, and supratidal SWACs. See Appendix C-5.4

Uncertainty Analysis 5: Original fish and invertebrate species groupings; sediment concentration is weighted average of intertidal and supratidal SWACs excluding the gunite channel portion of Patrick Bayou. See Appendix C-5.8

Uncertainty Analysis 6: Original fish and invertebrate species groupings; sediment concentration is weighted average of intertidal and supratidal SWACs based on hot spot analysis. See Appendix C-5.9

HQ - hazard quotient

COPC - chemical of potential concern

PCB - polychlorinated biphenyl

TEQ - toxic equivalent

PAH - polycyclic aromatic hydrocarbon

-- Not assessed

NA - Not applicable - Belted Kingfisher are not assumed to incidentally ingest sediment from the Site



Upper and Lower Bound Area Use Factor Estimates for Wildlife Receptors

Receptor	Upper Bound Home Range ^a	Lower Bound Home Range ^a	Site Habitat ^a	Units	AUF _{LOWER}	AUF _{UPPER}
Spotted Sandpiper	9.6	1.5	2.5	km	0.26	1.0
Carnivorous Bird	15	1.4	2.5	km	0.17	1.0
Belted Kingfisher	4.4	0.7	2.5	km	0.57	1.0
Raccoon	1.56	1.33	0.03	km²	0.019	0.023

^a Taken from Table 6-5.

Notes:

AUF - area use factor

km - kilometer

km² - square kilometers



Area Use Factor Uncertainty Analysis for Wildlife Receptors

conc	Spotted Sandpiper		Carnivor	ous Birds	Belted K	ingfisher	Raccoon	
COPC	AUF _{UPPER} HQ	AUF _{LOWER} HQ						
Lead	0.12	0.031	•		. 			
Mercury (Methyl)	0.83	0.22	0.29	0.050	1.0	0.58	0.0053	0.0044
Mercury (Inorganic)	0.011	0.0028	0.0039	0.00070	NA	NA	0.00028	0.00023
Total Mercury (Methyl & Inorganic)	0.84	0.22	0.29	0.051	1.0	0.58	0.0056	0.0046
Total PAH	0.00047	0.00012				1		
Dioxin/Furan Congener TEQ	0.16	0.042						
Hexachlorobenzene	0.34	0.09			0.14	0.080		
Hexachlorobutadiene	0.019	0.0049						
1,3-Dichlorobenzene	0.0023	0.00060			0.0033	0.0019		
1,4-Dichlorobenzene	0.0023	0.00060						
PCB Congener TEQ	3.1	0.80	1.42	0.24	2.5	1.4	0.17	0.14

Notes:

COPC - chemical of potential concern

AUF - area use factor

HQ - hazard quotient

PAH - polycyclic aromatic hydrocarbon

TEQ - toxic equivalent

PCB - polychlorinated biphenyl

AUF_{LOWER} HQ - See Appendix C-5.5

AUF_{UPPER} HQ - See Appendix C-5.6

-- Not assessed

NA - Not applicable - Belted Kingfisher are not assumed to incidentally ingest sediment from the Site

Table 6-12
PCB Uncertainty Analysis for Wildlife Receptors

Receptor	PCB TEQ HQ	Total PCB Congener HQ
Spotted Sandpiper	1.0	0.14
Carnivorous Bird	0.51	0.16
Belted Kingfisher	1.7	0.92
Raccoon	0.16	0.049

PCB - polychlorinated biphenyl

TEQ - toxic equivalent

HQ - hazard quotient

Total PCB Congener HQ - see Appendix C-5.7



Fish Groupings for Fish Exposure Point Concentration Calculations

			Fish	Group	
Receptor	СОРС	Group 1	Group 2	Group 3	Group 3 (All Species)
	Mercury	GM	GKF, STM, PNF, SAS	NA	GKF, GM, STM, PNF, SAS
C:-h	Total PCBs	GKF	GM, STM, PNF, SAS	NA	GKF, GM, STM, PNF, SAS
Fish	PCDD/PCDF TEQ (Fish)	GM	GKF, STM, PNF	SAS	GKF, GM, STM, PNF, SAS
	PCB Congener TEQ (Fish)	GKF	GM, STM, PNF, SAS	NA	GKF, GM, STM, PNF, SAS

Notes:

COPC - chemical of potential concern

GKF - Gulf killifish

GM - Gulf menhaden

STM - Striped mullet

PNF - Pinfish

SAS - Sand seatrout

TEQ - toxic equivalent

NA - Not applicable; grouping determined by ANOVA with post-hoc testing (see Section 7.1.1)



Tissue EPCs for Fish Bioaccumulation Pathways

		Chemical of Potential Concern ^{a,b}							
Prey Grouping	Mercury	PCB Congener TEQ (Fish)	Total PCBs	PCDD/PCDF TEQ	Selenium ^c				
GKF	NA	0.000072	5.8	NA	NA				
GM	0.15	NA	NA	0.000016	NA				
GM, STM, PNF, SAS	NA	0.000029	2.9	NA	NA				
GKF, STM, PNF, SAS	0.06	NA	NA	NA	NA				
GKF, STM, PNF	NA	NA	NA	0.0000090	NA				
SAS	NA	NA	NA	0.0000013 ^d	NA				
All Species	0.09	0.000053	4.3	0.000010	1.6				

^a - All units are mg/kg ww.

Notes:

GKF - Gulf killifish

GM - Gulf menhaden

STM - Striped mullet

PNF - Pinfish

SAS - Sand seatrout

PCB - polychlorinated biphenyl

TEQ - toxic equivalent

NA - Not applicable; prey grouping not indicated by ANOVA analysis (see Table 7-1)

^b - All totals and TEQs calculated using non-detects at 1/2 the detection limit.

^c - All COPC with exception of selenium based on empirical measurements in whole body tissue. Selenium EPC based on simple surface water bioaccumulatio model. See Section 7.1.1.1 for derivation.

^d - EPC based on maximum detected value.

Table 7-3
Fish LOED TRVs

СОРС	LOED (mg/kg ww)	Source
Mercury (Methyl)	3.8	ERED (2011)
Total PCBs	13	ERED (2011)
PCB Congener TEQ (Fish)	0.00090	ERED (2011) as 2,3,7,8-TCDD
PCDD/PCDF TEQ (Fish)	0.00090	ERED (2011) as 2,3,7,8-TCDD
Selenium	1.9	ERED (2011)

COPC - chemical of potential concern

LOED - lowest observable effects dose

mg/kg - milligram per kilogram

ww - wet weight

PCB - polychlorinated biphenyl



HQs for Fish Receptors

СОРС	HQ _{GKF}	HQ _{GM}	HQ _(GM, STM, PNF, SAS)	HQ _(GKF, STM, PNF, SAS)	HQ _(GKF, STM, PNF)	HQ _(SAS)	HQ _{All Fish}
Mercury (Methyl)	NA	0.038	NA	0.015	NA	NA	0.024
PCB Congener TEQ (Fish)	0.0081	NA	0.0033	NA	NA	NA	0.0059
PCDD/PCDF TEQ (Fish)	NA	0.018	NA	NA	0.010	0.0014	0.011
Selenium	NA	NA	NA	NA	NA	NA	0.82

Notes:

COPC - chemical of potential concern

GKF - Gulf killifish

GM - Gulf menhaden

STM - Striped mullet

PNF - Pinfish

SAS - Sand seatrout

PCB - polychlorinated biphenyl

Table 7-5
PCB Risk Characterization Uncertainty Assessment For Fish

СОРС	HQ _{GKF}	HQ _(GM, STM, PNF, SAS)	HQ _{All Fish}	
Total PCBs	0.43	0.21	0.32	

COPC - chemical of potential concern

GKF - Gulf killifish

GM - Gulf menhaden

STM - Striped mullet

PNF - Pinfish

SAS - Sand seatrout

PCB - polychlorinated biphenyl

Table 7-6
Biomagnification Uncertainty Analysis

COPC	LOGKow ^a	FCM _{TL2}	FCM _{TL4}	BMF ^c	EPC _{TL2} d	EPC _{TL4} e	TRV	HQ ^g
Mercury (methyl)	0.0763	1	1	1	0.09	0.09	3.8	0.02
PCB TEQ	7.02 ^b	1	26	26	5.3E-06	1.4E-04	9.0E-04	0.15
PCDD/PCDF TEQ	7.02 ^b	1	26	26	1.0E-05	2.6E-04	9.0E-04	0.29
Selenium	0.24	1	1	1	1.6	1.6	1.9	0.84

a Taken from TCEQ (2012)

b As 2,3,7,8-TCDD

c BMF = FCMTL4 / FCMTL2

d Table 7-2

e EPCTL4 = EPCTL2 * BMF

f Table 7-3

g HQ = EPCTL4 / TRV

FCM - food chain multiplier

TL - trophic level

BMF - biomagnification level

EPC - exposure point concentration

TRV - toxicity reference value

HQ - hazard quotient

Table 8-1
Sediment Toxicity Line of Evidence Summary

Station	Proportion Toxic
V	0%
2.5	0%
S	0%
E	25%
U	0%
3	31%
G	25%
4A	54%
5	25%
Т	0%
6A	69%
Q	25%

Toxicity

Low

<= 25% Toxic

Indeterminate Probable 25-50% Toxic > 50% Toxic

Table 8-2
Sediment Chemistry Line of Evidence Summary

Station	Relative Mean PEL-Q
V	-47%
2.5	-42%
S	38%
E	-16%
U	-94%
3	222%
G	-69%
4A	2286%
5	31%
Т	-35%
6A	419%
Q	90%

Mean PEL-Q

Low

Below Average

Indeterminate Probable Above Average Highest 15%

Table 8-3
Benthic Community Line of Evidence Summary

Station	ES Relative Benthic Index
V	3.51
2.5	4.89
S	5.25
E	6.95
U	3.18
3	6.82
G	6.12
4A	3.41
5	4.03
T	5.87
6A	0.18
Q	2.49

Low Indeterminate Probable

Benthic Index
Above Average
Below Average
Lowest 15%

Table 8-4
Weight of Evidence Risk Characterization Summary for Benthic Invertebrates

Station	Proportion Toxic	Relative Mean PEL-Q	Relative Benthic Index	WOE Risk Characterization
V	0%	-33%	3.51	Indeterminate
2.5	0%	-9%	4.89	Low
S	0%	60%	5.25	Low
E	25%	9%	6.95	Low
U	0%	-92%	3.18	Indeterminate
3	31%	233%	6.82	Indeterminate
G	25%	-64%	6.12	Low
4A	54%	2327%	3.41	Probable
5	25%	40%	4.03	Indeterminate
T	0%	-26%	5.87	Low
6A	69%	441%	0.18	Probable
Q	25%	103%	2.49	Probable

Toxicity Mean PEL-Q Benthic Index
Low <= 25% Toxic Below Average Above Average
Indeterminate 25-50% Toxic Above Average Below Average
Probable > 50% Toxic Highest 15% Lowest 15%

Table 8-5
Sediment Toxicity Sensitivity Analysis

Station	50% Survival	60% Survival (Baseline)	70% Survival	80% Survival	Relative Mean PEL-Q	Relative Benthic Index
V	0%	0%	25%	50%	-33%	3.51
2.5	0%	0%	0%	0%	-9%	4.89
S	0%	0%	25%	25%	60%	5.25
E	0%	25%	25%	25%	9%	6.95
U	0%	0%	25%	25%	-92%	3.18
3	31%	31%	31%	38%	233%	6.82
G	25%	25%	25%	25%	-64%	6.12
4A	46%	54%	62%	77%	2327%	3.41
5	25%	25%	25%	25%	40%	4.03
T	0%	0%	0%	50%	-26%	5.87
6A	69%	69%	85%	85%	441%	0.18
Q	25%	25%	25%	50%	103%	2.49

	Toxicity	Toxicity	Toxicity	Toxicity	Mean PEL-Q	Benthic Index
Low	<= 25% Toxic	<= 25% Toxic	<= 25% Toxic	<= 25% Toxic	Below Average	Above Average
Indeterminate	25-50% Toxic	25-50% Toxic	25-50% Toxic	25-50% Toxic	Above Average	Below Average
Probable	> 50% Toxic	> 50% Toxic	> 50% Toxic	> 50% Toxic	Highest 15%	Lowest 15%

Table 8-6
Sediment Chemistry Sensitivity Analysis

Station	Proportion Toxic	Relative Mean PEL- Q (Baseline)	Relative Mean PEL-Q	Relative Benthic Index
V	0%	-33%	-33%	3.51
2.5	0%	-9%	-9%	4.89
S	0%	60%	60%	5.25
E	25%	9%	9%	6.95
U	0%	-92%	-92%	3.18
3	31%	233%	233%	6.82
G	25%	-64%	-64%	6.12
4A	54%	2327%	2327%	3.41
5	25%	40%	40%	4.03
T	0%	-26%	-26%	5.87
6A	69%	441%	441%	0.18
Q	25%	103%	103%	2.49

Notes:

	Toxicity	Mean PEL-Q	Mean PEL-Q	Benthic Index
Low	<= 25% Toxic	Below Average	Below Average	Above Average
Indeterminate	25-50% Toxic	Above Average	Above Average	Below Average
Probable	> 50% Toxic	Highest 15%	Highest 25%	Lowest 15%

Table 8-7
Benthic Community Sensitivity Analysis

Station	Proportion Toxic	Relative Mean PEL-Q	Relative Benthic Index (Baseline)	Relative Benthic Index	TCEQ Benthic Index	TCEQ Benthic Index
V	0%	-33%	3.51	3.51	21.41	21.41
2.5	0%	-9%	4.89	4.89	16.25	16.25
S	0%	60%	5.25	5.25	22.93	22.93
E	25%	9%	6.95	6.95	22.01	22.01
U	0%	-92%	3.18	3.18	10.07	10.07
3	31%	233%	6.82	6.82	26.74	26.74
G	25%	-64%	6.12	6.12	19.24	19.24
4A	54%	2327%	3.41	3.41	18.15	18.15
5	25%	40%	4.03	4.03	11.07	11.07
T	0%	-26%	5.87	5.87	20.81	20.81
6A	69%	441%	0.18	0.18	11.33	11.33
Q	25%	103%	2.49	2.49	12.63	12.63

Notes:


	Toxicity	Mean PEL-Q	Benthic Index	Benthic Index	Benthic Index	Benthic Index
Low	<= 25% Toxic	Below Average	Above Average	Above Average	Above Average	Above Average
Indeterminate	25-50% Toxic	Above Average	Below Average	Below Average	Below Average	Below Average
Probable	> 50% Toxic	Highest 15%	Lowest 15%	Lowest 25%	Lowest 15%	Lowest 25%

C

FIGURES

C

0













Figure 2-1

Patrick Bayou Superfund Site Vicinity Map Baseline Ecological Risk Assessment Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas










Figure 3-2

Simplified Ecological Conceptual Site Model
Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas











Feet 275 550 825 1,100





Notes:

Multiple samples may be associated with some sample locations.

Station numbers from Patrick Bayou PSCR indicate length along channel in hundreds of feet. Aerial orthoimagery: December 2007.



Figure 4-3

BERA Dataset Invertebrate Tissue Sample Locations
Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas



ANCHOR OEA WW

Notes:


Multiple samples may be associated with some sample locations.

Station numbers from Patrick Bayou PSCR indicate length along channel in hundreds of feet. Aerial orthoimagery: December 2007.



Figure 4-4
BERA Dataset Fish Tissue Sample Locations

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas





Feet 50 700 1,050 1,400

C






Station numbers from Patrick Bayou PSCR indicate length along channel in hundreds of feet. Aerial orthoimagery: Bing Maps 2010

Feet 0 275 550 825 1,100



Figure 6-1
Patrick Bayou Tidal Zones
Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas



Key:	Toxicity	Mean PEL-Q	Benthic Index
Low	<= 25% Toxic	Below Average	Above Average
Indeterminate	> 25-50% Toxic	Above Average	Below Average
Probable	> 50% Toxic	Highest 15%	Lowest 15%






Figure 8-2
Relative Contribution of COPC to Optimized Mean PEL-Q by Station
Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas

APPENDIX A BERA DATASET

APPENDIX A-1 BERA SEDIMENT DATASET

Appendix A-1
BERA Sediment Dataset

Task: Location ID:	RI/FS 2009 PB001.1	RI/FS 2009 PB001.2	RI/FS 2009 PB001.3	RI/FS 2009 PB002	RI/FS 2009 PB004	RI/FS 2009 PB005	RI/FS 2009 PB007.1	RI/FS 2009 PB007.2	RI/FS 2009 PB009
-							PB007.1-1SS010-091028-N		
Sample Date:	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:	0 - 10 cm								
Sample Type:	N	N	N	N	N	N	N	N	N
X: Y:	3202382.3842 13836546.3402	3202624.0211 13836566.3557	3202754.9026 13836489.4987	3202606.9920 13836435.6794	3202330.4788 13836348.7826	3202262.1385 13836144.2614	3202065.5011 13836300.0036	3202037.3755 13836110.2965	3201747.2539 13836119.3947
Conventional Parameters (µmol/g)									
Sulfide, Acid Volatile/SEM	0.071 J		0.096 J		0.075 J		0.06 J		0.174 J
Acid volatile sulfide	53.8 J		14.7 J		52.8 J		103 J		13.2 J
Conventional Parameters (mg/kg)									
Ammonia	58	90	35	19	78	16	240	20	12
Conventional Parameters (pct)									
Specific gravity			2.58						-
Total organic carbon	1.64	1.46	1.09	1.11	1.47	1.15	2.01	1.26	1
Grain Size (pct)									
Granule	0		0.1 U		0.1 U		0		0.1 U
Cobbles	0		0.1 U		0.1 U		0		0.1 U
Gravel, Coarse	0		0.1 U		0.1 U		0		0.1 U
Gravel, Fine	0		0.1 U		0.1 U		0		0.1 U
Sand	46.6		45.3		13.7		42.5		42.4
Sand, Coarse	3.5		0.1		0.1 U		13.9		0.1
Sand, Medium	9.2		0.5		0.3		18		2.2
Sand, Fine	33.9		44.7		13.4		10.6		40.1
Silt, Fine	42.2		42.7		71.6		33.6		45.2
Clay, Fine	10.4		12		14.7		9.3		12.4
Fines (silt + clay)	52.6		54.7		86.3		42.9		57.6
Metals (mg/kg)									
Arsenic	3.99	4.32	2.52	3.34	4.24	3.52	6.35	3.21	1.95
Cadmium	0.495	0.696	0.296	0.535	0.776	0.641	1.17	0.597	0.424
Chromium	44.5 J	54.5 J	21.6 J	67.7 J	50.6 J	42.8 J	58.5 J	37.3 J	27.1 J
Copper	36.3 J	48.1 J	21.1 J	39.9 J	58.1 J	44.1 J	68 J	40.9 J	23.8 J
Lead	45 J	37.6 J	18.6 J	43.2 J	41 J	33.1 J	52.3 J	32.3 J	23.9 J
Mercury	4.48	3.93	1.73	7.96	5.24	4.32	5.95	3.84	2.35
Nickel	21.9 J	25 J	12.9 J	23.5 J	25.8 J	20 J	31.8 J	17.8 J	11.1 J
Selenium	1.2	1.13	0.478	0.816	1.53	1.25	1.58	1.22	0.935
Zinc	296 J	342 J	142 J	229 J	359 J	298 J	597 J	280 J	187 J
Metals (umol/g)									
Cadmium-SEM	0.00462 J		0.00212 J		0.00505 J		0.00862 J		0.00284 J
Copper-SEM	0.232 UJ		0.123 UJ		0.266 UJ		0.311 UJ		0.0839 J
Lead-SEM	0.179 J		0.0852 J		0.165 J		0.241 J		0.0963 J
Mercury-SEM	0.000367 UJ		0.000195 UJ		0.000422 UJ		0.000493 UJ		0.000114 UJ
Nickel-SEM	0.126 UJ		0.0666 UJ		0.144 UJ		0.191 J		0.0634 J
Silver-SEM	0.00341 UJ		0.00181 UJ		0.00392 UJ		0.00458 UJ		0.00106 UJ
Zinc-SEM	3.63 J		1.32 J		3.78 J		5.76 J		2.05 J
Volatile Organics (μg/kg)									
1,2-Dichlorobenzene	2.88 U	2.28 U	101 U	89 U	2.68 U	1.99 U	4.39 UJ	1.47 U	1.16 U
1,3-Dichlorobenzene	281	63.6	4090	382	22.9	18.1	244 J	28.4	25.4 J
1,4-Dichlorobenzene	114	62.8	1260	243	39.1	16.5	118 J	22.7	18.4 J
4-Isopropyltoluene (4-Cymene)	8.02	2.47	101 U	89 U	1.88 J	1.99 U	3.68 J	1.47 U	1.16 U
Benzene	2.88 U	2.28 U	101 U	89 U	2.68 U	1.99 U	2.22 J	1.47 U	1.16 U

Appendix A-1
BERA Sediment Dataset

Task	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID	•	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
Sample Date	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type		N N	N N	N N	N N	N	N N	N	FD
x	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
Y	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
Conventional Parameters (µmol/g)									
Sulfide, Acid Volatile/SEM		0.246 J		0.082 J			2.47 J		0.774 J
Acid volatile sulfide		8.6 J		71.1 J			2.53 J		1.64 J
Conventional Parameters (mg/kg)									
Ammonia	29	12 U	9.7 U	190	10 UJ		10 J	11 UJ	9.8 UJ
Conventional Parameters (pct)									
Specific gravity									
Total organic carbon	1.24	0.936	0.574	2	0.858		0.716	1.07	0.636
Grain Size (pct)									
Granule		0.1 U		0.1 U			0		0.1 U
Cobbles		0.1 U		0.1 U			0		0.1 U
Gravel, Coarse		0.1 U		0.1 U			0		0.1 U
Gravel, Fine		0.1 U		0.1 U			0		0.1 U
Sand		58.1		6			39.4		69.3
Sand, Coarse		0.3		0.1		-	4.7	-	0.3
Sand, Medium		3.1		0.6			10.8		1.9
Sand, Fine		54.7		5.3			23.9		67.1
Silt, Fine		32.3		63.3			21.2		26.6
Clay, Fine		9.6		30.7			3.2		4.1
Fines (silt + clay)		41.9		94			24.4		30.7
Metals (mg/kg)									
Arsenic	4.43	2.45	1.42	5.15	1.83		4.45	2.37	1.55
Cadmium	0.733	0.371	0.179	0.994	0.232		0.307	0.41	0.245
Chromium	116 J	28.9 J	14.4 J	68 J	20.9 J		29.7 J	29.1 J	21.8 J
Copper	61.7 J	50.6 J	16.9 J	73.3 J	22.9 J		26.8 J	31 J	24.1 J
Lead	220 J	26.9 J	13.9 J	52.8 J	57.2 J		234 J	23.9 J	26.9 J
Mercury	5.28	3.38	1.24	6.98	2.62 J		5.18 J	2.56 J	2.25 J
Nickel	25.7 J	16.3 J	8.29 J	31.7 J	10.4 J		17 J	14.6 J	11.2 J
Selenium	1.19	1.13	0.538	1.96	0.626		0.661	0.811	0.584
Zinc	325 J	203 J	103 J	527 J	137 J		530 J	237 J	151 J
Metals (umol/g)									
Cadmium-SEM		0.00215 J		0.00782 J			0.00366 J		0.00138 J
Copper-SEM		0.054 UJ		0.35 UJ			0.402 UJ		0.0986 J
Lead-SEM		0.0908 J		0.237 J			0.119 J		0.0507 J
Mercury-SEM		8.6E-05 UJ		0.000555 UJ			6.4E-05 UJ		7.5E-05 UJ
Nickel-SEM		0.0692 J		0.193 J			0.218 UJ		0.0411 J
Silver-SEM		0.0008 UJ		0.00515 UJ			0.00059 UJ		0.00069 UJ
Zinc-SEM		1.95 J		5.39 J		-	6.12 J		1.08 J
Volatile Organics (µg/kg)									
1,2-Dichlorobenzene	122 U	1.38 U	62 U	2.59 U	1.3 U		1.16 U	1.51 U	1.27 U
1,3-Dichlorobenzene	246	35.3	1030	72.8	10.1		38.6	10.2	10.8 J
1,4-Dichlorobenzene	594	31.5	1150	55.2	7.73		43.7	7.96	9.6 J
4-Isopropyltoluene (4-Cymene)	122 U	1.38 U	62 U	10.2	1.3 U		1.16 U	1.51 U	1.27 U
Benzene	122 U	1.38 U	62 U	2.59 U	0.836 J		1.16 U	1.51 U	1.27 U
: -			1				_:		

Appendix A-1
BERA Sediment Dataset

		D1/50 2000	D1/50 0000	D1/50 0000	D1/50 2000	DI /50 2000	DI /50 0000	DI /50 2000	D1/50 2000	DI /EC 2000
	Task:	RI/FS 2009								
	Location ID:	PB022	PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
	-						PB028-1SS010-091030-N			
	Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
	Depth:	0 - 10 cm								
	Sample Type:	N 2204474 4204	FD	N 2204245 4072	N 2201100 0567	N 2204224 F647	N 2201255 5225	N 2204250 5522	N 2201207 6450	N 2204200 4202
	X: Y:	3201174.1294 13835456.7389	3201315.1973 13835429.2411	3201315.1973 13835429.2411	3201180.8567 13835181.4775	3201331.5617 13835155.0673	3201355.6226 13834855.4834	3201368.6522 13834541.6071	3201387.6450 13834242.2769	3201399.4292 13833936.8351
Conventional Parameters (µmol/g)										
Sulfide, Acid Volatile/SEM		0.448 J			0.525 J		0.054 UJ		0.054 UJ	
Acid volatile sulfide		3.6 J			4.08 J		97.6 J		183 J	
Conventional Parameters (mg/kg)										
Ammonia		9.8 UJ	11 UJ	11 UJ	14 J	160 J	150 J	34 J	70 J	52 J
Conventional Parameters (pct)										
Specific gravity					2.59					
Total organic carbon		0.567	0.837	0.637	0.923	1.78	2.71	1.13	1.31	2.4
Grain Size (pct)										
Granule		0			0.1 U		0		0.1 U	
Cobbles		0			0.1 U		0		0.1 U	
Gravel, Coarse		0			0.1 U		0		0.1 U	
Gravel, Fine		0			0.1 U		0		0.1 U	
Sand		67.1			47.6		14.9		14	
Sand, Coarse		0.2			0.1		1.3		1	
Sand, Medium		1.4			1.5		6.4		2.8	
Sand, Fine		65.5			46		7.2		10.2	
Silt, Fine		28			44.7		74.3		73.6	
Clay, Fine		4.7			7.7		10.4		12.4	
Fines (silt + clay)		32.7			52.4		84.7		86	
Metals (mg/kg)										
Arsenic		1.56	3.01	2.73	1.92	8.04	6.8	3.81	6.98	5
Cadmium		0.268	0.411	0.263	0.389	1.47	0.35	0.418	2.14	1.14
Chromium		31.3 J	35.4 J	39.5 J	39.4 J	142 J	33.4 J	32 J	83.9 J	65.2 J
Copper		37.2 J	20.6 J	20.2 J	33.1 J	71.5 J	30.1 J	52 J	99.7 J	59.5 J
Lead		25.4 J	20.4 J	20.8 J	32.3 J	187 J	41.9 J	112 J	142 J	47.5 J
Mercury		2.52 J	2.55 J	2.87 J	5.44 J	64.2 J	2.54 J	24.7 J	9.11 J	5.4 J
Nickel		9.6 J	17.5 J	18.3 J	14.2 J	79.4 J	20.4 J	25.4 J	71.7 J	35 J
Selenium		0.641	0.675	0.585	0.996	1.29	1.28	0.634	1.67	1.54
Zinc		172 J	141	138	208	400	189	124	616	603
Metals (umol/g)										
Cadmium-SEM		0.00138 J			0.00238 J		0.00535 J		0.0167 J	
Copper-SEM		0.108 J			0.0601 UJ		R		0.299 UJ	
Lead-SEM		0.0567 J			0.101 J		0.23 J		0.432 J	
Mercury-SEM		7.3E-05 UJ			9.5E-05 UJ		0.000454 UJ		0.000473 UJ	
Nickel-SEM		0.0485 J			0.0689 J		0.247 J		0.528 J	
Silver-SEM		0.00068 UJ			0.00088 UJ		0.00422 UJ		0.0044 UJ	
Zinc-SEM		1.4 J			1.97 J		4.17 J		6.74 J	
Volatile Organics (μg/kg)										
1,2-Dichlorobenzene		1.23 U	1.25 U	1.18 U	1.16 U	360 U	3.84 U	1620 U	6.35 U	283 U
1,3-Dichlorobenzene		26.1 J	7.56	6.31	32.3 J	7880	106	1290 J	200	283
1,4-Dichlorobenzene		25.8 J	3.58	3.21	5.39 J	6940	43.2	1050 J	976	150 J
4-Isopropyltoluene (4-Cymene)		1.23 U	1.25 U	1.18 U	1.16 UJ	360 U	3.84 U	1620 U	19.9	283 U
Benzene		1.23 U	1.25 U	1.18 U	1.16 U	360 U	3.84 U	1620 U	4.5 J	283 U

Appendix A-1
BERA Sediment Dataset

Task	: RI/FS 2009	RI/FS 2009	RI/FS 2009	DI /EC 2000	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
	-	PB043	PB044	RI/FS 2009 PB047.1	PB047.2	PB048	PB049	PB049	PB053
Location ID									
I -		10/30/2009	10/30/2009	10/30/2009	PB047.2-1SS010-091030-N 10/30/2009	10/31/2009	PB049-1SS010-091031-D	10/31/2009	10/31/2009
Sample Date Depth		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	10/31/2009 0 - 10 cm	0 - 10 cm	0 - 10 cm
· ·		N N	N N	0 - 10 cm	N N	0 - 10 CIII	FD	N N	
Sample Type	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	N 3201421.7211
Ŷ	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
Conventional Parameters (μmol/g)									
Sulfide, Acid Volatile/SEM	0.12 J		0.079 J		0.684 J		0.354 J	0.25 J	
Acid volatile sulfide	42.1 J		62.1 J		6.7 J		10.4 J	16.2 J	
Conventional Parameters (mg/kg)									
Ammonia	32 J	28 J	53 J	18 J	21 J	37 J	25 J	20 J	97 J
Conventional Parameters (pct)									
Specific gravity	-						2.63	2.65	
Total organic carbon	1.35	1.04	1.26	0.978	1.14	1.69	1.49	1.16	2.06
Grain Size (pct)									
Granule	0		0.1 U		0.1 U		0	0	
Cobbles	0		0.1 U		0.1 U		0	0	
Gravel, Coarse	0		0.1 U	-	0.1 U		0	0	
Gravel, Fine	0		0.1 U	-	0.1 U	-	0	0	
Sand	56.5		37.8	-	14.8		57	54.9	
Sand, Coarse	1.7		1.1	-	0.1 U	-	2.6	2.3	
Sand, Medium	18.9		6.6	-	1.1	-	17.2	16.6	
Sand, Fine	35.9		30.1	-	13.7		37.2	36	
Silt, Fine	37.4		56.1		75.8		33.9	38	
Clay, Fine	5.9		6.1	-	9.4		6.5	4.2	
Fines (silt + clay)	43.3		62.2	-	85.2		40.4	42.2	
Metals (mg/kg)									
Arsenic	5.2	2.68	4.66	3.26	3.82	5.57	4.09	3.5	6.06
Cadmium	0.999	0.361	0.853	0.69	0.81	1.34	0.75	0.685	0.861
Chromium	48.4 J	28.4 J	62.4 J	50.3 J	55.2 J	72.1 J	83.6 J	72.4 J	322 J
Copper	47.4 J	25.3 J	53.5 J	32.8 J	55.4 J	73.2 J	45.3 J	42.6 J	74.3 J
Lead	52.4 J	24.6 J	37.7 J	24.9 J	34.6 J	43.2 J	38 J	34.9 J	70.9 J
Mercury	3.82 J	1.23 J	13.7 J	2.02 J	5.14 J	8.55 J	3.15 J	2.97 J	843 J
Nickel	52.3 J	16.8 J	73.3 J	17.6 J	32 J	52.2 J	20.8 J	19.4 J	55.4 J
Selenium	1.47	0.717	1.16	0.93	1.42	2.12	1.02	0.985	0.792
Zinc	1020	306	424	281	429	578 J	383 J	350 J	628 J
Metals (umol/g)									
Cadmium-SEM	0.00414 J		0.00482 J		0.00505 J		0.0029 J	0.00291 J	
Copper-SEM	0.278 J		0.213 UJ		0.378 J		0.163 J	0.15 J	
Lead-SEM	0.153 J		0.133 J		0.125 J		0.0811 J	0.0888 J	
Mercury-SEM	0.000133 UJ		0.000338 UJ	-	0.000127 UJ	-	0.000119 UJ	0.000112 UJ	
Nickel-SEM	0.292 J		0.702 J		0.219 J		0.106 J	0.112 J	
Silver-SEM	0.00124 UJ		0.00314 UJ	-	0.00118 UJ	-	0.0011 UJ	0.00104 UJ	
Zinc-SEM	4.3 J		4.09 J		3.86 J		3.31 J	3.69 J	
Volatile Organics (μg/kg)									
1,2-Dichlorobenzene	2.99 U	2.18 U	2.54 U	1.74 U	2.42 U	3.18 U	102 U	1.48 U	893 U
1,3-Dichlorobenzene	14.5	3.03	41.7	1.67 J	3.31	2.07 J	102 U	1.07 J	893 U
1,4-Dichlorobenzene	19.5	1.75 J	19.2	0.973 J	2.27 J	2.05 J	102 U	1.48 U	893 U
4-Isopropyltoluene (4-Cymene)	2.99 U	2.18 U	5.28	1.74 U	2.42 U	3.18 U	102 U	1.24 J	1160
Benzene	2.99 U	2.18 U	2.54 U	1.74 U	2.42 U	3.18 U	102 U	1.48 U	3410

Appendix A-1
BERA Sediment Dataset

Tas	*	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location I		PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
•			PB059.1-1SS010-091102-N						
Sample Date		10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Dept		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Typ		N	N	N	N	N	N	N	N
	X: 3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
	Y: 13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
Conventional Parameters (µmol/g)									
Sulfide, Acid Volatile/SEM	0.054 UJ		0.671 J		0.055 J		0.311 J		0.148 J
Acid volatile sulfide	127 J		4.48 J		57.1 J		5.25 J		15.8 J
Conventional Parameters (mg/kg)									
Ammonia	220 J	24 J	9.2 UJ	27 J	21 J	35 J	31 J	41 J	8.3 UJ
Conventional Parameters (pct)									
Specific gravity									2.61
Total organic carbon	1.79	1.03	1.09	1	1.51	0.977	0.549	0.71	2.9
Grain Size (pct)									
Granule	0		0		0		0		0
Cobbles	0		0		0		0		0
Gravel, Coarse	0		0		0		0		0
Gravel, Fine	0		0		0		0		0
Sand	39.3		84.3		71.6		56.1		68.3
Sand, Coarse	0.2		11.1		12		0.4		29.1
Sand, Medium	6.7		40.1		38.3		3.9		33.7
Sand, Fine	32.4		33.1		21.3		51.8		5.5
Silt, Fine	53.3		7.8		11.3		37.3		5.7
Clay, Fine	7.3		0.9		1.8		5.9		0.7
Fines (silt + clay)	60.6		8.7		13.1		43.2		6.4
Metals (mg/kg)									
Arsenic	3.68	2.86	6.11	3.4	6.37	3.62	4.07	2.3	9.61 J
Cadmium	0.56	0.458	0.537	0.577	0.406	0.562	0.236	0.285	1.05
Chromium	47 J	44.8 J	57.9 J	94.7 J	305 J	161 J	25.6 J	23.6 J	34.5 J
Copper	42 J	35.7 J	83.8 J	43.9 J	37.6 J	57 J	29.9 J	45.4 J	48.2 J
Lead	31.9 J	22.9 J	33.3 J	37 J	57 J	39 J	9.72 J	24.5 J	22.7 J
Mercury	2.08 J	1.13 J	0.795 J	1.12 J	0.567 J	0.769 J	0.124 J	0.48 J	1.32 J
Nickel	21.8 J	13.8 J	26.2 J	16.8 J	21.6 J	25.2 J	11.4 J	10.6 J	23.6 J
Selenium	0.891	0.785	0.563	0.766	0.669	0.919	0.299	0.27	0.426 J
Zinc	464 J	249 J	784 J	299 J	630 J	353 J	124 J	264 J	1750 J
Metals (umol/g)									
Cadmium-SEM	0.00438 J		0.00235 J		0.00294 J		0.00163 J		0.00324 UJ
Copper-SEM	0.258 UJ		0.224 UJ		0.0597 UJ		0.139 J		0.229 UJ
Lead-SEM	0.128 J		0.0776 J		0.0862 J		0.0335 J		0.117 J
Mercury-SEM	0.000409 UJ		7.1E-05 UJ		9.5E-05 UJ		0.0011 UJ		7.3E-05 UJ
Nickel-SEM	0.171 J		0.167 J		0.162 J		0.0584 J		0.208 J
Silver-SEM	0.0038 UJ		0.00066 UJ		0.00088 UJ		0.00103 UJ		0.00337 UJ
Zinc-SEM	5.14 J		2.76 J		2.87 J		1.4 J		2.02 J
Volatile Organics (μg/kg)									
1,2-Dichlorobenzene	2.38 U	1.59 U	1.32 U	1.47 U	93.8 U	1.44 U	1.08 U	1.33 U	0.629 J
1,3-Dichlorobenzene	3.15	1.59 U	1.32 U	0.847 J	93.8 U	0.979 J	1.08 U	1.33 U	1.02 U
1,4-Dichlorobenzene	1.34 J	1.59 U	1.32 U	1.47 U	93.8 U	1.44 U	1.08 U	1.33 U	0.822 J
4-Isopropyltoluene (4-Cymene)	2.77	1.59 U	2.19	1.89	144	1.14 J	1.08 U	1.6	5.95
Benzene	7.52	1.59 U	1.32 U	1.47 U	93.8 U	1.44 U	1.08 U	1.33 U	1.75

Appendix A-1
BERA Sediment Dataset

1	D. /50 2000	D1/50 2000	D1/50 2000	D1/50 2000	D1/50 0000	2014 1100	2044 1100	2044 1100
Task:	RI/FS 2009	2011 USC	2011 USC	2011 USC				
Location ID:	PB074	PB081	PB093	PB097	PB101	PB068	PB069.1	PB070A
· •		PB081-1SS010-091103-N				PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm				
Sample Type:	N	N	N	N	N	N	N	N
x: Y:	3200828.0183 13830463.2171	3200761.3787 13830005.1879	3200970.4291 13828894.7843	3201103.1242 13828566.1540	3201287.2658 13828181.2285	3201209.861 13831121.51	3201205.859 13830973.3	3201087.255 13830903.51
Conventional Parameters (μmol/g)								
Sulfide, Acid Volatile/SEM		0.054 UJ	0.159 J		0.244 J			
Acid volatile sulfide		96.3 J	6.04 J		3.97 J	-		
Conventional Parameters (mg/kg)								
Ammonia	13 J	11 UJ	9.1 UJ	50 J	9 UJ	-		
Conventional Parameters (pct)								
Specific gravity								
Total organic carbon	0.507	1.67	1.47	0.411	0.686			
Grain Size (pct)								
Granule		0	0		0			
Cobbles		0	0		0	-		
Gravel, Coarse		0	0		0			
Gravel, Fine		0	0		0	-		
Sand		47	53.8	-	76.5	1		
Sand, Coarse	-	21.2	35.1	-	31.9	•		
Sand, Medium	-	20.3	17.9 J	-	37.9	•		
Sand, Fine	-	5.5	0.8	-	6.7	1		
Silt, Fine	-	17.6	1.7	-	5.8			
Clay, Fine	-	2.2	0.1		0.6			
Fines (silt + clay)	-	19.8	1.8	-	6.4	1		
Metals (mg/kg)								
Arsenic	7.12	10.1	10.8	5.62	2.82			
Cadmium	0.265	0.537	0.246	0.214	0.19			
Chromium	22.6 J	242 J	23 J	99.9 J	13 J			
Copper	38.6 J	69.5	13.5	25	14.3			
Lead	26.3 J	335 J	36.3 J	13.1 J	172 J			
Mercury	7.54 J	0.659 J	0.026 J	0.042 J	0.0121 J			
Nickel	15 J	59	17.6	15	7.69			
Selenium	0.377	0.953 J	0.707 J	0.723 J	0.427 J	-		
Zinc	198 J	390 J	120 J	145 J	82.4 J	-		
Metals (umol/g)	_							
Cadmium-SEM		0.00234 J	0.00329 UJ		0.00322 UJ			
Copper-SEM		0.797 UJ	0.466 UJ		0.455 UJ			
Lead-SEM		0.226 J	0.0372 J		0.0472 J			
Mercury-SEM		0.000252 UJ	7.4E-05 UJ		7.2E-05 UJ			
Nickel-SEM		0.431 UJ	0.252 UJ		0.246 UJ			
Silver-SEM		0.00235 UJ	0.00343 UJ		0.00335 UJ			
Zinc-SEM		2.84 J	0.921 J		0.923 J			
Volatile Organics (μg/kg)								
1,2-Dichlorobenzene	1.33 U	R	1.07 U	1.04 U	0.816 U			
1,3-Dichlorobenzene	1.33 U	R	1.07 U	1.04 U	0.816 U			
1,4-Dichlorobenzene	0.842 J	R	1.07 U	1.04 U	0.816 U			
4-Isopropyltoluene (4-Cymene)	3.15	776 J	1.07 U	1.04 U	0.816 U			
								-

Appendix A-1 BERA Sediment Dataset

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:	PB076.1-1SS010-20110802-N	PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
Conventional Parameters (µmol/g)				
Sulfide, Acid Volatile/SEM				
Acid volatile sulfide				
Conventional Parameters (mg/kg)				
Ammonia				
Conventional Parameters (pct)				
Specific gravity				
Total organic carbon	-			
Grain Size (pct)				
Granule	-			-
Cobbles			-	-
Gravel, Coarse				
Gravel, Fine	-			
Sand	-			
Sand, Coarse				
Sand, Medium	1			
Sand, Fine				
Silt, Fine				
Clay, Fine				
Fines (silt + clay)				
Metals (mg/kg)				
Arsenic				
Cadmium				
Chromium				
Copper			-	
Lead				
Mercury			-	
Nickel				
Selenium				
Zinc				
Metals (umol/g)				
Cadmium-SEM				
Copper-SEM				
Lead-SEM			-	
Mercury-SEM			-	
Nickel-SEM			-	
Silver-SEM			-	-
Zinc-SEM			-	
Volatile Organics (µg/kg)				
1,2-Dichlorobenzene				
1,3-Dichlorobenzene			-	-
1,4-Dichlorobenzene				
4-Isopropyltoluene (4-Cymene)				
Benzene				-
Delizelle				

Appendix A-1
BERA Sediment Dataset

T1	DI /EC 2000	DI /50 2000	DI /50 2000	DI /FC 2000	DI /50 2000	DI /5C 2000	DI /EC 2000	DI /EC 2000	DI /50 2000
Task:	•	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB001.1	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
•	PB001.1-1SS010-091027-N			1				PB007.2-1SS010-091028-N	
Sample Date:		10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:		N 2202624 0244	N 2202754 0026	3202505 0020	N 2202220 4700	N 2202262 4205	N 2202005 F044	N 2202027 2755	N 2204747 2520
X:	3202382.3842 13836546.3402	3202624.0211 13836566.3557	3202754.9026 13836489.4987	3202606.9920	3202330.4788	3202262.1385 13836144.2614	3202065.5011	3202037.3755	3201747.2539 13836119.3947
T:				13836435.6794	13836348.7826		13836300.0036	13836110.2965	
Chlorobenzene	92.6	62.8	790	144	13	4.04	97.3 J	1.97	3.12 J
Ethylbenzene	9.01	2.86	101 U	89 U	2.68 U	1.99 U	5.44 J	1.47 U	1.16 U
Isopropylbenzene (Cumene)	272	24	101 U	89 U	2.51 J	1.17 J	42.8 J	1.18 J	1.22 J
m,p-Xylene	8.48	2.96 J	202 U	178 U	5.37 U	3.99 U	6.06 J	2.94 U	2.32 U
n-Butylbenzene	4.08	2.28 U	101 U	89 U	2.68 U	1.99 U	4.39 UJ	1.47 U	1.16 U
n-Propylbenzene	7.99	1.19 J	101 U	89 U	2.68 U	1.99 U	2.48 J	1.47 U	1.16 U
o-Xylene	3.67	2.28 U	101 U	89 U	2.68 U	1.99 U	4.39 U	1.47 U	1.16 U
sec-Butylbenzene	5.73	2.28 U	101 U	89 U	2.68 U	1.99 U	4.39 UJ	1.47 U	1.16 U
tert-Butylbenzene	17.7	1.3 J	101 U	89 U	2.68 U	1.99 U	2.5 J	1.47 U	1.16 U
Toluene Total Xylene (U = 1/2)	3.11 J	4.56 U	202 U 202 U	178 U	5.37 U	3.99 U	24 J 8.255 J	2.94 U	2.32 U 2.32 U
Aromatic Hydrocarbons (µg/kg)	12.15	4.1 J	202 0	178 U	5.37 U	3.99 U	8.235 J	2.94 U	2.32 0
2-Methylnaphthalene	526	356	266	382	180	289	103	218	554 J
Acenaphthene	1610	777	492	1340	332	854	260	748	2730 J
Acenaphthylene	2170	1530	890	3810	781	1480	541	1210	1370
Anthracene	1440	489	381	982	267	471	334	449	4040 J
Benzo(a)anthracene	1040	476	631	759	389	458	569	498	8520 J
Benzo(a)pyrene	1350	690	663	1710	544	742	796	896	7550 J
Benzo(b)fluoranthene	1050	624	628	832	610	639	737	745	7370 J
Benzo(e)pyrene	1100	663	564	1390	619	666	688	841	5270 J
Benzo(g,h,i)perylene	2350	1390	949	5540	1130	1370	1180	2330	6150 J
Benzo(k)fluoranthene	838	547	544	654	551	580	673	620	6510 J
C1-Chrysene	725	342	244	560	331	325	385	292	1780 J
C1-Fluoranthenes/Pyrenes	3250	1350	1020	3120	1110	1270	1180	1260	5620 J
C1-Fluorenes	1440	457	727	880	250	436	240	418	755 J
C1-Naphthalenes	881	458	343	526	244	390	143	304	636 J
C1-Phenanthrenes/Anthracenes	1830	537	375	851	339	469	449	452	2220 J
C2-Chrysenes	481	264	138	519	260	240	286	209	597 J
C2-Fluorenes	850	260	145	539	179	230	189	201	307 J
C2-Naphthalenes	1480	589	404	797	258	426	239	357	585 J
C2-Phenanthrenes/Anthracenes	1060	378	227	508	251	296	375	242	656 J
C3-Chrysenes	496	262	118	537	261	225	280	213	485 J
C3-Fluorenes	670	259	126	448	201	198	255	172	698 J
C3-Naphthalenes	4520	967	474	2930	650	807	988	729	715
C3-Phenanthrenes/Anthracenes	743	272	113	408	205	191	342	182	286 J
C4-Chrysenes	322	175	93.2	352	188	154	192	132	203
C4-Naphthalenes	1220	448	224	820	268	330	344	271	333
C4-Phenanthrenes/Anthracenes	671	286	105	435	291	224	392	198	217
Chrysene	1400	714	819	856	614	679	856	688	8150 J
Dibenzo(a,h)anthracene	155	97.5	99.1	80.4	103	98.8	126	114	1300 J
Fluoranthene	5400	2950	2500	6980	1900	2960	2170	3310	27300 J
Fluorene	981	620	590	932	284	593	184	535	2540 J
Indeno(1,2,3-c,d)pyrene	1080	671	553	1830	578	681	676	1010	5880 J
Naphthalene	1240	1430	3490	1670	568	1220	304	957	2540

Appendix A-1
BERA Sediment Dataset

Task	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	· · · · · · · · · · · · · · · · · · ·	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type	N	N	N	N	N	N	N	N	FD
x , x	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
Y	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
Chlorobenzene	1610	1.38 U	43.5 J	9.26	0.903 J		1.08 J	0.856 J	1.27 U
Ethylbenzene	122 U	1.38 U	62 U	1.32 J	1.3 U		1.16 U	1.51 U	1.27 U
Isopropylbenzene (Cumene)	83.2 J	1.38 U	62 U	4.65	1.3 U		0.865 J	1.51 U	1.27 U
m,p-Xylene	243 U	2.76 U	124 U	5.18 U	2.59 U		2.32 U	3.02 U	2.54 U
n-Butylbenzene	122 U	1.38 U	62 U	2.59 U	1.3 U		1.16 U	1.51 U	1.27 U
n-Propylbenzene	122 U	1.38 U	62 U	2.59 U	1.3 U		1.16 U	1.51 U	1.27 U
o-Xylene	122 U	1.38 U	62 U	2.59 U	1.3 U		1.16 U	1.51 U	1.27 U
sec-Butylbenzene	122 U	1.38 U	62 U	2.59 U	1.3 U		1.16 U	1.51 U	1.27 U
tert-Butylbenzene	122 U	1.38 U	62 U	2.59 U	1.3 U		1.16 U	1.51 U	1.27 U
Toluene	243 U	2.76 U	124 U	4.34 J	2.59 U		2.32 U	3.02 U	2.54 U
Total Xylene (U = 1/2)	243 U	2.76 U	124 U	5.18 U	2.59 U		2.32 U	3.02 U	2.54 U
Aromatic Hydrocarbons (μg/kg)									
2-Methylnaphthalene	301	204	115	152	218	-	256	200	187
Acenaphthene	1090	604	283	291	605	-	873	603	326
Acenaphthylene	1990	971	569	781	1200	-	1630	1140	806
Anthracene	702	300	161	342	393	-	588	355	300
Benzo(a)anthracene	610	411	242	379	481	-	2830	395	548
Benzo(a)pyrene	919	580	363	679	917	-	2990	701	804
Benzo(b)fluoranthene	658	606	353	506	659	-	2640	552	693
Benzo(e)pyrene	824	636	342	564	682	-	2050	570	625
Benzo(g,h,i)perylene	2280	1500	746	1190	1680	-	3100	1320	1040
Benzo(k)fluoranthene	490	533	298	423	561	-	2220	454	614
C1-Chrysene	424	250	130	343	285		895	231	318
C1-Fluoranthenes/Pyrenes	1930	1040	499	1180	1140	-	2740	954	968
C1-Fluorenes	664	385	262	266	371		562	402	289
C1-Naphthalenes	458	284	160	213	293		359	285	258
C1-Phenanthrenes/Anthracenes	616	355	187	411	456		674	381	375
C2-Chrysenes	334	169	82	275	173		438	154	196
C2-Fluorenes	390	146	74.3	193	194		263	154	139
C2-Naphthalenes	618	277	162	260	322		371	331	250
C2-Phenanthrenes/Anthracenes	458	180	105	309	257		528	218	198
C3-Chrysenes	346	182	73.1	286	164		359	158	190
C3-Fluorenes	315	141	69.5	205	175		234	138	139
C3-Naphthalenes	1290	554	240	933	758		1330	426	525
C3-Phenanthrenes/Anthracenes	376	115	67.3	246	174		317	139	137
C4-Chrysenes	253	118	57.9	199	106		188	112	113
C4-Naphthalenes	549	205	102	312	251		330	198	188
C4-Phenanthrenes/Anthracenes	396	164	76.5	347	182		232	145	151
Chrysene	814	606	379	560	677		3300	571	767
Dibenzo(a,h)anthracene	81.5	97	53.7	81	116		400	90.4	144
Fluoranthene	4220	2400	1290	1770	3250		8060	2520	2190
Fluorene	780	437	248	245	440		614	490	335
Indeno(1,2,3-c,d)pyrene	847	718	367	546	836		1980	652	694
Naphthalene	1920	838	960	477	936	-	2060	1530	606

Appendix A-1
BERA Sediment Dataset

	Task:	1	RI/FS 2009	RI/FS 2009	RI/FS 2009					
	Location ID:		PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
	•							PB032-1SS010-091030-N		
	Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
	Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm						
	Sample Type:	N	FD	N	N	N	N	N	N	N
	X:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
	Y:	13835456.7389	13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
Chlorobenzene		1.23 U	0.783 J	1.18 U	2.42	14400	14.7	1620 U	329	231 J
Ethylbenzene		1.23 U	1.25 U	1.18 U	1.16 U	361	3.84 U	1620 U	58.3	283 U
Isopropylbenzene (Cumene)		0.858 J	1.25 U	1.18 U	2.27	4780	7.04	1380 J	17.2	2330
m,p-Xylene		2.45 U	2.5 U	2.36 U	2.31 U	688 J	7.68 U	3250 U	22.6	566 U
n-Butylbenzene		1.23 U	1.25 U	1.18 U	1.16 UJ	201 J	3.84 U	1620 U	6.75	283 U
n-Propylbenzene		1.23 U	1.25 U	1.18 U	1.16 U	216 J	3.84 U	1620 U	28	283 U
o-Xylene		1.23 U	1.25 U	1.18 U	1.16 U	256 J	3.84 U	1620 U	5.48 J	283 U
sec-Butylbenzene		1.23 U	1.25 U	1.18 U	1.16 UJ	360 U	3.84 U	1620 U	7.24	283 U
tert-Butylbenzene		1.23 U	1.25 U	1.18 U	1.16 U	369	3.84 U	1620 U	6.35 U	283 U
Toluene		2.45 U	2.5 U	2.36 U	2.31 U	719 U	7.68 U	3250 U	12.7 U	566 U
Total Xylene (U = 1/2)		2.45 U	2.5 U	2.36 U	2.31 U	944 J	7.68 U	3250 U	28.08 J	566 U
Aromatic Hydrocarbons (μg/kg)										
2-Methylnaphthalene		289	101	110	252	3830	55.2	1320	266	164
Acenaphthene		316	331	278	595	6410	133	641	1690	439
Acenaphthylene		709	651	711	1520	6860	193	1420	2190	852
Anthracene		307	230	218	511	3570	99.3	1120	3290	812
Benzo(a)anthracene		668	217	215	677 J	1210	137	1430	1000	809
Benzo(a)pyrene		850	504	517	1020	2500	221	13600	803	909
Benzo(b)fluoranthene		712	271	261	766	697	153	2770	562	822
Benzo(e)pyrene		702	380	393	818	1160	197	6410	622	716
Benzo(g,h,i)perylene		951	1320	1420	1400	3350	343	29000	847	827
Benzo(k)fluoranthene		584	220	227	676 J	583	146	2440	519	650
C1-Chrysene		672	124	122	572	919	175	312	803	585
C1-Fluoranthenes/Pyrenes		1230	656	708	1630	5800	480	5390	5200	2340
C1-Fluorenes		280	286	311	483	4820	119	1380	3440	718
C1-Naphthalenes		334	138	147	372	6440	76.6	1620	1410	259
C1-Phenanthrenes/Anthracenes		879	203	210	732	5580	202 J	1150 J	4450 J	1250 J
C2-Chrysenes		324	91.7	86.1	338	713	138	205	490	393
C2-Fluorenes		218	104	114	265	2420	158	453	2120	587
C2-Naphthalenes		315	167	174	454	6810	106	1420	4810	837
C2-Phenanthrenes/Anthracenes		428	129	136	350	2950	188 J	642 J	1810 J	875 J
C3-Chrysenes		220	88.9	92.4	296	714	123	227	462	491
C3-Fluorenes		229	88.4	97.9	208	1430	151	332	1140	559
C3-Naphthalenes		502	426	480	1110	19000	594	2660	33600	3560
C3-Phenanthrenes/Anthracenes		167	78.5	83.4	213	1610	185 J	346 J	817 J	939 J
C4-Chrysenes		135	76.6	81.5	187	2550	92.2	1240	319	305
C4-Naphthalenes		172	157	174	379	5460	294	1080	4880	873
C4-Phenanthrenes/Anthracenes		160	87.1	101	233	1020	153 J	240 J	677 J	823 J
Chrysene		996	289	307	990	1320	221	1590	1170	1120
Dibenzo(a,h)anthracene		160	33.8	28.1	147	51.6	28.1	345	94.4	152
Fluoranthene		1980	1760	1830	3050	12700	536	28800	3880	2200
Fluorene		306	261	291	570	7570	81.9	1980	613	267
Indeno(1,2,3-c,d)pyrene		638	479	506	787	1050	163	9910	475	621
~-····\-,-,- 0,\\\p ₁ . 00		633	1250	1990	795	11300	133	4580	487	252

Appendix A-1
BERA Sediment Dataset

Task	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009				
Location ID:		PB043	PB044	PB047.1	PB047.2	PB048	PB049	PB049	PB053
			_		PB047.2-1SS010-091030-N		PB049-1SS010-091031-D		
Sample Date:	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm			
Sample Type		N N	N N	0 - 10 CIII	0 - 10 CIII	0 - 10 CIII	FD	N N	N N
Sample Type	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
γ.	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
Chlorobenzene	4.87	2.18 U	2.54 U	1.74 U	2.42 U	3.18 U	102 U	1.48 U	893 U
Ethylbenzene	2.99 U	2.18 U	2.07 J	1.74 U	2.42 U	3.18 U	102 U	1.48 U	4090
Isopropylbenzene (Cumene)	9.99	16.9	169	7.73	1.67 J	3.18 U	338 J	57.2 J	106000
m,p-Xylene	5.98 U	4.37 U	4.19 J	3.49 U	4.83 U	6.36 U	204 U	2.96 U	8960
n-Butylbenzene	2.99 U	2.18 U	4.12	1.74 U	2.42 U	3.18 U	102 U	1.01 J	1410
n-Propylbenzene	2.99 U	2.18 U	5.46	1.74 U	2.42 U	3.18 U	102 U	1.74	1280
o-Xylene	2.99 U	2.18 U	2.16 J	1.74 U	2.42 U	3.18 U	102 U	1.48 U	5210
sec-Butylbenzene	2.99 U	2.18 U	4.24	1.74 U	2.42 U	3.18 U	102 U	0.825 J	1320
tert-Butylbenzene	2.2 J	2.18 U	15.6	1.74 U	2.42 U	3.18 U	102 U	1.65	3300
Toluene	5.98 U	4.37 U	5.08 U	3.49 U	4.83 U	6.36 U	204 U	2.96 U	1780 U
Total Xylene (U = 1/2)	5.98 U	4.37 U	6.35 J	3.49 U	4.83 U	6.36 U	204 U	2.96 U	14170
Aromatic Hydrocarbons (μg/kg)									
2-Methylnaphthalene	231	52.5	137	106	55.6	46.1	108 J	52.9	32000
Acenaphthene	686	176	714	113	49.1	47.8	166 J	140	9770
Acenaphthylene	969	638	1010	1340	310	329	522 J	329	10900
Anthracene	1230	447	1370	453	128	136	604 J	327	15700
Benzo(a)anthracene	1370	995	988	1340	410	406	431 J	600	3780
Benzo(a)pyrene	1350	1160	1080	1440	637	625	400 J	703	2590
Benzo(b)fluoranthene	1330	1020	990	1160	638	638	358 J	629	1190
Benzo(e)pyrene	1030	823	821	1040	539	557	368	500	1470
Benzo(g,h,i)perylene	1010	890	826	988	624	646	308 J	517	944
Benzo(k)fluoranthene	1130	920	894	1050	603	579	282 J	541	1020
C1-Chrysene	800	537	611	880	301	340	440	351	2640
C1-Fluoranthenes/Pyrenes	2930	1690	2800	3030	783	956	1430	1140	22500
C1-Fluorenes	1220	397	1190	448	95.9	104	471 J	262	14600
C1-Naphthalenes	379	89.5	434	162	75.3	57.8	259 J	125	36400
C1-Phenanthrenes/Anthracenes	1620 J	600 J	1640 J	741 J	197 J	207	762 J	431	24500
C2-Chrysenes	595	311	326	542	223	254	286	209	1230
C2-Fluorenes	892	381	693	553	107	144	464 J	237	9030
C2-Naphthalenes	1200	452	1370	352	114	114	571 J	307	23200
C2-Phenanthrenes/Anthracenes	1240 J	451 J	854 J	1010 J	193 J	249	779 J	419	13900
C3-Chrysenes	543	286	312	567	231	253	288	207	1160
C3-Fluorenes	763	342	438	669	156	209	453 J	241	5960
C3-Naphthalenes	5700	4760	4820	2160	473	690	2120	985	41600
C3-Phenanthrenes/Anthracenes	911 J	374 J	481 J	1010 J	176 J	247	615 J	326	6960
C4-Chrysenes	325	149	176	326	134	195	190	124	698
C4-Naphthalenes	1210	639	861	657	169	217	649	344	15100
C4-Phenanthrenes/Anthracenes	782 J	354 J	424 J	988 J	262 J	358	570 J	317	4100
Chrysene	1660	1260	1350	1630	618	596	608 J	728	3760
Dibenzo(a,h)anthracene	239	217	188	227	126	121	75	120	200
Fluoranthene	3220	2460	2730	2740	1020	1140	984 J	1410	12100
Fluorene	236	181	150	119	66.2	56.8	119 J	77	21500
Indeno(1,2,3-c,d)pyrene	964	853	763	915	541	526	250 J	475	670
Naphthalene	448	70.8	269	174	109	125	187 J	103	45100

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009				
Location ID:	PB056	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
					PB063.1-1SS010-091031-N				
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm				
Sample Type:	N N	N N	N N	N N	N N	N N	N N	N N	N N
v.	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
Chlorobenzene	2.38 U	1.59 U	1.32 U	1.47 U	93.8 U	1.44 U	1.08 U	1.33 U	0.856 J
Ethylbenzene	2.06 J	1.59 U	0.933 J	1.47 U	76.1 J	1.44 U	1.08 U	1.33 U	34.1
Isopropylbenzene (Cumene)	25.7	1.59 U	37	2.52	7760	18.7	1.08 U	69.3	6530
m,p-Xylene	2.52 J	3.18 U	2.63 U	2.94 U	188 U	2.88 U	2.16 U	2.66 U	17
n-Butylbenzene	1.49 J	1.59 U	1.25 J	1.47 U	128	0.875 J	1.08 U	1.8	4.1
n-Propylbenzene	2.38 U	1.59 U	3.44	1.47 U	236	0.825 J	1.08 U	3.45	11.1
o-Xylene	2.38 U	1.59 U	0.725 J	1.47 U	93.8 U	1.44 U	1.08 U	0.793 J	20.1
sec-Butylbenzene	2.38 U	1.59 U	2.04	1.47 U	151	1.44 U	1.08 U	2.08	5.08
tert-Butylbenzene	1.48 J	1.59 U	9.23	1.47 U	448	0.969 J	1.08 U	5.44	17.8
Toluene	4.76 U	3.18 U	2.63 U	2.94 U	96.6 J	2.88 U	2.16 U	2.66 U	6.38 U
Total Xylene (U = 1/2)	3.71 J	3.18 U	2.04 J	2.94 U	188 U	2.88 U	2.16 U	2.123 J	37.1
Aromatic Hydrocarbons (μg/kg)									
2-Methylnaphthalene	78.6	23.5	324	280	486	49.1	560	55.8	2520
Acenaphthene	122	28.4	711	199	4470	54.1	672	66.5	6160
Acenaphthylene	333	89.4	3210	284	3200	162	125	330	9830
Anthracene	284	102	1510	281	4780	178	675	211	7040
Benzo(a)anthracene	1260	692	2600	312	5310	457	889	850	2130
Benzo(a)pyrene	1650	971	3060	401	4780	593	897	1020	1240
Benzo(b)fluoranthene	1740	1070	2070	384	3860	648	751	1070	689
Benzo(e)pyrene	1240	755	2070	321	3000	516	579	814	802
Benzo(g,h,i)perylene	1180	821	2840	311	2990	525	552	844	476
Benzo(k)fluoranthene	1540	861	1820	312	3580	592	759	950	585
C1-Chrysene	703	293	1830	248	1990	289	257	412	1730
C1-Fluoranthenes/Pyrenes	1600	741	6800	849	7740	943	680	1220	8730
C1-Fluorenes	163	50.1	2040	218	3000	119	72.6	279	6460
C1-Naphthalenes	117	29.3	530	378	1610	68.3	640	86.9	9040
C1-Phenanthrenes/Anthracenes	440	153	2580	294	4880	183	328	362	8950
C2-Chrysenes	432	168	1190	207	919	188	152	232	1070
C2-Fluorenes	217	108	1730	234	1900	225	77	265	3270
C2-Naphthalenes	254	53.1	1640	308	4060	142	268	207	9690
C2-Phenanthrenes/Anthracenes	545	202	2740	323	3150	296	135	396	4780
C3-Chrysenes	425	174	830	240	648	211	174	227	634
C3-Fluorenes	303	174	1740	279	1570	276	146	307	2680
C3-Naphthalenes	663	192	6990	675	6870	613	115	855	14800
C3-Phenanthrenes/Anthracenes	431	175	2480	341	2180	266	82.9	364	3480
C4-Chrysenes	230	102	348	184	240	171	116	122	287
C4-Naphthalenes	309	165	1960	399	1990	416	64.8	286	3270
C4-Phenanthrenes/Anthracenes	415	210	2360	405	1560	327	105	321	2920
Chrysene	1850	1050	2810	473	5160	726	866	1180	2290
Dibenzo(a,h)anthracene	288	201	470	71.5	827	130	133	204	126
Fluoranthene	2260	2020	5730	909	13000	1320	3010	2310	5090
Fluorene	99.6	34.7	443	220	1320	40.2	536	75.7	1800
Indeno(1,2,3-c,d)pyrene	1210	846	2130	279	3080	510	546	857	392
Naphthalene	94.3	34.3	605	622	967	31.2	90	93.8	4810

Appendix A-1
BERA Sediment Dataset

Taskil	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI/EC 2000	2014 LICC	2014 UCC	2014 LICC
Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009 PB097	RI/FS 2009	2011 USC PB068	2011 USC PB069.1	2011 USC PB070A
Location ID:	PB074	PB081	PB093		PB101			
•		PB081-1SS010-091103-N				PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm N	0 - 10 cm	0 - 10 cm N	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm
Sample Type:		3200761.3787	3200970.4291	3201103.1242	3201287.2658	N 3201209.861	N 320120F 8F0	3201087.255
Λ: Y:	3200828.0183 13830463.2171	13830005.1879	13828894.7843	13828566.1540	13828181.2285	13831121.51	3201205.859 13830973.3	13830903.51
Chlorobenzene	1.33 U	R	1.07 U	1.04 U	0.816 U			
Ethylbenzene	3.76	6290 J	1.07 U	1.04 U	0.816 U	-		
Isopropylbenzene (Cumene)	244	71400 J	1.07 U	1.04 U	5.81	-		
m,p-Xylene	3.6	9080 J	2.14 U	2.08 U	1.63 U			
n-Butylbenzene	3.65	2370 J	1.07 U	1.04 U	0.816 U	-		
n-Propylbenzene	10.8	2890 J	1.07 U	1.04 U	0.816 U	-		
o-Xylene	1.24 J	3790 J	1.07 U	1.04 U	0.555 J			
sec-Butylbenzene	3.83	618 J	1.07 U	1.04 U	0.816 U			
tert-Butylbenzene	8.26	665 J	1.07 U	1.04 U	0.816 U			
Toluene	2.66 U	7140 J	2.14 U	2.08 U	1.63 U			
Total Xylene (U = 1/2)	4.84 J	12870 J	2.14 U	2.08 U	1.37 J	-		
Aromatic Hydrocarbons (μg/kg)								
2-Methylnaphthalene	40.9	179000	9.61	4.27 J	12.9	1710	81.1	43.4
Acenaphthene	172	41900	29.8	6.4	55.1	8800	202	113
Acenaphthylene	283	88300	60.8	28.3	43.4	7240	1310	302
Anthracene	204	54200	23.9	18.2	73.4	6690	859	239
Benzo(a)anthracene	279	15100	121	106	415	1930	1190	384
Benzo(a)pyrene	250	8710	176	102	460	1570	1060	551 J
Benzo(b)fluoranthene	255	3530	201	145	515	1300	973	675 J
Benzo(e)pyrene	201	5130	140	100	368	1400	771	527
Benzo(g,h,i)perylene	172	2360	157	112	376	802	650	510 J
Benzo(k)fluoranthene	216	4120	154	111	412	959	846	513 J
C1-Chrysene	183	8190	59.2	40.8	163	1520	605	279
C1-Fluoranthenes/Pyrenes	664	60500	114	82.9	276	9530	2450	743
C1-Fluorenes	245	49100	18.8	5.82	19.9	7200	428	249
C1-Naphthalenes	90.2	201000	22.7	5.76	20.4	6480	107	76.4
C1-Phenanthrenes/Anthracenes	313	78500	46.2	30.9	171	11300	1060	906
C2-Chrysenes	116	698 U	59.1	21	82.1	930	359	212
C2-Fluorenes	198	17400	21.1	11.6	26.6	7690	732	481
C2-Naphthalenes	262	65700	34.1	9.22	28.7	10200	411	542
C2-Phenanthrenes/Anthracenes	329	16300	44.4	28.3	86	8740	1320	848
C3-Chrysenes	144	698 U	113	45.6	104	913	316	231
C3-Fluorenes	233	6510	25	4.36 U	76.1	6740	980	531
C3-Naphthalenes	539	121000	40.1	15.1	34.1	19000	1440	800
C3-Phenanthrenes/Anthracenes	307	4200	35.9	21.6	44.4	6430	1210	482
C4-Chrysenes	92.6	698 U	86	4.36 U	68.2	107 U	23.4 U	192
C4-Naphthalenes	206	10000	19.8	8.2	13.7	6960	414	611
C4-Phenanthrenes/Anthracenes	268	2200	27.9	8.94	25.7	4070	860	253
Chrysene	345	13400	185	138	551	2260	1300	696 J
Dibenzo(a,h)anthracene	43.2	668 J	36.7	26.3	96.3	176	165	116
Fluoranthene	856	36500	263	248	1070	5710	2650	1150
Fluorene	84.5	112000	13.4	4.38	33.6	10400	214	94.3
Indeno(1,2,3-c,d)pyrene	168	1830	158	118	400	749	615	505 J
Naphthalene	62.8	413000	32.8	5.14	11.7	6660	102	70

Appendix A-1 BERA Sediment Dataset

Task:		2011 USC	2011 USC	2011 USC
Location ID:		PB081.1	PB081.1	PB082
Sample ID:		PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:		8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
Chlorobenzene				
Ethylbenzene				
Isopropylbenzene (Cumene)				
m,p-Xylene				
n-Butylbenzene				
n-Propylbenzene				
o-Xylene				
sec-Butylbenzene				
tert-Butylbenzene				
Toluene				
Total Xylene (U = 1/2)				
Aromatic Hydrocarbons (µg/kg)				
2-Methylnaphthalene	63	53000	41000	44.8
Acenaphthene	293	23700	19800	548 J
Acenaphthylene	332	39400	32800	1460 J
Anthracene	896	31800	23600	1190 J
Benzo(a)anthracene	2050	5870	5060	508 J
Benzo(a)pyrene	2070	3100	2880	349 J
Benzo(b)fluoranthene	2400	1400	1180	254 J
Benzo(e)pyrene	1690	1840	1700	287 J
Benzo(g,h,i)perylene	1510	830	819	182 J
Benzo(k)fluoranthene	2020	1230	1220	193 J
C1-Chrysene	721	3320	2890	480 J
C1-Fluoranthenes/Pyrenes	2570	26200	22000	2260 J
C1-Fluorenes	539	23600	19500	1310 J
C1-Naphthalenes	79.2	74400	58700	123
C1-Phenanthrenes/Anthracenes	1030	36700	29100	1190 J
C2-Chrysenes	384	1310	1160	434 J
C2-Fluorenes	1230	10400	9020	1160 J
C2-Naphthalenes	404	36800	31200	1040 J
C2-Phenanthrenes/Anthracenes	1470	11600	9350	1790 J
C3-Chrysenes	326	924	699	380 J
C3-Fluorenes	1080	5790	4620	1160 J
C3-Naphthalenes	1120	76000	64000	3560 J
C3-Phenanthrenes/Anthracenes	836	5450	4190	1580 J
C4-Chrysenes	188	108 U	102 U	292
C4-Naphthalenes	911	7250	5860	835 J
C4-Phenanthrenes/Anthracenes	361	2900	2420	942 J
Chrysene	2830	5580	4810	622 J
Dibenzo(a,h)anthracene	413	227	202	34.8 J
Fluoranthene	5610	16800	13600	1410 J
Fluorene	132	43200	33900	78.3
Indeno(1,2,3-c,d)pyrene	1740	709	636	136
- (/ //- II /	93.3	96300	73800	114

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB001.1	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
	PB001.1-1SS010-091027-N							PB007.2 PB007.2-1SS010-091028-N	
•									
Sample Date:	10/27/2009 0 - 10 cm	10/27/2009 0 - 10 cm	10/27/2009	10/27/2009	10/27/2009	10/27/2009 0 - 10 cm	10/28/2009 0 - 10 cm	10/28/2009	10/28/2009 0 - 10 cm
Depth: Sample Type:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	N N	0 - 10 CIII	0 - 10 cm N	N N
Sample Type.	3202382.3842	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011	3202037.3755	3201747.2539
Λ. V.	13836546.3402	13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614	13836300.0036	13836110.2965	13836119.3947
Domilara .									
Perylene	285 3710	223 2570	178 2510	309 3960	187 1230	185 2530	226 1240	210 2360	2240 J 21500 J
Phenanthrene		5450				5450			
Pyrene Total 10 of 17 HPAH (U = 1/2)	9990 24653	13609.5	3620 11006.1	17800 37041.4	3260 9679	13657.8	3150 10933	6110 16321	22800 J 101530 J
					13321			22798	136804 J
Total 7 PAH (U = 1/2)	36330	21381.5 7772	19625.1	50117.4		21094.8	13899		
Total 7 of 17 LPAH (U = 1/2)	11677		8619	13076	3642	7437	2966	6477	35274 J
Total cPAH TEF (7 minimum) (U = 1/2)	1780.3	938.69	916.7	2134.1	773.24	994.47	1082.66	1201.58	10589.5 J
Semivolatile Organics (µg/kg) 2,2'-Oxybis (1-chloropropane)	528 U	488 U	321 U	366 U	514 U	406 U	766 U	360 U	313 U
Benzidine	1060 UJ	976 UJ	641 UJ	731 UJ	1030 UJ	812 UJ	766 U R	720 UJ	625 UJ
Bis(2-ethylhexyl) phthalate	5370	1460	693	2060	1030 03	1240	11800 J	1040	963 J
Cholesterol									302.1
Di-n-octyl phthalate	528 U	488 U	321 U	366 U	514 U	406 U	766 U	360 U	313 U
Coprostanol									
Hexachlorobenzene	7300	2400	150000	8900	520 J	1800 J	2900 J	1200	20000
Hexachlorobutadiene	19000	16000	310000	32000	1900 J	6800 J	1300 J	5400	27000
Hexachloroethane	860	1200	21000	1400	170 J	540 J	110 J	410	4100
PCB Congeners (ng/kg)	800	1200	21000	1400	1703	3403	1107	410	4100
PCB-001	11400	4270	2140	8390	1760	3480	1550	2940	2770
PCB-002	585	481	265	635	187	380	182	276	490
PCB-003	5970	3160	1420	6420	1420	2770	1350	2220	2360
PCB-004	40900 J	9320 J	4070 J	17900 J	7500 J	10900 J	6910 J	13600 J	6010 J
PCB-005	1070 J	276 J	138 J	434 J	134 J	247 J	213 J	281 J	260 J
PCB-006	14700 J	11400 J	3030 J	11100 J	2990 J	4300 J	2630 J	4220 J	2810 J
PCB-007	2480 J	687 J	313 J	1250 J	364 J	611 J	419 J	676 J	521 J
PCB-008	81500 J	21300 J	9420 J	38800 J	13200 J	20300 J	15400 J	21300 J	14300 J
PCB-009	3600 J	1040 J	456 J	1920 J	578 J	928 J	630 J	998 J	744 J
PCB-010	1400 J	247 J	113 J	366 J	148 J	206 J	188 J	264 J	157 J
PCB-011	841 J	680 J	8.93 U	499 J	315 J	464 J	314 J	8.47 U	390 J
PCB-012/013	6980 J	3900 J	1410 J	3970 J	1470 J	2200 J	1400 J	1960 J	1810 J
PCB-014	210 J	8.93 U	537 J	251 J	62.4 J	128 J	51.4 J	79.7 J	205 J
PCB-015	42700 J	14700 J	7240 J	24400 J	9260 J	13600 J	10700 J	13900 J	11800 J
PCB-016	129000 J	25600 J	8570	32500 J	15100	19100 J	25900 J	20800 J	12700
PCB-017	148000 J	39600 J	13500	62000 J	30800 J	36700 J	37800 J	38800 J	23600 J
PCB-018/030	17.9 UJ	86600 J	26800 J	119000 J	60300 J	75600 J	90900 J	71900 J	41900 J
PCB-019	33200 J	6290	2140	9130	4680	5950	6890	7320	3620
PCB-020/028	381000 J	105000 J	38800 J	160000 J	78100 J	96600 J	101000 J	106000 J	70500 J
PCB-021/033	219000 J	46400 J	19900 J	87700 J	44900 J	54800 J	44500 J	56300 J	35200 J
PCB-022	125000 J	25700 J	9590	36900 J	16200	22000 J	26900 J	22700 J	16400 J
PCB-023	8.93 U	40.3	43	54.3	9.62 U	31.4	46 J	42.6	7.58 U
PCB-024	8.93 U	8.93 U	194	746	413	393	469 J	461	317
PCB-025	19800 J	11300	2830	10200	4180	5050	5000	4870	3390
PCB-026/029	60100 J	24200 J	7510	27400 J	12600	15700	16000	16100	11300
PCB-027	21100 J	4690	1650	6560	3610	4160	5030	4850	2820

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	-	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N N	N N	N N	N N	N N	N N	N N	FD
y.	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
ν. γ.	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
Domilana									
Perylene	209	167	85	176	200		784	152	193
Phenanthrene	2990	1880	1060	1160	2090		2720	2120	1610
Pyrene	8050	4400	2220	3270	6110		11000	4520	3500
Total 10 of 17 HPAH (U = 1/2)	18969.5	11851	6311.7	9404	15287		38520	11775.4	10994
Total 17 PAH (U = 1/2)	28742.5	17085	9707.7	12852	21169		47261	18213.4	15164
Total 7 of 17 LPAH (U = 1/2)	9773	5234	3396	3448	5882	-	8741	6438	4170
Total cPAH TEF (7 minimum) (U = 1/2)	1195.79	822.56	498.16	878.1	1189.07		4030	921.05	1080.97
Semivolatile Organics (µg/kg)	252.11	205.11	262.11	670.11	254.11		247.11	20011	220.11
2,2'-Oxybis (1-chloropropane)	353 U	305 U	268 U	679 U	251 U		247 U	290 U	239 U
Benzidine	705 UJ	610 UJ	536 UJ	1360 UJ	R		R	R	R
Bis(2-ethylhexyl) phthalate	1660	1530	632	1280	949		959	991	812
Cholesterol	252.11				320 U		300 U	310 U	
Di-n-octyl phthalate	353 U	305 U	268 U	679 U	251 U		247 U	290 U	239 U
Coprostanol					1600 U	-	340 U	310 U	
Hexachlorobenzene	2700	10000	22000	1500 J	2400		8500 J	2500	5200
Hexachlorobutadiene	11000	7900	20000	3200 J	7000 J	-	4600 J	7200 J	2000 J
Hexachloroethane	960	450	1600	180 J	1000	-	470 J	2000	350
PCB Congeners (ng/kg)									
PCB-001	3830	1660	828	1640		2920	3710 J	3120	1340
PCB-002	391	232	204	229		238	444 J	362	218
PCB-003	3160	1250	781	1530		2050	2570	2350	1090
PCB-004	9010 J	5730 J	2750 J	5850 J		35500 J	8790 J	6170 J	4490 J
PCB-005	235 J	162 J	113 J	194 J		499 J	297 J	213 J	124 J
PCB-006	6800 J	2120 J	1090 J	2540 J		8220 J	3380 J	2610 J	1520 J
PCB-007	638 J	372 J	222 J	432 J		1320 J	668 J	471 J	263 J
PCB-008	21300 J	11400 J	5310 J	13900 J		54200 J	17400 J	13000 J	8090 J
PCB-009	1040 J	577 J	263 J	617 J		2040 J	1080 J	698 J	409 J
PCB-010	221 J	128 J	70.7 J	151 J		645 J	289 J	196 J	122 J
PCB-011	353 J	244 J	6.85 U	393 J		382 J	213 J	328 J	266 J
PCB-012/013	2690 J	1140 J	1000 J	1480 J		4320 J	1560 J	1640 J	1030 J
PCB-014	106 J	41.9 J	35 J	52.6 J		50.5 J	105 J	114 J	43.9 U
PCB-015	15400 J	8580 J	4260 J	11900 J		28900 J	11200 J	10900 J	7040 J
PCB-016	20600 J	11800	5500	20300 J		105000 J	18900 J	10900	11800
PCB-017	39300 J	22600 J	9670	35000 J		145000 J	26200 J	20100 J	20100 J
PCB-018/030	82300 J	45900 J	19900 J	73900 J		541000 J	75900 J	59000 J	52100 J
PCB-019	5340	3680	1640	5270		43100 J	4900	2860	3510
PCB-020/028	110000 J	68300 J	33000 J	102000 J		384000 J	62600 J	62000 J	59200 J
PCB-021/033	58100 J	33000 J	14500 J	43400 J		188000 J	31900 J	29100 J	25900 J
PCB-022	23900 J	15500	7130	23900 J		122000 J	15900 J	13800	14300
PCB-023	47.5	29	22.4	9.62 U		107	46.8	49.6	27.9
PCB-024	413	282	150	507		1490	428	261	251
PCB-025	6790	3050	1530	4550		15300	3030	2870	2660
PCB-026/029	19200 J	10100	4790	15000		54800 J	10000	9130	8550
PCB-027	4190	2620	1200	4410		20500 J	3290	2360	2490

Appendix A-1
BERA Sediment Dataset

	Taalu	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /FC 2000	DI /EC 2000	DI /FC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000
1	Task: Location ID:	RI/FS 2009 PB022	RI/FS 2009 PB023	RI/FS 2009 PB023	RI/FS 2009 PB024	RI/FS 2009 PB026	RI/FS 2009 PB028	RI/FS 2009 PB032	RI/FS 2009 PB034	RI/FS 2009 PB037
1										
1	•	PB022-1SS010-091101-N					PB028-1SS010-091030-N			
1	Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
1	Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
1	Sample Type:	N	FD	N	N	N	N	N	N	N
1	Х:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
	Y:	13835456.7389	13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
Perylene		185	94.7	97.9	219	274	976	1750	185	224
Phenanthrene		2080	1180	1370	2730	23700	356 J	12700 J	5750 J	1270 J
Pyrene		3320	3660	3820	5580	27500	1040	66300	8320	3490
Total 10 of 17 HPAH (U = 1/2)		10859	8753.8	9131.1	15093 J	50961.6	2988.1	156185	17670.4	11600
Total 17 PAH (U = 1/2)		15499	12757.8	14099.1	22066 J	114201.6	4039.5 J	179946 J	31956.4 J	15656 J
Total 7 of 17 LPAH (U = 1/2)		4640	4004	4968	6973	63240	1051.4 J	23761 J	14286 J	4056 J
Total cPAH TEF (7 minimum) (U = 1/2)		1136.16	628.97	643.78	1335.2 J	2872.36	285.92	15305.4	1079.74	1225.6
Semivolatile Organics (μg/kg)										
2,2'-Oxybis (1-chloropropane)		242 U	262 U	265 U	284 U	589 U	520 U	339 U	584 U	534 U
Benzidine		R	R	R	R	R	R	R	R	2180 J
Bis(2-ethylhexyl) phthalate		812	1010	1980	1920 J	1020 J	458 J	3320 J	1300 J	4780 J
Cholesterol		290 U	-							
Di-n-octyl phthalate		242 U	262 U	265 U	284 U	589 U	520 U	339 U	584 U	534 U
Coprostanol		290 U								
Hexachlorobenzene		7100	8200	17000	280 J	1200 J	200 J	200000	240	380
Hexachlorobutadiene		2500 J	11000 J	11000 J	1300 J	2000 J	200 J	300000 J	510 J	250 J
Hexachloroethane		410	3400 J	1800	140 J	170 UJ	150 UJ	100 U	170 U	630
PCB Congeners (ng/kg)										
PCB-001		1290 J	5150 J	2110 J	4440	178000 J	1020	23500 J	10400	3680
PCB-002		173 J	556 J	422 J	425	14700 J	125	12800	1940	281
PCB-003		983	2350 J	1610	3190	46900 J	567	17700	6700	2020
PCB-004		4800 J	7060 J	4110 J	8630 J	94800 J	1880 J	7700 J	5930 J	19500 J
PCB-005		101 J	235 J	136 J	307 J	937 J	73.3 J	172 J	300 J	702 J
PCB-006		1480 J	2310 J	1640 J	3610 J	7660 J	773 J	803 J	2520 J	8270 J
PCB-007		245 J	502 J	312 J	658 J	2510 J	143 J	563 J	583 J	1410 J
PCB-008		8380 J	12900 J	8920 J	21600 J	81000 J	4410 J	17600 J	15000 J	49600 J
PCB-009		383 J	721 J	504 J	1000 J	3570 J	238 J	599 J	819 J	2140 J
PCB-010		127 J	275 J	161 J	251 J	2180 J	66.9 J	136 J	214 J	631 J
PCB-011		172 J	255 J	187 J	322 J	3100 J	120 J	45 U	455 J	800 J
PCB-012/013		875 J	1400 J	1000 J	2060 J	18800 J	454 J	5420 J	1780 J	4940 J
PCB-014		24.3 J	269 J	247 J	59.2 J	897 J	23.3 J	267 J	138 J	39.4 J
PCB-015		7320 J	9940 J	7640 J	16300 J	89600 J	3530 J	22700 J	11500 J	27700 J
PCB-016		13100	15000 J	9160	22000 J	50700 J	6190	1240	23100 J	84000 J
PCB-017		22400 J	24100 J	15300 J	30900 J	75800 J	9280	1860	35300 J	108000 J
PCB-018/030		56800 J	70800 J	42100 J	76800 J	149000 J	23000 J	4560	86200 J	281000 J
PCB-019		3670	3430 J	1980	5430	12300 J	1420	405	5450	20800 J
PCB-020/028		62700 J	57700 J	54500 J	97800 J	72500 J	26900 J	5910	128000 J	359000 J
PCB-021/033		26900 J	21800 J	20700 J	43500 J	33200 J	11600	2370	51100 J	199000 J
PCB-022		14300	12800 J	12100	26000 J	18500 J	7030	1490	33100 J	102000 J
PCB-023		22.4	282 J	56.9	52.8	40 UJ	19.9	70.9	87.5	229
PCB-024		277	441 J	280	519	2120 J	158	47.6	582	1890
				2130	4500	4330 J	1290	597	6060	18400 J
PCB-025		2/00	2340 J	2130	1 4300	40001	1230			
PCB-025 PCB-026/029		2700 8460	2340 J 8250 J	7850	14200 J	18400 J	4190	1120	19300 J	55300 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	DI/EC 2000	DI/EC 2000	RI/FS 2009	DI/FC 2000	RI/FS 2009	RI/FS 2009
	PB041	PB043	PB044	RI/FS 2009 PB047.1	RI/FS 2009 PB047.2	PB048	RI/FS 2009 PB049	PB049	PB053
Location ID:									
-							PB049-1SS010-091031-D		
Sample Date:	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm			
Sample Type:	N N	N	N	N	N	N	FD	N	N
X:	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
Υ:	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
Perylene	359	282	272	308	183	195	98 J	178	344
Phenanthrene	974 J	1130 J	976 J	653 J	365 J	371	618 J	582	57300
Pyrene	4340	2890	3930	3990	1240	1490	1610 J	1650	27900
Total 10 of 17 HPAH (U = 1/2)	16613	12665	13739	15480	6457	6767	5306 J	7373	54154
Total 17 PAH (U = 1/2)	21387 J	15360.3 J	18365 J	18438 J	7539.9 J	7878.7	7630 J	8983.9	246424
Total 7 of 17 LPAH (U = 1/2)	4774 J	2695.3 J	4626 J	2958 J	1082.9 J	1111.7	2324 J	1610.9	192270
Total cPAH TEF (7 minimum) (U = 1/2)	1869.9	1573.1	1475.8	1925.5	874.98	857.96	545.68 J	946.78	3313.6
Semivolatile Organics (μg/kg)									
2,2'-Oxybis (1-chloropropane)	383 U	270 U	458 U	308 U	381 U	570 U	307 U	297 U	1530 U
Benzidine	R	R	R	339 J	846 J	R	R	R	R
Bis(2-ethylhexyl) phthalate	1450 J	1270 J	1880 J	1240 J	1530 J	1620	2460	2120	2300
Cholesterol									-
Di-n-octyl phthalate	383 U	270 U	458 U	308 U	381 U	570 U	307 U	297 U	1530 U
Coprostanol									
Hexachlorobenzene	580 U	390	220	48 J	450 J	170 UJ	92 U	88 U	2200 U
Hexachlorobutadiene	360 UJ	130 J	120 J	84 J	580 J	100 UJ	57 U	54 J	1400 U
Hexachloroethane	580 U	170	130 U	210	330 UJ	170 UJ	70 J	120	2200 U
PCB Congeners (ng/kg)									
PCB-001	3770	1370	6140	5370	411	943 J	803	815	415000 J
PCB-002	355	40.3	186	166	84.5	111 J	68	62.6	15100
PCB-003	2080	758	3570	1550	409	777	522	565	142000 J
PCB-004	22000 J	12500 J	25200 J	29100 J	2840 J	6470 J	6090 J	6090 J	1380000 J
PCB-005	592 J	385 J	862 J	844 J	92.7 J	150 J	197 J	227 J	45100 J
PCB-006	7820 J	5410 J	10000 J	9670 J	1210 J	2300 J	2410 J	2820 J	366000 J
PCB-007	1320 J	808 J	1680 J	1580 J	212 J	393 J	388 J	432 J	95400 J
PCB-008	46400 J	32800 J	59400 J	59600 J	7570 J	13200 J	15200 J	17700 J	1690000 J
PCB-009	1970 J	1290 J	2530 J	2480 J	328 J	621 J	603 J	742 J	142000 J
PCB-010	652 J	372 J	714 J	935 J	98.2 J	209 J	189 J	185 J	56300 J
PCB-011	700 J	425 J	701 J	703 J	314 J	446 J	643 J	661 J	8520 J
PCB-012/013	4150 J	2570 J	4550 J	4730 J	948 J	1380 J	3390 J	3660 J	120000 J
PCB-014	42.7 J	11.7 J	47.2 J	8.4 U	9.17 U	42 U	48.5 U	47.6 U	225 J
PCB-015	27800 J	18900 J	28300 J	36800 J	7190 J	11400 J	11300 J	12000 J	462000 J
PCB-016	83300 J	61200 J	82500 J	155000 J	13800	28500 J	29500 J	33200 J	968000 J
PCB-017	107000 J	78900 J	95600 J	181000 J	24000 J	46200 J	42600 J	48800 J	1100000 J
PCB-018/030	263000 J	205000 J	248000 J	494000 J	56900 J	113000 J	104000 J	121000 J	3200000 J
PCB-019	19400 J	14800	19700 J	35200 J	3360	6860	7730	8470	356000 J
PCB-020/028	340000 J	256000 J	286000 J	580000 J	91300 J	129000 J	138000 J	160000 J	3670000 J
PCB-021/033	169000 J	128000 J	156000 J	305000 J	35500 J	51800 J	59800 J	69400 J	2360000 J
PCB-022	99300 J	77000 J	87700 J	184000 J	22700 J	32600 J	40000 J	45600 J	1410000 J
PCB-023	114	116	144	250	39.1	40.6	63	75.7	2840
PCB-024	1450	1380	1740	2410	344	682	653	720	23600 J
PCB-025	14700	12000	13900	25000 J	3930	5370	6390	7230	186000 J
PCB-026/029	48300 J	38300 J	42200 J	87600 J	12800	18100 J	19900 J	22600 J	635000 J
PCB-027	13800	11000	12900	25900 J	2970	5700	5600	6400	144000 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB056	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
			PB059.1-1SS010-091102-N						
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N N	N N	N 10 cm	N 20 cm	N 20 cm	N N	N N	N N
X:	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
Perylene	410	251	624	101	1140	160	272	264	173
Phenanthrene	640	414	2030	623	8090	241	3170	675	9160
Pyrene	2220	1750	8800	1060	14400	1380	2330	2390	10200
Total 10 of 17 HPAH (U = 1/2)	15198	10282	32330	4512.5	56987	6881	10733	11675	23218
Total 17 PAH (U = 1/2)	16849.5	11008.3	41163	7021.5	80300	7636.6	16561	13182.8	64538
Total 7 of 17 LPAH (U = 1/2)	1651.5	726.3	8833	2509	23313	755.6	5828	1507.8	41320
Total cPAH TEF (7 minimum) (U = 1/2)	2272.3	1348.5	3997.1	541.58	6497.3	833.96	1213.46	1424.9	1655.1
Semivolatile Organics (μg/kg)									
2,2'-Oxybis (1-chloropropane)	441 U	302 U	937 U	319 U	237 U	313 U	260 U	241 U	826 U
Benzidine	R	R	R	R	R	R	R	R	R
Bis(2-ethylhexyl) phthalate	2230	1470	1030	2290	577	2390	198 J	922	1140
Cholesterol			-	-		-			
Di-n-octyl phthalate	441 U	302 U	937 U	319 U	237 U	313 U	260 U	241 U	826 U
Coprostanol				-					
Hexachlorobenzene	130 U	89 U	1300 U	90 U	700 U	91 UJ	74 UJ	360 UJ	5900 U
Hexachlorobutadiene	80 U	56 U	820 UJ	56 U	440 UJ	57 UJ	46 UJ	220 UJ	3700 UJ
Hexachloroethane	130 U	89 U	1300 U	90 U	700 UJ	49 J	74 UJ	360 UJ	5900 U
PCB Congeners (ng/kg)									
PCB-001	1480	1140 J	15800	962	46300 J	885	67.8 J	4110	28900 J
PCB-002	73.2	92.8 J	264	79.4	1270	119	36.8 UJ	310	749
PCB-003	782	399 J	6880	409	15900	482	28.9	1880	13700 J
PCB-004	12400 J	6580 J	129000 J	6740 J	219000 J	5880 J	580 J	21700 J	201000 J
PCB-005	415 J	137 J	3370 J	178 J	6780 J	167 J	20.6 J	507 J	4520 J
PCB-006	4970 J	1510 J	42200 J	2170 J	63300 J	1810 J	201 J	6890 J	70300 J
PCB-007	727 J	241 J	6200 J	322 J	13300 J	351 J	42.4 J	943 J	8200 J
PCB-008	29800 J	8230 J	276000 J	11000 J	343000 J	9780 J	965 J	37100 J	453000 J
PCB-009	1200 J	401 J	11100 J	545 J	19200 J	541 J	61.3 J	1630 J	18200 J
PCB-010	402 J	241 J	3680 J	252 J	7840 J	233 J	28.6 J	632 J	4950 J
PCB-011	605 J	664 J	1400 J	1010 J	1680 J	442 J	75.3 UJ	600 J	1600 J
PCB-012/013	3000 J	2800 J	15600 J	5420 J	18600 J	2370 J	171 J	3680 J	17300 J
PCB-014	44.2 U	43.5 UJ	90.1 U	50 U	49.5 U	46.3 U	36.8 U	34.8 J	44.2 J
PCB-015	18800 J	7530 J	145000 J	9610 J	122000 J	8960 J	902 J	21300 J	191000 J
PCB-016	62200 J	26900 J	620000 J	31000 J	393000 J	26900 J	2530	83700 J	693000 J
PCB-017	79900 J	36100 J	666000 J	40200 J	470000 J	30500 J	3530	93500 J	674000 J
PCB-018/030	209000 J	93800 J	2020000 J	108000 J	1300000 J	81100 J	8610	263000 J	1920000 J
PCB-019	15700	7210	139000 J	8790	106000 J	6700	666	21200 J	165000 J
PCB-020/028	245000 J	79600 J	1650000 J	111000 J	1170000 J	83700 J	8680	229000 J	1810000 J
PCB-021/033	112000 J	29600 J	933000 J	39300 J	613000 J	31600 J	2870	110000 J	1130000 J
PCB-022	72500 J	20900 J	529000 J	28000 J	379000 J	21100 J	2090	65500 J	637000 J
PCB-023	112	32.1	750	40.5	592	40.1	36.8 U	105	1070
PCB-024	1320	616	9030	620	8730	46.3 U	45.4 J	1360	12900 J
PCB-025	11800	3830	68800 J	5610	50600 J	3970	422	12200	87300 J
PCB-026/029	37600 J	12500	239000 J	17700	171000 J	13100	1300	35300 J	284000 J
PCB-027	11500	5310	93500 J	6330	66500 J	4880	514	13500	100000 J

Appendix A-1
BERA Sediment Dataset

TI.	DI /5C 2000	DI /50 2000	DI /50 2000	DI /50 2000	D1/FC 2000	2014 1100	2044 1100	2044 1100
Task:	· ·	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	2011 USC	2011 USC	2011 USC
Location ID:	PB074	PB081	PB093	PB097	PB101	PB068	PB069.1	PB070A
-		PB081-1SS010-091103-N				PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm
Sample Type:	N 2200220 0402	N 2200764 2707	N	N 2204402 4242	N 2204207.2650	N	N 2204205.050	N 2204007.255
X:	3200828.0183	3200761.3787	3200970.4291	3201103.1242	3201287.2658	3201209.861	3201205.859	3201087.255
Υ:	13830463.2171	13830005.1879	13828894.7843	13828566.1540	13828181.2285	13831121.51	13830973.3	13830903.51
Perylene	63.1	1170	47.2	37.8	119	278	275	155
Phenanthrene	268	250000	78.1	49.2	620	20000	1230	541 J
Pyrene	1020	81900	245	213	880	12700	3420	1240
Total 10 of 17 HPAH (U = 1/2)	3604.2	168118 J	1696.7	1319.3	5175.3	27980	12704	6224 J
Total 17 PAH (U = 1/2)	4719.4	1306518 J	1945.11	1435.19 J	6025.4	89480	16702.1	7626.7 J
Total 7 of 17 LPAH (U = 1/2)	1115.2	1138400	248.41	115.89 J	850.1	61500	3998.1	1402.7 J
Total cPAH TEF (7 minimum) (U = 1/2)	349.57	11368.8 J	244.92	154.01	649.34	1970.16	1339.1	708.096 J
Semivolatile Organics (µg/kg)	224.11	120011	224 11	242.11	22011			
2,2'-Oxybis (1-chloropropane)	221 U	1360 U	231 U	242 U	220 U			
Benzidine	R	R	R	R	R			
Bis(2-ethylhexyl) phthalate	1790	1220 J	245	266	132 J			
Cholesterol		6000 U	310 U	290 U	300 UJ			
Di-n-octyl phthalate	221 U	1360 U	231 U	242 U	220 U			
Coprostanol		6000 U	310 U	290 U	300 UJ	-		
Hexachlorobenzene	320 UJ	4000 U	67 U	68 U	63 U			
Hexachlorobutadiene	200 UJ	2500 U	42 U	43 U	40 U	-		
Hexachloroethane	320 UJ	4000 U	67 U	68 U	63 U			
PCB Congeners (ng/kg)	2670	427000	224	24.0	445			
PCB-001	2670	137000 J	331	34.9	115			
PCB-002	34.6	2540	94.4	54.3	46			
PCB-003	1180	82300 J	293	126 3720 J	81			
PCB-004 PCB-005	12800 J 297 J	183000 J 3810 J	14100 J 224 J	21.9 J	3170 J 90.5 J			
			15300 J	3060 J	5450 J			
PCB-006 PCB-007	4150 J 553 J	61900 J 8020 J	507 J	58.2 J	175 J			
PCB-007 PCB-008	22000 J	419000 J	13900 J	3740 J	3860 J			
PCB-009	929 J	14400 J	528 J	40.9 J	190 J			
PCB-010	366 J	3990 J	192 J	22.7 J	53.9 J			
PCB-010 PCB-011	340 J	849 J	2310 J	1080 J	1010 J			
PCB-012/013	1800 J	10900 J	7460 J	1750 J	2830 J			
PCB-012/013 PCB-014	36.8 U	46.3 U	9.39 J	45.9 U	48.1 U			
PCB-014 PCB-015	11500 J	194000 J	6280 J	1840 J	2220 J			
PCB-013	42900 J	281000 J	24000 J	4980	4710	 	 	
PCB-010 PCB-017	47100 J	377000 J	42400 J	37900 J	9560			
PCB-018/030	125000 J	779000 J	97300 J	34900 J	19900 J			
PCB-019	11000	97500 J	16100	17400	3410			
PCB-020/028	103000 J	905000 J	91000 J	110000 J	26000 J	 		
PCB-021/033	55300 J	606000 J	23800 J	4490	7460	 	 	
PCB-022	32800 J	295000 J	24000 J	16600	7320			
PCB-022 PCB-023	49.3	611	40.3	45.9 U	17.9			
PCB-023	569	6140	297 J	45.9 U	124			
PCB-025	6200	53600 J	21900 J	20100 J	7130			
PCB-025	15700 J	141000 J	26800 J	6210	9310			
PCB-020/029	6500	43900 J	5720	6700	1330			
1 00 041	0300	+3300 J	3/20	0700	1330		I	

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:	PB076.1-1SS010-20110802-N	PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
Perylene	548	367	339	61.2
Phenanthrene	1740	115000	89000	284 J
Pyrene	5790	37600	30400	2700 J
Total 10 of 17 HPAH (U = 1/2)	26020	73119	60605	6354 J
Total 17 PAH (U = 1/2)	29569.3	475519	374505	10073.1 J
Total 7 of 17 LPAH (U = 1/2)	3549.3	402400	313900	3719.1 J
Total cPAH TEF (7 minimum) (U = 1/2)	2691.83	3903.48	3572.41	439.422 J
Semivolatile Organics (μg/kg)				
2,2'-Oxybis (1-chloropropane)				
Benzidine				
Bis(2-ethylhexyl) phthalate				
Cholesterol				
Di-n-octyl phthalate				
Coprostanol				
Hexachlorobenzene				
Hexachlorobutadiene				
Hexachloroethane				
PCB Congeners (ng/kg)				
PCB-001				
PCB-002				
PCB-003				
PCB-004				
PCB-005				
PCB-006				
PCB-007				
PCB-008				
PCB-009				
PCB-010				
PCB-011				
PCB-012/013				
PCB-014				
PCB-015				
PCB-016				
PCB-017				
PCB-018/030			-	-
PCB-019			-	-
PCB-020/028				
PCB-021/033				
PCB-022				
PCB-023				-
PCB-024				-
PCB-025				
PCB-026/029				
PCB-027				
1 00 027		J		-=

Appendix A-1
BERA Sediment Dataset

Task	: RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID	*	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
	: PB001.1-1SS010-091027-N	PB001.2-1SS010-091027-N	PB001.3-1SS010-091027-N	PB002-1SS010-091027-N	PB004-1SS010-091027-N	PB005-1SS010-091027-N		PB007.2-1SS010-091028-N	
Sample Date	l .	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type		N N	N N	N N	N N	N N	N	N	N N
Sumple Type	3202382.3842	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011	3202037.3755	3201747.2539
Ý	13836546.3402	13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614	13836300.0036	13836110.2965	13836119.3947
PCB-031	465000 J	126000 J	46100 J	188000 J	96600 J	118000 J	115000 J	122000 J	80700 J
PCB-032	115000 J	26400 J	8740	34900 J	19700 J	24100 J	28900 J	27500 J	16000 J
PCB-034	2460	1150	430	2340	928	1040	710	926	636
PCB-035	7830	2680	1080	3280	1760	2290	2210	2420	1800
PCB-036	8.93 U	48.9 J	8.93 U	68.8	9.62 U	8.55 U	9.17 U	8.47 U	53.2
PCB-037	125000 J	29400 J	13500	45100 J	23900 J	30600 J	36000 J	32900 J	24000 J
PCB-038	8.93 U	279	1990	566	111	360	168	289	439
PCB-039	1800	583	208	878	466	532	9.17 U	589	403
PCB-040/071	252000 J	52800 J	17600	66000 J	45900 J	50600 J	78000 J	54900 J	36700 J
PCB-041	8.93 U	9490	2620	4370	9.62 U	6040	15300	6360	4740
PCB-042	172000 J	40900 J	13400	50700 J	36500 J	37000 J	54800 J	37800 J	25200 J
PCB-043	8.93 U	3820	1050	3010	9.62 U	2490	6970	2710	2230
PCB-044/047/065	558000 J	138000 J	45300 J	171000 J	119000 J	129000 J	186000 J	133000 J	89400 J
PCB-045	118000 J	20000 J	5840	24600 J	21100 J	17400 J	26200 J	18400 J	11400
PCB-046	43900 J	8910	2780	11400	7080	8230	11300	8970	5320
PCB-048	107000 J	20900 J	5780	20800 J	16100	20200 J	40200 J	23300 J	15500 J
PCB-049/069	311000 J	92400 J	30000 J	119000 J	81100 J	87800 J	117000 J	91000 J	62300 J
PCB-050/053	97500 J	22100 J	6820	27200 J	18000	20900 J	29000 J	22800 J	14400
PCB-051	8.93 U	7860	2390	9970	9.62 U	7920	9860	8800	5720
PCB-052	516000 J	137000 J	46900 J	169000 J	118000 J	130000 J	183000 J	134000 J	95000 J
PCB-054	1870	388	125	459	298	347	508	380	214
PCB-055	8.93 U	1130	469	9.8 U	9.62 U	992	9.17 U	8.47 U	799 J
PCB-056	8.93 U	53100 J	17700	57100 J	9.62 U	45800 J	86200 J	44400 J	34200 J
PCB-057	8.93 U	386	117	326	9.62 U	273	732	261	238
PCB-058	8.93 U	723	273	1050	9.62 U	568	1560 J	434	350
PCB-059/062/075	34000 J	9990	3200	12500	7510	9150	14000	9560	6370
PCB-060	8.93 U	22800 J	6620	14800	9.62 U	17500 J	45500 J	17600 J	15400 J
PCB-061/070/074/076	965000 J	242000 J	86200 J	300000 J	216000 J	241000 J	350000 J	232000 J	172000 J
PCB-063	18800 J	3420	1050	3080	2110	2570	6530	2530	2260
PCB-064	223000 J	49100 J	16200	59500 J	43300 J	46200 J	73400 J	48200 J	33400 J
PCB-066	558000 J	130000 J	45900 J	168000 J	132000 J	134000 J	187000 J	133000 J	95700 J
PCB-067	11900	2280	701	2210	1660	1870	3960	2020	1560
PCB-068	1420	1010	402	1440	831	799	715	679	551
PCB-072	3580	2380	872	3470	2100	2030	1730	1780	1490
PCB-073	8.93 U	8.93 U	8.93 U	409	9.62 U	8.55 U	9.17 U	280	172 J
PCB-077	62100 J	15000	8.93 U	18600	15000	16700	24300 J	16600	12300
PCB-078	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-079	8.93 U	1280	637	1520	9.62 U	1280	9.17 U	1260	712
PCB-080	8.93 U	186	8.93 U	9.8 U	9.62 U	176	356	8.47 U	7.58 U
PCB-081	8.93 U	438	132	344	378	404	865	461	361
PCB-082	76000 J	12300	5100	10900	11000	12500	24300 J	11500	9840
PCB-083	272000 J	8.93 U	2470	5550	77400 J	8.55 U	9280	8.47 U	5720
PCB-084	8.93 U	35100 J	13700	41500 J	34700 J	35100 J	48500 J	32600 J	26000 J
PCB-085/116	83200 J	17800	7300	15300	16200	17100	29500 J	16600	14000

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB011	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N N	0 - 10 CIII	N N	N N	0 - 10 CIII	0 - 10 CIII	N N	FD
Sample Type.	3201727.2504		IN 2201420 F706	3201223.8464	3201140.0114	IN 2201140 0114	IN 2201146 0FF0	3201281.6017	3201174.1294
Λ; ν.	13836239.7593	3201333.4751	3201420.5786	13836229.0694		3201140.0114	3201146.8550	13835711.9947	13835456.7389
Ti		13836187.7023	13836111.1432		13836081.5392	13836081.5392	13835702.1476		
PCB-031	135000 J	75700 J	36000 J	111000 J		489000 J	69000 J	66500 J	63900 J
PCB-032	24200 J	15400	7270	25900 J		125000 J	16500 J	14000	15500
PCB-034	1420	724	271	758		1940	719	580	437
PCB-035	2710	1530	1190	2330	-	6260	1240	1440	1270
PCB-036	50.3 J	27.7	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	43.9 U
PCB-037	32900 J	20900 J	10700	34100 J		126000 J	20200 J	20600 J	19300 J
PCB-038	305	130	126	178		698	150	453	170
PCB-039	596	383	222	584		76.9 U	39.1 U	216	43.9 U
PCB-040/071	58700 J	36900 J	18700 J	67200 J		315000 J	33500 J	30000 J	39600 J
PCB-041	5400	4340	3010	11600		67300 J	4820	3510	6500
PCB-042	46400 J	26700 J	13000	48900 J		202000 J	24700 J	21700 J	27700 J
PCB-043	2730	1960	1380	5160		29400 J	2320	1880	3330
PCB-044/047/065	151000 J	92100 J	46200 J	166000 J		680000 J	85200 J	77600 J	98500 J
PCB-045	19700 J	12900	6130	21200 J		112000 J	11900	8450	12400
PCB-046	8430	5510	2520	9180		53200 J	5380	3580	5450
PCB-048	18400 J	16000	8830	33900 J	-	150000 J	13800	12600	20600 J
PCB-049/069	109000 J	61300 J	31300 J	110000 J	-	393000 J	56500 J	55700 J	64300 J
PCB-050/053	23000 J	15100	7270	25500 J		134000 J	13600	11300	16100
PCB-051	9180	5120	2520	10200	-	42700 J	4590	4310	6030
PCB-052	155000 J	92000 J	47800 J	169000 J		663000 J	87700 J	83000 J	100000 J
PCB-054	326	215	117	385	-	2660	234	169	250
PCB-055	8.2 U	8.47 U	681	9.62 U	-	5900	477	475	776
PCB-056	45600 J	35100 J	17900 J	71100 J	-	250000 J	28300 J	26400 J	37700 J
PCB-057	288	202	167	559	-	2150	180	181	297
PCB-058	876	344	152	594		896	350	346	239
PCB-059/062/075	10900	6710	3390	12600	-	47800 J	6490	5670	7250
PCB-060	16000	12800	9070	33400 J		148000 J	9930	9810	17500
PCB-061/070/074/076	261000 J	167000 J	84500 J	319000 J		946000 J	132000 J	138000 J	169000 J
PCB-063	2580	1940	1320	5250		18100 J	1600	1580	2720
PCB-064	53000 J	34900 J	17900 J	67400 J		273000 J	31000 J	29400 J	38500 J
PCB-066	148000 J	91900 J	45400 J	176000 J		458000 J	68500 J	75600 J	84400 J
PCB-067	1860	1370	972	3360		10700	1260	1160	1830
PCB-068	1180	468	227	787		957	505	532	322
PCB-072	3000	1270	537	1900		2700	1220	1160	805
PCB-073	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	120
PCB-077	16200	11000	6570	22900 J		58800 J	8700	10100	10900
PCB-078	8.2 U	8.47 U	26.4 J	9.62 U		76.9 U	12.5	38.5 U	43.9 U
PCB-079	1440	576	380	989		3520	853	824	826 J
PCB-080	8.2 U	148	6.85 U	350		76.9 U	39.1 U	38.5 U	43.9 U
PCB-081	374	323	226	735		2680	221	255	335
PCB-082	11200	9590	6880	18700		72700 J	7750	6590	11700
PCB-083	5770	3670 J	3260	7390		26800 J	4040	3910	5690
PCB-084	40400 J	22600 J	13400	40800 J		145000 J	19500 J	16900 J	24000 J
				1	i				

Appendix A-1
BERA Sediment Dataset

	Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
	Location ID:	PB022	PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
					PB024-1SS010-091030-N					
	Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
	Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
	Sample Type:	N N	FD	N N	N N	N N	N N	0 - 10 CIII	O - 10 CIII	N N
	X:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
	λ. Υ:	13835456.7389	13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
DCD 034	1.									
PCB-031		67900 J	59900 J	57500 J	103000 J	74300 J	29400 J	5900	141000 J	431000 J
PCB-032		17000 J	18600 J	11700	23400 J	47600 J	7150	1360	27700 J	84700 J
PCB-034		460	620 J	574	810	784 J	233 593	45.7 1680	1030	1780
PCB-035		1230	1370 J	1100	2190	3130 J			2640	3710
PCB-036		37.6 U	231 UJ	183 U	6.94 U	596 J	9.17 U	295	83.9	8.13 U
PCB-037		20400 J	16800 J	13700 J	32000 J	21000 J	8060	3020	35300 J	89600 J
PCB-038		124	1660 J	1550	183	329 J	65.4	255	109	8.13 U
PCB-039		37.6 U	495 J	305	633	923 J	209	794	777	646
PCB-040/071		44400 J	56800 J	26600 J	44500 J	36700 J	14800	3160 809	76300 J	191000 J
PCB-041		7080	8190	4130	9780	3770	2870	2180	15300 J	31700 J
PCB-042		31900 J 3770	38000 J 4300	20200 J 2080	32200 J 3640	30300 J 2150 J	10400 1340	2180 318	54300 J 7810	141000 J 36100 J
PCB-043								9900		
PCB-044/047/065		108000 J	135000 J	73700 J	113000 J	117000 J	37000 J		193000 J	517000 J
PCB-045		14500	15500 J	9000	16400 J	13100 J	5010	1120	27800 J	73600 J
PCB-046		5810	6710	3590	6690	5370 J	1920	451	10600	35500 J
PCB-048		23200 J	23200 J	11100	22300 J	17800 J	7670	1800	41200 J	99600 J
PCB-049/069		70900 J	83700 J	49900 J	71900 J	98500 J	24000 J	5980	124000 J	294000 J
PCB-050/053		17300 J	17200 J	10600	18300 J	14300 J	5360	1380	31100 J	93900 J
PCB-051		6010	6050	3740	6180	4770 J	1720	429	10000	40900 J
PCB-052		107000 J	135000 J	80000 J	112000 J	128000 J	37900 J	14100	200000 J	377000 J
PCB-054		259 788	234	130	301	398 J	99.6	45 U 127	424	1410
PCB-055			2100 57400 J	1090 25500 J	1750 56200 J	718 J 30000 J	9.17 U 17900	3560	2770 91900 J	8.13 U 273000 J
PCB-056 PCB-057		37100 J 323	335	154	440	364 J	164	45 U		6150
PCB-057 PCB-058		265	480	416		821 J			701 765	
PCB-058 PCB-059/062/075		8030	10200	5370	441 8400	13400 J	124 2790	45 U 860	14400	8.13 U 30500 J
PCB-059/062/075 PCB-060		17600 J	18000 J	8300	24500 J	8390	9690	1960	47000 J	
PCB-061/070/074/076		167000 J	219000 J	117000 J	226000 J	151000 J	76900 J	19800 J	385000 J	123000 J 896000 J
PCB-061/070/074/076		2860	3370	1600	3920	2150 J	1480	45 U	7420	18700 J
PCB-064		43800 J	54000 J	27800 J	46000 J	37900 J	15200	4320	77700 J	216000 J
PCB-066		89000 J	120000 J	62300 J	123000 J	90100 J	40100 J	7760	198000 J	588000 J
PCB-0667		1870	2230	914	2520	2080 J	821	45 U	4230	10300
PCB-068		368	677	502	515	3810 J	197	192	1010	1450
PCB-072		842	1620	1120	1280	6060 J	506	45 U	2300	3120
PCB-072		37.6 U	231 U	183 U	6.94 U	40 UJ	9.17 U	45 U	7.63 U	117000 J
PCB-073		11500	14000	6860	15700 J	8600	4620	1190	22200 J	55900 J
PCB-077		17.7	231 U	183 U	6.94 U	40 UJ	9.17 U	278	7.63 U	8.13 U
PCB-079		747	987	470	819	1630 J	252	319	1240	5150
PCB-080		37.6 U 386	231 U 372	183 U 136	6.94 U 494	40 UJ	9.17 U 157	45 U 429	7.63 U 837	8.13 U 3870
PCB-081				6800		257	4660			
PCB-082		10400	17800 J		14100 J	6540		2310	26900 J	76600 J
PCB-083		4870	8320	4140	5290	4210	2270	1270	12700	8.13 U
PCB-084		21600 J	37100 J	18300 J	29200 J	30900 J	9130	5540	53500 J	169000 J
PCB-085/116		15600 J	22500 J	9150	18700 J	9960	6610	3260	37000 J	60700 J

Appendix A-1
BERA Sediment Dataset

Tas	k: RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location I		PB043	PB044	PB047.1	PB047.2	PB048	PB049	РВ049	PB053
			· -				PB049-1SS010-091031-D		
Sample Dat		10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Dept		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Typ		N N	N N	N 20 5	N N	N 20 cm	FD	N N	N N
Sumple Typ	X: 3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
	Y: 13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
DCD 024						144000 J			
PCB-031 PCB-032	406000 J 81800 J	309000 J 63800 J	331000 J 72100 J	729000 J 156000 J	101000 J 20000 J	36800 J	159000 J 35500 J	182000 J 40500 J	4750000 J 827000 J
					1				
PCB-034	1680	928	1620	2030	544	850	736	826	13800
PCB-035	8350	4820	5670	10900	2040	3140	3280	3550	157000 J
PCB-036	8.2 U	7.75 U	9.71 U	8.4 U	23.8	42 U	48.5 U	47.6 U	48.1 U
PCB-037	105000 J	79200 J	86500 J	165000 J	26900 J	40800 J	40400 J	45600 J	1240000 J
PCB-038	240	196	198	466	71.5	82	99.1	137	3560
PCB-039	2210	1740	1660	3910	614	42 U	48.5 U	47.6 U	48.1 U
PCB-040/071	216000 J	180000 J	170000 J	459000 J	55200 J	84900 J	88100 J	106000 J	2850000 J
PCB-041	53600 J	51200 J	45200 J	124000 J	11800	16100	24600 J	30500 J	829000 J
PCB-042	150000 J	122000 J	116000 J	304000 J	38900 J	56800 J	60700 J	72200 J	1810000 J
PCB-043	20900 J	19100 J	17100	46200 J	5850	7810	9940	11900	296000 J
PCB-044/047/065	524000 J	417000 J	401000 J	1080000 J	141000 J	206000 J	216000 J	257000 J	6420000 J
PCB-045	86100 J	72500 J	74000 J	202000 J	19700 J	29400 J	33700 J	40400 J	1190000 J
PCB-046	35700 J	28400 J	30600 J	84400 J	6790	10800	12900	15400	509000 J
PCB-048	122000 J	108000 J	97500 J	260000 J	32500 J	46200 J	52000 J	62600 J	1630000 J
PCB-049/069	321000 J	253000 J	232000 J	612000 J	92300 J	132000 J	134000 J	160000 J	3550000 J
PCB-050/053	90100 J	76100 J	73900 J	207000 J	23900 J	31600 J	36200 J	43100 J	1130000 J
PCB-051	28900 J	21700 J	21200 J	58900 J	7930	10200	11600	14000	343000 J
PCB-052	527000 J	405000 J	385000 J	1090000 J	150000 J	212000 J	215000 J	256000 J	5830000 J
PCB-054	1340	1060	1050	3080	278	577	618	686	16600
PCB-055	8290	9890	5120	14600	1900	3180	2640	4350	81700 J
PCB-056	247000 J	207000 J	183000 J	489000 J	69400 J	104000 J	106000 J	127000 J	2740000 J
PCB-057	2130	1910	1580	4340	566	784	967	1170	26100 J
PCB-058	1480	802	807	1810	352	521	498	594	8770
PCB-059/062/075	38900 J	31900 J	30700 J	77500 J	10400	15400	16400	19600 J	477000 J
PCB-060	145000 J	135000 J	103000 J	299000 J	38200 J	53700 J	56800 J	70400 J	1810000 J
PCB-061/070/074/076	1010000 J	805000 J	702000 J	1820000 J	279000 J	412000 J	402000 J	480000 J	9080000 J
PCB-063	19800 J	17500 J	14600	37900 J	5740	8070	8810	10500	233000 J
PCB-064	207000 J	169000 J	160000 J	417000 J	57700 J	84800 J	87000 J	104000 J	2500000 J
PCB-066	493000 J	401000 J	346000 J	853000 J	147000 J	220000 J	212000 J	251000 J	4380000 J
PCB-067	12300	10800	8940	23500 J	3290	4660	4860	5860	143000 J
PCB-068	1450	679	854	1350	430	605	570	617	8120
PCB-072	4160	2090	2350	4510	1080	1610	1490	1640	26700 J
PCB-073	8.2 U	7.75 U	9.71 U	8.4 U	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-077	63900 J	45900 J	39400 J	99100 J	18600 J	28900 J	23700 J	26900 J	541000 J
PCB-078	154	7.75 U	85.3	8.4 U	25.2	42 U	48.5 U	47.6 U	1860
PCB-079	3260	2350	1840	5240	874	1340	1200	1430	25600 J
PCB-080	1180	1070	890	2670	395	42 U	48.5 U	47.6 U	48.1 U
PCB-081	2830	2240	1700	4640	729	1050	1040	1220	29000 J
PCB-082	80100 J	59800 J	46300 J	169000 J	19200 J	25400 J	26500 J	32000 J	571000 J
PCB-083	31000 J	24400 J	16000	70000 J	8980	9810	10200	14600	230000 J
PCB-084	151000 J	111000 J	90800 J	339000 J	34500 J	47600 J	50000 J	57600 J	1090000 J
PCB-085/116	87300 J	63000 J	50000 J	166000 J	24900 J	33400 J	32500 J	38600 J	609000 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB056	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
Sample ID:	PB056-1SS010-091031-N	PB057-1SS010-091031-N	PB059.1-1SS010-091102-N	PB059.2-1SS010-091031-N	PB063.1-1SS010-091031-N	PB063.2-1SS010-091102-N	PB064-1SS010-091102-N	PB066-1SS010-091101-N	PB069-1SS010-091101-N
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	N
x:	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
PCB-031	284000 J	92700 J	2040000 J	124000 J	1390000 J	91300 J	8940	262000 J	1920000 J
PCB-032	66600 J	32100 J	546000 J	36000 J	368000 J	27000 J	2950	75700 J	502000 J
PCB-034	1190	410	5090	567	3570	391	58.9	1090	6040
PCB-035	5020	2100	22300 J	3390	23600 J	2250	219	3970	20100 J
PCB-036	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-037	79300 J	22100 J	535000 J	28900 J	398000 J	24600 J	2370	69300 J	583000 J
PCB-038	173	73.1	1950	104	910	121	36.8 U	274	1610
PCB-039	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	11900
PCB-040/071	174000 J	58500 J	1290000 J	106000 J	859000 J	76100 J	7520	200000 J	1340000 J
PCB-041	51800 J	13100	337000 J	29300 J	297000 J	19100 J	1200	46600 J	290000 J
PCB-042	116000 J	38400 J	826000 J	70100 J	581000 J	53300 J	5190	134000 J	846000 J
PCB-043	18300 J	6210	140000 J	11100	91500 J	8480	696	20700 J	139000 J
PCB-044/047/065	419000 J	140000 J	2860000 J	249000 J	1990000 J	181000 J	17700 J	449000 J	2940000 J
PCB-045	68500 J	22200 J	496000 J	37400 J	357000 J	28600 J	2520	76700 J	486000 J
PCB-046	28500 J	8630	208000 J	16400	150000 J	12100	1070	32500 J	209000 J
PCB-048	97700 J	32400 J	719000 J	56000 J	503000 J	42500 J	3550	107000 J	686000 J
PCB-049/069	248000 J	86800 J	1640000 J	146000 J	1140000 J	107000 J	10700	256000 J	1560000 J
PCB-050/053	69100 J	22700 J	503000 J	40900 J	351000 J	30700 J	2740	80300 J	499000 J
PCB-051	21500 J	7030	142000 J	14100	105000 J	9010	868	23500 J	139000 J
PCB-052	405000 J	138000 J	2690000 J	243000 J	1890000 J	170000 J	16400 J	413000 J	2620000 J
PCB-054	1200	513	9200	683	5810	501	44.9 J	1390	10200
PCB-055	6060	2590	28900 J	3910	28100 J	1410	108	3740	23400 J
PCB-056	202000 J	69000 J	1130000 J	116000 J	894000 J	61800 J	6320	155000 J	1110000 J
PCB-057	1840	645	11800	1020	9150	589	54.2 J	1560	11300
PCB-058	801	285	3360	466	3540	292	36.9 J	724	2920
PCB-059/062/075	32400 J	10400	214000 J	18400	152000 J	13200	1330	32900 J	203000 J
PCB-060	120000 J	37100 J	745000 J	63500 J	616000 J	30700 J	3000	88700 J	646000 J
PCB-061/070/074/076	756000 J	265000 J	4430000 J	425000 J	3200000 J	233000 J	24200 J	595000 J	4050000 J
PCB-063	16300	5610	90600 J	9190	78300 J	4670	472	12500	84400 J
PCB-064	167000 J	55300 J	1150000 J	96300 J	806000 J	69100 J	7060	172000 J	1080000 J
PCB-066	388000 J	139000 J	2090000 J	226000 J	1630000 J	126000 J	13500	302000 J	1990000 J
PCB-067	9850	3150	55500 J	5040	47100 J	2980	301	7940	58300 J
PCB-068	838	315	3140	493	2640	277	41.9	646	3290
PCB-072	2460	896	9440	1440	8740	803	114	1790	9630
PCB-073	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-077	42800 J	15500	280000 J	24500 J	183000 J	15700	1530	37200 J	236000 J
PCB-078	44.2 U	43.5 U	90.1 U	50 U	352	35.1 J	36.8 U	59 J	357 J
PCB-079	2060	936	15800	1280	9820	1070	117	1870 J	11500
PCB-080	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-081	1970	749	12800	1230	9950	659	65.9	1680	12000
PCB-082	49800 J	19400 J	345000 J	35200 J	206000 J	18300	1430	42600 J	276000 J
PCB-083	21800 J	8880	144000 J	13000	84300 J	6690	618	17100 J	101000 J
PCB-084	93100 J	33600 J	628000 J	62800 J	384000 J	34600 J	3250	83800 J	456000 J
PCB-085/116	58000 J	23700 J	386000 J	38400 J	221000 J	21800 J	1930	49500 J	303000 J

Appendix A-1
BERA Sediment Dataset

Task: Location ID: Sample ID: Sample Date: Depth: Sample Type: X:	11/1/2009 0 - 10 cm	RI/FS 2009 PB081 PB081-1SS010-091103-N 11/3/2009	RI/FS 2009 PB093	RI/FS 2009 PB097	RI/FS 2009 PB101	2011 USC PB068	2011 USC PB069.1	2011 USC
Sample ID: Sample Date: Depth:	PB074-1SS010-091101-N 11/1/2009 0 - 10 cm	PB081-1SS010-091103-N		1 5037				PB070A
Sample Date: Depth:	11/1/2009 0 - 10 cm			DR097_1SS010_091103_N	l	PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Depth:	0 - 10 cm		11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
•		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm
X: Y:	N	N N	N N	N N	N N	N	N N	N N
Y:	3200828.0183	3200761.3787	3200970.4291	3201103.1242	3201287.2658	3201209.861	3201205.859	3201087.255
	13830463.2171	13830005.1879	13828894.7843	13828566.1540	13828181.2285	13831121.51	13830973.3	13830903.51
PCB-031	123000 J	984000 J	91000 J	52100 J	26800 J			
PCB-032	37700 J	248000 J	34500 J	61200 J	8140		-	
PCB-034	444	3310	1080	1020	325		-	
PCB-035	1690	11500	2220	622	733		1	
PCB-036	36.8 U	46.3 U	36.8	45.9 U	48.1 U		-	
PCB-037	30500 J	350000 J	11900	5730	4200		-	
PCB-038	111	579	27.1 J	45.9 U	48.1 U		-	
PCB-039	36.8 U	46.3 U	397	300	136		-	
PCB-040/071	76600 J	575000 J	36700 J	63700 J	12600		-	
PCB-041	16700 J	173000 J	5550	1940	1300		-	
PCB-042	49900 J	365000 J	26300 J	48600 J	9160		-	-
PCB-043	7880	58900 J	3830	7240	1440	-		
PCB-044/047/065	180000 J	1230000 J	87100 J	167000 J	29700 J			
PCB-045	30000 J	228000 J	17700	8330	4990			
PCB-046	13600	105000 J	9350	18100	2710			
PCB-048	41200 J	327000 J	14400	8740	4680			
PCB-049/069	101000 J	681000 J	57200 J	151000 J	19900 J			
PCB-050/053	33300 J	229000 J	24000 J	60600 J	7490			
PCB-051	10500	82600 J	6650	24700 J	2330			
PCB-052	185000 J	1160000 J	86300 J	162000 J	29400 J			
PCB-054	672	3900	634	1650	179			
PCB-055	1430	20600 J	786	1430	566			
PCB-056	55900 J	541000 J	19700	25400 J	7070			
PCB-057	558	5780	419	452	147			
PCB-058	216	1460	256	590	89.4			
PCB-059/062/075	12300	102000 J	5860	7770	2080			
PCB-060	33400 J	362000 J	6780	6100	2350			
PCB-061/070/074/076	229000 J	1910000 J	54100 J	97500 J	26900 J			
PCB-063	4720	43900 J	2420	4990	887			
PCB-064	66700 J	510000 J	30200 J	29800 J	10600			
PCB-066	107000 J	913000 J	38800 J	79100 J	13600			
PCB-067	3020	28900 J	1240	807	426			
PCB-068	269	1550	289	1120	122			
PCB-072	707	4880	567	1710	237			
PCB-073	36.8 U	46.3 U	392	1800	151			
PCB-077	14400	101000 J	3630	7290	1400		-	
PCB-078	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-079	1300	4190	176	338	94			
PCB-080	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-081	583	6220	74.8	119	39.3			
PCB-082	19800 J	114000 J	3210	4320	1140			
PCB-083	10800	52300 J	1790	3760	577		-	
PCB-084	48600 J	223000 J	10700	12900	3790		-	
PCB-085/116	22200 J	117000 J	3910	5160	1090			

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:		PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
PCB-031				
PCB-032				
PCB-034				
PCB-035				
PCB-036				
PCB-037				
PCB-038				
PCB-039				
PCB-040/071				
PCB-041				
PCB-042				
PCB-043				
PCB-044/047/065				
PCB-045				
PCB-046				
PCB-048				
PCB-049/069				
PCB-050/053				
PCB-051				
PCB-052				
PCB-054				
PCB-055				
PCB-056				
PCB-057				
PCB-058			-	
PCB-059/062/075			-	
PCB-060			-	
PCB-061/070/074/076			-	
PCB-063				
PCB-064				
PCB-066			-	
PCB-067			-	
PCB-068			-	
PCB-072				
PCB-073				
PCB-077				-
PCB-078				
PCB-079				-
PCB-080				
PCB-081				
PCB-082				
PCB-083				
PCB-084				
PCB-085/116				

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009						
Location ID:	PB001.1	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
							PB007.1-1SS010-091028-N		
Sample Date:	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm						
Sample Type:	N N	N N	N 10 cm	N N	N N	N N	N 10 cm	N N	N N
y.	3202382.3842	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011	3202037.3755	3201747.2539
γ.	13836546.3402	13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614	13836300.0036	13836110.2965	13836119.3947
PCB-086/087/097/108/119/125	275000 J	58500 J	26200 J	61500 J	57300 J	63400 J	105000 J	60100 J	51700 J
PCB-088	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	21800 J	9.17 U	8.47 U	7.58 U
PCB-089	8.93 U	2310	697	2060	1780	2010	4310	1910	1560
PCB-090/101/113	328000 J	89700 J	39800 J	112000 J	93500 J	98400 J	139000 J	93600 J	79700 J
PCB-091	8.93 U	18900 J	6800	23900 J	16900	8.55 U	23800 J	17500 J	13800
PCB-092	8.93 U	17200	7630	18200	13800	14600	24600 J	12800	12200
PCB-093/100	17.9 U	1840	601	2080	19.2 U	1530	2640	1550	1200
PCB-094	8.93 U	721	238	607	591	542	1190	517	445
PCB-095	8.93 UJ	77200 J	32100 J	96300 J	74800 J	81000 J	104000 J	76200 J	64500 J
PCB-096	7880	1850	618	2280	1640	1940	2730	1810	1250
PCB-098	8.93 U	8.93 U	8.93 U	7580	9.62 U	6510	9.17 U	6670	7.58 U
PCB-099	8.93 UJ	47300 J	18000 J	55800 J	9.62 UJ	4400 J	66400 J	3830 J	35900 J
PCB-102	8.93 U	5720	1940	9.8 U	6460	8.55 U	8240	8.47 U	4130
PCB-103	8.93 U	1490 J	483	2420	1350	1390	1500	1310	1030
PCB-104	104 J	41.7	13.9 J	52.6	41.6	38.5	48.3	36.9	27.5 J
PCB-105	181000 J	32700 J	15600 J	32100 J	31400 J	34700 J	68000 J	33300 J	30100 J
PCB-106	8.93 U	8.93 U	8.93 U	48.2	9.62 U	80.2	168	86.8	70.9
PCB-107/124	12500	2670	1380	2480	2490	2910	5050	2780	2580
PCB-109	22900 J	5320	2490	6180	4610	5280	9740	5200	4620
PCB-110	332000 J	90000 J	40600 J	108000 J	88500 J	94300 J	142000 J	93500 J	81000 J
PCB-111	8.93 U	8.93 U	89.5	162	9.62 U	95	9.17 U	105	89.7
PCB-112	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	49500 J	9.17 U	48600 J	7.58 U
PCB-114	12000	1970	813	1420	1720	1950	4280	1790	1630
PCB-115	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-117	8.93 U	8.93 U	8.93 U	3330	9.62 U	2010	5670	8.47 U	7.58 U
PCB-118	294000 J	68000 J	33000 J	75900 J	66800 J	73900 J	121000 J	72600 J	62800 J
PCB-120	888	523	354	925	9.62 U	538	539	486	427
PCB-121	8.93 U	8.93 U	39.1	74.8	9.62 U	24.4 J	9.17 U	33.5	7.58 U
PCB-122	7550	1530	651	1450	1450	1670	3060	1520	1320
PCB-123	8790	1870	824	1750	2380	2140	3570	1830	1680
PCB-126	2040	382	247	381	498	416	630	386	364
PCB-127	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-128/166	25400 J	8210	5750	9920	8170	9030	12300	8260	9010
PCB-129/138/163	132000 J	49000 J	31600 J	68900 J	48700 J	54000 J	69600 J	48700 J	51700 J
PCB-130	12300	5320	3600	7740	4870	5290	6780	4760	4970
PCB-131	8.93 U	805	542	951	819	879	1150	841	856
PCB-132	8.93 U	17500	10800	23100 J	15800	18500 J	23700 J	16800	17900 J
PCB-133	8.93 U	1230	777	2020	1040	1050	1270	961	922
PCB-134	8700	3680	2230	9.8 U	3620	3590	5420	3660	3870
PCB-135/151	37300 J	16900	9060	26800 J	14700	15900	21200 J	14100	14600
PCB-136	17800	7590	4090	12500	7020	8290	9030	7330	7120
PCB-137	8300	1800	1370	1940	9.62 U	2140	2960	2330	2190
PCB-139/140	2340	1210	729	1700	1120	1210	1580	1210	1170
PCB-141	19500 J	7120	4510	9920	7490	8300	11300	7580	8050

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB011	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
•	10/28/2009	10/28/2009			11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Sample Date:			10/28/2009	10/28/2009					
Depth:	0 - 10 cm N	0 - 10 cm N	0 - 10 cm	0 - 10 cm	0 - 10 cm N	0 - 10 cm	0 - 10 cm	0 - 10 cm N	0 - 10 cm FD
Sample Type:	3201727.2504		IN 2201420 F796	3201223.8464	3201140.0114	IN 2201140 0114	N 2201146 0550	3201281.6017	3201174.1294
X.		3201333.4751	3201420.5786	13836229.0694		3201140.0114	3201146.8550	13835711.9947	13835456.7389
T:	13836239.7593	13836187.7023	13836111.1432		13836081.5392	13836081.5392	13835702.1476		
PCB-086/087/097/108/119/125	65200 J	46200 J	32400 J	88200 J		276000 J	39500 J	38200 J	55700 J
PCB-088	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	43.9 U
PCB-089	1860	1580	1030	3330		13400	1290	986	1860
PCB-090/101/113	110000 J	68600 J	45500 J	126000 J		312000 J	60100 J	60400 J	76100 J
PCB-091	22700 J	11900	6870	21700 J		65900 J	11400	10400	13100
PCB-092	17500 J	10600	7310	21000 J		56100 J	9740	9280	12900
PCB-093/100	2280	1100	702	2380		6450	1110	858	1140
PCB-094 PCB-095	660 96700 J	434 53600 J	281 32000 J	931 93200 J		3270 284000 J	373 48800 J	293 44200 J	533 56400 J
PCB-095	1970	1170	706	2380		284000 J 8600	48800 J 1100	936	1440
PCB-098	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	43.9 U
PCB-099	52800 J	33000 J	20600 J	63100 J		143000 J	26600 J	27300 J	32900 J
PCB-099	6210	3820	2320	7070		20700 J	2920	2760	3750
PCB-102 PCB-103	2030	842	459	1500		2620	935	748	659
PCB-105 PCB-104	44.4	22	10.9 J	40.8		117 J	33.5 J	21.3 J	28.3 J
PCB-104 PCB-105	31900 J	27500 J	21700 J	57700 J		170000 J	22100 J	22800 J	37700 J
PCB-103	8.2 U	8.47 U	38.7 J	9.62 U		301	39.1 U	38.5 U	43.9 U
PCB-100 PCB-107/124	2720	2150	1820	4400		12300	1790	1880	3030
PCB-107/124 PCB-109	5710	4090	3300	8670		20600 J	3490	3670	5190
PCB-109	116000 J	68900 J	46900 J	127000 J		328000 J	56000 J	55600 J	71700 J
PCB-111	73.8	43.7	41.2 J	9.62 U		77.4	71.6	56.8	32.2 J
PCB-112	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	43.9 U
PCB-114	1740	1520	1290	3310		11700	1170	1220	2400
PCB-115	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	43.9 U
PCB-117	2230 J	2290	1220	4150		9570	1710	1270	2500
PCB-118	75500 J	55200 J	42300 J	110000 J		277000 J	47700 J	51400 J	75200 J
PCB-120	605	314	204	555		76.9 U	409	373	220
PCB-121	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	19.5	43.9 U
PCB-122	1500	1270	946	2480		7350	919	896	1580
PCB-123	2040	1580	1150	3140		7950	1230	1250	1840
PCB-126	403	307	268	637		1140	246	282	345
PCB-127	8.2 U	8.47 U	6.85 U	9.62 U		76.9 U	39.1 U	38.5 U	43.9 U
PCB-128/166	11100	6900	5460	11000		25600 J	5870	5920	8720
PCB-129/138/163	62000 J	38900 J	29300 J	67400 J		146000 J	42900 J	41500 J	56600 J
PCB-130	5940	3880	2810	6580		13000	4020	3930	4970
PCB-131	957	662	494	1050		2980	626	579	888
PCB-132	21500 J	13900	10200	22100 J		55200 J	13800	12600	17800 J
PCB-133	1280	718	489	1330		1750	864	739	773
PCB-134	5130	3000	2160	4800		10600	2640	2310	3170
PCB-135/151	21400 J	11700	7890	20900 J		38600 J	12900	11000	13300
PCB-136	11200	5570	3620	8940		20800 J	7140	5750	6680
PCB-137	2650	1730	1430	2620		7640	1630	1740	2750
PCB-139/140	1470	907	640	1470		3090	846	812	1040
PCB-141	9490	6160	4770	10800		23900 J	6670	6220	8780

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	1	PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
Sample ID:				PB024-1SS010-091030-N					
Sample Date:		10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:		FD	0 - 10 CIII	N N	N N	N N	N N	0 - 10 CIII	N N
X:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
Λ· Υ:		13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
PCB-086/087/097/108/119/125	48600 J	75000 J	37200 J	62800 J	60500 J	21200 J	16700	148000 J	298000 J
PCB-088	37.6 U	231 U	183 U	6.94 U	40 U	9.17 U	45 U	7.63 U	8.13 U
PCB-089	1810	3330	1080	2390	1390	815	1000	4440	13700
PCB-090/101/113	64600 J	105000 J	62700 J	88000 J	144000 J	30700 J	24600 J	171000 J	373000 J
PCB-091	11500	19100 J	10100	14700 J	17700 J	5010	2260	26800 J	44100 J
PCB-092	10900	18300 J	10700	15200 J	24200 J	5450	4150	31400 J	67200 J
PCB-093/100	1050	2150	960	1440	4930	572	90.1 U	3110	16.3 U
PCB-094	499	778	341	623	495	229	70.7	1280	4240
PCB-095	47800 J	83600 J	50700 J	67100 J	101000 J	20900 J	17000	120000 J	273000 J
PCB-096 PCB-098	1420 37.6 U	2130 231 U	852 183 U	1720 192	1580 1760	621 76.6	183 45 U	2690 639	7670 8.13 U
		48400 J	26600 J		96900 J	15300	7710	78300 J	200000 J
PCB-099	32000 J			41200 J					
PCB-102	4070	6580	2690	4830	3600	1640	45 U	8210	30000 J
PCB-103	639	1320	942	839	8120	335	131	1740	2360
PCB-104	26.9 J	51.4	183 U	23.8	177	7.97 J	45 U	48.9	93.5
PCB-105	32100 J	41500 J	19400 J	39300 J	20700 J	13700	8120	72700 J	175000 J
PCB-106	37.6 U	231 U	183 U	6.94 U	40 U	9.17 U	388	603 J	8.13 U
PCB-107/124	2420	3350	1600	3040	2000	1060	966	5840	13100
PCB-109	4370	6720	3670	6030	9030	2220	1350	11300	32300 J
PCB-110	57400 J	97200 J	55500 J	82400 J	99500 J	27100 J	19600 J	155000 J	370000 J
PCB-111	32.9	231 U	72.3	48.3	2980	9.17 U	56	136	8.13 U
PCB-112	37.6 U	405	237	6.94 U	1330	9.17 U	45 U	749	8.13 U
PCB-114	2010	1910	798	2040	2020	761	1870	3950	11600
PCB-115	37.6 U	3270	1440	1970	910	864	409	7.63 U	8.13 U
PCB-117	37.6 U	2100	2010	2520	4300	813	1760	7.63 U	8.13 U
PCB-118	61400 J	77500 J	41800 J	73500 J	82100 J	25300 J	17600	121000 J	294000 J
PCB-120	208	490	403	289	8050	132	308	816	8.13 U
PCB-121	37.6 U	231 U	183 U	6.94 U	1790	9.17 U	96.1	7.63 U	8.13 U
PCB-122	1360	2310	936	1500	1200 J	556	429 J	3400 J	7450
PCB-123	1710	2390	979	1900	1100	629	557	4620	8.13 U
PCB-126 PCB-127	299	525 231 U	251	388 6.94 U	40 U 40 U	155	180 45 U	647 92.4	2340 8.13 U
PCB-127 PCB-128/166	37.6 U 5810	8800	183 U 6740	9360	8680	9.17 U 2680	3800	92.4 18000 J	28300 J
PCB-128/166 PCB-129/138/163	35400 J	64900 J	49400 J	59300 J		18000	27800 J	18000 J 104000 J	28300 J 154000 J
	33400 J	6810	49400 J 5070	59300 J	89500 J 19500 J	1710	27800 J 2500	9680	13100
PCB-130	607	969	646		911			1640	2910
PCB-131				892		9.17 U 5730	388 8790		
PCB-132	12000	21900 J	16100 J 1080	19900 J 892	42700 J	357		33700 J	55300 J
PCB-133	583	1340			23100 J		11100	1850	2100
PCB-134	2100	4270	3240 16700 I	3570	8390	1170	1560	6780	12100
PCB-135/151	9970	21200 J	16700 J	16700 J	58200 J	5570	8630	30500 J	43000 J
PCB-136	4740	9410	8150	7880	28400 J	2360	3640	12700	18700 J
PCB-137	1750	2090	1540 968	2160	1330	666	1530	3820	18600 J
PCB-139/140	731	1350		1080	6320	349	613	1980	3290
PCB-141	5990	10200	7880	9780	7860	2780	5650	16700 J	21100 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB041	PB043	PB044	PB047.1	PB047.2	PB048	PB049	PB049	PB053
			_				PB049-1SS010-091031-D		
Sample Date:	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N N	N N	N N	N N	N N	FD	N	N
v.	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
ν. •	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
DCD 00C/007/007/400/440/43F									
PCB-086/087/097/108/119/125 PCB-088	326000 J 8.2 U	236000 J 7.75 U	181000 J 9.71 U	671000 J 8.4 U	85500 J 9.17 U	109000 J 42 U	111000 J 48.5 U	129000 J 47.6 U	2060000 J 48.1 U
PCB-089	13900	11100	8520	29700 J	3380	4640	4970	5980	124000 J
PCB-089 PCB-090/101/113	421000 J	284000 J	229000 J	822000 J	113000 J	143000 J	144000 J	163000 J	2280000 J
PCB-091	74200 J	52400 J	41500 J	141000 J	18900 J	26400 J	25900 J	30700 J	524000 J
PCB-092	74600 J	50500 J	41300 J	152000 J	20200 J	25000 J	26400 J	29400 J	425000 J
PCB-093/100	7800	5570	4460	13300	2030	2520	2610	3010	56800 J
PCB-094 PCB-095	3570 315000 J	2820 218000 J	2170 181000 J	7380 682000 J	928 77900 J	1220 103000 J	1310 108000 J	1570 121000 J	29800 J 1970000 J
PCB-095	7980	6310	5480	18200 J	1900	3350	3200	3560	71800 J
PCB-098	8.2 U	7.75 U	9.71 U	8.4 U	6890	484	357	47.6 U	8830
PCB-099	202000 J	127000 J	107000 J	341000 J	9.17 U	69000 J	66300 J	76800 J	1100000 J
PCB-099	25500 J	19000 J	14600	49000 J	9.17 U	8780	8650	10900	191000 J
PCB-102 PCB-103	3890	2240	1870	5740	9.17 0	1260	1170	1330	20400 J
	115	75.5	55.9	186	27.2	44.3	36.7	32.7 J	740
PCB-104									
PCB-105	195000 J	151000 J	118000 J	394000 J	54300 J	73100 J	71400 J	84600 J	1220000 J
PCB-106 PCB-107/124	544	7.75 U 11000	270 9130	8.4 U 33300 J	126 4380	42 U 5720	145 5770	242 6500	3220 93300 J
	16100								
PCB-109	28500 J	26000 J	15800	51300 J	7810	9810	10100	11600	150000 J
PCB-110	383000 J	272000 J	207000 J	766000 J	99700 J	128000 J	129000 J	146000 J	2020000 J
PCB-111	8.2 U	7.75 U	9.71 U	1940	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-112	1680	6470	680	8.4 U	53200 J	796	592	672	10900
PCB-114	13400	11300	7690	27900 J	3520	4730	5230	5860	84300 J
PCB-115	10500	7.75 U	6570	26600 J	4200	5080	4370	5390	93100 J
PCB-117	18400 J	7.75 U	9590	33700 J	5580	5010	5440	6460	89800 J
PCB-118	351000 J	250000 J	207000 J	681000 J	96700 J	126000 J	135000 J	146000 J	1890000 J
PCB-120 PCB-121	1080	219	437	833	228	309	227	268	2600
	8.2 U 9970	7.75 U 6510	9.71 U 5480	8.4 U 19700 J	9.17 U 2630	42 U 3520	48.5 U 3660	47.6 U 4030	48.1 U 57900 J
PCB-122	10500	7.75 U	5590		2810	4000	3660	4280	70200 J
PCB-123				18100 J					
PCB-126	1780	1850	1030	3120	529	789	671	686	12200
PCB-127	8.2 U	83.8	9.71 U	447	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-128/166	31900 J	24100 J	25200 J	88100 J	9890	10900	12400	12800	118000 J
PCB-129/138/163	186000 J	127000 J	135000 J	434000 J	54100 J	64900 J	73200 J	70300 J	653000 J
PCB-130	16700 J	10900	11800	38800 J	4780	5710	6300	6070	65300 J
PCB-131	3500 67400 I	2850 46200 I	2260	8970 165000 I	930 18200	1020 20100 I	1250 23700 J	1240	16600 273000 J
PCB-132	67400 J	46200 J	43900 J 1650	165000 J 5080		20100 J 856		23500 J	
PCB-133	2590	1490			743		852	859 4790	8150
PCB-134	13800	12600	8460	33000 J	3540	4040	5080	4780	61200 J
PCB-135/151	50500 J	31900 J	30900 J	104000 J	14400	15500	16800	16800	182000 J
PCB-136	24800 J	15600 J	14600	57900 J	6390	8270	8570	8480	89400 J
PCB-137	9820	7.75 U	7160	23600 J	2720	3260	4190	3910	47400 J
PCB-139/140	4110	2620	2560	9170	1130	1260	1420	1450	18000
PCB-141	30100 J	19100 J	22000 J	70600 J	8890	10800	11800	11900	116000 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB056	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
Sample ID:	PB056-1SS010-091031-N	PB057-1SS010-091031-N	PB059.1-1SS010-091102-N	PB059.2-1SS010-091031-N	PB063.1-1SS010-091031-N	PB063.2-1SS010-091102-N	PB064-1SS010-091102-N	PB066-1SS010-091101-N	PB069-1SS010-091101-N
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	N
 X:	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
PCB-086/087/097/108/119/125	194000 J	77900 J	1310000 J	136000 J	762000 J	74100 J	6310	171000 J	1010000 J
PCB-088	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-089	9550	3670	65300 J	6710	41300 J	3450	271	8450	52500 J
PCB-090/101/113	240000 J	95900 J	1440000 J	160000 J	869000 J	87900 J	8210	201000 J	1080000 J
PCB-091	45300 J	18100 J	306000 J	31200 J	180000 J	16600	1550	38800 J	212000 J
PCB-092	43700 J	17200	259000 J	28800 J	159000 J	16300	1530	36300 J	196000 J
PCB-093/100	4660	1930	32400 J	3310	20300 J	1840	160	4150	21700 J
PCB-094	2330	937	15600	1640	10200	900	81.9	2120	12800 J
PCB-095	183000 J	68600 J	1200000 J	124000 J	728000 J	68200 J	6240	159000 J	875000 J
PCB-096	5940	2390	37600 J	4090	26300 J	2310	230	5580	33500 J
PCB-098	302	43.5 U	90.1 U	657	1960	46.3 U	36.8 U	37.3 U	60.6 U
PCB-099	110000 J	44900 J	695000 J	77700 J	417000 J	44400 J	4300	103000 J	530000 J
PCB-102	15500	6390	97200 J	10200	64500 J	5840	541	14300	92800 J
PCB-103	1880	763	11300	1250	7550	710	75.3	1580	8480
PCB-104	69.6	27.8	564 J	41.5	284	35.2 J	36.8 U	90.8 J	456 J
PCB-105	124000 J	50900 J	822000 J	94900 J	487000 J	49900 J	4290	112000 J	701000 J
PCB-106	44.2 U	43.5 U	2130	50 U	1420	93.3	36.8 U	250	1770
PCB-107/124	88.5 U	3900	63300 J	6920	37200 J	3640	322	8290	46200 J
PCB-109	16500	6810	103000 J	11700	61600 J	6070	551	13500	74800 J
PCB-110	216000 J	85200 J	1450000 J	150000 J	803000 J	78900 J	7330	183000 J	1130000 J
PCB-111	44.2 U	43.5 U	165 J	50 U	49.5 U	31.9 J	36.8 U	37.3 U	139 J
PCB-112	918	526	90.1 U	50 U	3670	46.3 U	36.8 U	37.3 U	60.6 U
PCB-114	8780	3450	70400 J	6360	38200 J	3250	263	7790	56000 J
PCB-115	8340	3330	90.1 U	5580	31200 J	46.3 U	36.8 U	37.3 U	60.6 U
PCB-117	9670	3640	56300 J	7150	42600 J	3420	285	7400	35300 J
PCB-118	211000 J	82300 J	1320000 J	146000 J	795000 J	86400 J	7690	193000 J	1020000 J
PCB-120	317	131	90.1 U	50 U	928	135	36.8 U	281	1080
PCB-121	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-122	6000	2550	40400 J	4560	24000 J	2260	170	4790	27100 J
PCB-123	6140	2520	45200 J	4600	25900 J	2620	237	5790	34800 J
PCB-126	1020	471	6230	793	4250	452	51.6	841	4910
PCB-127	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-128/166	18300 J	6550	101000 J	12900	53600 J	6960	667	14700	90900 J
PCB-129/138/163	103000 J	36800 J	544000 J	66800 J	289000 J	40100 J	3970	78900 J	410000 J
PCB-130	9050	3250	50200 J	6190	26200 J	3550	349	7230	31700 J
PCB-131	1840	685	12700	1390	6130	780	63.4	1680	8320
PCB-132	34800 J	12600	206000 J	25600 J	108000 J	14100	1210	30200 J	147000 J
PCB-133	1170	405	6120	798	3210	478	52.9	981	4900
PCB-134	7010	2510	40200 J	5160	22500 J	2700	242	5300	29600 J
PCB-135/151	23600 J	8340	132000 J	15800	68800 J	10300	960	20500 J	106000 J
PCB-136	13800	4670	79200 J	8670	40300 J	5010	446	10600	67700 J
PCB-137	5820	2200	35100 J	3910	17700	2370	211	4940	24400 J
PCB-139/140	2050	776	12800	1550	6540	843	75.9	1830	9130
PCB-141	17500	6130	92500 J	11400	49600 J	7010	617	14100	71200 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	2011 USC	2011 USC	2011 USC				
Location ID:	PB074	PB081	PB093	PB097	PB101	2011 03C PB068	PB069.1	2011 03C PB070A
		PB081-1SS010-091103-N				PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	
-								PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm				
Sample Type:	N 2200220 0492	N 2200764 2707	N 2200070 4204	N 2201102 1242	N 2201207 2650	N 2201200 001	N 2201205 850	N 2201007.255
x: Y:	3200828.0183 13830463.2171	3200761.3787 13830005.1879	3200970.4291 13828894.7843	3201103.1242 13828566.1540	3201287.2658 13828181.2285	3201209.861 13831121.51	3201205.859 13830973.3	3201087.255 13830903.51
PCB-086/087/097/108/119/125	99100 J	457000 J	12700	20100 J	4570			
PCB-088	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U	-		
PCB-089	3140	21600 J	1010	854	375	-		
PCB-090/101/113	136000 J	559000 J	17400	28000 J	6260			
PCB-091	19600 J	100000 J	4960	10300	1710			
PCB-092	25300 J	99900 J	3890	10900	1380			
PCB-093/100	1630	9620	611	1690	229			
PCB-094	915	5170	318	1110	123			
PCB-095	109000 J	463000 J	19800	21900 J	6650			
PCB-096	2410	12900	870	1240	302	-		-
PCB-098	36.8 U	2110	98.8	323	48.1 U			
PCB-099	52600 J	238000 J	8230	15500	3220			
PCB-102	6210	34000 J	1770	2570	642			
PCB-103	815	4060	251	928	92.2			
PCB-104	36.8 U	104	8.77 J	64	48.1 U			
PCB-105	54900 J	334000 J	4920	12200	2140			
PCB-106	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U	-		
PCB-107/124	4930	21900 J	389	959	144			
PCB-109	7440	32700 J	984	3020	399	-		
PCB-110	116000 J	503000 J	20200	27700 J	6720			
PCB-111	36.8 U	46.3 U	52.1 U	43.6	48.1 U			
PCB-112	36.8 U	2010	52.1 U	219	48.1 U			
PCB-114	3630	23800 J	310	721	136			
PCB-115	36.8 U	21800 J	52.1 U	1060	48.1 U			
PCB-117	5070	26200 J	52.1 U	3350	368			
PCB-118	109000 J	490000 J	11100	25600 J	4970			
PCB-120	175	403	29.3 J	65.3	48.1 U			
PCB-121	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-122	2220	13600	281	676	119			
PCB-123	2620	13900	255	771	125			
PCB-126	801	2200	47.5	119	48.1 U			
PCB-127	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-128/166	16900 J	44600 J	2160	1590	286	-		-
PCB-129/138/163	93800 J	264000 J	10800	8650	1510	-		-
PCB-130	8580	22000 J	983	799	148	-		-
PCB-131	1820	5240	151	152	32.6			-
PCB-132	35400 J	103000 J	3990	3120	717			
PCB-133	1160	3180	149	184	22.1	-		-
PCB-134	6130	21700 J	897	924	155			
PCB-135/151	21500 J	83800 J	2810	2680	493			
PCB-136	13000	44900 J	1400	1290	262			
PCB-137	5470	12600	463	461	80.5			
PCB-139/140	1880	5250	192	213	36.6			
PCB-141	15300 J	52500 J	1430	1270	241			
		·			i			

Task:		2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:	PB076.1-1SS010-20110802-N	PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	. N	FD	N	N
X:		3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
PCB-086/087/097/108/119/125				-
PCB-088				
PCB-089				
PCB-090/101/113				
PCB-091				
PCB-092				-
PCB-093/100				
PCB-094				
PCB-095				
PCB-096				-
PCB-098				
PCB-099				
PCB-102				<u></u>
PCB-103				
PCB-104				
PCB-105				
PCB-106				
PCB-107/124				
PCB-109				
PCB-110				
PCB-111				
PCB-112				
PCB-114				
PCB-115				
PCB-117				
PCB-118				
PCB-120				
PCB-121		-	-	-
PCB-122			-	-
PCB-123				
PCB-126				-
PCB-127			-	-
PCB-128/166				-
PCB-129/138/163			-	-
PCB-130				-
PCB-131				
PCB-132				-
PCB-133		-		-
PCB-134				
PCB-135/151				
PCB-136				
PCB-137				
PCB-139/140				
PCB-139/140	I			

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB001.1	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
	PB001.1-1SS010-091027-N								
Sample Date:	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N N	N 20 CIII	N N	N N	N N	N N	N N	N N
Y-	3202382.3842	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011	3202037.3755	3201747.2539
Y:	13836546.3402	13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614	13836300.0036	13836110.2965	13836119.3947
PCB-142	43600 J	8.93 U	8.93 U	9.8 U	47.4	44	9.17 U	47.2	56.3
PCB-143	2770	8.93 U	8.93 U	3060	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-144	6060	2170	1340	3450	2260	2530	3110	2360	2380
PCB-145	8.93 U	8.93 U	29.6	52.5	37.1	38.8	53.1	37	35.9
PCB-146	15800	8.93 U	5000	9.8 U	7820	7770	9630	7030	6980
PCB-147/149	88300 J	39200 J	21300 J	65900 J	38200 J	42500 J	47000 J	39000 J	38000 J
PCB-148	8.93 U	8.93 U	135	282	139	116	154	121	103
PCB-150	8.93 U	204	119	385	213	200	167	189	158
PCB-152	8.93 U	74.8	40.9	69.8	51.3	61.7	103	56.9	54.8
PCB-153/168	83100 J	35700 J	20400 J	61900 J	38400 J	41700 J	49700 J	37900 J	38400 J
PCB-154	2630	1670	709	2730	1610	1410	1600	1290	1140
PCB-155	8.93 U	182	142	333	174	186	198	186	128
PCB-156/157	21000 J	6390	4690	7250	6240	6880	9530	6490	7200
PCB-158	14400	5010	3520	6860	5160	5610	7170	5200	5490
PCB-159	443 J	229 J	173 J	510 J	256 J	285 J	321 J	236 J	248 J
PCB-160	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-161	8.93 U	7990	8.93 U	13600	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-162	631	299	8.93 U	259	279	183	247 J	212	189
PCB-164	7600	3820	2250	5560	5560	4170	4800	3480	3870
PCB-165	8.93 U	8.93 U	92	164	9.62 U	8.55 U	9.17 U	8.47 U	48.1
PCB-167	6110	2160	1660	2950	2240	2420	3280	2290	2460
PCB-169	8.93 U	8.93 U	8.93 U	9.8 U	9.62 U	8.55 U	9.17 U	8.47 U	7.58 U
PCB-170	13600 J	6570 J	3410 J	12900 J	7210 J	7520 J	8480 J	6720 J	7270 J
PCB-171/173	5740	2950	1840	5910	3170	3360	3910	2990	3030
PCB-172	3100	1740	1220	3560	1880	1850	2300	1720	1730
PCB-174	16800	7320	3420	15200	8830	8870	10600	7440	7870
PCB-175	1060	700	638	1400	629	690	745	651	634
PCB-176	2200	1140	668	2560	1190	1250	1450	1090	1070
PCB-177	9500	5100	2600	10400	5360	5640	6560	4750	4910
PCB-178	3720	2260	1380	4600	2230	2140	2650	1840	1840
PCB-179	6830	3490	1810	7350	3630	3700	4470	2990	3080
PCB-180/193	42800 J	21500 J	10600	45100 J	24800 J	25500 J	30700 J	21900 J	22400 J
PCB-181	270	195	218	289	207	193	207	182	183
PCB-182	8.93 U	130 J	113 J	225 J	95.8 J	117 J	142 J	123 J	106 J
PCB-183	10600 J	5630 J	2800 J	12200 J	6600 J	6590 J	7930 J	5610 J	5810 J
PCB-184	168	191	210	455	177	218	169	221	163
PCB-185	8.93 U	621	448	1230	9.62 U	716	882	1010	799
PCB-186	8.93 U	8.93 U	81.8	9.8 U	9.62 U	8.55 U	9.17 U	18.5	7.58 U
PCB-187	18500 J	10600 J	4940 J	21300 J	11200 J	11100 J	13400 J	9830 J	9470 J
PCB-188	116	139	201	186	86.9	83.4	97.2	82.6	71.6
PCB-189	1020	565	700	1090	579	576	651	563	533
PCB-190	3770 J	1750 J	1120 J	3530 J	1980 J	1960 J	2120 J	1750 J	1730 J
PCB-191	839	512	511	1010	535	561	635	496	480
PCB-192	8.93 U	56.7	147	107	9.62 U	8.55 U	9.17 U	59.9	43.4

Appendix A-1
BERA Sediment Dataset

Task	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	· · · · · · · · · · · · · · · · · · ·	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:		N N	0 - 10 CIII	N N	N N	0 - 10 CIII	N N	N N	FD
Sample Type.	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
A. V.	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
1000 443									
PCB-142	8.2 U	41.2	6.85 U	52		102	34.8	22.2	35.4
PCB-143	8.2 U	8.47 U	6.85 U	9.62 U		533	39.1 U	69.4	190
PCB-144	2950	1820	1280	3070		6570	1860	1630	2200
PCB-145	42.6	30.5	19.9	55.5		132	33.6	31.8	41.2
PCB-146	9560	5330	3640	9870		15000	6060	5680	6120
PCB-147/149	53300 J	28300 J	19300 J	49300 J	-	88500 J	31000 J	26800 J	31500 J
PCB-148	191	8.47 U	52	167	-	132	127	93	62.6
PCB-150	345	103	63.7	200		189	205	129	103
PCB-152	81.9	49.9	34	80.8		266	55.4	53.2	71.4
PCB-153/168	50900 J	28500 J	20000 J	51200 J		92600 J	36300 J	34500 J	40900 J
PCB-154	2110	870	532	1780		1540	1200	943	893
PCB-155	410	80.9	69.8	221	-	74.5	207	175	33.8
PCB-156/157	8330	5190	4410	8990		21000 J	4910	5430	8140
PCB-158	6380	4010	3180	6930		15800 J	4150	4170	6060
PCB-159	331 J	200 J	142 J	309 J		657	320	277	375
PCB-160	8.2 U	8.47 U	6.85 U	9.62 U	-	76.9 U	39.1 U	38.5 U	43.9 U
PCB-161	8.2 U	8.47 U	6.85 U	9.62 U	-	76.9 U	39.1 U	38.5 U	43.9 U
PCB-162	223	144	110	208 J	-	427	147	142	164
PCB-164	4590	2820	1980	4910		9430	3100	2990	3680
PCB-165	63.8	8.47 U	6.85 U	9.62 U	-	50.6	53.3 J	38.5 U	26.2
PCB-167	2930	1760	1460	3050		6150	1850	1880	2600
PCB-169	8.2 U	8.47 U	6.85 U	9.62 U	-	76.9 U	39.1 U	38.5 U	43.9 U
PCB-170	10000 J	5300 J	3810 J	9430 J		14700	7060	5960	9670
PCB-171/173	4210	2190	1550	4020	-	5790	2710	2220	3090
PCB-172	2630	1270	924	2440		3080	1700	1460	2290
PCB-174	11300	6040	4090	11000	-	15600 J	6740	5280	7730
PCB-175	885	439	315	816		1040	557	422	565
PCB-176	1790	809	512	1450		2190	1220	1050	1200
PCB-177	7120	3630	2440	6820		9250	4370	3550	4870
PCB-178	2850	1340	863	2610		3560	2050	1800	2490
PCB-179	5120	2400	1520	4470		6790	3400	2870	3720
PCB-180/193	34400 J	16600	12100	32400 J		43900 J	23500 J	18800 J	35200 J
PCB-181	271	145	88	224		291	162	122	143
PCB-182	150 J	86.4 J	6.85 U	144 J		119	118	77.5	77.8
PCB-183	9070 J	4270 J	3050 J	8350 J		10000	4800	3810	5930
PCB-184	341	73.9	65.7	175		65.7	237	178	49.5
PCB-185	731	479	244 J	686		1110	835	636	850
PCB-186	8.2 U	8.47 U	6.85 U	9.62 U		18	39.1 U	38.5 U	43.9 U
PCB-187	15200 J	7480 J	4980 J	14200 J		16900 J	8660	7070	11500
PCB-188	163	44	36	98.7		50.5	78.9	88.2	57.1
PCB-189	790	367	296	730		807	534	438	695
PCB-190	2630 J	1350 J	938 J	2470 J		2860	2100	1640	2700
PCB-191	698	341	254	611		781	39.1 U	404	581
PCB-192	55.5	23.3 J	18.3 J	9.62 U	-	76.9 U	79.1	38.5 U	43.9 U

Appendix A-1
BERA Sediment Dataset

Tools	: RI/FS 2009	RI/FS 2009	RI/FS 2009	DI/FC 2000	RI/FS 2009	RI/FS 2009	DI /EC 2000	RI/FS 2009	RI/FS 2009
Task		PB023	PB023	RI/FS 2009 PB024	PB026	PB028	RI/FS 2009 PB032	PB034	PB037
Location ID	_								
-	: PB022-1SS010-091101-N								
Sample Date	1 ' '	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
Depth		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type		FD	N N	N N	N	N N	N	N	N
X		3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
Y		13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
PCB-142	28.8	231 U	33.2	40.1	420	9.17 U	3090	97.9	8.13 U
PCB-143	118	231 U	183 U	189	188	9.17 U	467	275	8.13 U
PCB-144	1590	3090	2390	2600	2760	775	1460	4490	7110
PCB-145	27.9	79.5	183 U	35.5	308	9.17 U	45 U	65.2	124
PCB-146	4480	9910	7950	7160	59700 J	2710	3390	13800	17600 J
PCB-147/149	23000 J	48100 J	38300 J	38600 J	118000 J	12100	18200 J	65800 J	97500 J
PCB-148	58.6	226	133	79.8	9220	54.7	95.1	249	154
PCB-150	70	268	180	86.3	2460	50.2	112	198	194
PCB-152	52.7	113	48.2	72.2	200	22.4	22.9 J	118	257
PCB-153/168	26800 J	48700 J	39400 J	41400 J	120000 J	14100	19400 J	73400 J	103000 J
PCB-154	629	1590	1280	867	24600 J	454	1750	2080	2090
PCB-155	37.5	354	264	41.7	11200	83.2	105	207	80.9
PCB-156/157	5100	6450	4820	7550	7680	2080	3650	12900	22400 J
PCB-158	3870	6020	4560	5980	5670	1760	3070	10600	14100
PCB-159	242	231 U	183 U	287	1050	81.7	246	468	461
PCB-160	37.6 U	231 U	183 U	6.94 U	1160	9.17 U	1500	7.63 U	8.13 U
PCB-161	37.6 U	231 U	183 U	6.94 U	469	9.17 U	238	7.63 U	8.13 U
PCB-162	120	173	183 U	178	1300	53.5	170	402	639
PCB-164	2470	5160	3730	4270	12300	1260	2170	7520	8.13 U
PCB-165	37.6 U	231 U	183 U	6.94 U	4070	9.17 U	155	126	8.13 U
PCB-167	1640	2480	1860	2360	2790	713	1450	4650	6250
PCB-169	37.6 U	231 U	183 U	6.94 U	40 U	9.17 U	45 U	7.63 U	8.13 U
PCB-170	5040	8520	8600	7990	11100	2490	3580 J	14500	14100
PCB-171/173	1930	4460	3740	3630	13500	1060	2790	5710	6390
PCB-172	1200	2470	1940	1980	8650	646	1400	3750	3550
PCB-174	5340	11800	9750	9780	29900 J	2950	5320	16000 J	18500 J
PCB-175	375	970	723	658	5180	225	844	1350	1050
PCB-176	763	1880	1540	1220	10700	375	813	2200	2140
PCB-177	3250	7850	6270	5870	28900 J	1910	3590	9830	10300
PCB-178	1370	3100	2600	2030	40000 J	756	1370	4170	3140
PCB-179	2260	5180	4590	3690	19600 J	1270	2140	6610	6670
PCB-180/193	17000 J	34400 J	28500 J	27300 J	69000 J	8940	17200	47100 J	47100 J
PCB-181	108	512	187	181	1180	69.7	1330	307 J	8.13 U
PCB-182	55.8	231 U	143	110	11300	9.17 U	45 U	223	8.13 U
PCB-183	3740	8960	6900	6530	18200 J	2120	4360	10800	12000
PCB-184	47.1	337	172	60.4 J	9310	52.2	245	187	8.13 U
PCB-185	559	1070	925	1300	2560	314	3330	1690	8.13 U
PCB-186	7.01 J	231 U	183 U	10.2	253	9.17 U	83	7.63 U	8.13 U
PCB-187	6480	15100	12400 J	11100	75200 J	3970	6480	20500 J	22300 J
PCB-188	29.1	218	93	36.7	8150	32.8	799 J	200	8.13 U
PCB-189	328	863	560	528	3850	168	919	1180	950
PCB-190	1400	2550	2390	2280	5580	743	2360	3970	3580
PCB-191	321	777	593	556	3060	147	896	1010	913
PCB-192	17.3 J	186	183 U	22.9	937	9.17 U	472	7.63 U	8.13 U

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009						
Location ID:	PB041	PB043	PB044	PB047.1	PB047.2	PB048	PB049	PB049	PB053
•			10/30/2009			10/31/2009	PB049-1SS010-091031-D	10/31/2009	
Sample Date:	10/30/2009	10/30/2009		10/30/2009	10/30/2009		10/31/2009		10/31/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm						
Sample Type:	N	N	N	N	N	N	FD	N	N
X:	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
Y	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
PCB-142	204	7.75 U	85.3	257	40.2	38.8	60.2	68.4	741
PCB-143	670	7.75 U	395	1640	198	246	130	310	4070
PCB-144	8190	5700	5230	19200 J	2380	2520	2930	2860	34100 J
PCB-145	165	109	89.9	339	34.1 J	67.2	52.2	61.5	892
PCB-146	20900 J	12600	13800	40900 J	6100	7410	7640	7600	67600 J
PCB-147/149	118000 J	72900 J	72900 J	244000 J	33400 J	36300 J	39400 J	38700 J	412000 J
PCB-148	188	76.8	91	234	55	62	48.2	51.8	538
PCB-150	395	136	155	407	95.1	128	80.7	75.8	830
PCB-152	307	212	174	672	79.1	109	104	118	1710
PCB-153/168	121000 J	76500 J	82900 J	239000 J	36100 J	42600 J	44300 J	42900 J	372000 J
PCB-154	3140	1590	1330	3590	806	900	667	673	6910
PCB-155	37.6	7.75 U	12.4 J	8.4 U	17.5	24.8 J	48.5 U	47.6 U	48.1 U
PCB-156/157	26000 J	19700 J	21300 J	69200 J	7510	8970	10900	10300	89800 J
PCB-158	19800 J	14100	14300	48900 J	5690	6730	8060	7620	74200 J
PCB-159	538	320	471	798	203	42 U	48.5 U	47.6 U	48.1 U
PCB-160	8.2 U	7.75 U	9.71 U	8.4 U	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-161	8.2 U	7.75 U	9.71 U	8.4 U	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-162	636	564	473	1440	197	212	235	260	2290
PCB-164	12700	14800	8840	29200 J	3580	4120	4400	4350	38900 J
PCB-165	8.2 U	7.75 U	9.71 U	8.4 U	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-167	7910	5640	6230	19800 J	2350	2890	3190	3200	27300 J
PCB-169	8.2 U	7.75 U	9.71 U	8.4 U	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-170	18900 J	11700	18800	35700 J	6440	7840	7100	7340	37600 J
PCB-171/173	7490	4650	7100	15000	2510	3190	2830	2930	16800
PCB-172	4310	2560	4020	7430	1530	2030	1620	1700	9730
PCB-174	19700 J	12800	17600	33300 J	6920	8330	6980	7930	46000 J
PCB-175	1270	774	1080	2340	475	575	485	522	3090
PCB-176	2560	1550	2050	4340	822	1240	1080	1150	6900
PCB-177	12100	7470	10900	20700 J	4330	5220	4320	4750	24800 J
PCB-178	4490	2390	3420	6180	1440	2280	1830	1990	11400
PCB-179	8030	4540	5950	11600	2580	3810	3140	3440	23000 J
PCB-180/193	58900 J	33100 J	51300 J	91100 J	20600 J	27400 J	21500 J	23400 J	120000 J
PCB-181	632	257	391	1050	135	167	174	171	1190
PCB-182	181	7.75 U	135	297	60.7	64.2	68	58.9	493
PCB-183	13200	8480	11800	21800 J	4900	6110	5150	5690	35100 J
PCB-184	45.7	7.75 U	27.9 J	48.3	20.8	32.7	48.5 U	15.2 J	48.1 U
PCB-185	2050	7.75 U	1050	3210	701	673	866	781	3920
PCB-186	8.2 U	7.75 U	9.71 U	31.2	9.17 U	42 U	48.5 U	47.6 U	48.1 U
PCB-187	24500 J	13600	19700 J	33700 J	8830	10700	8440	9180	60800 J
PCB-188	64.2	7.75 U	32.1 J	46.4	21.6	39.8	18.3	14.5 J	116
PCB-189	1310	699	1130	2290	396	548	510	497	2860
PCB-190	4910	2780	4250	8180	1540	2080	1840	1870	8300
PCB-191	1090	628	956	1890	370	487	413	431	2140
PCB-192	8.2 U	7.75 U	9.71 U	8.4 U	9.17 U	42 U	48.5 U	47.6 U	48.1 U

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	-	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
			PB059.1-1SS010-091102-N		PB063.1-1SS010-091031-N		PB064-1SS010-091102-N		PB069-1SS010-091101-N
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	N
x: 'X	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
PCB-142	83.8	43.5 U	90.1 U	51.5	243	46.3 U	36.8 U	55 J	289
PCB-143	397	179	2650	371	1570	46.3 U	36.8 U	404	2030
PCB-144	4090	1460	25300 J	2930	12700	1800	151	3730	19900 J
PCB-145	91.5	35.3	658	74	351	46.3 U	36.8 U	69.3	519
PCB-146	10400	3750	52300 J	6580	27500 J	4060	423	7900	41100 J
PCB-147/149	55900 J	19600 J	311000 J	37800 J	161000 J	23300 J	2170	47400 J	245000 J
PCB-148	58.3	23.1	345	41	157	27.6	36.8 U	57.9	234
PCB-150	118	41.3	649	78.5	313	45.6	36.8 U	88	486
PCB-152	186	70.8	1250	130	641	72	36.8 U	155	973
PCB-153/168	61700 J	21600 J	335000 J	39600 J	171000 J	26600 J	2730	50700 J	275000 J
PCB-154	904	344	4740	628	2400	386	45.9	776	2930
PCB-155	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-156/157	15000	5190	92500 J	9960	45900 J	6170	561	12700	74400 J
PCB-158	11300	4030	62300 J	7470	32700 J	4340	415	9180	47600 J
PCB-159	44.2 U	43.5 U	2050	50 U	49.5 U	191	17.1	343	60.6 U
PCB-160	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-161	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-162	364	123	1770	221	936	140	15.1	264	1160
PCB-164	6310	2220	34300 J	4350	17600	2510	256	5070	24800 J
PCB-165	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	35.5	60.6 U
PCB-167	4450	1570	25000 J	2980	12100	1830	183	3650	20300 J
PCB-169	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-170	11300	3040	45600 J	5790	25700 J	4090	468	7330	47300 J
PCB-171/173	4190	1240	18300 J	2160	9840	1520	167	2740	17200 J
PCB-172	2310	741	9290	1190	5300	894	102	1520	8720
PCB-174	10700	3250	48000 J	5730	26300 J	4430	432	7550	52600 J
PCB-175	719	219	3050	375	1660	274	26.2	467	2660
PCB-176	1500	537	6310	863	3890	608	57.2	1050	7010
PCB-177	6490	2040	27300 J	3430	15800	2520	264	4230	29000 J
PCB-178	2540	898	8900	1390	6190	1010	119	1680	10200
PCB-179	4670	1560	18500 J	2640	12100	1930	191	3130	22800 J
PCB-180/193	31400 J	9680	134000 J	16500	73700 J	13300	1480	21900 J	149000 J
PCB-181	198	67.2	1110	116	520	70.6	36.8 U	138	836
PCB-182	82.7	35	342	51.9	191	25.7	36.8 U	47.4	259
PCB-183	7930	2400	29900 J	4010	18000	3120	322	5020	29400 J
PCB-184	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	44
PCB-185	850	246	4340	474	2400	327	57.1	691	5770
PCB-186	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	8.67	27.7
PCB-187	12900	4020	49000 J	6950	31000 J	5200	576	37.3 U	57700 J
PCB-188	44.2 U	43.5 U	90.1 U	50 U	35.3	46.3 U	36.8 U	37.3 U	43.9
PCB-189	680	221	2790	389	1700	247	29.8	441	2710
PCB-190	2910	772	10500	1440	6500	999	118	1850	11000
PCB-191	577	171	2230	299	1290	214	22.7 J	367	2230
PCB-192	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U

Appendix A-1
BERA Sediment Dataset

	Task: ocation ID:	RI/FS 2009	RI/FS 2009						2011 USC
		PB074	PB081	RI/FS 2009 PB093	RI/FS 2009 PB097	RI/FS 2009 PB101	2011 USC PB068	2011 USC PB069.1	PB070A
	Sample ID:		PB081-1SS010-091103-N				PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Su.	nple Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
	Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm
Sar	nple Type:	N	N N	N N	N N	N N	N	N	N
54.	χ.	3200828.0183	3200761.3787	3200970.4291	3201103.1242	3201287.2658	3201209.861	3201205.859	3201087.255
	Y:	13830463.2171	13830005.1879	13828894.7843	13828566.1540	13828181.2285	13831121.51	13830973.3	13830903.51
PCB-142		36.8 U	158	52.1 U	45.9 U	48.1 U		-	
PCB-143		189	922	52.1 U	35.7	7.59 J	-		
PCB-144		3910	14300	367	355	63			
PCB-145		65.4	205	7.55 J	45.9 U	48.1 U			
PCB-146		9230	27600 J	1200	962	188			
PCB-147/149		52800 J	182000 J	6430	5820	1090			
PCB-148		56.1	150	10.4	28	48.1 U			
PCB-150		101	265	8.44 J	57.6	48.1 U			
PCB-152		141	440	13.5	63.1	48.1 U			
PCB-153/168		57200 J	183000 J	5530	5330	1010	-	-	
PCB-154		833	2250	88.6	217	30.3	-	ı	-
PCB-155		36.8 U	46.3 U	52.1 U	6.05 J	48.1 U		-	-
PCB-156/157		13200	37300 J	678	1170	189	-	-	
PCB-158		11000	29100 J	1070	853	151	-	ı	-
PCB-159		203 J	46.3 U	52.1 U	36.2	48.1 U			
PCB-160		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-161		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U	-		
PCB-162		290	697	35.6	26.7	48.1 U			
PCB-164		6130	17900	805	612	107			
PCB-165		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U	-		
PCB-167		3600	10800	301	368	61.1			
PCB-169		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-170		6610	34600 J	1070	738	126	-		
PCB-171/173		2590	14600	466	287	50.9			
PCB-172		1250	8150	251	179	29.4	-		
PCB-174		5370	44100 J	1150	921	161			
PCB-175		382	2550	78.4	60.8	12.9		-	-
PCB-176		781	5780	152	122	25.6			
PCB-177		3120	25500 J	719	525	98.8			
PCB-178		1030	9010	247	280	51.2			
PCB-179		1890	19000 J	469	513	92.6			
PCB-180/193		15200 J	117000 J	3130	2550	458			
PCB-181		173	522	19.8	17.6	48.1 U			
PCB-182		66.5 J	196	11.5	11.4	48.1 U			
PCB-183		3850	28700 J	787	613	115			
PCB-184		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-185		234	4890	69.5 J	117	16.2 J			
PCB-186		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-187		4610	49300 J	1360	1450	270			
PCB-188		36.8 U	46.3 U	52.1 U	21.6	48.1 U			
PCB-189		370	2120	64.5	47.4	48.1 U			
PCB-190		1470	8510	233	182	29.9			
PCB-191		333	1920	68.1	37.6	48.1 U			
PCB-192		36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			

Appendix A-1
BERA Sediment Dataset

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:		PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
PCB-142				
PCB-143				
PCB-144				
PCB-145				
PCB-146	-		1	
PCB-147/149				
PCB-148			1	
PCB-150	-			
PCB-152			-	-
PCB-153/168				
PCB-154				
PCB-155				
PCB-156/157				
PCB-158				
PCB-159				
PCB-160				
PCB-161				
PCB-162				
PCB-164				
PCB-165				
PCB-167				
PCB-169				
PCB-170				
PCB-171/173				
PCB-172				
PCB-174				
PCB-175				
PCB-176				
PCB-177				
PCB-178				
PCB-179				
PCB-180/193				
PCB-181				
PCB-182				
PCB-183				
PCB-184				
PCB-164 PCB-185				
PCB-186				
PCB-180	 			
PCB-187				
PCB-188 PCB-189	 			
PCB-190				
PCB-191				
PCB-192				

Appendix A-1
BERA Sediment Dataset

Task	DI /EC 2000	DI/EC 2000	DI /EC 2000	DI /EC 2000	DI/FC 2000	DI/FC 2000	DI /EC 2000	DI /EC 2000	DI/EC 2000
Task: Location ID:	RI/FS 2009 PB001.1	RI/FS 2009 PB001.2	RI/FS 2009 PB001.3	RI/FS 2009 PB002	RI/FS 2009 PB004	RI/FS 2009 PB005	RI/FS 2009 PB007.1	RI/FS 2009 PB007.2	RI/FS 2009 PB009
							PB007.1-1SS010-091028-N		
Sample Date:	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	0 - 10 CIII	0 - 10 CIII	0 - 10 CIII	0 - 10 Cili	0 - 10 CIII	N N	0 - 10 CIII		0 - 10 Cili
Sample Type:	3202382.3842	N 2202624 0211	N 22027F4 0026	3202000 0020	N 2202220 4700		2202065 5011	N 2202027 27FF	3201747.2539
A:	13836546.3402	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011 13836300.0036	3202037.3755	
1:		13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614		13836110.2965	13836119.3947
PCB-194	6420 J	4070 J	2240 J	8920 J	4470 J	4350 J	5610 J	4160 J	3820 J
PCB-195	2760	1700	999	3640	1910	1790	2260	1770	1630
PCB-196	4670	2480	1840	5120	2600	2600	3030	2380	2180
PCB-197	901	812	855	1940	814	975	675	923	705
PCB-198/199	8860	5070	3230	10900	5290	4960	6610	4640	4290
PCB-200	1290	772	718	1430	841	703	1080	734	710
PCB-201	1850	1400	1990	2460	1150	1180	1370	1120	1030
PCB-202	3520	4950	15900	7390	2630	2750	3000	2500	2450
PCB-203	4470	2440	1680	4910	2600	2570	3120	2380	2130
PCB-204	170	147	262	273	115	135	119	145	117
PCB-205	1210	875	2320	1830	740	914	734	742	721
PCB-206	18700 J	11700 J	24600 J	30400 J	10700 J	12600 J	11700 J	11600 J	16800 J
PCB-207	5940	5280	10500	9770	4680	5750	5600	5200	5110
PCB-208	8340	8760	20300 J	14800	6330	7180	8530	6660	6460
PCB-209	1450000 J	884000 J	1410000 J	1710000 J	1080000 J	1210000 J	1130000 J	994000 J	2110000 J
Total PCB Congener (U = 1/2)	10254611.615 J	3549333.095 J	2528236.635 J	5263006.6 J	3372482.64 J	3829972.525 J	4684077.285 J	3587434.36 J	4106808.25 J
Total PCB Congener minus 209 (U = 1/2)	8804611.615 J	2665333.095 J	1118236.635 J	3553006.6 J	2292482.64 J	2619972.525 J	3554077.285 J	2593434.36 J	1996808.25 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	3333.950165 J	836.836415 J	40.599855 J	1007.3988 J	842.2608 J	922.147635 J	1373.970595 J	919.635065 J	692.07152 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	19.03205575 J	4.19749825 J	1.58810475 J	4.549545 J	4.7360355 J	4.56504375 J	7.06428425 J	4.41502675 J	3.7627045 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	226.0628895 J	43.375 J	26.5926065 J	43.884 J	54.89847 J	47.19643 J	72.13638 J	44.09124 J	41.04409 J
Monochlorobiphenyls (total)	17900	7910	3830	15400	3360	6630	3080	5430	5630
Dichlorobiphenyls (total)	196000	63600	26700	101000	36000	53900	38900	57300	38900
Trichlorobiphenyls (total)	1850000	562000	204000	827000	414000	514000	544000 J	537000	346000
Tetrachlorobiphenyls (total)	4060000	1090000	361000	1320000	884000	1040000	1550000	1050000	746000 J
Pentachlorobiphenyls (total)	1920000 J	593000 J	260000	692000	607000	632000	958000	601000	510000
Hexachlorobiphenyls (total)	556000	225000	137000	341000	222000	244000	303000	223000	229000
Heptachlorobiphenyls (total)	141000	73100	39100	151000	80400 J	82600	98200	72000	73200
Octachlorobiphenyls (total)	36100	24700	32000	48800	23200	22900	27600	21500	19800
Nonachlorobiphenyls (total)	33000	25800	55400	55100	21700	25500	25900	23500	28300
PCB Aroclors (μg/kg)									
Aroclor 1260									
Aroclor 1254									
Aroclor 1268									
Aroclor 1221									
Aroclor 1232									
Aroclor 1248									
Aroclor 1016									
Aroclor 1262									
Aroclor 1242									
Total PCB Aroclors (U = 1/2)									
Pesticides (μg/kg)									
2,4'-DDD (o,p'-DDD)	23.7 J	11.4 J	2.06 J	12.8 J	2.07 U	12.3 J	3.11 U	1.42 U	1.27 U
2,4'-DDE (o,p'-DDE)	2.07 U	1.97 UJ	1.28 U	1.44 U	2.07 U	1.65 U	3.11 U	1.42 U	1.27 U
2,4'-DDT (o,p'-DDT)	2.07 U	1.97 UJ	1.28 U	1.44 U	2.07 U	1.65 U	3.11 U	1.42 U	1.27 U

Appendix A-1
BERA Sediment Dataset

Task:	-	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:		PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
•		PB013.1-1SS010-091028-N							
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	FD
X:	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
Y:	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
PCB-194	7470 J	2830 J	2070 J	6320 J		5360	3670	3120	9750
PCB-195	3040	1180	864	2490	-	2220	1540	1260	3430
PCB-196	3730	1730	1220	3490		3400	2520	1890	5130
PCB-197	2670	392	280	855		350	979	758	472
PCB-198/199	7460	3480	2260	7790		7260	4960	4130	9910
PCB-200	1210	553	384	1090		1320	726	575	1260
PCB-201	1700	684	486	1420		1240	1040	870	1420
PCB-202	6620	1150	944	3000		2040	1560	1850	1920
PCB-203	3740	1670	1170	3420		3300	2610	2020	5460
PCB-204	161	76.5	52.7	132 J		59.8	183	140	45
PCB-205	970	432	344	871		508	1130	742	766
PCB-206	15900 J	6240 J	4940 J	21600 J		5330	14900	10800	5890
PCB-207	8910	2530	1900	6020		1980	4650	4480	1660
PCB-208	12200	3330	2430	10100		2850	6030	5830	2260
PCB-209	1770000 J	563000 J	680000 J	1160000 J		359000 J	1340000 J	883000 J	318000 J
Total PCB Congener (U = 1/2)	4748056.5 J	2365084.895 J	1716917.05 J	4414730.6 J		11089736.2 J	3080039.85 J	2505587.5 J	2306206.55 J
Total PCB Congener minus 209 (U = 1/2)	2978056.5 J	1802084.895 J	1036917.05 J	3254730.6 J		10730736.2 J	1740039.85 J	1622587.5 J	1988206.55 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	892.7137 J	617.014305 J	381.095485 J	1290.37401 J		3345.22752 J	485.05069 J	562.21393 J	618.6493 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	4.438355 J	3.26229675 J	2.47320125 J	6.7773405 J		15.3949575 J	2.6089475 J	2.9705525 J	3.6264725 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	45.8521 J	34.81746 J	29.80573 J	71.9624 J		136.67571 J	28.50762 J	32.39654 J	40.20625 J
Monochlorobiphenyls (total)	7380	3140	1810	3400		5210	6730 J	5830	2650
Dichlorobiphenyls (total)	57800	30500	15100	37400		136000	45100	36400	23300
Trichlorobiphenyls (total)	567000	332000	155000	504000		2370000	361000	316000	302000
Tetrachlorobiphenyls (total)	1170000	736000	378000	1390000		5070000	636000	615000	775000 J
Pentachlorobiphenyls (total)	692000 J	446000 J	304000	845000		2360000	382000	374000	515000 J
Hexachlorobiphenyls (total)	297000	173000	126000	299000		609000	191000	178000	229000
Heptachlorobiphenyls (total)	110000	54700	38100 J	103000		139000	70900	57800	93400
Octachlorobiphenyls (total)	38800	14200	10100	30900 J		27100	20900	17300	39600
Nonachlorobiphenyls (total)	37000	12100	9260	37700		10200	25600	21100	9820
PCB Aroclors (µg/kg)	37000	12100	3200	37700		10200	25000	21100	3020
Aroclor 1260									
Aroclor 1254									
Aroclor 1268									
Aroclor 1208 Aroclor 1221			 						
Aroclor 1221 Aroclor 1232			 			 			
Aroclor 1248			 						
Aroclor 1248 Aroclor 1016									
Aroclor 1016 Aroclor 1262									
Aroclor 1242						<u></u>			
Total PCB Aroclors (U = 1/2)									
Pesticides (µg/kg)		<u></u>	<u></u>						
	10 7 1	1.24 U	4.06 J	9.28 J	1.08 U		0.991 UJ	1 15 11	1 01 11
2,4'-DDD (o,p'-DDD) 2,4'-DDE (o,p'-DDE)	18.7 J 1.41 U	1.24 U	1.07 U	2.74 UJ	1.08 U		0.991 UJ	1.15 U 1.15 U	1.01 U 1.01 U
2,4'-DDT (o,p'-DDT)	1.41 U	1.24 U	1.07 U	2.74 UJ	1.08 U		0.991 UJ	1.15 U	1.01 U

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB022	PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
•	PB022-1SS010-091101-N			PB024-1SS010-091030-N					
Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N	N	N	N	N	N
X:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
Y:	13835456.7389	13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
PCB-194	3000	5790	5400	4960	23900 J	1600	4550	9390	7980
PCB-195	1120	2800	2070	1770	8210	570	2700	3530	3120
PCB-196	1620	4010	2880	2600	33700 J	873	6300	4880	3740
PCB-197	235	1230	858	348	56400 J	210	2970	937	8.13 U
PCB-198/199	3330	8060	5810	8070	204000 J	2100	9220	11300	8470
PCB-200	550	1230	1000	940	16600 J	314	2120	1650	1950
PCB-201	622	2140	1220	939	76200 J	391	3570	2190	1530
PCB-202	986	2620	1980	1400	154000 J	780	2350	3700	2030
PCB-203	1680	4730	2890	2840	23900 J	975	11100	5480	3820
PCB-204	35.3	403	191	70.9	14700	38.9	1470	231	54.6
PCB-205	324	2370	1080	495	22400 J	204	4900	1330	563
PCB-206	4410	50700 J	20700 J	38800 J	737000 J	4120	289000 J	22800 J	6270
PCB-207	1750	9660	7460	3040	724000 J	1610	38700 J	9130	1990
PCB-208	2300	9770	7580	7330	1300000 J	2440	32600 J	14300	2910
PCB-209	346000 J	6010000 J	3700000 J	649000 J	117000000 J	423000 J	57600000 J	1770000 J	315000 J
Total PCB Congener (U = 1/2)	2183986.11 J	8604078.4 J	5288205.6 J	3161184.91 J	124178512 J	1216484.14 J	58574618.05 J	5868663.58 J	9941384.695 J
Total PCB Congener minus 209 (U = 1/2)	1837986.11 J	2594078.4 J	1588205.6 J	2512184.91 J	7178512 J	793484.14 J	974618.05 J	4098663.58 J	9626384.695 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	648.09058 J	795.63383 J	384.74529 J	878.87535 J	461.6584 J	264.126785 J	121.99176	1268.673315 J	3439.91610565 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	3.36038 J	4.88224 J	2.36466 J	4.3930635 J	1.6907 J	1.53248425 J	1.405455	6.97869075 J	21.776223575 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	34.85844 J	61.46939 J	30.67831 J	44.43764 J	7.1443 J	17.44718 J	19.94768	73.91555 J	256.17907195 J
Monochlorobiphenyls (total)	2450 J	8060 J	4130 J	8050	240000 J	1710	54100	19000	5980
Dichlorobiphenyls (total)	23900	35800 J	24900	54800	305000 J	11700	56000	39200	116000
Trichlorobiphenyls (total)	321000	320000 J	255000	488000	598000 J	138000	33300	601000	1860000
Tetrachlorobiphenyls (total)	822000	1040000	554000	969000	829000 J	321000	82500	1650000	4280000
Pentachlorobiphenyls (total)	441000	691000	372000	583000	755000 J	199000	140000 J	1110000 J	2530000
Hexachlorobiphenyls (total)	153000	286000	222000	248000	680000	77700	137000	438000	644000
Heptachlorobiphenyls (total)	51600	111000	92600	86800	376000	28200	60300 J	151000	153000
Octachlorobiphenyls (total)	13500	35400	25400	24400	635000	8050	51200	44600	33300
Nonachlorobiphenyls (total)	8470	70100	35700	49200	2760000	8180	360000	46200	11200
PCB Aroclors (μg/kg)									
Aroclor 1260									
Aroclor 1254									
Aroclor 1268									
Aroclor 1221									
Aroclor 1232									
Aroclor 1248									
Aroclor 1016									
Aroclor 1262									
Aroclor 1242									
Total PCB Aroclors (U = 1/2)									
Pesticides (μg/kg)								·	
2,4'-DDD (o,p'-DDD)	1 UJ	9.22 J	6 J	1.07 U	42.7 J	2.1 U	10.5 J	8.81 J	2.04 U
2,4'-DDE (o,p'-DDE)	1 UJ	3.51	3.46	1.07 U	15.2 J	2.1 U	1.45 U	2.31 UJ	2.04 U
2,4'-DDT (o,p'-DDT)	1 UJ	4.9 J	1.06 U	1.07 U	2.4 U	2.1 UJ	1.45 UJ	2.31 UJ	2.04 UJ

Appendix A-1
BERA Sediment Dataset

Task:	DI /EC 2000	DI/EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI/EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000
	1	RI/FS 2009 PB043	RI/FS 2009 PB044	RI/FS 2009 PB047.1	RI/FS 2009 PB047.2	RI/FS 2009 PB048	RI/FS 2009 PB049	RI/FS 2009 PB049	RI/FS 2009 PB053
Location ID:									
•							PB049-1SS010-091031-D		
Sample Date:	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:		N	N N	N	N N	N	FD	N	N
X	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
Y	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
PCB-194	10700	4750	7080	10300	3370	5250	3810	3940	16800
PCB-195	3490	1950	2830	4260	1350	1910	1390	1460	6540
PCB-196	5070	2600	3670	5360	1620	2740	1910	2130	12700
PCB-197	476	169	368	378	216	288	178	184	1130
PCB-198/199	14000	5410	7650	11500	3710	5980	4860	5230	30100 J
PCB-200	1800	996	1170	1930	546	987	756	826	5430
PCB-201	1790	956	1260	1860	638	929	675	789	6290
PCB-202	2610	1140	1630	2330	828	1330	948	1010	9100
PCB-203	5920	2770	3830	5750	1760	2880	2110	2340	13800
PCB-204	29.3 J	7.75 U	9.71 U	17	20	22 J	48.5 U	47.6 U	51
PCB-205	755	295	478	623	243	364	230	246	965
PCB-206	11100	2090	4270	5510	3200	3680	3320	3390	10500
PCB-207	1360	470	1140	801	796	1150	445	487	2610
PCB-208	2910	745	1560	1620	1180	1710	973	1000	5270
PCB-209	103000 J	24100 J	177000 J	38200 J	93100 J	140000 J	33100 J	36800 J	48200 J
Total PCB Congener (U = 1/2)	10321975.8 J	7705686.85 J	7481713.97000001 J	19864911.9 J	2819075.49 J	3980062.4 J	3966069.4 J	4575212.1 J	95025849.8499999 J
Total PCB Congener minus 209 (U = 1/2)	10218975.8 J	7681586.85 J	7304713.97 J	19826711.9 J	2725975.49 J	3840062.4 J	3932969.4 J	4538412.1 J	94977649.85 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	3683.1513 J	2724.76730375 J	2259.903355 J	5787.3261 J	1063.360145 J	1638.93538 J	1366.30085 J	1547.23957 J	31329.33765 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	19.730805 J	17.151908125 J	11.77494275 J	33.89166 J	5.70765925 J	8.46224 J	7.3956625 J	8.004875 J	146.5235025 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	203.5156 J	203.52853625 J	118.60385 J	359.7967 J	60.14383 J	89.34214 J	77.4062 J	80.01211 J	1385.0553 J
Monochlorobiphenyls (total)	6200	2170	9890	7090	904	1830 J	1390	1440	571000
Dichlorobiphenyls (total)	113000	75500	134000	146000	20800	36500	40400	44500	4370000
Trichlorobiphenyls (total)	1760000	1340000	1550000	3140000	419000	663000	693000	796000	21100000
Tetrachlorobiphenyls (total)	4440000	3600000	3270000	8690000	1220000	1780000	1820000	2180000	48500000
Pentachlorobiphenyls (total)	2870000	2010000	1620000	5730000	755000	980000	1000000	1140000	17200000
Hexachlorobiphenyls (total)	782000	519000	534000	1760000	224000	260000	288000	281000	2780000
Heptachlorobiphenyls (total)	186000	108000	162000	300000	64600	82800	68400	73900	414000
Octachlorobiphenyls (total)	46700 J	21000	30000	44300	14300	22700	16900	18200	103000
Nonachlorobiphenyls (total)	15300	3300	6960	7930	5180	6530	4740	4880	18400
PCB Aroclors (μg/kg)									
Aroclor 1260									
Aroclor 1254									
Aroclor 1268									
Aroclor 1221									
Aroclor 1232									
Aroclor 1248									
Aroclor 1016									
Aroclor 1262									
Aroclor 1242						-			
Total PCB Aroclors (U = 1/2)									
Pesticides (μg/kg)									
2,4'-DDD (o,p'-DDD)	1.55 U	1.1 UJ	1.71 U	1.16 U	1.57 U	2.39 U	1.34 U	1.2 U	1.47 U
2,4'-DDE (o,p'-DDE)	1.55 U	1.1 UJ	1.71 U	1.16 U	1.57 U	2.39 U	1.34 U	1.2 U	1.47 U
2,4'-DDT (o,p'-DDT)	1.55 UJ	1.1 UJ	1.71 UJ	1.16 UJ	1.57 UJ	2.39 U	1.34 U	1.2 U	1.47 U

Appendix A-1
BERA Sediment Dataset

	D1/50 2000	DI /50 2000	DI /50 2000	D1/50 2000	DI /50 2000	DI /50 2000	D1/50 2000	DI /50 0000	D1/50 2000
Task:	1	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:		PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
•			PB059.1-1SS010-091102-N						
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:		0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:		N 2201624 7257	N 2201521 (202	N 2201640 7220	N 22014074002	N 2204552 0022	N 2201620 7105	N 2201424 2755	N 22011C0 2007
X:	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
1:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
PCB-194	4820	1560	16500	2540	11200	1800	234	3040	22400 J
PCB-195	1890	584	6980	942	4490	675	90.3	1230	9070
PCB-196	2920	937	9280	1460	5800	990	112	1570	12600 J
PCB-197	276	96.9	699	139	450	83	33.8	120	644
PCB-198/199	6280	2140	20000 J	3450	12600	2150	258	3280	27300 J
PCB-200	999	374	3600	528	2260	385	36.4	621	3670
PCB-201	1010	364	3460	574	2310	402	43.1 J	593	3510
PCB-202	1310	468	4980	787	3100	596	66.6	867	5440
PCB-203	3150	1040	10100	1650	6570	1030	132	1730	14900 J
PCB-204	44.2 U	43.5 U	90.1 U	50 U	49.5 U	46.3 U	36.8 U	37.3 U	60.6 U
PCB-205	316	90	1050	154	659	111	14 J	209	1350
PCB-206	2510	918	6150	1340	4130	910	143	1230	7480
PCB-207	563	252	947	276	634	209	109	220	1010
PCB-208	984	371	1940	545	1340	363	108	391	2060
PCB-209	42200 J	20900 J	4380	7260	2760	7150	4800	3490	3300
Total PCB Congener (U = 1/2)	7107162.55 J	2553728.4 J	46631784.2 J	4165038.3 J	31807916.55 J	2646980.85 J	262392.45 J	6585980.37 J	43592345.9 J
Total PCB Congener minus 209 (U = 1/2)	7064962.55 J	2532828.4 J	46627404.2 J	4157778.3 J	31805156.55 J	2639830.85 J	257592.45 J	6582490.37 J	43589045.9 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	2456.0228 J	903.84186 J	16015.46495 J	1439.98669 J	10635.48175 J	902.96612 J	88.861198	2127.39646 J	13584.9484 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	12.216355 J	5.0113425 J	77.4417025 J	8.357195 J	51.5552375 J	4.9127425 J	0.511139	10.4427875 J	63.697565 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	118.6355 J	53.91173 J	727.5282 J	90.82467 J	489.2015 J	52.17471 J	6.282384	98.94463 J	576.3853 J
Monochlorobiphenyls (total)	2340	1630 J	22900	1450	63400	1490	103 J	6310	43300
Dichlorobiphenyls (total)	72400	28300 J	634000	37200	814000	30500	3050	95000	970000
Trichlorobiphenyls (total)	1290000	466000	10100000	590000	6910000	449000	45800 J	1340000	10600000
Tetrachlorobiphenyls (total)	3470000	1180000	22100000	2010000	16000000	1300000	128000	3260000 J	21300000
Pentachlorobiphenyls (total)	1690000	670000	11000000	1170000	6540000	641000	57900	1470000	8390000
Hexachlorobiphenyls (total)	409000	145000	2260000	273000	1180000	166000	15800	334000 J	1760000
Heptachlorobiphenyls (total)	102000	31100	420000	53800	242000	40700	4430 J	60200	456000
Octachlorobiphenyls (total)	23000	7650	76800	12200	49500	8220	1020 J	13300	101000
Nonachlorobiphenyls (total)	4050	1540	9040	2160	6110	1480	360	1840	10500
PCB Aroclors (μg/kg)									
Aroclor 1260									
Aroclor 1254									
Aroclor 1268									
Aroclor 1221									
Aroclor 1232									
Aroclor 1248	-								
Aroclor 1016									
Aroclor 1262	-								
Aroclor 1242									
Total PCB Aroclors (U = 1/2)									
Pesticides (µg/kg)									
2,4'-DDD (o,p'-DDD)	1.71 U	1.23 U	0.951 U	1.26 U	0.998 U	1.28 U	1.04 U	0.962 U	0.832 U
2,4'-DDE (o,p'-DDE)	1.71 U	1.23 U	0.951 U	1.26 U	0.998 U	1.28 U	1.04 U	0.962 U	0.832 U
2,4'-DDT (o,p'-DDT)	1.71 U	1.23 U	0.951 U	1.26 U	0.998 U	1.28 U	1.04 U	0.962 U	0.832 U

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	2011 USC	2011 USC	2011 USC				
Location ID:	PB074	PB081	PB093	PB097	PB101	PB068	PB069.1	PB070A
					PB101-1SS010-091103-N	PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm				
Sample Type:	N N	0 - 10 CIII	N N	0 - 10 CIII	0 - 10 CIII	0 - 10 CIII	NI NI	0 - 10 cm
Sample Type.	3200828.0183	3200761.3787	3200970.4291	3201103.1242	3201287.2658	3201209.861	3201205.859	3201087.255
Λ. Υ:	13830463.2171	13830005.1879	13828894.7843	13828566.1540	13828181.2285	13831121.51	13830973.3	13830903.51
PCB-194	983	16300	317	403	61.1			
PCB-195	409	7030	132	165	19.1 J			
PCB-196	589	9750	206	292	41.6	-		
PCB-197	45.7	672	19.1	23.3	48.1 U	-		
PCB-198/199	1050	19400 J	382	716	110			
PCB-200	211	3550	72.6	125	21.6			
PCB-201	194	3260	69.7	139	20.2			
PCB-202	280	3790	104	241	34.9			
PCB-203	597	9960	216	363	52.8			
PCB-204	36.8 U	46.3 U	52.1 U	45.9 U	48.1 U			
PCB-205	60.3 J	1030	21.2 J	15.6 J	48.1 U			
PCB-206	340	4570	129	269	40.2			
PCB-207	54	768	45.8	57.2	12.5			
PCB-208	124	1230	61	122	25.8			
PCB-209	592 U	11700	482	331	379			
Total PCB Congener (U = 1/2)	3275873.3 J	21717830.3 J	1302666.6 J	1653530.6 J	405981.24 J			
Total PCB Congener minus 209 (U = 1/2)	3275577.3 J	21706130.3 J	1302184.6 J	1653199.6 J	405602.24 J			
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	866.7473 J	5936.70135 J	194.464055	389.999914 J	76.6573515			
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	6.67402 J	28.7707575 J	0.727345	1.5890345 J	0.31932825			
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	87.8865 J	260.0181 J	6.445795	14.579522 J	3.5076445			
Monochlorobiphenyls (total)	3890	221000	719	215	242			
Dichlorobiphenyls (total)	54800	899000	60800	15300	19000			
Trichlorobiphenyls (total)	639000	5180000	514000	380000	137000			
Tetrachlorobiphenyls (total)	1280000	9770000	541000	990000	193000			
Pentachlorobiphenyls (total)	864000	400000	130000	218000	47300			
Hexachlorobiphenyls (total)	380000	1170000	41900	37300	6890			
Heptachlorobiphenyls (total)	49400 J	376000	10300	8670	1540 J			
Octachlorobiphenyls (total)	4420 J	74800	1540 J	2480 J	361 J			
Nonachlorobiphenyls (total)	518	6570	236	448	78.5			
PCB Aroclors (µg/kg)	310	0370	230	110	70.5			
Aroclor 1260						1320 U	108 U	37.2 U
Aroclor 1254						15700	2630	604
Aroclor 1268						1320 U	108 U	37.2 U
Aroclor 1221						1320 U	108 U	37.2 U
Aroclor 1232						1320 U	108 U	37.2 U
Aroclor 1248						56000	6440	1720
Aroclor 1016						1320 U	108 U	37.2 U
Aroclor 1262						1320 U	108 U	37.2 U
Aroclor 1242						1320 U	108 U	37.2 U
Total PCB Aroclors (U = 1/2)			 			76320	9448	2454.2
Pesticides (µg/kg)	·-		·-			,0320	3-1-10	E-73716
2,4'-DDD (o,p'-DDD)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
2,4'-DDE (0,p'-DDE)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
2,4'-DDT (0,p'-DDT)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
ביא יומה. לומח. (חים י ליומר ליים)	0.030 0	J.33 U	4.07 0	4.09 0	4.31 0	-		

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:		PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
PCB-194				
PCB-195				
PCB-196				
PCB-197				
PCB-198/199				
PCB-200				
PCB-201			-	
PCB-202			-	
PCB-203			-	
PCB-204			-	
PCB-205			-	
PCB-206			-	
PCB-207				
PCB-208				
PCB-209				
Total PCB Congener (U = 1/2)			-	-
Total PCB Congener minus 209 (U = 1/2)				
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)				
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)				
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)			-	-
Monochlorobiphenyls (total)				<u></u>
Dichlorobiphenyls (total)				
Trichlorobiphenyls (total)				
Tetrachlorobiphenyls (total)				
Pentachlorobiphenyls (total)				
Hexachlorobiphenyls (total)				
Heptachlorobiphenyls (total)			-	
Octachlorobiphenyls (total)				
Nonachlorobiphenyls (total)				
PCB Aroclors (μg/kg)				
Aroclor 1260	47.2 U	1530 U	1490 U	251 U
Aroclor 1254	1010	20500 J	23200 J	4300
Aroclor 1268	47.2 U	1530 U	1490 U	251 U
Aroclor 1221	47.2 U	1530 U	1490 U	251 U
Aroclor 1232	47.2 U	1530 U	1490 U	251 U
Aroclor 1248	2830	80600 J	83600 J	15000
Aroclor 1016	47.2 U	1530 U	1490 U	251 U
Aroclor 1262	47.2 U	1530 U	1490 U	251 U
Aroclor 1242	47.2 U	1530 U	1490 U	251 U
Total PCB Aroclors (U = 1/2)	4005.2	106455 J	112015 J	20178.5
Pesticides (μg/kg)				
2,4'-DDD (o,p'-DDD)			-	
2,4'-DDE (o,p'-DDE)			-	
2,4'-DDT (o,p'-DDT)				

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB001.1	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
	PB001.1-1SS010-091027-N								
Sample Date:	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N 20 cm	N 20 CIII	N N	N 20 cm	N N	N N	N 10 cm	N N
y.	3202382.3842	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011	3202037.3755	3201747.2539
Y:	13836546.3402	13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614	13836300.0036	13836110.2965	13836119.3947
4,4'-DDD (p,p'-DDD)	58.8 J	30.6 J	4.98 J	35.1 J	28.3 J	29.4 J	46.2 J	24.8 J	17.8 J
4,4'-DDE (p,p'-DDE)	12.5 J	11.1 J	1.6 J	11.6 J	10.1 J	10.2 J	16.9 J	7.97 J	6.31
4,4'-DDT (p,p'-DDT)	62.7 J	68.1 J	1.28 U	109 J	59.4 J	48.4 J	3.11 UJ	64	1.27 U
Sum 2,4 DDT, DDE, DDD (U=1/2)	25.77 J	13.37 J	3.34 J	14.24 J	2.07 U	13.95 J	3.11 U	1.42 U	1.27 U
Sum 4,4 DDT, DDE, DDD (U = 1/2)	134 J	109.8 J	7.22 J	155.7 J	97.8 J	88 J	64.655 J	96.77 J	24.745 J
Sum DDD (U=1/2)	82.5 J	42 J	7.04 J	47.9 J	29.335 J	41.7 J	47.755 J	25.51 J	18.435 J
Sum DDE (U=1/2)	13.535 J	12.085 J	2.24 J	12.32 J	11.135 J	11.025 J	18.455 J	8.68 J	6.945
Sum DDT (U=1/2)	63.735 J	69.085 J	1.28 U	109.72 J	60.435 J	49.225 J	3.11 UJ	64.71	1.27 U
Total DDx (U = 1/2)	159.77 J	123.17 J	10.56 J	169.94 J	100.905 J	101.95 J	69.32 J	98.9 J	26.65 J
Aldrin	1.04 U	0.986 U	0.64 U	0.718 U	1.03 U	0.823 U	1.55 U	0.712 U	0.634 U
alpha-Chlordane (cis-Chlordane)	34.1	8.89 J	10.7 J	16 J	12.1	41.9	24.8	39.5	54.4 NJ
Endosulfan-alpha (I)	66.8 NJ	18.2 NJ	7.44 NJ	35.4 NJ	25.8 NJ	51.5 NJ	46.6 J	70.9	27.5 NJ
Endosulfan-beta (II)	1.04 U	0.986 UJ	0.64 U	0.718 U	1.03 U	0.823 U	1.55 U	0.712 U	0.634 U
Endrin	1.04 U	0.986 UJ	5.86 NJ	19.7 NJ	1.03 U	18.7 NJ	1.55 U	47.9 NJ	0.634 U
gamma-BHC (Lindane)	1.04 U	0.986 UJ	0.64 U	0.718 U	27.4 NJ	57.7 NJ	1.55 U	0.712 U	0.634 U
gamma-Chlordane	1.04 U	0.986 UJ	0.64 U	0.718 U	1.03 U	0.823 U	1.55 U	0.712 U	0.634 U
Heptachlor epoxide	1.04 U	0.986 U	0.64 U	0.718 U	1.03 U	0.823 U	1.55 U	0.712 U	0.634 U
Methoxychlor	1.04 U	0.986 UJ	0.64 U	0.718 U	1.03 U	0.823 U	1.55 U	0.712 U	0.634 U
Dioxin Furans (ng/kg)									
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	21.3	27	9.46	46.9	26	21.5	23.7	17.2	14.3
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	3.85	3.76	3.24 J	4.21 J	4.8 J	3.9	4 J	3.57 J	3.34
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	5.46 J	6.5	4.57	7.62	9.36	5.72	8.5	4.13 J	4.27 J
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	19.1	18.8	11.8	26.4	18.8	14.5 J	23.8	12.6 J	13.7
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	10.3	12	8.8 J	14.8	18.2 J	13.2	14.4 J	8.42 J	6.02 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	501	414	280	555	460	398	682	349	367
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	5570	5100	2690	5910	5510	4890	8640	4100	4550
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	104	94.8	39.9	192	121	114	105	95.1	81.4
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	56.2	79.8	45.4	160	186	110	81.9	103	72.5
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	52.7	72.1	43.6	124	151	80.3	68.1	78	61.4
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	103	155	228	327	349 J	201	144	177	135
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	46.4	65.4	139	144	124	88.1	64	81.4	59.7
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	23.8	36.9 J	38.7	63.4	122 J	53.4	36.2 J	42.3	32.7 J
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	26.8	34.2	75.4	62.7	69.9	41.9	34.3	38.5	29.6
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	1150	1730	7320 J	3990	1450	2240	1410	1880	1620
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	102	147	604	356	133	221	113	184	127
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	45500 J	67500 J	253000 J	149000 J	48500 J	81000 J	49600 J	60100 J	63300 J
Total Tetrachlorodibenzo-p-dioxin (TCDD)	58 J	61.1 J	23 J	89.1 J	65 J	53.7 J	66.7 J	43.4 J	31.8 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	68 J	53.8 J	34.7 J	82.5 J	64.7 J	54.3 J	60.3 J	43.9 J	41.8 J
Total Hexachlorodibenzo-p-dioxin (HxCDD)	238 J	233 J	147 J	297 J	286 J	210 J	289 J	182 J	170 J
Total Heptachlorodibenzo-p-dioxin (HpCDD)	1590	1160	680	1390	1350	1130	1990	1020	1110
Total Tetrachlorodibenzofuran (TCDF)	459 J	419 J	240 J	791 J	550 J	460 J	471 J	403 J	326 J
Total Pentachlorodibenzofuran (PeCDF)	345 J	412 J	411 J	765 J	740 J	490 J	438 J	468 J	363 J
Total Hexachlorodibenzofuran (HxCDF)	871	1500 J	2090 J	2170	1650 J	1290 J	1150 J	1200 J	1030 J
Total Heptachlorodibenzofuran (HpCDF)	1950	2900	10300	6050	2390	3470	2430	2930	2460

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB011	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
		PB013.1-1SS010-091028-N					PB018-1SS010-091101-N		
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N N	N N	N	N N	N N	N N	N N	N	FD
χ.	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
Y:	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
4,4'-DDD (p,p'-DDD)	33 J	1.24 U	9.56 J	26.4 J	1.08 U		0.991 UJ	14 J	1.01 U
4,4'-DDE (p,p'-DDE)	12.8 J	6.08	4.42 J	6.98 J	7.97	-	3.77 J	7.94	3.87
4,4'-DDT (p,p'-DDT)	1.41 U	1.24 U	19.6 J	2.74 UJ	53.5 J	-	0.991 UJ	20.7 J	21.4
Sum 2,4 DDT, DDE, DDD (U=1/2)	20.11 J	1.24 U	5.13 J	12.02 J	1.08 U		0.991 UJ	1.15 U	1.01 U
Sum 4,4 DDT, DDE, DDD (U = 1/2)	46.505 J	7.32	33.58 J	34.75 J	62.01 J		4.761 J	42.64 J	25.775
Sum DDD (U=1/2)	51.7 J	1.24 U	13.62 J	35.68 J	1.08 U		0.991 UJ	14.575 J	1.01 U
Sum DDE (U=1/2)	13.505 J	6.7	4.955 J	8.35 J	8.51		4.2655 J	8.515	4.375
Sum DDT (U=1/2)	1.41 U	1.24 U	20.135 J	2.74 UJ	54.04 J		0.991 UJ	21.275 J	21.905
Total DDx (U = 1/2)	66.615 J	9.18	38.71 J	46.77 J	63.63 J		6.2475 J	44.365 J	27.29
Aldrin	0.707 U	0.622 U	0.534 U	1.37 U	0.541 U	-	0.495 UJ	0.576 U	0.507 UJ
alpha-Chlordane (cis-Chlordane)	7.72 NJ	5.17 J	13.2 J	14.5 J	3.91		1.66 J	13.1 J	7.2 J
Endosulfan-alpha (I)	0.707 U	14.7 NJ	18.2 NJ	29.5 NJ	29.3 NJ		0.495 UJ	19.3 NJ	17.4 NJ
Endosulfan-beta (II)	0.707 U	0.622 U	0.534 U	1.37 UJ	0.541 U		0.495 UJ	0.576 U	0.507 UJ
Endrin	0.707 U	0.622 U	0.534 U	1.37 UJ	0.541 U		0.495 UJ	0.576 U	0.507 UJ
gamma-BHC (Lindane)	0.707 U	39.5 NJ	0.534 U	27.2 NJ	43.5 NJ		164 NJ	0.576 U	14.4 NJ
gamma-Chlordane	0.707 U	27.5 NJ	0.534 U	1.37 UJ	0.541 U		0.495 UJ	0.576 U	0.507 UJ
Heptachlor epoxide	0.707 U	0.622 U	0.534 U	1.37 U	11.2 NJ		0.495 UJ	0.576 U	0.507 UJ
Methoxychlor	0.707 U	0.622 U	0.534 U	1.37 UJ	0.541 U		0.495 UJ	0.576 U	0.507 UJ
Dioxin Furans (ng/kg)	0.1.0.1.0	***************************************			0.0.12.0		0.100 01	0.0.0	0.007
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	27.9	15.8 J	5.09	30	9.11 J		12.7	11.6 J	10.3
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	3.62 J	2.26 J	1.02 J	4.72 J	3.39 U		3.58 U	2.46 J	5.34 U
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	5.59 J	3.03 J	2.18	9.2	1.83 J		7.75 U	3.13 J	6.63 U
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	19.7	12.3	6.31	24.1	10.3		9.31	17.2	11.1
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	10.1 J	8.49	4.03 J	19.3	6.29 J		5.38 J	7.12	5.34 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	403	278	121	680	208		276	355	195
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	4290	3420	1340	7840	2460		2460	3260	2100
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	138	114	31.7	156	74		81.6	81.6	66.8
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	145	147	30.3	105	117		90.8	57	59.8
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	111	112	23.1	84.3	89.3		78.4	45.8	49.1
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	262	239	70.7	168	160	-	208	97.6	99
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	106	81.6	31.7	76.5	62.4		86.7	54.5	36.3
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	68.2	66.6	14.8 J	37 J	41.5		51.9	22.5 J	28.2 J
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	47.6	38.2	14.9	43.3	32.4		37.3	25.3 J	17.8
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	1920	1220	1250	1720	893		1630	1670	654
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	187	138	72.1	145	100		156	133	60.9
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	65700 J	35100 J	27600 J	59700 J	37000 J		99300 J	58700 J	19100 J
Total Tetrachlorodibenzo-p-dioxin (TCDD)	52.7 J	44.6 J	13.8 J	76.2 J	22 J		26.4 J	30.1 J	21.3 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	59 J	31.6 J	18.1 J	73.4 J	25.8 J		51.1 J	46.8 J	21.7 J
Total Hexachlorodibenzo-p-dioxin (HxCDD)	231 J	146 J	78.8 J	319 J	132 J		151 J	187 J	116 J
Total Heptachlorodibenzo-p-dioxin (HpCDD)	1170	755	361	1910	546		731	955	523
Total Tetrachlorodibenzofuran (TCDF)	555 J	437 J	157 J	671 J	334 J		338 J	366 J	296 J
Total Pentachlorodibenzofuran (PeCDF)	603	541 J	159 J	537	442 J		457 J	302 J	252 J
Total Hexachlorodibenzofuran (HxCDF)	1370 J	979	493 J	1290 J	758 J		1580	944 J	494 J
	2980	1950	1670	2890	1450		2710	2490	1030

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	· ·	PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
Sample ID:			PB023-1SS010-091030-N			PB028-1SS010-091030-N		PB034-1SS010-091030-N	
Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N 20 cm	FD	N 10 cm	N N	N	N N	N 10 cm	N 20 cm	N 20 0
X:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
Y:	13835456.7389	13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
4,4'-DDD (p,p'-DDD)	1 UJ	12.1 J	11.1 J	10.4 J	94.1 J	11.6 J	33.2 J	45.6 J	2.04 U
4,4'-DDE (p,p'-DDE)	3.34 J	13.8 J	12.8 J	7.2 J	38 J	4.78	7.75 J	23.7 J	2.04 U
4,4'-DDT (p,p'-DDT)	24 J	16.8 J	1.06 U	1.07 U	80 J	2.1 UJ	44.8 J	2.31 UJ	2.04 UJ
Sum 2,4 DDT, DDE, DDD (U=1/2)	1 UJ	17.63 J	9.99 J	1.07 U	59.1 J	2.1 UJ	11.95 J	11.12 J	2.04 UJ
Sum 4,4 DDT, DDE, DDD (U = 1/2)	27.84 J	42.7 J	24.43 J	18.135 J	212.1 J	17.43 J	85.75 J	70.455 J	2.04 UJ
Sum DDD (U=1/2)	1 UJ	21.32 J	17.1 J	10.935 J	136.8 J	12.65 J	43.7 J	54.41 J	2.04 U
Sum DDE (U=1/2)	3.84 J	17.31 J	16.26 J	7.735 J	53.2 J	5.83	8.475 J	24.855 J	2.04 U
Sum DDT (U=1/2)	24.5 J	21.7 J	1.06 U	1.07 U	81.2 J	2.1 UJ	45.525 J	2.31 UJ	2.04 UJ
Total DDx (U = 1/2)	29.34 J	60.33 J	34.42 J	19.74 J	271.2 J	20.58 J	97.7 J	81.575 J	2.04 UJ
Aldrin	0.5 U	0.565 UJ	0.531 UJ	0.534 UJ	1.2 UJ	1.05 UJ	0.723 UJ	1.16 UJ	1.02 UJ
alpha-Chlordane (cis-Chlordane)	6.48	4.06 J	19.4 NJ	7.48 J	1.2 UJ	2.07 J	0.723 UJ	10.7 J	66.9 J
Endosulfan-alpha (I)	0.5 U	0.565 UJ	0.531 UJ	21.2 J	1.2 UJ	1.05 UJ	0.723 UJ	27 J	87.7 J
Endosulfan-beta (II)	0.5 U	0.565 UJ	0.531 UJ	0.534 UJ	57.5 NJ	1.05 UJ	53.3 J	1.16 UJ	133 NJ
Endrin	0.5 U	16.2 NJ	41.4 NJ	0.534 UJ	54.6 NJ	4 J	0.723 UJ	1.16 UJ	1.02 UJ
gamma-BHC (Lindane)	19.4 NJ	76.3 NJ	0.531 UJ	34.8 NJ	1.2 UJ	24 NJ	86.9 J	1.16 UJ	6.56 J
gamma-Chlordane	0.5 U	5.05 J	15 NJ	0.534 UJ	1.2 UJ	1.05 UJ	74.9 J	1.16 UJ	1.02 UJ
Heptachlor epoxide	21.5 NJ	0.565 UJ	0.531 UJ	0.534 UJ	1.2 UJ	1.05 UJ	0.723 UJ	1.16 UJ	1.02 UJ
Methoxychlor	0.5 UJ	0.565 UJ	0.531 UJ	0.534 UJ	1.2 UJ	1.05 UJ	0.723 UJ	1.16 UJ	1.02 UJ
Dioxin Furans (ng/kg)									
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	9.44	10 J	14.3	19	830	9.3	158	63.6	23
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	3.08 U	3.08 J	4.45 J	3.11 J	9.66 J	1.29 J	4.06	6.65 J	5.63 J
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	2.13 J	4.23 J	4.31	4.89 J	7.74 J	3.38 U	3.55 J	9.6 J	8.15 J
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	8.54 J	11.6	11	16.4	49.6	3.77 U	18	74.7	30.1
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	4.86 J	7.59	6.79 J	9.6 J	17.9 J	3.77 U	8.83 J	26.2	16.3 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	195	208	188	364	1130	238	603	1800	773
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	2150	2290	2050	3790	8750	3680	4370	11400 J	8830
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	61.9	69.7	71.8	101	2860 J	40.4	364	298	128
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	54	74.2	77.6	114	154	26.5	51.6	143	111
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	46.3	62.5	51.3	91.6	180	27.3	48.1	125	94.8
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	85.2	206	120	206	556	73.3 J	116	292	165
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	39.8	99.9	72.1	85.7	170	26.9 J	37.4	120	67.5
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	19 J	32.4	34.4 J	52.4	130	13.1 J	25.2 J	64.3	37.2
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	14.8 J	45.7	30.7	36.8	164	16.6	23	93.1	35
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	638	2410	2690	1480	13100 J	455	7010 J	4510	594
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	58.9	234	252	155	1390	42.2	256 J	422	77.6
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	17800 J	77900 J	110000 J	32200 J	2570000 J	17400 J	1460000 J	250000 J	13100 J
Total Tetrachlorodibenzo-p-dioxin (TCDD)	25.6 J	42.9 J	35.8 J	41 J	976 J	25.4 J	181 J	130 J	71.5 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	25.8 J	56.7 J	43.8 J	43.3 J	252 J	21.3 J	22.5 J	117 J	74.1 J
Total Hexachlorodibenzo-p-dioxin (HxCDD)	122 J	148 J	134 J	189 J	1100 J	106 J	209 J	658 J	304 J
Total Heptachlorodibenzo-p-dioxin (HxCDD) Total Heptachlorodibenzo-p-dioxin (HpCDD)	555	558	510	972	3350	638	1340	4500	1910
Total Pentaghlaradibangaliyan (PaCDE)	271 J	321 J	385 J	438 J	5140 J	169 J	738 J	1030 J	599 J
Total Pentachlorodibenzofuran (PeCDF)	248 J	453 J	447 J	463 J	1860 J	165 J	248 J	954	536 J
Total Hexachlorodibenzofuran (HxCDF)	467 J	1610	2640 J	890 J	74600 J	540 J	527 J	3960 J	763 J
Total Heptachlorodibenzofuran (HpCDF)	1040	3670	5820	2210	163000	933	8820 J	10600	1170

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009						
Location ID:	PB041	PB043	PB044	PB047.1	PB047.2	PB048	PB049	PB049	PB053
							PB049-1SS010-091031-D		
Sample Date:	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm						
Sample Type:	O-10 CIII	N N	N N	O - 10 Cili	O-10 CIII	O - 10 CIII	FD	N N	N N
v.	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
Λ. V.	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
1.									
4,4'-DDD (p,p'-DDD)	16.6 J	11.3 J	16.9 J	32.4 J	15.2 J	26.7 J	1.34 U	1.2 U	142 J
4,4'-DDE (p,p'-DDE)	12.2 J	1.1 UJ	11.4	1.16 U	7.62	9.02	10.4	9.94	1.47 U
4,4'-DDT (p,p'-DDT)	1.55 UJ	1.1 UJ	1.71 UJ	1.16 UJ	1.57 UJ	2.39 U	33.1 J	32.7 J	1.47 U
Sum 2,4 DDT, DDE, DDD (U=1/2)	1.55 UJ	1.1 UJ	1.71 UJ	1.16 UJ	1.57 UJ	2.39 U	1.34 U	1.2 U	1.47 U
Sum 4,4 DDT, DDE, DDD (U = 1/2)	29.575 J	12.4 J	29.155 J	33.56 J	23.605 J	36.915 J	44.17 J	43.24 J	143.47 J
Sum DDD (U=1/2)	17.375 J	11.85 J	17.755 J	32.98 J	15.985 J	27.895 J	1.34 U	1.2 U	142.735 J
Sum DDE (U=1/2)	12.975 J	1.1 UJ	12.255	1.16 U	8.405	10.215	11.07	10.54	1.47 U
Sum DDT (U=1/2)	1.55 UJ	1.1 UJ	1.71 UJ	1.16 UJ	1.57 UJ	2.39 U	33.77 J	33.3 J	1.47 U
Total DDx (U = 1/2)	31.9 J	14.05 J	31.72 J	35.3 J	25.96 J	40.5 J	46.18 J	45.04 J	145.675 J
Aldrin	0.777 UJ	0.55 UJ	0.857 UJ	0.578 UJ	0.784 UJ	1.2 U	0.67 U	0.6 U	0.736 U
alpha-Chlordane (cis-Chlordane)	0.777 UJ	64.4 J	10.6 J	30.9 J	39.8 J	25.8	40.7	26.1	0.736 U
Endosulfan-alpha (I)	51.9 NJ	76.3 J	42.9 NJ	121 NJ	60.1 J	42.2 J	56.5 NJ	72.2 NJ	0.736 U
Endosulfan-beta (II)	0.777 UJ	0.55 UJ	0.857 UJ	0.578 UJ	0.784 UJ	1.2 U	0.67 U	112 NJ	0.736 U
Endrin	35.4 J	33.6 NJ	0.857 UJ	0.578 UJ	0.784 UJ	1.2 U	0.67 U	0.6 U	0.736 U
gamma-BHC (Lindane)	13.6 NJ	2.42 J	7.54 NJ	5.73 NJ	3.56 NJ	3.95	1.4 J	1.98	26.8
gamma-Chlordane	57.8 NJ	0.55 UJ	0.857 UJ	0.578 UJ	0.784 UJ	1.2 U	0.67 U	0.6 U	0.736 U
Heptachlor epoxide	38.3 J	0.55 UJ	0.857 UJ	0.578 UJ	0.784 UJ	1.2 U	0.67 U	0.6 U	0.736 U
Methoxychlor	0.777 UJ	0.55 UJ	0.857 UJ	0.578 UJ	0.784 UJ	1.2 U	0.67 U	0.6 U	0.736 U
Dioxin Furans (ng/kg)									
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	10	3.89	12.2	6.17	12	14.6	4.84	4.99	48.7
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	4.18 J	2.21 J	4.28	4.41	4.01	4.72	4.42 J	4.15	7.53 J
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	7.06	3.56	7.3	5.85	7.47	9.03	6.3	6.19	12.4
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	17.8	10.8	20.9	18.1	20	23.6	22.2	21.6	76.4
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	11.5	7.11	11.8	10.9	12.8	15.2	13.8	12.6	27.1
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	381	326	531	449	490	588	557	516	1790
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	4510	3740	7170	4850	5890	7600	5650	5910	19100 J
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	130	47.7	621	59.7	178	130	49.2	46.7	399
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	174	22.9	118	71.4	70	72.2	43.9	41.9	573
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	134	25	90.8	67.3	55.4	59	43.2	39.2	462
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	284	33.9	132	148	90.4	109	69.9	65.4	2000
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	82.7	14.9	46.5	55.9	37.7	45.7	33.5	30.8	487
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	58.9	7.32	27.2	25.1	19.3	23.2	14.5	14.4	358
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	40.1	9.7	24.8	22.9	21.4	25.2	18.2	17.2	156
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	377	135	331	544	326	387	211	200	1370
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	76.7	16.1	51.9	70.8	42.5	53	25.7	24.7	614
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	5550	1580	5190	4210	5720	6410	1720	1680	9300
Total Tetrachlorodibenzo-p-dioxin (TCDD)	41.9 J	24.5 J	48.5 J	38.5 J	50.2 J	59.7 J	43.3 J	40.9 J	191 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	48.4 J	26 J	51.2 J	48.3 J	53.1 J	63.8 J	64 J	61.2 J	85 J
Total Hexachlorodibenzo-p-dioxin (HxCDD)	194 J	107	223	185	220	271	211	199 J	415
Total Heptachlorodibenzo-p-dioxin (HpCDD)	972	754	1480	1170	1340	1670	1400	1320	3500
Total Tetrachlorodibenzofuran (TCDF)	525 J	203 J	1200 J	372 J	469 J	476 J	326 J	300 J	2000 J
Total Pentachlorodibenzofuran (PeCDF)	660 J	149 J	447 J	416 J	316 J	356 J	297 J	273 J	2030 J
Total Hexachlorodibenzofuran (HxCDF)	785 J	175	481	540	391	482 J	339	321	4050
Total Heptachlorodibenzofuran (HpCDF)	748	277	683	911	613	754	423	405	3350

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009				
Location ID:	PB056	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
					PB063.1-1SS010-091031-N				
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm				
Sample Type:	0 - 10 CIII	0 - 10 CIII	N N	0 - 10 CIII	0 - 10 CIII	N N	N N	N N	0 - 10 CIII
Sample Type.	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
4,4'-DDD (p,p'-DDD)	1.71 U	1.23 U	0.951 U	1.26 U	0.998 U	1.28 U	1.04 U	0.962 U	0.832 U
4,4'-DDE (p,p'-DDE)	6.16	4.29	0.951 U	8.54	0.998 U	5.96	1.65	0.962 U	0.832 U
4,4'-DDT (p,p'-DDT)	43.9 J	20.8 J	0.951 U	28.2 J	0.998 U	37.9	1.04 U	0.962 U	0.832 U
Sum 2,4 DDT, DDE, DDD (U=1/2)	1.71 U	1.23 U	0.951 U	1.26 U	0.998 U	1.28 U	1.04 U	0.962 U	0.832 U
Sum 4,4 DDT, DDE, DDD (U = 1/2)	50.915 J	25.705 J	0.951 U	37.37 J	0.998 U	44.5	2.69	0.962 U	0.832 U
Sum DDD (U=1/2)	1.71 U	1.23 U	0.951 U	1.26 U	0.998 U	1.28 U	1.04 U	0.962 U	0.832 U
Sum DDE (U=1/2)	7.015	4.905	0.951 U	9.17	0.998 U	6.6	2.17	0.962 U	0.832 U
Sum DDT (U=1/2)	44.755 J	21.415 J	0.951 U	28.83 J	0.998 U	38.54	1.04 U	0.962 U	0.832 U
Total DDx (U = 1/2)	53.48 J	27.55 J	0.951 U	39.26 J	0.998 U	46.42	4.25	0.962 U	0.832 U
Aldrin	0.854 U	0.615 U	0.476 U	0.631 U	0.499 U	0.641 U	0.519 U	0.481 U	0.416 U
alpha-Chlordane (cis-Chlordane)	10.5	13.3 J	19.2	9.19	146	2.89	0.519 U	124	0.416 U
Endosulfan-alpha (I)	34.6 NJ	24.1 NJ	124 NJ	33.5 NJ	165 NJ	19.7 NJ	2.79 NJ	133 J	292 NJ
Endosulfan-beta (II)	0.854 U	0.615 U	0.476 U	0.631 U	0.499 U	0.641 U	0.519 U	0.481 U	0.416 U
Endrin	0.854 U	0.615 U	24.1 NJ	0.631 U	0.499 U	0.641 U	0.519 U	0.481 U	0.416 U
gamma-BHC (Lindane)	2.72 NJ	0.924	8.33 NJ	0.631 U	0.499 U	0.641 U	0.519 U	2.76	0.416 U
gamma-Chlordane	0.854 U	0.615 U	0.476 U	0.631 U	0.499 U	0.641 U	0.519 U	0.481 U	0.416 U
Heptachlor epoxide	0.854 U	0.615 U	0.476 U	0.631 U	0.499 U	0.641 U	0.519 U	0.481 U	0.416 U
Methoxychlor	0.854 U	0.615 U	0.476 U	0.631 U	0.499 U	0.641 U	0.519 U	0.481 U	0.416 U
Dioxin Furans (ng/kg)									
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	4.92	1.77	3.35	1.69	2.92	1.74	0.468 J	1.18 J	5.04
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	4.48	2.4 J	5.15 J	3.19	2.49 J	4.43	1.51 J	2.43	4.17
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	6.42	3.8	6.11	5	3.14	7.44	2.37 J	2.48 J	4.18
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	20.2	11.7	26.3	16.9	14.6	29.9	10.3	9.63	23.2
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	12.9	7.48	14	11	7.44	17.1	5.35	5.8	10
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	495	445	720	448	391	757	338	255	690
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	5410	3980	7090	4360	3950	7850	3710	2450	8430
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	34.2	12.8	60.1	13.1	42.1	12.2	1.79	11.4	87.4
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	34.3	10.5	48.8	11	15.9	10.1	1.23 J	9.75	31.7
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	29.6	16.7	106	12.7	45.7	11.6	1.78 J	12.8	109
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	49.3	39.7	158	26	31.8	31.5	3.26	27.2	82.1
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	25.7	15.3	72.3	15.2	14.4	21	3.96	15.9	35.3
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	10.4	7.33	26.2	5.79	7.47	6.82	1.04 J	5.17 J	19.3
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	16	9.49	35.9	10.3	11.9	15.1	3.71	8.19	26.8
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	228	138	525	142	106	240	56.2	153	220
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	25.1	21.6	59.6	16.7	12.8	30.4	6.83	17.2	30.3
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	2060	1130	1170	612	308	822	299	690	537
Total Tetrachlorodibenzo-p-dioxin (TCDD)	44.7 J	20.3 J	38.1 J	22.8 J	27.8 J	23.1 J	7.57 J	15.6 J	45.4 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	54.6 J	25.7 J	47.5 J	31.1 J	28.8 J	41.1 J	13.3 J	21.1 J	43.1 J
Total Hexachlorodibenzo-p-dioxin (HxCDD)	208	123	185	151	118	246	109	87.1 J	159
Total Heptachlorodibenzo-p-dioxin (HpCDD)	1330	1080	1440	1060	890	1750	942	622	1450
Total Tetrachlorodibenzofuran (TCDF)	275 J	108 J	723 J	149 J	397 J	132 J	18.8 J	127 J	816 J
Total Pentachlorodibenzofuran (PeCDF)	271 J	106 J	638 J	114 J	242 J	122	15.5 J	108 J	520 J
Total Hexachlorodibenzofuran (HxCDF)	292 J	178 J	746	180	204	275 J	57 J	177 J	425
Total Heptachlorodibenzofuran (HpCDF)	449	279	933	290	260	530	179	354	472

Appendix A-1
BERA Sediment Dataset

1	5./50.000		5: /50 0000	-: /	5./50.000			
Task:	RI/FS 2009	2011 USC	2011 USC	2011 USC				
Location ID:	PB074	PB081	PB093	PB097	PB101	PB068	PB069.1	PB070A
•			PB093-1SS010-091103-N			PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 8 cm	0 - 10 cm				
Sample Type:	N	N	N	N	N	N	N	N
X: Y:	3200828.0183 13830463.2171	3200761.3787 13830005.1879	3200970.4291 13828894.7843	3201103.1242 13828566.1540	3201287.2658 13828181.2285	3201209.861 13831121.51	3201205.859 13830973.3	3201087.255 13830903.51
4,4'-DDD (p,p'-DDD)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
4,4'-DDE (p,p'-DDE)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
4,4'-DDT (p,p'-DDT)	22.8	5.33 U	4.67 U	4.69 U	4.51 U			
Sum 2,4 DDT, DDE, DDD (U=1/2)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
Sum 4,4 DDT, DDE, DDD (U = 1/2)	23.698	5.33 U	4.67 U	4.69 U	4.51 U			
Sum DDD (U=1/2)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U	-		
Sum DDE (U=1/2)	0.898 U	5.33 U	4.67 U	4.69 U	4.51 U			
Sum DDT (U=1/2)	23.249	5.33 U	4.67 U	4.69 U	4.51 U	-		
Total DDx (U = 1/2)	25.045	5.33 U	4.67 U	4.69 U	4.51 U			
Aldrin	0.449 U	5.33 U	4.67 U	4.69 U	4.51 U			
alpha-Chlordane (cis-Chlordane)	48.2 NJ	126	4.67 U	4.69 U	4.51 U	-		
Endosulfan-alpha (I)	63.3 NJ	5.33 U	4.67 U	4.69 U	4.51 U	-		
Endosulfan-beta (II)	80.9 NJ	291	4.67 U	4.69 U	4.51 U	-		
Endrin	0.449 U	5.33 U	4.67 U	4.69 U	4.51 U	-		
gamma-BHC (Lindane)	2.98	5.33 U	4.67 U	4.69 U	4.51 U			
gamma-Chlordane	0.449 U	5.33 U	4.67 U	4.69 U	4.51 U			
Heptachlor epoxide	0.449 U	5.33 U	4.67 U	4.69 U	4.51 U			
Methoxychlor	0.449 U	1780 NJ	4.67 U	4.69 U	4.51 U			
Dioxin Furans (ng/kg)								
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	0.341 J	4.58	0.217 J	0.137 J	0.156 U			
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	0.692 J	4.81 J	0.394 J	0.332 J	0.743 UJ	-		
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	1.4 J	4.72	0.33 U	0.366 J	0.304 U			
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	3.63	14.6	0.604 J	1.4 J	0.538 J	-		
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	2.66	7.52	0.606 J	1.01 J	0.613 J			
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	102	520 J	11.1 J	33	12.8	-		
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	1520	4600	102	380	361			
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	6.65	68.5	1.02	1.21	0.374 J			
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	3.01	17.6	0.253 J	0.638 J	0.558 U			
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	7.8	39	0.704 J	1.73 J	0.404 U	-		
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	6.63	28.6	0.629 J	1.4 J	0.659 J	-		
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	4.07	13.9	0.723 J	1.02 J	0.435 J			
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	1.61 J	36.1	0.294 J	0.281 J	0.272 J			
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	3.75	10.7	0.765 J	0.93 J	0.421 J			
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	25.6	125	3.24	8.57	3.56			
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	3.27	15	0.261 J	0.804 J	0.387 J			
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	83.8	2020	7.01	21.6	8.17			
Total Tetrachlorodibenzo-p-dioxin (TCDD)	10 J	54.9 J	3.34 J	3.17 J	3.14 J			
Total Pentachlorodibenzo-p-dioxin (PeCDD)	14.7 J	36.1 J	2.25 J	3.53 J	0.743 UJ			
Total Hexachlorodibenzo-p-dioxin (HxCDD)	51.3	110 J	6.48	14.4 J	7.28 J			
Total Heptachlorodibenzo-p-dioxin (HpCDD)	603	1010 J	27.4 J	67.7	32.6			
			13.6 J	20.1 J	5.93 J			
	72.8 J	I 358 J	I T2.0 1	20.1 J				
Total Tetrachlorodibenzofuran (TCDF)	72.8 J 58.5 J	358 J 150 J						
	72.8 J 58.5 J 51.4 J	150 J 204 J	9.53 J 10.6 J	19.4 J 15.4 J	3.86 J 5.29 J			

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:	PB076.1-1SS010-20110802-N	PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
4,4'-DDD (p,p'-DDD)				
4,4'-DDE (p,p'-DDE)				
4,4'-DDT (p,p'-DDT)				
Sum 2,4 DDT, DDE, DDD (U=1/2)				
Sum 4,4 DDT, DDE, DDD (U = 1/2)			1	
Sum DDD (U=1/2)				
Sum DDE (U=1/2)				
Sum DDT (U=1/2)			-	-
Total DDx (U = 1/2)			-	-
Aldrin				-
alpha-Chlordane (cis-Chlordane)				
Endosulfan-alpha (I)				
Endosulfan-beta (II)				
Endrin				
gamma-BHC (Lindane)				
gamma-Chlordane				
Heptachlor epoxide				
Methoxychlor				
Dioxin Furans (ng/kg)				
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)				
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)				
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)				
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)				
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)				
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)				
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)				
2,3,7,8-Tetrachlorodibenzofuran (TCDF)				
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)				
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)				
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)				
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)				
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)				
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)				
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)				
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)				
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)				
Total Tetrachlorodibenzo-p-dioxin (TCDD)				
Total Pentachlorodibenzo-p-dioxin (PeCDD)				
Total Hexachlorodibenzo-p-dioxin (HxCDD)				
Total Heptachlorodibenzo-p-dioxin (HyCDD)				
Total Tetrachlorodibenzofuran (TCDF)				
Total Heyarblaradibenzafuran (PeCDF)				
Total Hexachlorodibenzofuran (HxCDF)			-	
Total Heptachlorodibenzofuran (HpCDF)				

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB001.1	PB001.2	PB001.3	PB002	PB004	PB005	PB007.1	PB007.2	PB009
Sample ID:	PB001.1-1SS010-091027-N	PB001.2-1SS010-091027-N	PB001.3-1SS010-091027-N	PB002-1SS010-091027-N	PB004-1SS010-091027-N	PB005-1SS010-091027-N	PB007.1-1SS010-091028-N	PB007.2-1SS010-091028-N	PB009-1SS010-091028-N
Sample Date:	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/27/2009	10/28/2009	10/28/2009	10/28/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	N
X:	3202382.3842	3202624.0211	3202754.9026	3202606.9920	3202330.4788	3202262.1385	3202065.5011	3202037.3755	3201747.2539
Y:	13836546.3402	13836566.3557	13836489.4987	13836435.6794	13836348.7826	13836144.2614	13836300.0036	13836110.2965	13836119.3947
Total Dioxin/Furan (U = 1/2)	53295.91 J	75497.26 J	264541.87 J	160984.03 J	57253.06 J	89496.52 J	61052.9 J	67274.22 J	70477.93 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	227.092 J	262.947 J	255.1655 J	504.451 J	412.057 J	304.488 J	260.679 J	266.6735 J	218.9645 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	100.662 J	134.692 J	194.455 J	254.148 J	214.881 J	151.924 J	125.313 J	133.2792 J	108.6892 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	109.383 J	141.834 J	240.506 J	272.385 J	201.539 J	160.329 J	133.129 J	136.595 J	115.969 J

Appendix A-1
BERA Sediment Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB011	PB013.1	PB013.2	PB015	PB016	PB016	PB018	PB019	PB022
Sample ID:	PB011-1SS010-091028-N	PB013.1-1SS010-091028-N	PB013.2-1SS010-091028-N	PB015-1SS010-091028-N	PB016-1SS010-091101-N	PB016-1SS010-091102-N	PB018-1SS010-091101-N	PB019-1SS010-091101-N	PB022-1SS010-091101-D
Sample Date:	10/28/2009	10/28/2009	10/28/2009	10/28/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	FD
X:	3201727.2504	3201333.4751	3201420.5786	3201223.8464	3201140.0114	3201140.0114	3201146.8550	3201281.6017	3201174.1294
Y:	13836239.7593	13836187.7023	13836111.1432	13836229.0694	13836081.5392	13836081.5392	13835702.1476	13835711.9947	13835456.7389
Total Dioxin/Furan (U = 1/2)	73444.71 J	40996.28 J	30618.93 J	70842.42 J	41266.825 J		104489.755 J	64543.81 J	22499.625 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	373.3585 J	320.1335 J	93.9611 J	346.715 J	230.3425 J		251.09685 J	192.7715 J	163.25475 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	181.115 J	149.0829 J	51.3994 J	153.518 J	109.7999 J		131.0964 J	90.2692 J	73.2659 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	180.986 J	140.308 J	54.694 J	162.212 J	103.825 J		140.2885 J	100.573 J	71.7385 J

Task:	RI/FS 2009								
Location ID:	PB022	PB023	PB023	PB024	PB026	PB028	PB032	PB034	PB037
Sample ID:	PB022-1SS010-091101-N	PB023-1SS010-091030-D	PB023-1SS010-091030-N	PB024-1SS010-091030-N	PB026-1SS010-091030-N	PB028-1SS010-091030-N	PB032-1SS010-091030-N	PB034-1SS010-091030-N	PB037-1SS010-091030-N
Sample Date:	11/1/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009
Depth:	0 - 10 cm								
Sample Type:	N	FD	N	N	N	N	N	N	N
X:	3201174.1294	3201315.1973	3201315.1973	3201180.8567	3201331.5617	3201355.6226	3201368.6522	3201387.6450	3201399.4292
Y:	13835456.7389	13835429.2411	13835429.2411	13835181.4775	13835155.0673	13834855.4834	13834541.6071	13834242.2769	13833936.8351
Total Dioxin/Furan (U = 1/2)	21189.41 J	83668.9 J	115678.75 J	38729.5 J	2599498.9 J	22055.35 J	1473096.74 J	269448.15 J	24096.28 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	150.2969 J	226.8535 J	217.1475 J	285.8815 J	4403.638 J	101.53985 J	820.4205 J	645.597 J	305.0205 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	66.163 J	126.8989 J	120.7359 J	139.768 J	1590.81 J	48.7757 J	448.7933 J	294.809 J	132.671 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	65.017 J	134.345 J	136.493 J	135.076 J	2223.629 J	50.827 J	755.637 J	355.57 J	130.15 J

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB041	PB043	PB044	PB047.1	PB047.2	PB048	PB049	PB049	PB053
Sample ID:	PB041-1SS010-091030-N	PB043-1SS010-091030-N	PB044-1SS010-091030-N	PB047.1-1SS010-091030-N	PB047.2-1SS010-091030-N	PB048-1SS010-091031-N	PB049-1SS010-091031-D	PB049-1SS010-091031-N	PB053-1SS010-091031-N
Sample Date:	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/30/2009	10/31/2009	10/31/2009	10/31/2009	10/31/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm			
Sample Type:	N	N	N	N	N	N	FD	N	N
X:	3201410.6060	3201451.0315	3201663.5416	3201454.2006	3201656.8574	3201635.5547	3201435.3433	3201435.3433	3201421.7211
Y:	13833634.6932	13833347.5606	13833285.4731	13833042.2356	13833055.7842	13832859.6311	13832751.7292	13832751.7292	13832443.5506
Total Dioxin/Furan (U = 1/2)	11848.94 J	5986.09 J	14390.68	10619.53	12996.98	15569.45	8487.66 J	8635.83	36781.13 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	349.755 J	91.038 J	770.48	178.9765	280.4795	244.4465	125.238 J	117.3175	1303.194 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	152.697 J	33.0401 J	131.453	86.693	82.389	90.532	56.296 J	52.909	667.635 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	134.171 J	34.153 J	149.257	80.913	85.505	91.762	56.445 J	53.33	609.87 J

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB056	PB057	PB059.1	PB059.2	PB063.1	PB063.2	PB064	PB066	PB069
Sample ID:	PB056-1SS010-091031-N	PB057-1SS010-091031-N	PB059.1-1SS010-091102-N	PB059.2-1SS010-091031-N	PB063.1-1SS010-091031-N	PB063.2-1SS010-091102-N	PB064-1SS010-091102-N	PB066-1SS010-091101-N	PB069-1SS010-091101-N
Sample Date:	10/31/2009	10/31/2009	11/2/2009	10/31/2009	10/31/2009	11/2/2009	11/2/2009	11/1/2009	11/1/2009
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	N	N	N	N	N	N	N	N
X:	3201402.6901	3201634.7257	3201531.6282	3201640.7239	3201497.4803	3201553.0822	3201628.7185	3201431.2755	3201160.2007
Y:	13832184.9773	13832087.9098	13831850.4179	13831857.0149	13831523.9045	13831552.2764	13831275.6688	13831275.6811	13830971.1845
Total Dioxin/Furan (U = 1/2)	8466.52	5853.57 J	10126.81 J	5710.57	4967.66 J	9868.33	4446.798 J	3677.13 J	10345.49
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	92.356	45.509 J	218.0805 J	41.5602	104.4088 J	45.1312	8.9937 J	37.5023 J	230.6607
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	45.079	25.5108 J	107.035 J	23.4752	41.5122 J	29.0452	6.9267 J	20.3788 J	92.5267
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	46.543	27.834 J	97.179 J	26.9076	39.2574 J	36.9346	10.9399 J	21.5135 J	83.7821

Task:	RI/FS 2009	2011 USC	2011 USC	2011 USC				
Location ID:	PB074	PB081	PB093	PB097	PB101	PB068	PB069.1	PB070A
Sample ID:	PB074-1SS010-091101-N	PB081-1SS010-091103-N	PB093-1SS010-091103-N	PB097-1SS010-091103-N	PB101-1SS010-091103-N	PB068-1SS010-20110804-N	PB069.1-1SS008-20110804-N	PB070A-1SS010-20110802-N
Sample Date:	11/1/2009	11/3/2009	11/3/2009	11/3/2009	11/3/2009	8/4/2011	8/4/2011	8/2/2011
Depth:	0 - 10 cm	0 - 8 cm	0 - 10 cm					
Sample Type:	N	N	N	N	N	N	N	N
X:	3200828.0183	3200761.3787	3200970.4291	3201103.1242	3201287.2658	3201209.861	3201205.859	3201087.255
Y:	13830463.2171	13830005.1879	13828894.7843	13828566.1540	13828181.2285	13831121.51	13830973.3	13830903.51
Total Dioxin/Furan (U = 1/2)	1776.913 J	7530.63 J	129.985 J	454.428 J	390.3115 J			
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	18.31338 J	131.296 J	2.733301 J	4.1361 J	1.395567 J	-		
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	8.33598 J	47.2882 J	1.419361 J	2.1635 J	0.938547 J	-		-
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	8.29314 J	48.668 J	1.489103 J	2.31306 J	1.143091 J			

Task:	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB076.1	PB081.1	PB081.1	PB082
Sample ID:	PB076.1-1SS010-20110802-N	PB081.1-1SS010-20110803-D	PB081.1-1SS010-20110803-N	PB082-1SS010-20110803-N
Sample Date:	8/2/2011	8/3/2011	8/3/2011	8/3/2011
Depth:	0 - 10 cm	0 - 10 cm	0 - 10 cm	0 - 10 cm
Sample Type:	N	FD	N	N
X:	3200744.033	3200772.123	3200772.123	3200751.881
Y:	13830426.95	13829983.6	13829983.6	13829914.32
Total Dioxin/Furan (U = 1/2)				
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	1		-	1
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	1		-	-
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	1			-

Notes:

Bold = Detected result

J = Estimated value

U = Compound analyzed, but not detected above detection limit

UJ = Compound analyzed, but not detected above estimated detection limit

R = Rejected

-- Results not reported or not applicable

μg/kg = micrograms per kilogram

mg/kg = milligrams per kilogram

ng/kg = nanograms per kilogram

μmol/g = micromoles per gram pct = percent

N = Normal Field Sample FD = Field Duplicate

TEQ values as of 2005, World Health Organization

APPENDIX A-2 BERA SURFACE WATER DATASET

Appendix A-2
BERA Surface Water Dataset

	Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
	Location ID:	PB006	PB006	PB006	PB006	PB006	PB006	PB031	PB031
		PB006-1SWMID-091104-D				PB006-1SWNBT-091104-N		PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
	ample Date:	11/4/2009	11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
	ample Type:	FD FD	N	FD	N	N	N	N	N
-	X:	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
	Y:	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
Conventional Parameters (mg/L)									
Alkalinity, as Calcium carbonate (CaCO3)		91	94	88	86	110	96	200	130
Chloride (total)		1400	1500	1800	1800	1600	1900	1700	1900
Cyanide		0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.0025 J	0.005 U	0.003 J
Fluoride		0.29	0.3	0.32	0.31	0.32	0.31	0.46	0.33
Nitrate as Nitrogen		1.1	1.2	1.1	1	1.4	1.2	4.1	1.9
Phosphorus		0.431	0.416	0.39	0.371	0.482	0.523	0.863	0.531
Sulfate		260	280	300	260	330	290	520	370
Total Kjeldahl nitrogen		1.4	1.6	1.1	1	1.4	1.5	2	1.2
Conventional Parameters (pct)									
Dissolved organic carbon		0.00069	0.00074	0.00061	0.0006	0.00074	0.00066	0.0011	0.00079
Total dissolved solids		0.27	0.27	0.31	0.32	0.28	0.34	0.35	0.36
Total organic carbon		0.00065	0.00064	0.00058	0.00056	0.00069	0.00058	0.00096	0.00074
Total Suspended Solids		0.0025	0.003	0.0024	0.0022	0.0053	0.0024	0.0028	0.002
Metals (dissolved) (μg/L)									
Mercury		0.01 U	0.01 U	0.01 U	0.01 U	0.0051 J	0.01 U	0.01 U	0.01 U
Selenium		11	11	14	14	11	15	3.8 J	12
Metals (μg/L)									
Mercury		0.155 J	0.153 J	0.12 J	0.128 J	0.24 J	0.149 J	0.2 J	0.181 J
Selenium		14	16	16	15	11	12	4.5 J	10
Volatile Organics (μg/L)									
1,1-Dichloroethane		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1-Dichloroethene		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1,1-Trichloroethane		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1,2-Trichloroethane		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1,2-Trichlorotrifluoroethane		2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
1,1,2,2-Tetrachloroethane		2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
1,2-Dibromo-3-chloropropane		5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
1,2-Dibromoethane (Ethylene dibromide)		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,2-Dichlorobenzene		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,2-Dichloroethane		3.26	3.59	2.49	1.97	3.03	2.56	2.75	3.54
1,2-Dichloroethene, cis-		1.58	1.55	1.02	0.72 J	1.36	1.03	1.54	1.82
1,2-Dichloroethene, trans- 1,2-Dichloropropane		1 U 1 U	1 U 1 U	1 U 1 U	1 U 1 U	1 U 1 U	1 U 1 U	1 U 1 U	1 U 1 U
		5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
1,2,3-Trichlorobenzene 1,2,4-Trichlorobenzene		5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
1,2,4,5-Tetrachlorobenzene		0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
1,3-Dichlorobenzene		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,3-Dichloropenzene 1,3-Dichloropropene, cis-		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,3-Dichloropropene, trans-		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,4-Dichlorobenzene		1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
TIT DICHIOLONGHEETIC		10	5 U	5 U	5 U	5 U	10	5 U	5 U

Appendix A-2
BERA Surface Water Dataset

	Task: RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Locatio		PB031	PB059	PB059	PB076	PB076	PB076	PB076
Samp			PB059-1SWMID-091104-N		PB076-1SWMID-091104-N		PB076-1SWNBT-091104-N	
Sample I		11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample 1		N N	N N	N N	N N	N	N N	N N
•	X: 3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
	Y: 13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
Conventional Parameters (mg/L)								
Alkalinity, as Calcium carbonate (CaCO3)	170	130	220	210	260	220	260	230
Chloride (total)	2000	2100	1700	1800	2400	2200	2400	2300
Cyanide	0.005 U	0.005 U	0.005 U	0.005 U	0.0038 J	0.005 U	0.005 U	0.005 U
Fluoride	0.41	0.37	0.56	0.5	0.35	0.47	0.34	0.43
Nitrate as Nitrogen	3.4	1.8	6.9	6.8	7.7	8.6	7.6	8.5
Phosphorus	0.835	0.548	1.54	1.44	1.57	1.88	1.57	1.95
Sulfate	510	390	840	660	280	600	280	630
Total Kjeldahl nitrogen	1.5	1.2	1.8	1.5	1.7	1.6	1.6	1.6
Conventional Parameters (pct)								
Dissolved organic carbon	0.0012	0.00078	0.0014	0.0013	0.0013	0.0013	0.0014	0.0014
Total dissolved solids	0.38	0.39	0.4	0.38	0.42	0.43	0.44	0.44
Total organic carbon	0.00097	0.00073	0.0012	0.0013	0.0013	0.0014	0.0013	0.0013
Total Suspended Solids	0.0029	0.0027	0.0037	0.0012	0.0011	0.00077	0.001	0.0012
Metals (dissolved) (μg/L)								
Mercury	0.01 U	0.01 U	0.01 U	0.01 U	0.01 U	0.01 U	0.01 U	0.01 U
Selenium	8	14	5 U	5 U	5 U	5 U	5 U	5 U
Metals (μg/L)								
Mercury	0.209 J	0.215 J	0.097 J	0.025 J	0.011 J	0.01 J	0.046 J	0.012 J
Selenium	8	11	5 U	5 U	5 U	5 U	5 U	5 U
Volatile Organics (μg/L)								
1,1-Dichloroethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1-Dichloroethene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1,1-Trichloroethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1,2-Trichloroethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,1,2-Trichlorotrifluoroethane	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
1,1,2,2-Tetrachloroethane	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
1,2-Dibromo-3-chloropropane	5 U	5 U 1 U	5 U	5 U 1 U	5 U 1 U	5 U	5 U	5 U
1,2-Dibromoethane (Ethylene dibromide) 1,2-Dichlorobenzene	1 U 1 U	1 U	1 U 1 U	1 U	1 U	1 U 1 U	1 U 1 U	1 U 1 U
1,2-Dichloroethane	3.72	3.9	1 U	1 U	1 U	1 U	1 U	1 U
1,2-Dichloroethane 1,2-Dichloroethene, cis-	1.91	2.27	1 U	1 U	1 U	1 U	1 U	1 U
1,2-Dichloroethene, trans-	1.91 1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,2-Dichloropropane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,2,3-Trichlorobenzene	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
1,2,4-Trichlorobenzene	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
1,2,4,5-Tetrachlorobenzene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
1,3-Dichlorobenzene	1 U	1 U	1 U	0.837 J	1 U	1.11	1 U	0.989 J
1,3-Dichloropropene, cis-	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,3-Dichloropropene, trans-	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
1,4-Dichlorobenzene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
2-Butanone (MEK)	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U

Appendix A-2 BERA Surface Water Dataset

			I		
Task		2011 USC	2011 USC	2011 USC	2011 USC
Location ID	•	PB080	PB080	PB082.1	PB101C
Sample ID		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date		8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type		FD	N	N	N
X	•	3200769.755	3200769.755	3200777.403	3201307.665
Υ	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
Conventional Parameters (mg/L)					
Alkalinity, as Calcium carbonate (CaCO3)					
Chloride (total)					
Cyanide					
Fluoride					
Nitrate as Nitrogen					
Phosphorus					
Sulfate					
Total Kjeldahl nitrogen					
Conventional Parameters (pct)					
Dissolved organic carbon					
Total dissolved solids					
Total organic carbon	0.0016	0.0016	0.0015	0.0017	0.002
Total Suspended Solids	19	26	22	16	16
Metals (dissolved) (μg/L)	ļ				
Mercury					
Selenium					
Metals (μg/L)	_				
Mercury				-	
Selenium				-	
Volatile Organics (μg/L)					
1,1-Dichloroethane				-	
1,1-Dichloroethene					
1,1,1-Trichloroethane					
1,1,2-Trichloroethane					
1,1,2-Trichlorotrifluoroethane				-	
1,1,2,2-Tetrachloroethane					
1,2-Dibromo-3-chloropropane					
1,2-Dibromoethane (Ethylene dibromide)					
1,2-Dichlorobenzene					
1,2-Dichloroethane					
1,2-Dichloroethene, cis-					
1,2-Dichloroethene, trans-					
1,2-Dichloropropane					
1,2,3-Trichlorobenzene					
1,2,4-Trichlorobenzene					
1,2,4,5-Tetrachlorobenzene					
1,3-Dichlorobenzene					
1,3-Dichloropropene, cis-					
1,3-Dichloropropene, trans-					
1,4-Dichlorobenzene					
2-Butanone (MEK)					

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	DI/EC 2000
Location ID:	PB006	PB006	PB006	PB006	PB006	PB006	PB031	RI/FS 2009 PB031
	PB006-1SWMID-091104-D				PB006-1SWNBT-091104-N		PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
Sample Date:	11/4/2009	11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Date:	FD	11/4/2003 N	FD	11/3/2003 N	11/4/2003 N	11/3/2003 N	11/4/2003 N	11/3/2003 N
X:	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
γ. γ:	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
2-Hexanone (Methyl butyl ketone)	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
Acetone	5 U	5.28	5 U	5 U	4.58 J	5 U	16.2	6.13
Benzene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromochloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromodichloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromoform	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromomethane	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
Carbon disulfide	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Carbon tetrachloride (Tetrachloromethane)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chlorobenzene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chloroethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chloroform	0.934 J	0.921 J	1	0.695 J	0.822 J	0.863 J	0.622 J	0.803 J
Chloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Cyclohexane	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Dibromochloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Dichlorodifluoromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Dichloromethane (Methylene chloride)	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Ethylbenzene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Isopropylbenzene (Cumene)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Methyl acetate	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Methyl isobutyl ketone (4-Methyl-2-pentanone or (MIBK))	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
Methyl tert-butyl ether (MTBE)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Methylcyclohexane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
m,p-Xylene	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
o-Xylene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Styrene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Tetrachloroethene (PCE)	1.06	1.18	0.704 J	0.596 J	1.2	1.04	1.01	1.72
Total Yulong (H = 1/2)	1 U	1 U 2 U	1 U	1 U	1 U	1 U	1 U 2 U	1 U
Total Xylene (U = 1/2) Trichloroethene (TCE)	2 U 0.739 J	0.799 J	2 U 1 U	2 U 0.529 J	2 ∪ 0.94 J	2 U 0.762 J	0.792 J	2 U 1.24
Trichlorofluoromethane (Fluorotrichloromethane)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1.24 1 U
Vinyl chloride	1 U	0.598 J	1 U	1 U	0.56 J	1 U	0.707 J	0.705 J
Aromatic Hydrocarbons (µg/L)	10	0.3363	10	10	0.30 1	10	0.7073	0.7033
Total 10 of 17 HPAH (U = 1/2)	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Total 17 PAH (U = 1/2)	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Total 7 of 17 LPAH (U = 1/2)	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Total cPAH TEF (7 minimum) (U = 1/2)	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
2-Methylnaphthalene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Acenaphthene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Acenaphthylene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Anthracene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Benzo(a)anthracene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U

Appendix A-2
BERA Surface Water Dataset

Tools	DI/EC 2000	DI/EC 2000	DI/EC 2000	DI/EC 2000	DI/EC 2000	DI/FC 2000	DI/EC 2000	DI/EC 2000
Task:	RI/FS 2009 PB031	RI/FS 2009 PB031	RI/FS 2009 PB059	RI/FS 2009 PB059	RI/FS 2009 PB076	RI/FS 2009 PB076	RI/FS 2009 PB076	RI/FS 2009 PB076
Location ID:			1				PB076 PB076-1SWNBT-091104-N	
· ·				PB059-1SWMID-091105-N				
Sample Date:	11/4/2009 N	11/5/2009 N	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009 N	11/5/2009 N
Sample Type:	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
X: v.	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
2-Hexanone (Methyl butyl ketone)	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
	17.7	9.73	18.7	18.8	16.9	16.2	17.7	18.8
Acetone	1 U	1 U	10.7 1 U	1 U	10.9 1 U	10.2 1 U	17.7 1 U	1 U
Bromochloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromodichloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromoform	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Bromomethane	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
Carbon disulfide	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Carbon distince Carbon tetrachloride (Tetrachloromethane)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chlorobenzene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chloroethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chloroform	0.704 J	0.964 J	1 U	1 U	1 U	1 U	1 U	1 U
Chloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Cyclohexane	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Dibromochloromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Dichlorodifluoromethane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Dichloromethane (Methylene chloride)	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Ethylbenzene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Isopropylbenzene (Cumene)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Methyl acetate	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Methyl isobutyl ketone (4-Methyl-2-pentanone or (MIBK))	5 U	5 U	5 U	5 U	5 U	5 U	5 U	5 U
Methyl tert-butyl ether (MTBE)	1 U	1 U	1 U	1 U	0.579 J	1 U	0.68 J	0.623 J
Methylcyclohexane	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
m,p-Xylene	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
o-Xylene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Styrene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Tetrachloroethene (PCE)	1.09	1.8	1 U	1 U	1 U	1 U	1 U	1 U
Toluene	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Total Xylene (U = 1/2)	2 U	2 U	2 U	2 U	2 U	2 U	2 U	2 U
Trichloroethene (TCE)	0.862 J	1.4	1 U	1 U	1 U	1 U	1 U	1 U
Trichlorofluoromethane (Fluorotrichloromethane)	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Vinyl chloride	0.904 J	0.94 J	1 U	1 U	1 U	1 U	1 U	1 U
Aromatic Hydrocarbons (μg/L)								
Total 10 of 17 HPAH (U = 1/2)	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Total 17 PAH (U = 1/2)	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Total 7 of 17 LPAH (U = 1/2)	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Total cPAH TEF (7 minimum) (U = 1/2)	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
2-Methylnaphthalene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Acenaphthene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Acenaphthylene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Anthracene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Benzo(a)anthracene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U

Appendix A-2 BERA Surface Water Dataset

Task:		2011 USC	2011 USC	2011 USC	2011 USC
Location ID:		PB080	PB080	PB082.1	PB101C
<u> </u>	PB066-1SWMID-20110808-N	PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N		PB101C-1SWMID-20110808-N
Sample Date:		8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type:		FD	N	N	N
X:		3200769.755	3200769.755	3200777.403	3201307.665
Y:		13830051.11	13830051.11	13829942.54	13828208.48
2-Hexanone (Methyl butyl ketone)					
Acetone					
Benzene					
Bromochloromethane					
Bromodichloromethane					
Bromoform					
Bromomethane		-			
Carbon disulfide		-			
Carbon tetrachloride (Tetrachloromethane)					
Chlorobenzene					
Chloroethane					
Chloroform					
Chloromethane					
Cyclohexane					
Dibromochloromethane					
Dichlorodifluoromethane					
Dichloromethane (Methylene chloride)					
Ethylbenzene					
Isopropylbenzene (Cumene)					
Methyl acetate					
Methyl isobutyl ketone (4-Methyl-2-pentanone or (MIBK))					
Methyl tert-butyl ether (MTBE)					
Methylcyclohexane					
m,p-Xylene					
o-Xylene					
Styrene					
Tetrachloroethene (PCE)					
Toluene					
Total Xylene (U = 1/2)					
Trichloroethene (TCE)					
Trichlorofluoromethane (Fluorotrichloromethane)					
Vinyl chloride					
Aromatic Hydrocarbons (μg/L)					
Total 10 of 17 HPAH (U = 1/2)					
Total 17 PAH (U = 1/2)					
Total 7 of 17 LPAH (U = 1/2)					
Total cPAH TEF (7 minimum) (U = 1/2)					
2-Methylnaphthalene					
Acenaphthene					
Acenaphthylene					
Anthracene					
Benzo(a)anthracene					

Appendix A-2
BERA Surface Water Dataset

Tools	B1/EC 2000	DI/EC 2000	DI/EC 2000	BI /EC 2000	DI/EC 2000	BI /EC 2000	DI/EC 2000	DI/EC 2000
Task:	1	RI/FS 2009 PB006	RI/FS 2009 PB006	RI/FS 2009 PB006	RI/FS 2009 PB006	RI/FS 2009 PB006	RI/FS 2009	RI/FS 2009 PB031
Location ID:	PB006-1SWMID-091104-D						PB031	
-					PB006-1SWNBT-091104-N		PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
Sample Date:	11/4/2009 FD	11/4/2009 N	11/5/2009 FD	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:		3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
X: Y:		13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
•••								
Benzo(a)pyrene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Benzo(b)fluoranthene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Benzo(g,h,i)perylene Benzo(k)fluoranthene	0.556 U 0.556 U	0.568 U 0.568 U	0.5 U 0.5 U	0.595 U 0.595 U	0.556 U 0.556 U	0.625 U 0.625 U	0.543 U 0.543 U	0.446 U 0.446 U
Chrysene Dibenzo(a,h)anthracene	0.556 U 0.556 U	0.568 U 0.568 U	0.5 U 0.5 U	0.595 U 0.595 U	0.556 U 0.556 U	0.625 U 0.625 U	0.543 U 0.543 U	0.446 U 0.446 U
Fluoranthene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Fluorene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Indeno(1,2,3-c,d)pyrene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Naphthalene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Phenanthrene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Pyrene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Semivolatile Organics (µg/L)	0.550 0	0.500 0	0.5 0	0.555 0	0.550 0	0.023 0	0.543 0	0.440 0
2,2'-Oxybis (1-chloropropane)	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
2,3,4,6-Tetrachlorophenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2,4,5-Trichlorophenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2,4,6-Trichlorophenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2,4-Dichlorophenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2,4-Dimethylphenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2,4-Dinitrophenol	2.22 UJ	2.27 UJ	2 UJ	2.38 UJ	2.22 UJ	2.5 UJ	2.17 UJ	1.78 UJ
2,4-Dinitrotoluene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
2,6-Dinitrotoluene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
2-Chloronaphthalene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
2-Chlorophenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2-Methylphenol (o-Cresol)	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
2-Nitroaniline	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
2-Nitrophenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
3,3'-Dichlorobenzidine	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
3-Nitroaniline	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
4-Bromophenyl-phenyl ether	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
4-Chloro-3-methylphenol	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
4-Chloroaniline	0.556 UJ	0.568 UJ	0.5 U	0.595 U	0.556 UJ	0.625 U	0.543 UJ	0.446 U
4-Chlorophenyl phenyl ether	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
4-Methylphenol (p-Cresol)	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
4-Nitroaniline	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
4-Nitrophenol	1.11 UJ	1.14 UJ	1 UJ	1.19 UJ	1.11 UJ	1.25 UJ	1.09 UJ	0.893 UJ
Acetophenone	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Atrazine	0.3 J	0.284 J	0.31 J	0.31 J	0.289 J	0.325 J	0.543 U	0.268 J
Biphenyl (1,1'-Biphenyl)	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
bis(2-Chloroethoxy)methane	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
bis(2-Chloroethyl)ether	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Bis(2-ethylhexyl) phthalate	0.556 U	0.568 U	0.5 U	0.595 U	0.311 J	0.625 U	0.37 J	0.446 U

Appendix A-2
BERA Surface Water Dataset

Task	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000
Task:	·	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB031	PB031	PB059	PB059	PB076	PB076	PB076	PB076
Sample ID:					PB076-1SWMID-091104-N		PB076-1SWNBT-091104-N	PB076-1SWNBT-091105-N
Sample Date:	11/4/2009 N	11/5/2009 N	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009 N	11/5/2009 N
Sample Type:		3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
X: v·	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
Panzalalnurana	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Benzo(a)pyrene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Benzo(b)fluoranthene Benzo(g,h,i)perylene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Benzo(k)fluoranthene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Chrysene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Dibenzo(a,h)anthracene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Fluoranthene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Fluorene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Indeno(1,2,3-c,d)pyrene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Naphthalene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Phenanthrene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Pyrene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Semivolatile Organics (μg/L)								
2,2'-Oxybis (1-chloropropane)	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
2,3,4,6-Tetrachlorophenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
2,4,5-Trichlorophenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
2,4,6-Trichlorophenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
2,4-Dichlorophenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
2,4-Dimethylphenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.37 J	0.588 UJ	0.658 UJ
2,4-Dinitrophenol	2.3 UJ	2.02 UJ	2.3 UJ	2 UJ	2.17 UJ	2 UJ	2.35 UJ	2.63 UJ
2,4-Dinitrotoluene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
2,6-Dinitrotoluene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
2-Chloronaphthalene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
2-Chlorophenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
2-Methylphenol (o-Cresol)	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
2-Nitroaniline	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
2-Nitrophenol	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
3,3'-Dichlorobenzidine	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
3-Nitroaniline	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
4-Bromophenyl-phenyl ether	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
4-Chloro-3-methylphenol	0.575 UJ	0.505 UJ 0.505 U	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ 0.588 UJ	0.658 UJ
4-Chloroaniline 4-Chlorophenyl phenyl ether	0.575 UJ 0.575 U	0.505 U	0.575 UJ 0.575 U	0.5 U 0.5 U	0.543 UJ 0.543 U	0.5 U 0.5 U	0.588 U	0.658 U 0.658 U
4-Methylphenol (p-Cresol)	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
4-Nitroaniline	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
4-Nitrophenol	1.15 UJ	1.01 UJ	1.15 UJ	1 UJ	1.09 UJ	1 UJ	1.18 UJ	1.32 UJ
Acetophenone	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Atrazine	0.575 U	0.283 J	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Biphenyl (1,1'-Biphenyl)	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
bis(2-Chloroethoxy)methane	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
bis(2-Chloroethyl)ether	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Bis(2-ethylhexyl) phthalate	0.356 J	0.505 U	0.621	0.5 U	201 J	0.37 J	0.376 J	0.355 J

Appendix A-2 BERA Surface Water Dataset

Task:		2011 USC	2011 USC	2011 USC	2011 USC
Location ID:		PB080	PB080	PB082.1	PB101C
Sample ID:		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date:		8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type:		FD	N	N	N
X:		3200769.755	3200769.755	3200777.403	3201307.665
Y:	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
Benzo(a)pyrene					
Benzo(b)fluoranthene					
Benzo(g,h,i)perylene					
Benzo(k)fluoranthene					
Chrysene					
Dibenzo(a,h)anthracene					
Fluoranthene					
Fluorene					
Indeno(1,2,3-c,d)pyrene					
Naphthalene					
Phenanthrene					
Pyrene					
Semivolatile Organics (μg/L)					
2,2'-Oxybis (1-chloropropane)					
2,3,4,6-Tetrachlorophenol					
2,4,5-Trichlorophenol					
2,4,6-Trichlorophenol					
2,4-Dichlorophenol					
2,4-Dimethylphenol					
2,4-Dinitrophenol					
2,4-Dinitrotoluene					
2,6-Dinitrotoluene					
2-Chloronaphthalene					
2-Chlorophenol					
2-Methylphenol (o-Cresol)					
2-Nitroaniline					
2-Nitrophenol					
3,3'-Dichlorobenzidine					
3-Nitroaniline					
4-Bromophenyl-phenyl ether					
4-Chloro-3-methylphenol					
4-Chloroaniline					
4-Chlorophenyl phenyl ether					
4-Methylphenol (p-Cresol)					
4-Nitroaniline					
4-Nitrophenol					
Acetophenone					
Atrazine					
Biphenyl (1,1'-Biphenyl)					
bis(2-Chloroethoxy)methane					
bis(2-Chloroethyl)ether					
Bis(2-ethylhexyl) phthalate					

Appendix A-2
BERA Surface Water Dataset

7.1	DI /50 2000	D1/FC 2000	DI /50 2000	DI /50 2000	DI /50 2000	DI /50 2000	DI /50 2000	DI /50 2000
Task:	· ·	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:		PB006	PB006	PB006	PB006	PB006	PB031	PB031
•	PB006-1SWMID-091104-D						PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
Sample Date:		11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:		N 2202127 4160	FD 2202427 4450	N 2202427 4460	N 2202427.4460	N 2202427.4460	N 22042CC 020E	N 2204266 0205
X: V.	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
T:	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
Diethyl phthalate	0.556 U	0.568 U	0.37 J	0.595 U	0.556 U	0.625 U	0.543 U	0.393 J
Butylbenzyl phthalate	0.556 U	0.295 J	0.5 U	0.595 U	0.356 J	0.625 U	0.337 J	0.688 U
Dimethyl phthalate	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Caprolactam	0.467 J	0.568 UJ	0.66 J	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.598 J
Di-n-butyl phthalate	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.438 J
Carbazole	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Dinitro-o-cresol (4,6-Dinitro-2-methylphenol)	2.22 UJ	2.27 UJ	2 UJ	2.38 UJ	2.22 UJ	2.5 UJ	2.17 UJ	1.78 UJ
Di-n-octyl phthalate	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Hexachlorobenzene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Hexachlorobutadiene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Dibenzofuran	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Hexachlorocyclopentadiene	0.556 UJ	0.568 UJ	0.5 UJ	0.595 UJ	0.556 UJ	0.625 UJ	0.543 UJ	0.446 UJ
Hexachloroethane	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Isophorone	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Nitrobenzene	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
N-Nitrosodi-N-propylamine	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
N-Nitrosodiphenylamine	0.556 U	0.568 U	0.5 U	0.595 U	0.556 U	0.625 U	0.543 U	0.446 U
Pentachlorophenol	2.22 UJ	2.27 UJ	2 UJ	2.38 UJ	2.22 UJ	2.5 UJ	2.17 UJ	1.78 UJ
PCB Congeners (ng/L)								
PCB-001	0.0127 J	0.0141 J	0.0118	0.00819 J	0.0226 J	0.00902 J	0.0439	0.017
PCB-002	0.012 UJ	0.00221 J	0.00191 J	0.00935 U	0.00286 J	0.012 U	0.00222 J	0.0103 U
PCB-003	0.00807 J	0.00774 J	0.00632 J	0.00572 J	0.0148 J	0.00571 J	0.0103 J	0.00702 J
PCB-004	0.494 J	0.507 J	0.403 J	0.435 J	0.923 J	0.529 J	1.89 J	1.04 J
PCB-005	0.00496 J	0.0118 UJ	0.0106 U	0.00935 U	0.00559 J	0.012 U	0.0131 J	0.0103 U
PCB-006	0.0517 J	0.0547 J	0.0268 J	0.0255 J	0.0716 J	0.0317 J	0.195 J	0.0685 J
PCB-007	0.00773 J	0.00649 J	0.00354 J	0.00339 J	0.00901 J	0.0038 J	0.0137 J	0.0103 U
PCB-008	0.145 J	0.147 J	0.0837 J	0.0829 J	0.242 J	0.0986 J	0.635 J	0.192 J
PCB-009	0.015 J	0.0154 J	0.00672 J	0.00716 J	0.0191 J	0.01 J	0.0484 J	0.017 J
PCB-010	0.0204 J	0.0169 J	0.0164 J	0.0154 J	0.0329 J	0.0207 J	0.0651 J	0.0373 J
PCB-011 PCB-012/013	0.0273 UJ 0.0183 J	0.0353 UJ 0.0181 J	0.0259 UJ 0.0126 J	0.0196 UJ 0.0139 J	0.0341 UJ 0.0257 J	0.0385 UJ 0.014 J	0.0445 UJ 0.0359 J	0.0301 UJ 0.0156 J
	0.0183 J 0.012 U	0.0181 J		0.0139 J 0.00935 U		0.014 J	0.0339 J 0.0132 U	
PCB-014		0.0118 UJ 0.267 J	0.0106 U		0.0111 U	0.012 U		0.0103 U
PCB-015 PCB-016	0.242 J 0.92	0.267 J 0.918 J	0.204 J 0.847	0.221 J 0.911	0.393 J 2.01 J	0.239)	0.749 J 3.58	0.326 J 1.87
							3.82	
PCB-017 PCB-018/030	1.24 4.7	1.25 J 4.58 J	1.12 3.46	1.19 3.66	2.44 J 8.81 J	1.31 4.18	13.9	2.3 7.43
,	0.372							
PCB-019 PCB-020/028		0.384	0.345	0.356 2.8	0.692 J	0.431	1.29 7.95	0.791
	2.77	2.89 J	2.88		4.63 J	3.28		4.87
PCB-021/033	0.818	0.827 J	0.613	0.595	1.38 J 0.874 J	0.713	2.86	1.37
PCB-022	0.561	0.584 J	0.603	0.583		0.65	1.78	0.887
PCB-023	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.00445 J	0.0103 U
PCB-024	0.0219	0.0233 J	0.0243	0.0186 J	0.0473 J	0.0289	0.0743	0.0438

Appendix A-2
BERA Surface Water Dataset

Tada	DI /FC 2000	DI /EC 2000	DI /FC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /FC 2000
Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB031	PB031	PB059	PB059	PB076	PB076	PB076	PB076
•	PB031-1SWNBT-091104-N							
Sample Date:	11/4/2009 N	11/5/2009 N	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009 N	11/5/2009 N
Sample Type:	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
X: v.	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
Disthist white state								
Diethyl phthalate	0.575 U	0.273 J	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Butylbenzyl phthalate	0.575 U	0.525 U	0.299 J	0.5 U	0.304 J	0.5 U	0.294 J	0.658 U
Dimethyl phthalate	0.575 U 0.575 UJ	0.505 U 0.434 J	0.575 U 0.575 UJ	0.5 U	0.543 U 0.543 UJ	0.5 U	0.588 U 0.588 UJ	0.658 U 0.658 UJ
Caprolactam Di-n-butyl phthalate	0.575 U	0.434 J 0.505 U	0.575 U	0.5 UJ 0.5 U	0.543 U	0.5 UJ 0.5 U	0.588 U	0.658 U
Carbazole	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Dinitro-o-cresol (4,6-Dinitro-2-methylphenol)	2.3 UJ	2.02 UJ	2.3 UJ	2 UJ	2.17 UJ	2 UJ	2.35 UJ	2.63 UJ
Di-n-octyl phthalate	0.575 U	0.505 U	0.575 U	0.5 U	0.348 J	0.5 U	0.588 U	0.658 U
Hexachlorobenzene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Hexachlorobutadiene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Dibenzofuran	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Hexachlorocyclopentadiene	0.575 UJ	0.505 UJ	0.575 UJ	0.5 UJ	0.543 UJ	0.5 UJ	0.588 UJ	0.658 UJ
Hexachloroethane	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Isophorone	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Nitrobenzene	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
N-Nitrosodi-N-propylamine	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
N-Nitrosodiphenylamine	0.575 U	0.505 U	0.575 U	0.5 U	0.543 U	0.5 U	0.588 U	0.658 U
Pentachlorophenol	2.3 UJ	2.02 UJ	2.3 UJ	2 UJ	2.17 UJ	2 UJ	2.35 UJ	2.63 UJ
PCB Congeners (ng/L)								
PCB-001	0.064 J	0.0179 J	0.181 J	0.0666 J	0.0401	0.0366 J	0.0503 J	0.0423 J
PCB-002	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.00516 J	0.00267 J	0.00445 J	0.0122 UJ
PCB-003	0.0119	0.00728 J	0.0229 J	0.0103 J	0.0156	0.0108 J	0.0178 J	0.0118 J
PCB-004	1.84 J	1.39 J	3.41 J	1.26 J	0.434 J	0.458 J	0.581 J	0.528 J
PCB-005	0.0159 J	0.0106 UJ	0.0394 J	0.0112 J	0.013 U	0.0102 UJ	0.00943 J	0.0122 UJ
PCB-006	0.209 J	0.0771 J	0.498 J	0.195 J	0.204 J	0.156 J	0.234 J	0.155 J
PCB-007	0.0196 J	0.0106 UJ	0.0581 J	0.0157 J	0.0144 J	0.0102 UJ	0.0183 J	0.00858 J
PCB-008	0.751 J	0.209 J	2.06 J	0.608 J	0.321 J	0.265 J	0.432 J	0.289 J
PCB-009	0.0517 J	0.0208 J	0.128 J	0.038 J	0.0212 J	0.017 J	0.0263 J	0.0184 J
PCB-010	0.0614 J	0.0472 J	0.107 J	0.0418 J	0.0143 J	0.0159 J	0.0183 J	0.0176 J
PCB-011	0.0416 UJ	0.0336 UJ	0.0541 UJ	0.0487 J	0.0536 UJ	0.0417 UJ	0.0639 UJ	0.0509 J
PCB-012/013	0.0422 J	0.0178 J	0.156 J	0.071 J	0.0783 J	0.0605 J	0.0879 J	0.0682 J
PCB-014	0.0115 U	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-015	0.744 J	0.42 J	1.82 J	0.609 J	0.276 J	0.201 J	0.359 J	0.275 J
PCB-016	3.62	2.53 J	6.89 J	2.15 J	0.835	0.717 J	1.23 J	1.13 J
PCB-017	3.95	3.02 J	7.12 J	2.38 J	1.11	0.945 J	1.61 J	1.33 J
PCB-018/030	13.7	9.9 J	24 J	7.59 J	3.48	2.8 J	5.06 J	4.11 J
PCB-019	1.27	0.988 J	2.24 J	0.935 J	0.451	0.43 J	0.595	0.573 J
PCB-020/028	8.56	5.89 J	16.5 J	5.75 J	3.1	2.33 J	4.33 J	2.99 J
PCB-021/033	3.07	1.55 J	5.96 J	1.61 J	0.78	0.55 J	1.09 J	0.758 J
PCB-022	1.97	1.17 J	4.61 J	1.4 J	0.714	0.528 J	0.992 J	0.669 J
PCB-023	0.0115 U	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-024	0.0673	0.0643 J	0.115 J	0.0458 J	0.0192	0.0176 J	0.0261 J	0.0231 J

Appendix A-2
BERA Surface Water Dataset

	_				
Task:	2011 USC	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB066B	PB080	PB080	PB082.1	PB101C
Sample ID:		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date:		8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type:	N	FD	N	N	N
X:		3200769.755	3200769.755	3200777.403	3201307.665
Y:	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
Diethyl phthalate					
Butylbenzyl phthalate					
Dimethyl phthalate					
Caprolactam					
Di-n-butyl phthalate					
Carbazole					
Dinitro-o-cresol (4,6-Dinitro-2-methylphenol)					
Di-n-octyl phthalate					
Hexachlorobenzene					
Hexachlorobutadiene					
Dibenzofuran					
Hexachlorocyclopentadiene					
Hexachloroethane					
Isophorone					
Nitrobenzene					
N-Nitrosodi-N-propylamine					
N-Nitrosodiphenylamine					
Pentachlorophenol					
PCB Congeners (ng/L)					
PCB-001	0.0539	0.07	0.0547	0.0626	0.00741 U
PCB-002	0.00263 U	0.00247 U	0.00366 U	0.00308 U	0.00268 U
PCB-003	0.00183 U	0.00633 J	0.00253 U	0.00231 U	0.0046 J
PCB-004	0.992	1.14	1.17	0.788	0.0379
PCB-005	0.0156 J	0.0229 J	0.0153	0.0123	0.00311 U
PCB-006	0.209	0.242	0.243	0.199	0.0209
PCB-007	0.0162	0.0228	0.0192 J	0.0183	0.00678 J
PCB-008	0.35	0.422	0.437	0.259	0.0253
PCB-009	0.749	0.965	0.89	0.647	0.0286 U
PCB-010	0.0549	0.0686	0.0626	0.049	0.00272 U
PCB-011	0.0803	0.0911	0.0831	0.0517	0.00364 J
PCB-012/013	0.0343 U	0.0416 U	0.0544 U	0.0382 U	0.0351 U
PCB-014	0.0227 J	0.0384	0.0272	0.0279	0.0121 J
PCB-015	0.00277 U	0.00265 U	0.00425 U	0.00364 U	0.00244 U
PCB-016	1.69	1.96	2.12	1.23	0.0418
PCB-017	2.44	2.9	3.04	1.79	0.0742
PCB-018/030	6.92	8.04	8.54	5.01	0.153
PCB-019	0.663	0.73	0.769	0.493	0.0283
PCB-020/028	5.52	7.69	7.76	4.38	0.233
PCB-021/033	1.96	2.7	2.7	1.51	0.051
PCB-022	1.39	1.98	1.91	1.15	0.0591
PCB-023	0.00303 J	0.0039 J	0.00455 J	0.00298 J	0.00117 U
PCB-024	0.0658	0.0822 J	0.0682	0.0433	0.00221 J

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	· ·	PB006	PB006	PB006	PB006	PB006	PB031	PB031
	PB006-1SWMID-091104-D			PB006-1SWMID-091105-N	PB006-1SWNBT-091104-N		PB031-1SWMID-091104-N	
Sample Date:		11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:		N	FD FD	N	N	N	N	N
X:		3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
Y:		13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
PCB-025	0.144	0.151 J	0.132	0.127	0.223 J	0.145	0.407	0.205
PCB-026/029	0.554	0.553 J	0.537	0.527	0.921 J	0.615	1.59	0.995
PCB-027	0.196	0.198 J	0.196	0.208	0.384 J	0.229	0.584	0.386
PCB-031	3.35	3.45 J	3.33	3.3	5.82 J	3.9	9.94	6.44
PCB-032	1.08	1.11 J	1.07	1.14	2.16 J	1.23	3.24	2.07
PCB-034	0.0213	0.0209 J	0.0181	0.0182	0.032 J	0.0204	0.0417	0.0278
PCB-035	0.0586	0.0489 J	0.0106 U	0.0421	0.0913 J	0.0507	0.158	0.0103 U
PCB-036	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-037	0.68	0.679 J	0.625	0.651	1.28 J	0.708	1.89	1.08
PCB-038	0.012 U	0.00913 J	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-039	0.012 U	0.0118 UJ	0.0232	0.0205	0.0111 UJ	0.0254	0.0132 U	0.0355
PCB-040/071	2.46 J	2.34 J	2.55	2.63	4.44 J	2.88	6.72	0.0206 U
PCB-041	0.469 J	0.511 J	0.382	0.527	0.983 J	0.545	1.54	5.99
PCB-042	1.66 J	1.67 J	1.56	1.67	3.09 J	1.81	4.53	2.65
PCB-043	0.256 J	0.255 J	0.297	0.318	0.493 J	0.365	0.755	0.505
PCB-044/047/065	6.23 J	6.11 J	6.42	6.79	11.2 J	7.37	16.5	10.4
PCB-045	1.09 J	1.05 J	1.22	1.26	1.82 J	1.4	3.12	2
PCB-046	0.434 J	0.425 J	0.448	0.453	0.709 J	0.508	1.18	0.772
PCB-048	1.21 J	1.2 J	1.21	1.27	2.19 J	1.4	3.44	2.16
PCB-049/069	3.79 J	3.69 J	3.86	4.11	6.73 J	4.47	9.66	6.19
PCB-050/053	1.25 J	1.19 J	1.32	1.33	2.05 J	1.5	3.46	2.25
PCB-051	0.366 J	0.332 J	0.38	0.32 J	0.605 J	0.416	0.863	0.626
PCB-052	6.25 J	6.06 J	6.7	7.03	10.8 J	7.67	16.3	11.1
PCB-054	0.0213	0.0209 J	0.0197	0.0214	0.0356 J	0.0218	0.0548	0.0361
PCB-055	0.0437 J	0.0443 J	0.0106 U	0.00935 U	0.0772 J	0.012 U	0.13	0.0103 U
PCB-056	1.82 J	1.84 J	1.87	2.07	3.55 J	2.19	5.08	3.14
PCB-057	0.0231 J	0.0211 J	0.0106 U	0.0203	0.0326 J	0.0193 J	0.0495	0.0301
PCB-058 PCB-059/062/075	0.0104 J 0.451 J	0.0151 J 0.443 J	0.0106 U 0.492	0.00935 U 0.513	0.0111 UJ 0.799 J	0.012 U 0.562	0.0294 1.18	0.0103 U 0.809
PCB-059/062/075 PCB-060	0.451 J 0.905 J	0.443 J 0.914 J	0.492	1.01	1.86 J	1.1	2.81	1.74
PCB-061/070/074/076	7.44 J	7.26 J	7.61	8.3	14 J	8.92	19.4	12.6
PCB-063	0.143 J	0.14 J	0.148	0.154	0.282 J	0.169	0.403	0.253
PCB-064	2.39 J	2.35 J	2.45	2.59	4.35 J	2.81	6.29	4.04
PCB-066	3.77 J	3.74 J	3.93	4.27	7.15 J	4.5	9.85	6.62
PCB-067	0.0898 J	0.0897 J	0.0863	0.0886	0.16 J	0.0989	0.243	0.147
PCB-068	0.0182 J	0.0192 J	0.0461	0.0458	0.0409 J	0.0439	0.0323	0.0533
PCB-072	0.0401 J	0.0408 J	0.0448	0.0452	0.0688 J	0.0484	0.0707	0.0538
PCB-073	0.012 UJ	0.0118 UJ	0.0451	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0884
PCB-077	0.405 J	0.416 J	0.419	0.439	0.818 J	0.478	1.1	0.682
PCB-078	0.012 UJ	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-079	0.0468 J	0.0351 J	0.0211 J	0.0583	0.0507 J	0.012 U	0.1	0.0399
PCB-080	0.012 UJ	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U

Appendix A-2
BERA Surface Water Dataset

Task	: RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID	1	PB031	PB059	PB059	PB076	PB076	PB076	PB076
	: PB031-1SWNBT-091104-N			PB059-1SWMID-091105-N	PB076-1SWMID-091104-N			
Sample Date		11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type		11/5/2003 N	N	N	11/4/2003 N	N	N	N
x	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
Ŷ	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
PCB-025	0.435	0.243 J	0.898 J	0.351 J	0.341	0.227 J	0.425 J	0.251 J
PCB-026/029	1.65	1.15 J	2.86 J	1.07 J	0.637	0.46 J	0.423 J	0.576 J
PCB-027	0.594	0.511 J	1.09 J	0.434 J	0.189	0.18 J	0.854 J	0.264 J
PCB-031	10.5	7.5 J	18.6 J	6.12 J	3.2	2.32 J	4.37 J	2.91 J
PCB-032	3.25	2.75 J	5.8 J	2.15 J	1.12	0.953 J	1.58 J	1.38 J
PCB-034	0.0115 U	0.0314 J	0.0119 UJ	0.0248 J	0.0201	0.0138 J	0.0225 J	0.0169 J
PCB-035	0.168	0.092 J	0.264 J	0.0943 J	0.0564	0.0437 J	0.0673 J	0.0632 J
PCB-036	0.0115 U	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-037	1.96	1.34	4.25 J	1.15 J	0.544	0.37	0.78 J	0.614 J
PCB-038	0.0172	0.0106 UJ	0.0276 J	0.00705 J	0.0087 J	0.0102 UJ	0.00896 J	0.0122 UJ
PCB-039	0.0115 U	0.0498 J	0.0119 UJ	0.0366 J	0.013 U	0.0141 J	0.0125 UJ	0.0253 J
PCB-040/071	6.29	5.5 J	12.3 J	4.08 J	2.27	1.69 J	3.16 J	2.86 J
PCB-041	1.47	1.28 J	2.89 J	0.904 J	0.434	0.289 J	0.57 J	0.526 J
PCB-042	4.24	3.46 J	8.17 J	2.51 J	1.56	1.05 J	2.11 J	1.74 J
PCB-043	0.714	0.667 J	1.41 J	0.504 J	0.26	0.0102 UJ	0.356 J	0.334 J
PCB-044/047/065	15.3	13.6 J	29.7 J	10.1 J	5.91	4.28 J	8.2 J	6.79 J
PCB-045	2.94	2.37 J	5.47 J	1.95 J	0.955	0.782 J	1.34 J	1.19 J
PCB-046	1.11	0.943 J	2.19 J	0.858 J	0.423	0.345 J	0.548 J	0.538 J
PCB-048	3.27	2.89 J	6.42 J	2 J	1.05	0.745 J	1.44 J	1.25 J
PCB-049/069	9.09	8.07 J	17.2 J	5.73 J	3.59	2.59 J	4.99 J	3.85 J
PCB-050/053	3.22	2.69 J	5.87 J	2.22 J	1.23	0.915 J	1.56 J	1.33 J
PCB-051	0.816	0.847 J	1.55 J	0.663 J	0.375	0.274 J	0.428 J	0.435 J
PCB-052	15.1	14.2 J	28.3 J	10.2 J	5.7	4.39 J	7.88 J	6.57 J
PCB-054	0.0542	0.0455 J	0.0942 J	0.0376 J	0.0211 J	0.0187 J	0.0299	0.0264 J
PCB-055	0.115	0.0106 UJ	0.244 J	0.01 UJ	0.0377	0.0102 UJ	0.0506 J	0.0122 UJ
PCB-056	4.9	4.28 J	9.94 J	2.84 J	1.7	1.22 J	2.52 J	2.15 J
PCB-057	0.0474	0.0418 J	0.0961 J	0.0277 J	0.0181 J	0.0102 UJ	0.027 J	0.0122 UJ
PCB-058	0.0211	0.0237 J	0.039 J	0.01 UJ	0.00883 J	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-059/062/075	1.11	1.01 J	2.17 J	0.769 J	0.429	0.324 J	0.577 J	0.494 J
PCB-060	2.72	2.29 J	5.67 J	1.45 J	0.885	0.6 J	1.31 J	0.954 J
PCB-061/070/074/076	19.2	16.7 J	37.1 J	10.8 J	6.49	4.66 J	9.78 J	7.8 J
PCB-063	0.383	0.342 J	0.767 J	0.218 J	0.137	0.101 J	0.208 J	0.148 J
PCB-064	5.92	5.27 J	11.3 J	3.53 J	2.1	1.5 J	2.96 J	2.37 J
PCB-066	9.62	8.59 J	18.8 J	5.79 J	3.57	2.76 J	5.39 J	4.58 J
PCB-067	0.231	0.193 J	0.471 J	0.133 J	0.0844	0.0573 J	0.12 J	0.0928 J
PCB-068	0.0331	0.055 J	0.0561 J	0.0151 J	0.0172	0.0102 UJ	0.0249 J	0.0234 J
PCB-072	0.0717	0.059 J	0.12 J	0.0423 J	0.0282	0.022 J	0.0444 J	0.0334 J
PCB-073	0.0115 U	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 U	0.151 J	0.0125 UJ	0.0655 J
PCB-077	1.08	0.893	2.04 J	0.585 J	0.399	0.246 J	0.623 J	0.519
PCB-078	0.0115 U	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-079	0.0932	0.047 J	0.155 J	0.0265 J	0.0298 J	0.0157 J	0.0491 J	0.0328 J
PCB-080	0.0115 U	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ

Appendix A-2
BERA Surface Water Dataset

	2011 USC	2014 UCC	2014 LICC	2014 LICC	2014 LICC
Task		2011 USC	2011 USC	2011 USC	2011 USC
Location ID		PB080	PB080	PB082.1	PB101C
Sample ID		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date		8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type		FD	N	N	N
X		3200769.755	3200769.755	3200777.403	3201307.665
Y		13830051.11	13830051.11	13829942.54	13828208.48
PCB-025	0.253	0.355	0.338	0.224	0.0384
PCB-026/029	0.935	1.19	1.22	0.706	0.0508
PCB-027	0.383	0.44	0.466	0.285	0.0138
PCB-031	6.41	8.65	8.89	4.95	0.21
PCB-032	1.93	2.31	2.42	1.41	0.0647
PCB-034	0.0211	0.0281	0.0288	0.0174	0.00179 J
PCB-035	0.0609	0.066	0.0655	0.0391	0.00693 J
PCB-036	0.00311 U	0.0045 U	0.00604 U	0.013	0.0027 U
PCB-037	0.561	0.739	0.767	0.415	0.031
PCB-038	0.00718 J	0.00457 U	0.00613 U	0.00337 U	0.00274 U
PCB-039	0.00295 U	0.00427 U	0.00572 U	0.00315 U	0.00255 U
PCB-040/071	2.77	4.2	3.93	2.16	0.102
PCB-041	0.61	0.904	0.804	0.417	0.0209
PCB-042	1.58	2.42	2.26	1.27	0.0646
PCB-043	0.283	0.466	0.408	0.228	0.0108
PCB-044/047/065	6.76	10.4	9.42	5.75	0.701
PCB-045	1.71	2.41	2.18	1.5	0.421
PCB-046	0.236	0.363	0.3565	0.183	0.01395
PCB-048	1.31	2.05	1.85	1.07	0.0332
PCB-049/069	3.95	6.17	5.86	3.24	0.183
PCB-050/053	1.34	1.85	1.75	0.936	0.0435
PCB-051	0.236	0.363	0.3565	0.183	0.01395
PCB-052	7.92	12.2	11.4	6.43	0.341
PCB-054	0.0226	0.0352	0.0287	0.0188 J	0.0016 J
PCB-055	0.0648	0.118	0.102	0.0563 J	0.00441 J
PCB-056	2.18	3.49	3.33	1.79	0.088
PCB-057	0.0176 J	0.0297	0.0274 J	0.0155	0.00208 U
PCB-058	0.0111 J	0.024	0.0248	0.00855 J	0.0021 U
PCB-059/062/075	0.486	0.749	0.652	0.408	0.023 J
PCB-060	1.1	1.8	1.7	0.934	0.0482
PCB-061/070/074/076	8.13	12.7	12.2	6.59	0.321
PCB-063	0.184	0.3	0.274	0.149	0.00906 J
PCB-064	2.95	4.66	4.3	2.37	0.12
PCB-066	4.57	7.22	6.97	3.78	0.195
PCB-067	0.102	0.162	0.149	0.0807	0.00622 J
PCB-068	0.319	0.399	0.363	0.473	0.418
PCB-072	0.0275	0.0445	0.0431 J	0.0242 J	0.002 U
PCB-073	0.071	0.000266 U	0.00088 U	0.00226 U	0.00271 J
PCB-077	0.306	0.447	0.461	0.231	0.018
PCB-078	0.00298 U	0.00602 U	0.00491 U	0.00471 U	0.00217 U
PCB-079	0.00251 U	0.0493 J	0.046 J	0.0211	0.00183 U
PCB-080	0.00269 U	0.00542 U	0.00443 U	0.00424 U	0.00195 U

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB006	PB006	PB006	PB006	PB006	PB006	PB031	PB031
	PB006-1SWMID-091104-D				PB006-1SWNBT-091104-N		PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
Sample Date:	11/4/2009	11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	FD	N	FD	N	N	N	N	N
X:		3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
Y:		13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
PCB-081	0.0159 J	0.0196 J	0.0106 U	0.0172	0.0343 J	0.012 U	0.0454	0.0319
PCB-082	0.432	0.468 J	0.481	0.508	0.907 J	0.552	1.2	0.77
PCB-083	0.216	0.167 J	0.23	0.247	0.337 J	0.283	0.465	0.33
PCB-084	1.01	1.02 J	1.03	1.12	1.89 J	1.22	2.49	1.66
PCB-085/116	0.575	0.615 J	0.683	0.719	1.19 J	0.802	1.56	1.02
PCB-086/087/097/108/119/125	2.03	2.14 J	2.19	2.32	4.14 J	2.56	5.2	3.47
PCB-088	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.947
PCB-089	0.0911	0.0915 J	0.101	0.112	0.181 J	0.114	0.254	0.167
PCB-090/101/113	2.73	2.81 J	2.88	3.09	5.2 J	3.38	6.32	4.49
PCB-091	0.538	0.546 J	0.585	0.645	0.917 J	0.696	1.16	0.0103 U
PCB-092	0.477	0.493 J	0.522	0.554	0.94 J	0.596	1.13	0.797
PCB-093/100	0.0607	0.0584 J	0.0749	0.0821	0.112 J	0.0898	0.132	0.119
PCB-094	0.0273	0.0282 J	0.0293	0.03	0.0499 J	0.0334	0.0648	0.0455
PCB-095	2.44	2.32 J	2.49	2.67	4.22 J	2.95	5.36	3.81
PCB-096	0.0622	0.0663 J	0.0641	0.0689	0.123 J	0.0764	0.171	0.111
PCB-098	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0257	0.0103 U
PCB-099	1.26	1.37 J	1.39	1.44	2.6 J	1.61	3.06	2.05
PCB-102	0.172	0.182 J	0.203	0.216	0.336 J	0.237	0.406	0.318
PCB-103	0.0344	0.0333 J	0.0385	0.0365	0.0544 J	0.0393	0.0593	0.0441 J
PCB-104	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 U	0.012 U	0.00429 J	0.0103 U
PCB-105	1.16	1.26 J	1.25	1.37	2.5	1.46	3.18	2.12
PCB-106	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-107/124	0.0945	0.0979 J	0.0983	0.105	0.195 J	0.116	0.244	0.159
PCB-109 PCB-110	0.16 2.59	0.177 J 2.78 J	0.175 2.75	0.183 2.91	0.324 J 5.18 J	0.212 3.23	0.403	0.269 4.3
PCB-110 PCB-111	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	6.13 0.0132 U	0.0103 U
PCB-112	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-114	0.077	0.0812 J	0.076	0.0852	0.161	0.096	0.205	0.134
PCB-115	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0972	0.0103 U
PCB-117	0.101	0.0893 J	0.0106 U	0.00935 U	0.188 J	0.012 U	0.212	0.125
PCB-118	2.11	2.3 J	2.33	2.51	4.49 J	2.68	5.42	3.67
PCB-120	0.00876 J	0.0105 J	0.00942 J	0.0101	0.0111 UJ	0.0128	0.0119 J	0.0103 U
PCB-121	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-122	0.0514	0.0601 J	0.068	0.0706	0.11 J	0.0746	0.142	0.109
PCB-123	0.0633	0.0657 J	0.0675	0.0771	0.137	0.0785	0.161	0.106
PCB-126	0.0123 J	0.0127 J	0.0133 J	0.00935 U	0.0248 J	0.0128	0.0303	0.02
PCB-127	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-128/166	0.219 J	0.241 J	0.232 J	0.25 J	0.416 J	0.256	0.472	0.339
PCB-129/138/163	1.21 J	1.33 J	1.37 J	1.4 J	2.44 J	1.55	2.54	1.96
PCB-130	0.111 J	0.119 J	0.126 J	0.133 J	0.219 J	0.152	0.214	0.168
PCB-131	0.0151 J	0.0213 J	0.0205 J	0.0245 J	0.0386 J	0.0261	0.0412	0.0103 U
PCB-132	0.413 J	0.454 J	0.461 J	0.484 J	0.811 J	0.532	0.857	0.642

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
	PB031	-	PB059	-		· ·	PB076	· ·
Location ID:		PB031		PB059	PB076	PB076		PB076
-	PB031-1SWNBT-091104-N			PB059-1SWMID-091105-N	PB076-1SWMID-091104-N		PB076-1SWNBT-091104-N	
Sample Date:	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	N 2204266 0205	N 2204266 0205	N	N 2204545 5205	N 2222240 7772	N 2200040 7772	N	N
X:	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
Y:	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
PCB-081	0.0502	0.032 J	0.101 J	0.0261 J	0.0194	0.0102 U	0.0323 J	0.0248 J
PCB-082	1.19	1.05	2.48 J	0.653 J	0.445	0.296 J	0.683 J	0.612 J
PCB-083	0.537	0.417	1.02 J	0.256 J	0.185	0.129 J	0.274 J	0.209 J
PCB-084	2.5	2.26	4.92 J	1.46 J	0.907	0.689 J	1.37 J	1.15 J
PCB-085/116	1.55	1.45	3.03 J	0.871 J	0.624	0.0204 UJ	0.976 J	0.0244 UJ
PCB-086/087/097/108/119/125	5.24	4.66	10.3 J	2.77 J	2	1.34 J	3.09 J	2.34 J
PCB-088	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.362 J	0.0125 UJ	0.599 J
PCB-089	0.25	0.225	0.5 J	0.138 J	0.0876	0.0694 J	0.126 J	0.119 J
PCB-090/101/113	6.3	6	12 J	3.38 J	2.38	1.67 J	3.69 J	2.77 J
PCB-091	1.18	1.16	2.23 J	0.69 J	0.408	0.0102 UJ	0.605 J	0.0122 UJ
PCB-092	1.14	1.06	2.17 J	0.606 J	0.446	0.314 J	0.679 J	0.501 J
PCB-093/100	0.138	0.138	0.266 J	0.0887 J	0.0572	0.0448 J	0.0783 J	0.0244 UJ
PCB-094	0.0688	0.063	0.129 J	0.0406 J	0.0248	0.0168 J	0.0336 J	0.0332 J
PCB-095	5.26	5.18	9.9 J	2.98 J	1.82	1.4 J	2.63 J	2.26 J
PCB-096	0.165	0.137	0.318 J	0.098 J	0.0656	0.0452 J	0.092 J	0.0766 J
PCB-098	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-099	3.07	2.82	5.95 J	1.66 J	1.24	0.851 J	1.91 J	1.4 J
PCB-102	0.426	0.447	0.813 J	0.262 J	0.147	0.119 J	0.216 J	0.213 J
PCB-103	0.0591	0.0593	0.11 J	0.0355 J	0.0251	0.0175 J	0.029 J	0.0285 J
PCB-104	0.00345 J	0.0106 U	0.00639 J	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-105	3.16	2.77	6.24 J	1.7 J	1.4	0.804 J	2.19 J	1.61
PCB-106	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-107/124	0.251	0.21	0.489 J	0.13 J	0.108	0.0616 J	0.168 J	0.134 J
PCB-109	0.405	0.361	0.797 J	0.207 J	0.179	0.104 J	0.28 J	0.196 J
PCB-110	6.4	5.67	12.1 J	3.26 J	2.45	1.63 J	3.81 J	2.89 J
PCB-111	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-112	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-114	0.207	0.185	0.418 J	0.104 J	0.0854	0.051 J	0.138 J	0.107 J
PCB-115	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-117	0.262	0.0106 U	0.482 J	0.01 UJ	0.0942	0.682 J	0.131 J	1.19 J
PCB-118	5.45	4.79	10.5 J	2.89 J	2.42	1.41 J	3.74 J	2.78 J
PCB-120	0.0115 U	0.0137 J	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.00676 J	0.0122 UJ
PCB-121	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-122	0.139	0.141	0.278 J	0.0956 J	0.0586	0.0424 J	0.0984 J	0.0861 J
PCB-123	0.165	0.136	0.311 J	0.0945 J	0.0764	0.0394 J	0.117 J	0.0958 J
PCB-126	0.0276	0.0227 J	0.0547 J	0.01 UJ	0.0168	0.00983 J	0.025 J	0.0122 UJ
PCB-127	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-128/166	0.483 J	0.413 J	0.797 J	0.21 J	0.197 J	0.12 J	0.334 J	0.225 J
PCB-129/138/163	2.49 J	2.45 J	4.14 J	1.13 J	0.996 J	0.56 J	1.56 J	1.09 J
PCB-130	0.209 J	0.211 J	0.345 J	0.101 J	0.0904 J	0.0486 J	0.138 J	0.0974 J
PCB-131	0.0426 J	0.0106 UJ	0.0844 J	0.023 J	0.0191 J	0.0137 J	0.0266 J	0.0228 J
PCB-132	0.869 J	0.84 J	1.48 J	0.38 J	0.307 J	0.194 J	0.486 J	0.387 J

Appendix A-2
BERA Surface Water Dataset

	Task:	2011 USC	2011 USC	2011 USC	2011 USC	2011 USC
	Location ID:	PB066B	PB080	PB080	PB082.1	PB101C
	Sample ID:		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	
	Sample Date:	8/8/2011	8/8/2011	8/8/2011	8/8/2011	8/8/2011
	Sample Type:	N	FD	0,0,2011 N	N	N
	X:	3201384.326	3200769.755	3200769.755	3200777.403	3201307.665
	Y:	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
PCB-081		0.013	0.0215	0.0216 J	0.0129	0.00209 U
PCB-082		0.392	0.624	0.608	0.303	0.0176
PCB-083		0.14	0.234	0.22	0.108	0.0118
PCB-084		0.829	1.28	1.23	0.654	0.0505
PCB-085/116		0.393	0.631	0.617	0.312	0.023
PCB-086/087/097/108/119/125		1.66	2.63	2.6	1.32	0.107
PCB-088		0.236	0.363	0.3565	0.183	0.01395
PCB-089		0.0801	0.123	0.119	0.0626	0.00383 J
PCB-090/101/113		1.75	2.78	2.75	1.4	0.135
PCB-091		0.236	0.363	0.3565	0.183	0.01395
PCB-092		0.325	0.536	0.526	0.266	0.0259
PCB-093/100		0.0424	0.0685	0.0687	0.0333	0.00271 J
PCB-094		0.026	0.0383	0.0385	0.0198	0.0028 U
PCB-095		1.75	2.69	2.6	1.37	0.113
PCB-096		0.0514	0.0812	0.0792	0.0433	0.00265 J
PCB-098		0.07	0.1095	0.108	0.058	0.00346 J
PCB-099		0.98	1.61	1.61	0.803	0.0661
PCB-102		0.07	0.1095	0.108	0.058	0.00346 J
PCB-103		0.019	0.0264	0.0284	0.0145	0.00239 U
PCB-104		0.00117 J	0.00116 U	0.00208 U	0.0013 U	0.000919 U
PCB-105		0.808	1.28	1.31	0.618	0.0475
PCB-106		0.00251 U	0.00495 U	0.00584 U	0.00318 U	0.00183 U
PCB-107/124		0.0633	0.1	0.109	0.051	0.00513 J
PCB-109		0.121	0.194	0.205	0.105	0.00859 J
PCB-110		1.095	1.74	1.735	0.86	0.073
PCB-111		0.00159 U	0.00284 U	0.00307 U	0.00244 U	0.00193 U
PCB-112		0.0149	0.0151	0.00327 U	0.00259 U	0.00205 U
PCB-114		0.0552	0.0816	0.0835	0.0425	0.00335 J
PCB-115		1.095	1.74	1.735	0.86	0.073
PCB-117		0.197	0.315	0.309	0.156	0.011
PCB-118		1.39	2.11	2.21	1.08	0.104
PCB-120		0.00344 J	0.00496 J	0.00629 J	0.00236 U	0.00187 U
PCB-121		0.0016 U	0.00286 U	0.0031 U	0.00246 U	0.00195 U
PCB-122		0.033	0.0531	0.0514	0.0253	0.00193 U
PCB-123		0.0394	0.0643	0.0624	0.0295	0.00586 J
PCB-126		0.00663 J	0.00976 J	0.013 J	0.00332 U	0.00196 U
PCB-127		0.00248 U	0.00489 U	0.00577 U	0.00314 U	0.00181 U
PCB-128/166		0.108	0.168	0.176	0.0854	0.0127 U
PCB-129/138/163		0.593	0.94	0.952	0.476	0.0841 U
PCB-130		0.0413	0.0696	0.0662	0.0334	0.00578 U
PCB-131		0.0113	0.0166	0.0198	0.00801 J	0.00174 U
PCB-132		0.211	0.328	0.338	0.164	0.0289 U

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB006	PB006	PB006	PB006	PB006	PB006	PB031	PB031
-		PB006-1SWMID-091104-N				PB006-1SWNBT-091105-N		PB031-1SWMID-091105-N
Sample Date:	11/4/2009	11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:		N	FD	N	N	N	N	N
X:	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
Y:	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
PCB-133	0.0232 J	0.0234 J	0.0237 J	0.0273 J	0.0435 J	0.0298	0.0327	0.0329
PCB-134	0.0756 J	0.092 J	0.0988 J	0.107 J	0.164 J	0.111	0.172	0.139
PCB-135/151	0.384 J	0.412 J	0.438 J	0.449 J	0.714 J	0.489	0.703	0.582
PCB-136	0.17 J	0.192 J	0.191 J	0.198 J	0.345 J	0.208	0.34	0.254
PCB-137	0.0562 J	0.0634 J	0.0554 J	0.0531 J	0.12 J	0.0764	0.127	0.0878
PCB-139/140	0.0237 J	0.0307 J	0.0281 J	0.0307 J	0.0479 J	0.0342	0.0514	0.0206 U
PCB-141	0.201 J	0.225 J	0.22 J	0.225 J	0.413 J	0.243	0.421	0.322
PCB-142	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-143	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.00771 J	0.0103 U
PCB-144	0.0593 J	0.0614 J	0.061 J	0.066 J	0.11 J	0.0704	0.111	0.0849
PCB-145	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-146	0.172 J	0.189 J	0.193 J	0.196 J	0.333 J	0.218	0.286	0.237
PCB-147/149	0.851 J	0.952 J	0.968 J	1.02 J	1.62 J	1.1	1.56	1.32
PCB-148	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.00277 J	0.0103 U
PCB-150	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.00633 J	0.012 U	0.00548 J	0.0103 U
PCB-152	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.00483 J	0.012 U	0.00616 J	0.0103 U
PCB-153/168	0.888 J	0.966 J	1.01 J	1.05 J	1.78 J	1.14	1.64	1.38
PCB-154	0.0295 J	0.0282 J	0.0365 J	0.034 J	0.0523 J	0.0378	0.0369	0.0368
PCB-155	0.0034 J	0.0118 U	0.00474 J	0.00355 J	0.00677 J	0.012 U	0.0132 U	0.0103 U
PCB-156/157	0.157	0.173 J	0.167	0.176	0.324 J	0.209	0.348	0.26
PCB-158	0.123 J	0.134 J	0.137 J	0.141 J	0.249 J	0.156	0.264	0.196
PCB-159	0.0123 J	0.00994 J	0.0106 UJ	0.00899 J	0.0111 UJ	0.0115 J	0.0178	0.0103 U
PCB-160	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-161	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-162	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00351 J	0.0111 UJ	0.00542 J	0.00754 J	0.0103 U
PCB-164	0.0821 J	0.0757 J	0.099 J	0.104 J	0.162 J	0.0981	0.158	0.134
PCB-165	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-167	0.0508	0.0562 J	0.0545	0.0613	0.102 J	0.0645	0.106	0.0828
PCB-169	0.012 UJ	0.0118 UJ	0.0106 UJ	0.00935 UJ	0.012 J	0.012 U	0.00566 J	0.0103 U
PCB-170	0.166	0.186 J	0.182	0.192	0.329 J	0.204	0.286	0.251
PCB-171/173	0.0751	0.0732 J	0.0763	0.0902	0.131 J	0.0903	0.112	0.106
PCB-172	0.042	0.0393 J	0.0463	0.0518	0.0712 J	0.056	0.0622	0.0588
PCB-174	0.188	0.205 J	0.203	0.23	0.342 J	0.231	0.301	0.273
PCB-175	0.0163	0.0151 J	0.0184 J	0.0192	0.0268 J	0.0229	0.0203	0.0201
PCB-176	0.0277	0.0293 J	0.0314	0.0301	0.0524 J	0.0367	0.0461	0.041
PCB-177	0.121	0.127 J	0.135	0.154	0.224 J	0.157	0.184	0.175
PCB-178	0.0569	0.0593 J	0.058	0.0589	0.0928 J	0.0616	0.0824	0.0713
PCB-179	0.0898	0.0928 J	0.0983	0.0989	0.166 J	0.107	0.152	0.127
PCB-180/193	0.558	0.592 J	0.613	0.639	1.04 J	0.685	0.893	0.819
PCB-181	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.00833 J	0.00563 J	0.00533 J	0.0103 U
PCB-182	0.012 U	0.0118 UJ	0.0106 U	0.00448 J	0.00434 J	0.012 U	0.0132 U	0.0103 U
PCB-183	0.142	0.151 J	0.148	0.166	0.265 J	0.173	0.221	0.201

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB031	PB031	PB059	PB059	PB076	PB076	PB076	PB076
	PB031-1SWNBT-091104-N			PB059-1SWMID-091105-N	PB076-1SWMID-091104-N			
Sample Date:	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Date:	11/4/2003 N	11/3/2003 N	11/4/2003 N	11/3/2003 N	11/4/2003 N	N	N	N
y.	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
γ. γ·	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
PCB-133	0.0366 J	0.04 J	0.0563 J	0.0134 J	0.0116 J	0.0102 UJ	0.0212 J	0.0122 UJ
PCB-133	0.182 J	0.174 J	0.296 J	0.0134 J 0.0877 J	0.0118 J	0.0102 01	0.103 J	0.0122 03 0.0874 J
PCB-135/151	0.182 J 0.681 J	0.718 J	1.11 J	0.0877 J	0.223 J	0.172 J	0.358 J	0.0874 J
PCB-133/131 PCB-136	0.347 J	0.718 J	0.569 J	0.144 J	0.223 J 0.111 J	0.1723 0.0713 J	0.175 J	0.133 J
PCB-130 PCB-137	0.126 J	0.313 J	0.244 J	0.0617 J	0.0577 J	0.0713 J	0.0908 J	0.0676 J
PCB-137/	0.120 J	0.0518 J	0.0966 J	0.0017 J	0.0204 J	0.032 J	0.0308 J	0.0254 J
PCB-133/140	0.43 J	0.413 J	0.712 J	0.19 J	0.164 J	0.0143 J	0.26 J	0.184 J
PCB-141	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-142	0.0115 UJ	0.0106 UJ	0.024 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.00474 J	0.0122 UJ
PCB-144	0.113 J	0.108 J	0.195 J	0.0497 J	0.0369 J	0.0102 03	0.0635 J	0.0481 J
PCB-145	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-146	0.287 J	0.294 J	0.437 J	0.115 J	0.107 J	0.0627 J	0.174 J	0.122 J
PCB-147/149	1.54 J	1.63 J	2.43 J	0.652 J	0.487 J	0.347 J	0.795 J	0.607 J
PCB-148	0.00377 J	0.00299 J	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-150	0.00561 J	0.0106 UJ	0.00502 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-152	0.00662 J	0.0106 UJ	0.0092 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-153/168	1.63 J	1.66 J	2.5 J	0.704 J	0.607 J	0.361 J	0.984 J	0.673 J
PCB-154	0.0412 J	0.0481 J	0.0485 J	0.01 UJ	0.0115 J	0.00883 J	0.0219 J	0.0122 UJ
PCB-155	0.0115 U	0.0106 U	0.0119 UJ	0.01 UJ	0.013 U	0.0102 U	0.0125 UJ	0.0122 UJ
PCB-156/157	0.351	0.338	0.632 J	0.16 J	0.156	0.0823 J	0.24 J	0.167 J
PCB-158	0.26 J	0.246 J	0.445 J	0.114 J	0.106 J	0.0598 J	0.169 J	0.113 J
PCB-159	0.0182 J	0.0106 UJ	0.0233 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.0102 J	0.0122 UJ
PCB-160	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-161	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-162	0.00829 J	0.0106 UJ	0.0127 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.00675 J	0.0122 UJ
PCB-164	0.162 J	0.166 J	0.239 J	0.0693 J	0.0563 J	0.0354 J	0.0901 J	0.0615 J
PCB-165	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-167	0.112	0.103	0.182 J	0.0467 J	0.048	0.0249 J	0.0758 J	0.054 J
PCB-169	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-170	0.284 J	0.31	0.4 J	0.104 J	0.103 J	0.0499 J	0.161 J	0.099 J
PCB-171/173	0.118 J	0.127	0.162 J	0.0437 J	0.0432 J	0.02 J	0.0663 J	0.0425 J
PCB-172	0.0612 J	0.074	0.0868 J	0.0256 J	0.0231 J	0.0133 J	0.0373 J	0.0281 J
PCB-174	0.328 J	0.339	0.42 J	0.106 J	0.105 J	0.0548 J	0.167 J	0.11 J
PCB-175	0.0206 J	0.0268	0.0286 J	0.00745 J	0.00677 J	0.0102 UJ	0.0151 J	0.0122 UJ
PCB-176	0.0445 J	0.0466	0.0524 J	0.0136 J	0.011 J	0.00737 J	0.02 J	0.0131 J
PCB-177	0.201 J	0.217	0.255 J	0.064 J	0.0617 J	0.041 J	0.104 J	0.0689 J
PCB-178	0.0782 J	0.0776	0.104 J	0.0264 J	0.0261 J	0.0107 J	0.0388 J	0.0241 J
PCB-179	0.158 J	0.156	0.205 J	0.0483 J	0.0394 J	0.0256 J	0.0713 J	0.0505 J
PCB-180/193	0.899 J	0.98	1.24 J	0.327 J	0.317 J	0.149 J	0.5 J	0.326 J
PCB-181	0.0065 J	0.00786 J	0.0082 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-182	0.0115 UJ	0.0106 U	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-183	0.245 J	0.239	0.31 J	0.0787 J	0.0851 J	0.0448 J	0.134 J	0.0838 J

Appendix A-2
BERA Surface Water Dataset

Task	2011 UCC	2011 UCC	2011 LICC	2011 LICC	2011 LICC
Task:		2011 USC	2011 USC	2011 USC	2011 USC
Location ID:		PB080	PB080	PB082.1	PB101C
Sample ID:		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date:		8/8/2011 FD	8/8/2011	8/8/2011	8/8/2011
Sample Type:			N 2200760 755	N 2200777 402	N 2201207 665
X:		3200769.755	3200769.755	3200777.403	3201307.665
Υ:		13830051.11	13830051.11	13829942.54	13828208.48
PCB-133	0.00711 J	0.0125	0.0123	0.00611 J	0.00157 U
PCB-134	0.0339	0.0555	0.0574	0.0283 J	0.00506 U
PCB-135/151	0.146	0.244	0.246	0.123	0.0237 U
PCB-136	0.0758	0.12	0.124	0.0639	0.0116 U
PCB-137	0.0352	0.0562	0.0568	0.0282	0.00417 U
PCB-139/140	0.0137 J	0.02 J	0.0207 J	0.0107 J	0.00147 U
PCB-141	0.0916	0.151	0.152	0.0745	0.0145 U
PCB-142	0.00166 U	0.00177 U	0.00275 U	0.0023 U	0.00166 U
PCB-143	0.00307 J	0.00379 J	0.00507 J	0.00219 U	0.00158 U
PCB-144	0.0229	0.0382	0.0399	0.0197	0.00404 U
PCB-145	0.000728 U	0.000886 U	0.00114 U	0.00184 U	0.000647 U
PCB-146	0.0695	0.109	0.117	0.0555	0.0124 U
PCB-147/149	0.353	0.57	0.567	0.29	0.0568 U
PCB-148	0.000907 U	0.0011 U	0.00141 U	0.0023 U	0.000806 U
PCB-150	0.000702 U	0.00161 J	0.00193 J	0.00177 U	0.000623 U
PCB-152	0.00172 J 0.367	0.00224 J 0.592	0.0024 J 0.608	0.00176 U 0.292	0.000617 U 0.0626 U
PCB-153/168	 				
PCB-154	0.00588 J	0.0101 J	0.00926 J	0.00492 J	0.00117 J
PCB-155 PCB-156/157	0.000627 U 0.0709	0.000945 J	0.000982 U	0.00159 U	0.00076 J 0.009 U
	0.0709	0.112	0.121	0.0555	
PCB-158 PCB-159	0.0014 0.00168 U	0.1 0.00188 U	0.101 0.0025 U	0.0476 0.00176 U	0.00833 U 0.00111 U
PCB-159	0.00168 U	0.00188 U	0.0023 U		0.00111 U
PCB-161	0.0012 U	0.00128 U	0.00198 U	0.00166 U 0.00159 U	0.0012 U
PCB-162	0.00113 U	0.00122 0	0.00362 J	0.00139 U	0.00113 U
PCB-164	0.0387	0.064	0.0616	0.0313	0.00100 U
PCB-165	0.00127 U	0.00135 U	0.00209 U	0.00175 U	0.00022 U
PCB-167	0.0214	0.0346	0.0362	0.0176	0.00127 U
PCB-169	0.00134 U	0.0016 U	0.00202 U	0.00145 U	0.000936 U
PCB-170	0.0743	0.116	0.117	0.0534	0.0128 U
PCB-171/173	0.0229	0.0372	0.0386	0.0189 J	0.005 U
PCB-172	0.0125	0.0231	0.0207	0.00955 J	0.00239 J
PCB-174	0.0672	0.101	0.105	0.0507	0.015 U
PCB-175	0.00307 J	0.00536 J	0.00511 J	0.00276 J	0.000755 U
PCB-176	0.0089 J	0.0151	0.0152	0.00726 J	0.00246 U
PCB-177	0.0401	0.0673	0.0675	0.0313	0.00836 U
PCB-178	0.0146	0.0239	0.0247	0.0126 J	0.0037 U
PCB-179	0.031	0.049	0.0479	0.0239	0.00734 U
PCB-180/193	0.139	0.22	0.232	0.108	0.0309 U
PCB-181	0.00138 U	0.0016 U	0.00233 J	0.00278 U	0.000851 U
PCB-182	0.00118 U	0.00223 U	0.00123 U	0.00197 U	0.000776 U
PCB-183	0.02345	0.0367	0.04	0.0186	0.00585 U

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB006	PB006	PB006	PB006	PB006	PB006	PB031	PB031
	PB006-1SWMID-091104-D		PB006-1SWMID-091105-D				PB031-1SWMID-091104-N	
Sample Date:	11/4/2009	11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	FD	N	FD	N	N	N	N	N
x:	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
Y:	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
PCB-184	0.012 U	0.0118 UJ	0.0106 U	0.00308 J	0.00615 J	0.012 U	0.0132 U	0.0103 U
PCB-185	0.021	0.0269 J	0.0106 U	0.0324	0.0258 J	0.0206 J	0.0231	0.0247 J
PCB-186	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-187	0.251	0.276 J	0.281	0.328	0.437 J	0.324	0.366	0.355
PCB-188	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.00412 J	0.00283 J	0.0132 U	0.0103 U
PCB-189	0.0129	0.0133 J	0.00915 J	0.0122	0.02 J	0.0154	0.0184	0.0144
PCB-190	0.0496	0.0523 J	0.0469 J	0.0574	0.0804 J	0.0564	0.0697	0.0722
PCB-191	0.00789 J	0.0117 J	0.0106 U	0.00935 U	0.0203 J	0.0115 J	0.0137	0.0171
PCB-192	0.012 U	0.0118 UJ	0.0106 U	0.00935 U	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-194	0.102 J	0.107 J	0.0999	0.111 J	0.172 J	0.12	0.142	0.137
PCB-195	0.038 J	0.0509 J	0.043	0.0502 J	0.0718 J	0.0509	0.0599	0.0522
PCB-196	0.0557 J	0.0559 J	0.0519 J	0.0499 J	0.0896 J	0.0641	0.0741	0.0755
PCB-197	0.0158 J	0.015 J	0.016	0.015 J	0.0233 J	0.0146 J	0.011 J	0.0149
PCB-198/199	0.124 J	0.116 J	0.136	0.117 J	0.2 J	0.142	0.178	0.166
PCB-200	0.0188 J	0.0208 J	0.0183 J	0.0198 J	0.0324 J	0.0228	0.0282	0.0209 J
PCB-201	0.0283 J	0.0286 J	0.0309	0.0291 J	0.0462 J	0.0314 J	0.0317	0.0279
PCB-202	0.0656	0.0667 J	0.064 J	0.0708	0.0936	0.0729	0.0465	0.0523
PCB-203	0.0568 J	0.0589 J	0.0665	0.0514 J	0.101 J	0.0634	0.0812	0.0723
PCB-204	0.012 UJ	0.0118 UJ	0.0106 U	0.00935 UJ	0.0111 UJ	0.012 U	0.0132 U	0.0103 U
PCB-205	0.0137 J	0.0178 J	0.0106 U	0.00935 UJ	0.0173 J	0.0141	0.0116 J	0.0125 J
PCB-206	0.17 J	0.188 J	0.184	0.179 J	0.241 J	0.189	0.125	0.16
PCB-207	0.125 J	0.131 J	0.123	0.102 J	0.149 J	0.118	0.0515	0.0889
PCB-208	0.183	0.184 J	0.176	0.171	0.219	0.173	0.0797	0.127
PCB-209	7.78 J	8.14 J	8	7.97 J	9.52 J	7.6 J	2.1	4.72
Total PCB Congener (U = 1/2)	96.4	97.5	98.1	103	174	112	234	156
Total PCB Congener minus 209 (U = 1/2)	88.6	89.4	90.1	95.1	164	104	232	151
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	0.0232	0.0242	0.0230	0.0243	0.0472	0.0260	0.0630	0.0396
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	0.000128	0.000135	0.000131	9.76E-05	0.000262	0.000138	0.000332	0.000216
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	0.0015642	0.00161	0.00165	0.000786	0.00316	0.00165	0.00361	0.00242
Monochlorobiphenyls (total)	0.0208 J	0.024 J	0.0201 J	0.0139 J	0.0403 J	0.0147 J	0.0564 J	0.024
Dichlorobiphenyls (total)	1.03	1.07 J	0.783	0.823	1.76	0.985	3.69	1.73
Trichlorobiphenyls (total)	17.5	17.7 J	15.8	16.2 J	31.8 J	18.5	53.1	30.8
Tetrachlorobiphenyls (total)	43.1 J	42.3 J	44.4	47.3 J	78.3 J	51.3	115	75
Pentachlorobiphenyls (total)	18.6 J	19.3 J	19.8 J	21.2	36.5 J	23.2	45.3	31.2 J
Hexachlorobiphenyls (total)	5.33 J	5.85 J	5.99 J	6.25 J	10.5 J	6.82	10.5	8.26
Heptachlorobiphenyls (total)	1.83	1.95 J	1.95 J	2.17 J	3.35 J	2.26 J	2.86	2.63 J
Octachlorobiphenyls (total)	0.519 J	0.538 J	0.526 J	0.514 J	0.848 J	0.596 J	0.664	0.632 J
Nonachlorobiphenyls (total)	0.479 J	0.503 J	0.483	0.453 J	0.609 J	0.481	0.256	0.376

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB031	PB031	PB059	PB059	PB076	PB076	PB076	PB076
		PB031-1SWNBT-091105-N				PB076-1SWMID-091105-N		
Sample Date:	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	N	N	N	N	N	N	N	N
X:	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
Y:	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
PCB-184	0.0115 UJ	0.0106 U	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-185	0.0316 J	0.0357	0.0511 J	0.0117 J	0.0109 J	0.00532 J	0.0195 J	0.0107 J
PCB-186	0.0115 UJ	0.0106 U	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-187	0.416 J	0.445	0.516 J	0.141 J	0.138 J	0.0854 J	0.226 J	0.157 J
PCB-188	0.0115 U	0.0106 U	0.0119 U	0.01 UJ	0.013 U	0.0102 UJ	0.0125 U	0.0122 UJ
PCB-189	0.0175 J	0.0172 J	0.024 J	0.00589 J	0.00461 J	0.0102 UJ	0.0109 J	0.0122 UJ
PCB-190	0.0675 J	0.0838	0.0981 J	0.0257 J	0.0264 J	0.0137 J	0.0363 J	0.0212 J
PCB-191	0.0127 J	0.0106 U	0.0219 J	0.0051 J	0.013 UJ	0.0102 UJ	0.00969 J	0.0122 UJ
PCB-192	0.0115 UJ	0.0106 U	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-194	0.135 J	0.159 J	0.183 J	0.0446 J	0.0465 J	0.0224 J	0.078 J	0.0487 J
PCB-195	0.0677 J	0.0635 J	0.0764 J	0.0167 J	0.0247 J	0.0104 J	0.033 J	0.0233 J
PCB-196	0.063 J	0.0834 J	0.0975 J	0.0267 J	0.0209 J	0.00964 J	0.0313 J	0.0222 J
PCB-197	0.00849 J	0.0127 J	0.0127 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.00518 J
PCB-198/199	0.147 J	0.19 J	0.207 J	0.0594 J	0.0506 J	0.0214 J	0.0749 J	0.0474 J
PCB-200	0.0296 J	0.0305 J	0.0338 J	0.0104 J	0.0108 J	0.0102 UJ	0.0114 J	0.0122 UJ
PCB-201	0.0313 J	0.0337 J	0.035 J	0.0117 J	0.00832 J	0.0102 UJ	0.0168 J	0.013 J
PCB-202	0.0479	0.054	0.0561 J	0.0169 J	0.0137	0.00626 J	0.0231 J	0.0148 J
PCB-203	0.0662 J	0.0801 J	0.102 J	0.0308 J	0.0269 J	0.0103 J	0.0361 J	0.0258 J
PCB-204	0.0115 UJ	0.0106 UJ	0.0119 UJ	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-205	0.0113 J	0.0106 UJ	0.011 J	0.01 UJ	0.013 UJ	0.0102 UJ	0.0125 UJ	0.0122 UJ
PCB-206	0.113 J	0.157 J	0.089 J	0.0246 J	0.0243 J	0.0102 UJ	0.0327 J	0.0312 J
PCB-207	0.0499 J	0.0639 J	0.0275 J	0.0075 J	0.013 UJ	0.0102 UJ	0.0103 J	0.0122 UJ
PCB-208	0.0781 J	0.0999	0.0456 J	0.0126 J	0.00989 J	0.0102 UJ	0.0186 J	0.0113 J
PCB-209	2.31 J	3.52 J	0.542 J	0.163 J	0.0916 J	0.245 J	0.144 J	0.685 J
Total PCB Congener (U = 1/2)	230	197	431	135	81.0	58.9	117	94.0
Total PCB Congener minus 209 (U = 1/2)	228	194	430	135	80.9	58.6	117	93.3
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	0.0622	0.0505	0.11841512	0.0326	0.0238	0.0139	0.0372	0.0293
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	0.000319	0.000261	0.000620	0.000122	0.000155	8.86E-05	0.000236	0.000119
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	0.00334	0.00278	0.00643	0.000866	0.00205	0.00123	0.00295	0.000997
Monochlorobiphenyls (total)	0.0758 J	0.0251 J	0.204 J	0.0769 J	0.0608	0.05 J	0.0726 J	0.0541 J
Dichlorobiphenyls (total)	3.77	2.21 J	8.33 J	2.9 J	1.42	1.21 J	1.83 J	1.41 J
Trichlorobiphenyls (total)	54.8	38.8 J	101 J	33.3 J	16.6	12.9 J	23.3 J	17.7 J
Tetrachlorobiphenyls (total)	109	96.4 J	211 J	68 J	39.7	29 J	56.3 J	46.7 J
Pentachlorobiphenyls (total)	45.6	41.5 J	87.7 J	24.5 J	17.7	12.2 J	27.2 J	21.4 J
Hexachlorobiphenyls (total)	10.5 J	10.3 J	17.1 J	4.57 J	3.88 J	2.37 J	6.22 J	4.44 J
Heptachlorobiphenyls (total)	2.99 J	3.18 J	3.98 J	1.03 J	1 J	0.521 J	1.62 J	1.04 J
Octachlorobiphenyls (total)	0.608 J	0.707 J	0.814 J	0.217 J	0.202 J	0.0803 J	0.304 J	0.2 J
Nonachlorobiphenyls (total)	0.241 J	0.321 J	0.162 J	0.0447 J	0.0342 J	0 UJ	0.0616 J	0.0425 J

Appendix A-2
BERA Surface Water Dataset

1			I		
Task:	2011 USC	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB066B	PB080	PB080	PB082.1	PB101C
•		PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date:	8/8/2011	8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type:	N	FD	N	N	N
X:	3201384.326	3200769.755	3200769.755	3200777.403	3201307.665
Y:	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
PCB-184	0.000847 U	0.00161 U	0.000883 U	0.00142 U	0.000559 U
PCB-185	0.02345	0.0367	0.04	0.0186	0.00585 U
PCB-186	0.000932 U	0.00177 U	0.000972 U	0.00156 U	0.000615 U
PCB-187	0.0737	0.119	0.132	0.056	0.0189 U
PCB-188	0.00105 U	0.00197 U	0.00111 U	0.00165 U	0.000688 U
PCB-189	0.00327 J	0.00513 J	0.00559 J	0.00375 J	0.00116 J
PCB-190	0.0146	0.0227	0.0231	0.0107 J	0.00316 J
PCB-191	0.0031 J	0.00442 J	0.00543 J	0.0023 U	0.000705 U
PCB-192	0.00117 U	0.00136 U	0.00158 U	0.00236 U	0.000722 U
PCB-194	0.0257	0.0382	0.0454	0.0222	0.00657 U
PCB-195	0.00951 J	0.0133 U	0.0189	0.00825 J	0.0028 U
PCB-196	0.0145	0.0225	0.0233	0.0108 J	0.00401 U
PCB-197	0.00279 J	0.004665 J	0.004425 J	0.001975 J	0.000815 J
PCB-198/199	0.0328	0.0529	0.0586	0.0258	0.00879 U
PCB-200	0.00279 J	0.004665 J	0.004425 J	0.001975 J	0.000815 J
PCB-201	0.00441 J	0.00718 J	0.00776 J	0.00351 J	0.00159 U
PCB-202	0.00765 J	0.0128	0.0139	0.00583 J	0.00288 J
PCB-203	0.0154	0.0271	0.0258	0.0114 J	0.00511 U
PCB-204	0.00058 U	0.0011 U	0.0013 U	0.001 U	0.000642 U
PCB-205	0.002 U	0.00908 U	0.00488 U	0.00213 U	0.00116 U
PCB-206	0.0179	0.0431	0.0356	0.0181 J	0.00402 J
PCB-207	0.00855 J	0.0108 J	0.0131	0.00645 J	0.00187 U
PCB-208	0.00926 J	0.0138 J	0.0162 J	0.00662 J	0.00234 J
PCB-209	0.353	1.07	0.965	0.267	0.00943 U
Total PCB Congener (U = 1/2)	100 J	147 J	143 J	79.9 J	5.65 J
Total PCB Congener minus 209 (U = 1/2)					
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	0.02	0.03	0.03	0.01	0.001
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)					
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	0.000789 J	0.00116 J	0.0015 J	0.00027 J	0.000119 J
Monochlorobiphenyls (total)	0.06	0.08	0.06	0.07	0.01
Dichlorobiphenyls (total)	2.51	3.03	2.98	2.07	0.14
Trichlorobiphenyls (total)	31.22	39.87	41.12	23.67	1.06
Tetrachlorobiphenyls (total)	49.28	75.953	71.167	40.32	3.21
Pentachlorobiphenyls (total)	13.98	22.01	21.86	11.03	0.936
Hexachlorobiphenyls (total)	2.391	3.829	3.904	1.93	0.187
Heptachlorobiphenyls (total)	0.558	0.89	0.925	0.43	0.068
Octachlorobiphenyls (total)	0.117	0.18	0.21	0.09	0.02
Nonachlorobiphenyls (total)	0.04	0.1	0.1	0.03	0.01

Appendix A-2
BERA Surface Water Dataset

Task	DI /FC 2000	DI /FC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000	DI /EC 2000
Task:	RI/FS 2009 PB006	RI/FS 2009	RI/FS 2009	RI/FS 2009 PB006	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:		PB006	PB006		PB006	PB006	PB031	PB031
-	PB006-1SWMID-091104-D				PB006-1SWNBT-091104-N		PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
Sample Date:	11/4/2009 FD	11/4/2009 N	11/5/2009 FD	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:			3202137.4168	3202137.4168	IN 2202127 4160	3202137.4168	3201366.8205	3201366.8205
X: v.	13836272.1183	3202137.4168 13836272.1183	13836272.1183	13836272.1183	3202137.4168 13836272.1183	13836272.1183	13834640.6333	13834640.6333
Destinidad (m./l)	13630272.1163	15050272.1105	150502/2.1105	13630272.1163	150502/2.1105	13630272.1163	13634040.0333	13634040.0333
Pesticides (µg/L)	0.00400.11	0.0011111	0.004.11	0.00110.11	0.00440.111	0.00422.11	0.0044411	0.004.11
2,4'-DDD (o,p'-DDD)	0.00109 U	0.00111 U	0.001 U	0.00119 U	0.00119 UJ	0.00122 U	0.00114 U	0.001 U
2,4'-DDE (o,p'-DDE)	0.00109 U	0.00111 U	0.001 U	0.00119 U	0.00119 UJ	0.00122 U	0.00114 U	0.001 U
2,4'-DDT (o,p'-DDT)	0.00109 U	0.00111 U	0.001 U	0.00119 U	0.00119 UJ	0.00122 U	0.00114 U	0.001 U
4,4'-DDD (p,p'-DDD)	0.00109 U	0.00111 U	0.001 U	0.00119 U	0.00119 UJ	0.00122 U	0.00114 U	0.001 U
4,4'-DDE (p,p'-DDE) 4,4'-DDT (p,p'-DDT)	0.00109 U 0.00246 J	0.00111 U 0.00317 J	0.001 U 0.001 U	0.00119 U 0.00119 U	0.00119 UJ 0.00187 J	0.00122 U 0.00122 U	0.00114 U 0.00254 J	0.001 U 0.001 U
Sum 2,4 DDT, DDE, DDD (U=1/2)	0.00246 J 0.00109 U	0.00317 J		0.00119 U		0.00122 U		
Sum 4,4 DDT, DDE, DDD (U=1/2)	0.00109 U	0.00111 U	0.001 U 0.001 U	0.00119 U	0.00119 UJ 0.00306 J	0.00122 U	0.00114 U 0.00368 J	0.001 U 0.001 U
Sum DDD (U=1/2)	0.00333 J	0.00428 J	0.001 U	0.00119 U	0.00306 J	0.00122 U	0.00368 J	0.001 U
Sum DDE (U=1/2)	0.00109 U	0.00111 U	0.001 U	0.00119 U	0.00119 UJ	0.00122 U	0.00114 U	0.001 U
Sum DDT (U=1/2)	0.003005 J	0.003725 J	0.001 U	0.00119 U	0.00119 03 0.002465 J	0.00122 U	0.00114 U	0.001 U
Total DDx (U = 1/2)	0.005185 J	0.005725 J	0.001 U	0.00119 U	0.002403 J	0.00122 U	0.00539 J	0.001 U
Aldrin	0.0051833	0.00261	0.001 U	0.00013 U	0.00299 J	0.00122 U	0.0423 NJ	0.001 U
alpha-BHC	0.00659 J	0.00527	0.0063 0 0.00453 NJ	0.00369 NJ	0.0064	0.00576	0.0449 NJ	0.0005 U
beta-BHC	0.00054 U	0.00327 0.00056 U	0.0005 U	0.0006 U	0.0004 0.0006 U	0.00061 U	0.00057 U	0.0005 U
alpha-Chlordane (cis-Chlordane)	0.00226	0.00255	0.00221	0.0018	0.00251 J	0.00319	0.00226 NJ	0.00895 NJ
delta-BHC	0.00054 U	0.00056 U	0.00606	0.0056	0.00744	0.00531	0.0384	0.0172
Dieldrin	0.00582 NJ	0.00579 J	0.00416	0.00423	0.00637 J	0.00061 U	0.0244 NJ	0.0115 NJ
Endosulfan-alpha (I)	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.00202 J	0.00061 U	0.00057 U	0.0005 U
Endosulfan-beta (II)	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.0006 U	0.00061 U	0.00057 U	0.0005 U
Endosulfan sulfate	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.0006 U	0.00061 U	0.00057 U	0.0005 U
Endrin	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.0006 U	0.00061 U	0.00057 U	0.0005 U
Endrin aldehyde	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.0006 U	0.00061 U	0.00057 U	0.0005 U
Endrin ketone	0.00054 U	0.00388 NJ	0.0005 U	0.0006 U	0.00271 J	0.00061 U	0.00057 U	0.0005 U
gamma-BHC (Lindane)	0.00054 U	0.00331 NJ	0.00552 NJ	0.00576 NJ	0.0006 U	0.00586 NJ	0.00057 U	0.0005 U
gamma-Chlordane	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.0006 U	0.00061 U	0.00057 U	0.0005 U
Heptachlor	0.0066 NJ	0.00056 U	0.0005 U	0.00324	0.00693 NJ	0.00644 J	0.04 NJ	0.0005 U
Heptachlor epoxide	0.00113 NJ	0.00103 NJ	0.00103 NJ	0.00093 NJ	0.00153 NJ	0.00132 NJ	0.00874 NJ	0.00327 NJ
Methoxychlor	0.00054 U	0.00056 U	0.0005 U	0.0006 U	0.0006 U	0.00061 U	0.00057 U	0.0005 U
Toxaphene	0.0272 U	0.0278 U	0.025 U	0.0298 U	0.0298 U	0.0305 U	0.0284 U	0.025 U
Dioxin Furans (ng/L)								
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	0.00227 U	0.00367 UJ	0.00206 U	0.00102 J	0.00247 U	0.00326 U	0.00148 J	0.00403 U
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	0.00272 U	0.00158 UJ	0.00289 U	0.00397 UJ	0.00368 U	0.00251 U	0.00202 U	0.00475 U
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	0.00286 UJ	0.00295 UJ	0.00583 UJ	0.00294 UJ	0.00351 UJ	0.00516 UJ	0.00189 U	0.00449 UJ
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	0.00279 UJ	0.00304 UJ	0.0058 UJ	0.00309 UJ	0.00373 UJ	0.00501 UJ	0.00204 U	0.00523 UJ
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	0.00312 UJ	0.00346 UJ	0.00661 UJ	0.00349 UJ	0.00394 UJ	0.00582 UJ	0.00197 U	0.00593 UJ
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	0.0186 J	0.0197 J	0.0178 J	0.018 J	0.0273 J	0.0167 J	0.0213 J	0.0178 J
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	0.19 J	0.206 J	0.176 J	0.2 J	0.257 J	0.193 J	0.145 J	0.186 J
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	0.00239 J	0.00285 J	0.000637 U	0.00225 J	0.00417 J	0.00167 U	0.0031 J	0.0031 J
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	0.00139 J	0.0027 J	0.00247 U	0.00152 J	0.003 J	0.00314 U	0.0024 U	0.00303 J
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	0.00192 U	0.002 UJ	0.00234 U	0.00326 U	0.0016 J	0.000897 J	0.00228 U	0.00317 U

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009	RI/FS 2009
Location ID:	PB031	PB031	PB059	PB059	PB076	PB076	PB076	PB076
	PB031-1SWNBT-091104-N			PB059-1SWMID-091105-N			PB076-1SWNBT-091104-N	
Sample Date:	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	N	N	N N	N N	N	N N	N N	N N
x:	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
Y:	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
Pesticides (μg/L)								
2,4'-DDD (o,p'-DDD)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
2,4'-DDE (o,p'-DDE)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
2,4'-DDT (o,p'-DDT)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
4,4'-DDD (p,p'-DDD)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
4,4'-DDE (p,p'-DDE)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
4,4'-DDT (p,p'-DDT)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Sum 2,4 DDT, DDE, DDD (U=1/2)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Sum 4,4 DDT, DDE, DDD (U = 1/2)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Sum DDD (U=1/2)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Sum DDE (U=1/2)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Sum DDT (U=1/2)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Total DDx (U = 1/2)	0.00133 UJ	0.001 U	0.00115 U	0.001 U	0.0012 U	0.001 U	0.00111 U	0.00122 U
Aldrin	0.0283 NJ	0.0005 U	0.00058 U	0.0005 U	0.0255 NJ	0.028 NJ	0.00056 U	0.0327 NJ
alpha-BHC	0.0346 J	0.0219	0.0885 NJ	0.0665 J	0.0993 NJ	0.0992 NJ	0.0985 J	0.116 NJ
beta-BHC	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
alpha-Chlordane (cis-Chlordane)	0.00209 NJ	0.0091	0.0254 NJ	0.0202 NJ	0.0432 NJ	0.0295 NJ	0.0379 NJ	0.00061 U
delta-BHC	0.0426 J	0.0005 U	0.0898	0.0496	0.123	0.116 J	0.14	0.00061 U
Dieldrin	0.0222 NJ	0.012 NJ	0.0309	0.0178 J	0.0361 NJ	0.0147	0.0344 NJ	0.00061 U
Endosulfan-alpha (I)	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Endosulfan-beta (II)	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Endosulfan sulfate	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Endrin	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Endrin aldehyde	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Endrin ketone	0.00067 UJ	0.0005 U	0.00516	0.00343	0.0006 U	0.00837 J	0.00701	0.00061 U
gamma-BHC (Lindane)	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
gamma-Chlordane	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Heptachlor	0.037 NJ	0.0005 U	0.055 J	0.0337	0.0622	0.061	0.0565	0.00061 U
Heptachlor epoxide	0.00762	0.00326 NJ	0.00997 NJ	0.00654 NJ	0.0148 NJ	0.00835 NJ	0.0152 NJ	0.0123 NJ
Methoxychlor	0.00067 UJ	0.0005 U	0.00058 U	0.0005 U	0.0006 U	0.0005 U	0.00056 U	0.00061 U
Toxaphene	0.0333 UJ	0.025 U	0.0287 U	0.025 U	0.0301 U	0.025 U	0.0278 U	0.0305 U
Dioxin Furans (ng/L)								
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	0.00148 U	0.00216 UJ	0.00247 UJ	0.00402 UJ	0.00219 U	0.00331 UJ	0.00176 U	0.00251 UJ
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	0.00281 U	0.0032 UJ	0.00242 UJ	0.00256 UJ	0.0027 U	0.00345 UJ	0.003 UJ	0.00278 UJ
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	0.00511 UJ	0.00358 UJ	0.0042 UJ	0.00581 UJ	0.00674 UJ	0.0188 UJ	0.00314 UJ	0.00739 UJ
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	0.00559 UJ	0.00344 UJ	0.00436 UJ	0.00566 UJ	0.00683 UJ	0.0189 UJ	0.00335 UJ	0.00704 UJ
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	0.00631 UJ	0.00377 UJ	0.00453 UJ	0.00664 UJ	0.0072 UJ	0.0209 UJ	0.00363 UJ	0.00769 UJ
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	0.0173 J	0.0289 J	0.0208 J	0.0114 J	0.00665 UJ	0.0305 UJ	0.00689 J	0.0224 UJ
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	0.193 J	0.192 J	0.187 J	0.0554 J	0.0323 J	0.105 UJ	0.0561 J	0.0622 J
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	0.00339 J	0.004 J	0.00218 UJ	0.00203 UJ	0.00179 U	0.00335 UJ	0.00131 U	0.00174 UJ
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	0.00275 J	0.00251 J	0.00363 UJ	0.00256 UJ	0.00217 U	0.00422 UJ	0.0016 UJ	0.00254 UJ
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	0.00155 J	0.00218 UJ	0.00352 UJ	0.00215 UJ	0.00211 U	0.00454 UJ	0.0017 UJ	0.00238 UJ

Appendix A-2
BERA Surface Water Dataset

Task:		2011 USC	2011 USC	2011 USC	2011 USC
Location ID:		PB080	PB080	PB082.1	PB101C
Sample ID:	1	PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date:		8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type:		FD	N 2202752 755	N	N 2224227.555
X		3200769.755	3200769.755	3200777.403	3201307.665
Y:	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
Pesticides (µg/L)					
2,4'-DDD (o,p'-DDD)					
2,4'-DDE (o,p'-DDE)					
2,4'-DDT (o,p'-DDT)					
4,4'-DDD (p,p'-DDD)					
4,4'-DDE (p,p'-DDE)					
4,4'-DDT (p,p'-DDT)					
Sum 2,4 DDT, DDE, DDD (U=1/2)					
Sum 4,4 DDT, DDE, DDD (U = 1/2)					
Sum DDD (U=1/2)					
Sum DDE (U=1/2)					
Sum DDT (U=1/2)	-				
Total DDx (U = 1/2)					
Aldrin					
alpha-BHC					
beta-BHC					
alpha-Chlordane (cis-Chlordane)					
delta-BHC					
Dieldrin					
Endosulfan-alpha (I)	-				
Endosulfan-beta (II)	-				
Endosulfan sulfate	-				
Endrin	-				
Endrin aldehyde	-				
Endrin ketone					
gamma-BHC (Lindane)	-				
gamma-Chlordane					
Heptachlor					
Heptachlor epoxide					
Methoxychlor					
Toxaphene					
Dioxin Furans (ng/L) 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)					
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)					
1,2,3,4,7,8-Pentachiorodibenzo-p-dioxin (PecDD) 1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)			 		
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)					
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)					
					
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	- -				
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)					
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	- -				
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)					
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)					

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009							
Location ID:	PB006	PB006	PB006	PB006	PB006	PB006	PB031	PB031
Sample ID:	PB006-1SWMID-091104-D	PB006-1SWMID-091104-N	PB006-1SWMID-091105-D	PB006-1SWMID-091105-N	PB006-1SWNBT-091104-N	PB006-1SWNBT-091105-N	PB031-1SWMID-091104-N	PB031-1SWMID-091105-N
Sample Date:	11/4/2009	11/4/2009	11/5/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	FD	N	FD	N	N	N	N	N
X:	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3202137.4168	3201366.8205	3201366.8205
Y:	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13836272.1183	13834640.6333	13834640.6333
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	0.00252 J	0.00345 J	0.00281 J	0.00311 J	0.00395 J	0.00311 J	0.00322 J	0.00379 J
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	0.00134 J	0.00164 UJ	0.000563 U	0.000968 UJ	0.00275 J	0.00114 UJ	0.00183 J	0.00299 U
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	0.00299 UJ	0.00276 UJ	0.000846 UJ	0.00158 UJ	0.003 UJ	0.00189 UJ	0.00233 U	0.00514 UJ
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	0.00229 UJ	0.00196 UJ	0.0006 UJ	0.00122 UJ	0.00228 UJ	0.00135 UJ	0.00173 UJ	0.00347 UJ
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	0.0215 J	0.0262 J	0.0164 J	0.0177 J	0.0269 J	0.0206 J	0.0129 J	0.0166 J
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	0.00262 UJ	0.00651 UJ	0.000881 UJ	0.00225 UJ	0.00592 UJ	0.00355 UJ	0.00262 UJ	0.00376 UJ
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	0.358 J	0.416 J	0.33 J	0.373 J	0.429 J	0.318 J	0.0747 J	0.226 J
Total Tetrachlorodibenzo-p-dioxin (TCDD)	0.00369 J	0.00434 J	0.00463 J	0.00509 J	0.00282	0.00335 J	0.0075 J	0.00403 U
Total Pentachlorodibenzo-p-dioxin (PeCDD)	0.00272 U	0.00158 UJ	0.00289 U	0.00397 UJ	0.00368 U	0.00251 U	0.00266 J	0.00475 U
Total Hexachlorodibenzo-p-dioxin (HxCDD)	0.00422 J	0.00472 J	0.00239 J	0.00464 J	0.00606 J	0.00423 J	0.00474 J	0.00523 J
Total Heptachlorodibenzo-p-dioxin (HpCDD)	0.05 J	0.0518 J	0.0497 J	0.0485 J	0.0717 J	0.043 J	0.0522 J	0.0502 J
Total Tetrachlorodibenzofuran (TCDF)	0.00935 J	0.0124 J	0.0062 J	0.00635 J	0.0201 J	0.00806 J	0.0257 J	0.0108 J
Total Pentachlorodibenzofuran (PeCDF)	0.00139	0.0027 J	0.00158 J	0.00284 J	0.0133 J	0.00345 J	0.00176 J	0.00303 J
Total Hexachlorodibenzofuran (HxCDF)	0.0178 J	0.0197 J	0.0161 J	0.0215 J	0.028 J	0.0171 J	0.0159 J	0.0162 J
Total Heptachlorodibenzofuran (HpCDF)	0.0347 J	0.0399 J	0.0338 J	0.0336 J	0.0504 J	0.0356 J	0.0219 J	0.0295 J
Total Dioxin/Furan (U = 1/2)	0.60753 J	0.691685 J	0.5587735 J	0.627984 J	0.769935 J	0.569557 J	0.27317 J	0.4778 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	0.00717695 J	0.0080464 J	0.005210505 J	0.0080634 J	0.0107769 J	0.0060406 J	0.00789932 J	0.0110157 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	0.00486005 J	0.00521195 J	0.00527548 J	0.00553935 J	0.00647785 J	0.00561945 J	0.00466092 J	0.0078701 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	0.0047307 J	0.00510465 J	0.004686555 J	0.00528015 J	0.0063324 J	0.00515825 J	0.00460201 J	0.0074943 J

Appendix A-2
BERA Surface Water Dataset

Task:	RI/FS 2009							
Location ID:	PB031	PB031	PB059	PB059	PB076	PB076	PB076	PB076
Sample ID:	PB031-1SWNBT-091104-N	PB031-1SWNBT-091105-N	PB059-1SWMID-091104-N	PB059-1SWMID-091105-N	PB076-1SWMID-091104-N	PB076-1SWMID-091105-N	PB076-1SWNBT-091104-N	PB076-1SWNBT-091105-N
Sample Date:	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009	11/4/2009	11/5/2009
Sample Type:	N	N	N	N	N	N	N	N
X:	3201366.8205	3201366.8205	3201646.6295	3201646.6295	3200819.7773	3200819.7773	3200819.7773	3200819.7773
Y:	13834640.6333	13834640.6333	13831831.4316	13831831.4316	13830437.4855	13830437.4855	13830437.4855	13830437.4855
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	0.00294 J	0.00463 J	0.00241 UJ	0.00151 UJ	0.00174 UJ	0.00294 UJ	0.000799 UJ	0.00248 UJ
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	0.00201 J	0.00197 J	0.0023 UJ	0.00137 UJ	0.00149 UJ	0.00281 UJ	0.000744 UJ	0.00254 UJ
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	0.0032 UJ	0.00229 UJ	0.00326 UJ	0.00225 UJ	0.00239 UJ	0.00524 UJ	0.00118 UJ	0.0041 UJ
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	0.00259 UJ	0.00166 UJ	0.00256 UJ	0.00161 UJ	0.00194 UJ	0.00409 UJ	0.000944 UJ	0.00297 UJ
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	0.0129 J	0.015 J	0.00802 J	0.00202 J	0.00317 UJ	0.00965 UJ	0.00356 J	0.00342 UJ
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	0.0036 UJ	0.00885 UJ	0.00261 UJ	0.0014 UJ	0.00461 UJ	0.0219 UJ	0.0078 UJ	0.00637 UJ
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	0.07 J	0.162 J	0.0401 J	0.0125 J	0.0346 UJ	0.0458 J	0.0234 UJ	0.0293 J
Total Tetrachlorodibenzo-p-dioxin (TCDD)	0.00396 J	0.00258 J	0.00214 J	0.00185 J	0.00587 J	0.00196 J	0.00619 J	0.00373 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	0.00221 J	0.0032 UJ	0.00382 J	0.00256 UJ	0.00323 J	0.00345 UJ	0.00517 J	0.00235 J
Total Hexachlorodibenzo-p-dioxin (HxCDD)	0.00567 UJ	0.00444 J	0.00735 J	0.00317 J	0.00692 UJ	0.0195 UJ	0.00337 UJ	0.00735 UJ
Total Heptachlorodibenzo-p-dioxin (HpCDD)	0.0562 J	0.0726 J	0.0687 J	0.0289 J	0.00665 UJ	0.0305 UJ	0.028 J	0.0116 J
Total Tetrachlorodibenzofuran (TCDF)	0.0286 J	0.0181 J	0.00879 J	0.00203 UJ	0.00179 U	0.00335 UJ	0.00152 J	0.00174 UJ
Total Pentachlorodibenzofuran (PeCDF)	0.0141 J	0.00694 J	0.00104 J	0.00235 UJ	0.000439 J	0.00437 UJ	0.00165 UJ	0.00246 UJ
Total Hexachlorodibenzofuran (HxCDF)	0.0151 J	0.0204 J	0.00687 J	0.00165 UJ	0.00185 UJ	0.00359 UJ	0.00416 J	0.00294 UJ
Total Heptachlorodibenzofuran (HpCDF)	0.0251 J	0.0276 J	0.0163 J	0.00567 J	0.00378 UJ	0.0142 UJ	0.00908 J	0.00455 UJ
Total Dioxin/Furan (U = 1/2)	0.321185 J	0.426575 J	0.276145 J	0.101105 J	0.07646 J	0.1756 J	0.0937285 J	0.130675 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	0.0088063 J	0.00943225 J	0.00649306 J	0.00639594 J	0.005491335 J	0.01008233 J	0.00451337 J	0.00611025 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	0.0055391 J	0.0055976 J	0.00522796 J	0.00583864 J	0.005251835 J	0.01054008 J	0.00396927 J	0.00594195 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	0.0050654 J	0.00546875 J	0.00468683 J	0.00515647 J	0.00438708 J	0.00831554 J	0.00357769 J	0.005026 J

Appendix A-2
BERA Surface Water Dataset

Task:	2011 USC	2011 USC	2011 USC	2011 USC	2011 USC
Location ID:	PB066B	PB080	PB080	PB082.1	PB101C
Sample ID:	PB066-1SWMID-20110808-N	PB080-1SWMID-20110808-D	PB080-1SWMID-20110808-N	PB082.1-1SWMID-20110808-N	PB101C-1SWMID-20110808-N
Sample Date:	8/8/2011	8/8/2011	8/8/2011	8/8/2011	8/8/2011
Sample Type:	N	FD	N	N	N
X:	3201384.326	3200769.755	3200769.755	3200777.403	3201307.665
Y:	13831305.66	13830051.11	13830051.11	13829942.54	13828208.48
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	-	-	-	1	
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	1	1	1	1	
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	1	1	1	1	
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	1	1	1	1	
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)					
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)					
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)					
Total Tetrachlorodibenzo-p-dioxin (TCDD)					
Total Pentachlorodibenzo-p-dioxin (PeCDD)					
Total Hexachlorodibenzo-p-dioxin (HxCDD)					
Total Heptachlorodibenzo-p-dioxin (HpCDD)					
Total Tetrachlorodibenzofuran (TCDF)					
Total Pentachlorodibenzofuran (PeCDF)					
Total Hexachlorodibenzofuran (HxCDF)					
Total Heptachlorodibenzofuran (HpCDF)					
Total Dioxin/Furan (U = 1/2)		-	-		
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)					
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)		-	-		
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)					

Appendix A-2 BERA Surface Water Dataset

Notes:

Bold = Detected result

J = Estimated value

U = Compound analyzed, but not detected above detection limit

UJ = Compound analyzed, but not detected above estimated detection limit

-- Results not reported or not applicable

μg/L = micrograms per liter

mg/L = milligrams per liter

ng/L = nanograms per liter

pct = percent

N = Normal Field Sample FD = Field Duplicate

TEQ values as of 2005, World Health Organization

TISSUE DATA CAN BE FOUND IN ATTACHMENT 1

APPENDIX B SURFACE SEDIMENT INTERPOLATION AND AVERAGING

TABLE OF CONTENTS

1 SPATIAL	INTERPOLATION OF COPCS IN SURFACE SEDIMENT1
1.1 Meth	nod Selection1
1.1.1 Iı	nverse-Distance Weighting3
1.1.1.1	IDW Optimization5
1.1.1.2	Data Transformation
2 SURFACE	WEIGHTED AVERAGING8
3 RESULTS	9
4 REFEREN	ICES10
List of Table	es e
Table B-1	Comparison of IDW and Kriging SWACs for PCBs
Table B-2	Transformation Parameters Selected for Each COPC
List of Figur	es
Figure B-1	Lead: Kriging and Inverse Distance Weighted
Figure B-2	Hexachlorobenzene: Kriging and Inverse Distance Weighted
Figure B-3	Total PCB Congener: Kriging and Inverse Distance Weighted
Figure B-4	Total PCB Congener TEQ (Avian): Kriging and Inverse Distance Weighted
Figure B-5	Total PCB Congener TEQ (Mammal): Kriging and Inverse Distance Weighted
Figure B-6	Total PCB Congener TEQ (Fish): Kriging and Inverse Distance Weighted
Figure B-7	Lead Variogram
Figure B-8	Hexachlorobenzene Variogram
Figure B-9	Total PCB Congener Variogram
Figure B-10	Total PCB Congener TEQ (Avian) Variogram
Figure B-11	Total PCB Congener TEQ (Mammal) Variogram
Figure B-12	Total PCB Congener TEQ (Fish) Variogram
Figure B-13	Lead Search Neighborhood Comparison
Figure B-14	Hexachlorobenzene Search Neighborhood Comparison
Figure B-15A	1,3 Dichlorobenzene Histogram
Figure B-15B	1,4 Dichlorobenzene Histogram

- Figure B-15C Bis(2-ethylhexyl) phthalate Histogram
- Figure B-15D Hexachlorobenzene Histogram
- Figure B-15E Hexachlorobutadiene Histogram
- Figure B-15F Lead Histogram
- Figure B-15G Mercury Histogram
- Figure B-15H Tetrachlorobiphenyl Histogram
- Figure B-15I Total Dioxin/Furan TEQ (Avian) Histogram
- Figure B-15J Total Dioxin/Furan TEQ (Mammal) Histogram
- Figure B-15K Total LPAH 6 of 15 Histogram
- Figure B-15L Total PAH 15 Histogram
- Figure B-15M Total PCB Congener Histogram
- Figure B-15N Total PCB Congener TEQ (Avian) Histogram
- Figure B-15O Total PCB Congener TEQ (Fish) Histogram
- Figure B-15P Total PCB Congener TEQ (Mammal) Histogram
- Figure B-15Q Trichlorobiphenyl Histogram
- Figure B-16A 1,3-Dichlorobenzene Interpolation
- Figure B-16B 1,4-Dichlorobenzene Interpolation
- Figure B-16C Bis(2-ethylhexyl) phthalate Interpolation
- Figure B-16D Hexachlorobenzene Interpolation
- Figure B-16E Hexachlorobutadiene Interpolation
- Figure B-16F Lead Interpolation
- Figure B-16G Mercury Interpolation
- Figure B-16H Tetrachlorobiphenyl Interpolation
- Figure B-16I Total Dioxin/Furan TEQ (Avian) Interpolation
- Figure B-16J Total Dioxin/Furan TEQ (Mammal) Interpolation
- Figure B-16K Total LPAH 6 of 15 Interpolation
- Figure B-16L Total PAH 15 Interpolation
- Figure B-16M Total PCB Congener Interpolation
- Figure B-16N Total PCB Congener TEQ (Avian) Interpolation
- Figure B-16O Total PCB Congener TEQ (Fish) Interpolation
- Figure B-16P Total PCB Congener TEQ (Mammal) Interpolation
- Figure B-16Q Trichlorobiphenyl Interpolation

LIST OF ACRONYMS AND ABBREVIATIONS

μg/kg microgram/kilogram

BERA Baseline Ecological Risk Assessment

COPC chemical of potential concern

IDW inverse distance weighting

mg/kg milligram/kilogram

RMSE root mean squared error

SWAC surface weighted averaging concentration

1 SPATIAL INTERPOLATION OF COPCs IN SURFACE SEDIMENT

As described in the *Baseline Ecological Risk Assessment Work Plan* (BERA Work Plan), mobile (e.g., fish, wildlife) receptors are expected to average their exposure within a defined foraging area. Sessile receptors (benthic invertebrates) are generally exposed to sediment contaminants within a small radius (e.g., centimeters) which may not be well characterized by a sample collected several meters away. Spatial interpolation¹ of the 2009 and 2011 bulk surface sediment data (Anchor QEA 2010, 2012) for selected chemical of potential concern (COPC) was performed to provide a basis to calculate surface weighted average concentrations (SWACs) across the Site and to characterize exposure of stationary receptors at unsampled locations. The following subsections describe the spatial interpolation methods and results for ecological COPCs at the Site. This information was used to support the exposure assessment for ecological receptors for the BERA.

1.1 Method Selection

Two commonly used interpolation methods were assessed for their ability to accurately predict the surface concentrations for the compounds within the COPC list. The two methods evaluated were inverse distance weighting (IDW) and Kriging. IDW assumes that each measured point has a local influence that diminishes with distance. It gives greater weight to the points closer to the prediction location than to those farther away, hence the name inverse distance weighting (QEA 2007). An important principle of IDW is that it is a deterministic exact interpolator that honors all data points. This means that the interpolation at sample points will be equal and the range of interpolated values will not exceed the sample observations. In addition, because IDW is deterministic, there is no way to evaluate the quality of the predictions without collecting confirmatory samples. Kriging is a probabilistic predictor, producing for each location an estimate of the parameter of interest and the uncertainty of that estimate (i.e., the prediction error). The estimate has the property of having the minimum variance among all estimates that are linear functions of the data. Similar to IDW, Kriging assumes that closer observations have more weight on an estimated value but also considers the spatial pattern of the observations (i.e., autocorrelation).

¹ Spatial interpolation is a general method of mapping a surface using limited sample data and estimation of values from unsampled locations.

To determine the most appropriate method for application to the surface sediment observations, two COPCs were interpolated by both methods and the results compared. The two compounds interpolated by both IDW and Kriging methods were lead and hexachlorobenzene. They were chosen for comparison because they: 1) had different concentration distributions, and 2) were generally representative of the apparent distribution of several other COPC. The spatial distribution of lead observations were relatively heterogeneous (i.e., variable) with several relatively higher loci and demonstrated a linear decrease in concentration with distance. This distribution type and others like it were classified as "linear distribution". The spatial distribution of hexachlorobenzene results also showed several relatively higher loci, however the decrease in concentration was exponential with distance; this distribution type and others like it were classified as an "exponential" distribution. The lead and hexachlorbenzene results at each sample location along with the IDW and Kriging interpolations are shown in Figure B-1 and Figure B-2, respectively.

Based on visual analysis of the interpolated results for lead; the IDW and Kriging methods produced similar interpolation results. The mean IDW interpolated surface lead concentration was 47 milligram/kilogram (mg/kg), the mean Kriging interpolated surface lead concentration was 65 mg/kg, and the mean measured data point concentration was 57 mg/kg. Based on visual analysis of the interpolated results for hexachlorobenzene, the IDW and Kriging methods did not produce similar results. While the IDW produced a "bull's eye" effect at several locations, the Kriging produced a smoothed dataset with less influence from higher observations, thus creating much lower concentration than actual at those locations (i.e., under predicting at sampled locations). The mean hexachlorobenzene concentration was 3214 μ g/kg from the IDW interpolated surface, 1700 μ g/kg from the Kriging interpolated surface, and 9397 μ g/kg from the measured data points.

Based on the results of the IDW and Kriging comparisons for lead, its seems either approach would produce a reasonable interpolation for data with a "linear distribution", however based on the results from the IDW and Kriging comparisons for hexachlorobenzene; the smoothing done by the Kriging approach is not well suited for this Site for COPCs demonstrating exponential changes in concentration within neighboring data points. For the reasons mentioned above and for the sake of consistency, IDW was the method selected to

generate the interpolated surfaces needed for the SWAC calculations. An additional practical consideration for selecting IDW over Kriging is the advantage of the IDW algorithms being more fully defined; leading to highly reproducible results. Kriging requires much more professional judgment to define the interpolation and is therefore much more subjective than IDW.

In addition, total PCB Congeners, PCB TEQ (Avian), PCB TEQ (Mammalian), and PCB TEQ (Fish) were interpolated using Kriging and IDW. These COPCs were selected for comparison due to their expected significance in characterizing risk (i.e., risk drivers) to receptors at the Site. Kriging and IDW interpolations for these COPC are shown in Figures B-3 through B-6. A comparison of the Site-wide SWAC based on the two methods is provided in Table B-1. Relative differences between IDW and Kriging range between 14-26% for PCB and PCB TEQ. In all cases, IDW estimates higher SWACs for PCB and PCB TEQ. Generally, the differences are considered acceptable for this type of analysis.

1.1.1 Inverse-Distance Weighting

IDW predicts concentration at a location as a weighted average of the surrounding measurements. The weight applied to each measurement depends on how close it is to the prediction location, with greater weight given to the closer measurements. The basic equation for IDW is:

$$\hat{Z}(s_0) = \sum_{i=1}^{N} \lambda_i Z(s_i)$$
 (B-1)

where:

 $\hat{Z}(s_0)$ = the value being predicted at location s_0

N = the number of measured sample points in the search neighborhood of the location to be estimated

 λi = the weights assigned to measured points in the search neighborhood

 $Z(s_i)$ = the observed value at the location s_i

The formula to determine the weights is:

$$\lambda_{i} = d_{i0}^{-p} / \sum_{j=1}^{N} d_{j0}^{-p}$$
where $\sum_{i=1}^{N} \lambda_{i} = 1$ (B-2)

where:

p = the power parameter that controls how much influence a measurement has on the interpolation

 $di\theta$ = the distance between the predicted location, $s\theta$, and the measured

 d_{i0} = the distance between the predicted location, s_0 , and the observed location s_i within the search neighborhood

As the distance between the predicted location and a measured location increases, the weight of the measured point will decrease exponentially according to the power parameter, *p*. The predicted value is the sum of the product of the data points within the search neighborhood and their assigned weights. The weights for the measured locations are scaled so that their sum is equal to 1.

The search neighborhood is defined using two common methods which are "Nearest neighbors" and "elliptical neighborhood". "Nearest neighbors" incorporates a preset number of data points that are closest to the predicted location into the interpolation and uses a variable search radius to obtain the set number of data points. "Elliptical neighborhood" incorporates all data points within a prescribed elliptical area. The three parameters that define an elliptical neighborhood are:

- 1. Azimuth the orientation of the ellipse
- 2. Minor semiaxis half the width of the ellipse

locations, si

3. Major semiaxis - half the length of the ellipse

The anisotropy ratio is defined as the ratio of the major semiaxis to the minor semiaxis. During interpolation, this parameter governs how much influence the interpolator gives to data points along the orientation of the ellipse relative to data points across the orientation of the ellipse (QEA 2007).

1.1.1.1 IDW Optimization

1.1.1.1.1 Power Parameter

The power parameter controls the significance of surrounding points on the interpolated value. A higher power results in less influence from distant points. The default value used by the ArcMap IDW interpolation tool is 2 (Esri). IDW interpolations of lead and hexachlorobenzene were "previewed²" using power parameters ranging from 0 to 3 at increments of 0.5. The default value of 2 was reasonable for both the lead and hexachlorobenzene IDW interpolations whereby it did not overly smooth the data (large p value) nor create large "bull's eyes" (low p value), and was therefore used for all of the COPC IDW interpolations.

1.1.1.1.2 Search Neighborhood

IDW interpolations were performed for lead and hexachlorobenzene using both an "elliptical neighborhood" and "natural neighbors" search neighborhood. The purpose of using both methods was to assess if there were major differences in the interpolated results based on search neighborhood method. Since the natural neighbor's method found in the ArcMap spatial analyst tool only has a few parameters to select when defining search neighborhood it is an easily reproducible process for performing numerous interpolations. The elliptical neighborhood method is based on correlation between data points and thus requires manual selection of parameter's using the ArcMap Geostatistical Analyst tool. The elliptical neighborhood has to be optimized for each of the 19 COPCs. While using an elliptical search radius provides a more robust analysis of the data correlation it introduces inconsistencies and analyst subjectivity between IDW methodologies for each compound. To test the significance of the difference between the two search neighborhood methods the IDW for lead and hexachlorobenzene were performed using both approaches (nearest neighbors and elliptical) and the results compared.

The parameters selected for the elliptical search neighborhood were based on cross validation results. Cross validation is a technique used to assess the accuracy of an interpolation model. The cross validation method leaves one point out and uses the rest to predict a value at that

² The ArcMap 10 Geostatistical Analyst tool allows a preview of the IDW interpolated surface based on selected parameters.

location. The point is then added back into the dataset, and a different one is removed. This is done for all samples in the dataset and provides pairs of predicted and known values that can be compared to assess the interpolation performance (Esri). The goal of the cross validation is to find the elliptical neighborhood which produces the smallest mean error and root mean squared error. The mean error is the averaged difference between the measured and the predicted values without regard to sign (Esri). The root mean squared error (RMSE) is the difference between predicted and measured values which are each squared and then averaged. Finally, the square root of the average is taken. Since the errors are squared before they are averaged, the RMSE gives a relatively high weight to large errors (Eumetcal). A small mean error and RMSE indicate a good model fit with the actual data, the greater the difference between the mean error and RMSE, the greater the variance in the individual errors. The optimized major range and anisotropic ratios for lead, hexachlorobenzene, PCB congeners, and PCB TEQ are shown in Figure B-7 through Figure B-12.

Parameters available using the spatial analyst tool includes the maximum neighbors and the optional max search distance. For the purposes of this analysis the max search distance was not entered and the maximum neighbors were set to 15. The maximum neighbors value was based on the optimum number of neighbors selected during the elliptical neighborhood analysis and the results of the cross validation.

The results of the lead and hexachlorobenzene search neighborhood comparison are shown in Figure B-13 and Figure B-14, respectively. The visual interpretation of the results for both analyses indicates little difference between the two methods. For the reason mentioned above and for the sake of minimizing inconsistencies in IDW methodologies between COPCs, the nearest neighbors approach was chosen as the appropriate search neighborhood method for the COPC IDW interpolations.

1.1.1.2 Data Transformation

Typically, the distribution for environmental data is skewed; characterized by a few relatively elevated sample results and many low concentration results. However, spatial interpolations produce the most accurate results when performed on normally distributed data. For this reason the data were transformed before interpolation.

1.1.1.2.1 Selecting Transformations for COPC Result Values

In an effort to normalize the data distribution, the data were transformed either by a natural log transformation (equation B-3) or Box-Cox transformation (equation B-4). The transformation changes the original variable (*X*) into the transformed variable (*Y*) (QEA 2007):

$$Y = \ln(X) \text{ for } \lambda = 0 \tag{B-3}$$

$$Y = \frac{X^{\lambda} - 1}{\lambda} \text{ for } \lambda \neq 0$$
 (B-4)

where:

 λ = the transformation parameter

A value for λ was selected independently for each COPC so as to produce transformed data that were approximately normally distributed. Normality was optimized by minimizing skewness, kurtosis, and the standard deviation in the same order of importance. Histograms showing untransformed and transformed data for each COPC are presented in Figure B-15A through Figure B-15Q. Transformation parameters selected for each COPC are presented in Table B-2.

2 SURFACE WEIGHTED AVERAGING

Once the appropriate transformation had been determined, values were transformed according to the values in Table B-1 and interpolated using the Spatial Analysis Tool in ArcGIS. IDW was the selected interpolation method and parameterized using the following setting:

- Search Radius = Variable
- Number of nearest sample points to interpolate = 15
- Cell size = 1 foot

The resulting intermediate raster using the transformed values was translated back into the original scale of measurement using equation B-3 for natural log transformations and B-4 for Box-Cox transformations. The transformation changes the transformed variable (Y) back into the original variable (X) (QEA 2007):

$$X = e^{Y} \tag{B-5}$$

$$X = \left(\lambda Y + 1\right)^{\frac{1}{\lambda}} \tag{B-6}$$

The back-transformation is valid because the transformation (and its inverse) are monotonic and hence preserve the percentile's rankings.

Finally, the Zonal Stats tool in Spatial Analyst was used to calculate summary statistics for desired polygons representing areas of interest (e.g., Site-wide, intertidal zone, etc.). The mean value within the polygon was taken as the SWAC of the area of interest.

3 RESULTS

Rasters of the interpolated values (normal scale) for each COPC in Table B-2 are provided in Figures B-16A through B-16Q. For the purposes of mapping interpolated values the 10th percentile ranges were selected based on professional judgment.

4 REFERENCES

- Anchor QEA, 2010. *Patrick Bayou Sediment and Surface Water Contaminant of Potential Concern Delineation Data Report.* Patrick Bayou Superfund Site, Deer Park, Texas.
- Anchor QEA, 2012. *Upstream Patrick Bayou Characterization Data Report.* Patrick Bayou Superfund Site, Deer Park, Texas.
- Esri, (n.d.) ArcMap Version 10: ArcMap Help Geostatistical Analyst. Retrieved February 1, 2012.
- Eumetcal, (n.d.). *Mean absolute error (MAE) and root mean squared error (RMSE)*. Retrieved February 3, 2012, from Eumetcal.org:

 http://www.eumetcal.org/resources/ukmeteocal/verification/www/english/msg/ver_c ont_var/uos3/uos3_ko1.htm
- QEA, 2007. Phase 2 dredge area delineation report: Hudson River PCBs Superfund Site.

TABLES

Table B-1
Comparison of IDW and Kriging SWACs for PCBs

СОРС	Units	IDW SWAC	Kriging SWAC
Total PCB Congeners	mg/kg	7.87	6.69
PCB TEQ (Fish)	μg/kg	0.011	0.0085
PCB TEQ (Avian)	μg/kg	2.08	1.68
PCB TEQ (Mammals)	μg/kg	0.11	0.096


Notes:

IDW – Inverse distance weighting SWAC – Surface weighted average mg/kg - milligrams per kilogram µg/kg - micrograms per kilogram


Table B-2
Transformation Parameters Selected for Each COPC

Chemical Name	Transformation Type	Transformation Parameter	
1,3-Dichlorobenzene	Box-Cox	(-) 0.06	
1,4-Dichlorobenzene	Box-Cox	(-) 0.1	
Bis(2-ethylhexyl) phthalate	Natural Log	0	
Hexachlorobenzene	Box-Cox	(-) 0.1	
Hexachlorobutadiene	Box-Cox	(-) 0.1	
Lead	Box-Cox	(-) 0.3	
Mercury	Natural Log	0	
Tetrachlorobiphenyls (total)	Box-Cox	(-) 0.09	
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	Box-Cox	0.13	
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	Box-Cox	0.11	
Total LPAH (6 of 15) (U = 1/2)	Box-Cox	(-) 0.12	
Total PAH (15) (U = 1/2)	Box-Cox	(-) 0.06	
Total PCB Congener (U = 1/2)	Natural Log	0	
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	Natural Log	0	
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	Box-Cox	(-) 0.05	
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	Natural Log	0	
Trichlorobiphenyls (total)	Box-Cox	(-) 0.08	


FIGURES
























































Figure B-10






Figure B-11








Figure B-12













Patrick Bayou Superfund Site Remedial Alternatives, Deer Park, Texas





































































Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas











































































Station numbers from Patrick Bayou PSCR indicate length along channel in hundreds of feet. Aerial orthoimagery: Bing Maps 2010


























Feet

550

825








Feet


550

825














Feet

550

825











































APPENDIX C WILDLIFE RISK ASSESSMENT SUPPORTING INFORMATION

APPENDIX C-1 TISSUE DATA ANALYSIS

APPENDIX C-1.1 PREY GROUP ANALYSIS

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Hexachlorobenzene Size_Class_Code=FISH (<15 CM)



Oneway Anova Summary of Fit

Rsquare	0.173362
Adj Rsquare	0.099883
Root Mean Square Error	0.572538
Mean of Response	6.028825
Observations (or Sum Wgts)	50

Analysis of Variance


Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	4	3.093561	0.773390	2.3593	0.0676
Error	45	14.750992	0.327800		
C. Total	49	17.844553			

Means for Oneway Anova


Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Gulf killifish	25	5.84569	0.11451	5.6151	6.0763
Gulf menhaden	10	6.06552	0.18105	5.7009	6.4302
Pinfish	4	6.42302	0.28627	5.8464	6.9996
Sand seatrout	2	5.63223	0.40485	4.8168	6.4476
Striped mullet	9	6.40969	0.19085	6.0253	6.7941

Std Error uses a pooled estimate of error variance


Oneway Analysis of LN Adj Result By Common_Name chemical_name=Hexachlorobenzene Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Lead Size_Class_Code=INVERTEBRATES(2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Mercury Size_Class_Code=FISH (<15 CM)



Oneway Anova Summary of Fit

Rsquare	0.392097
Adj Rsquare	0.338062
Root Mean Square Error	0.448351
Mean of Response	-2.90355
Observations (or Sum Wgts)	50

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	4	5.834577	1.45864	7.2563	0.0001*
Error	45	9.045852	0.20102		
C. Total	49	14.880429			

Means for Oneway Anova

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Gulf killifish	25	-3.0420	0.08967	-3.223	-2.861
Gulf menhaden	10	-2.2332	0.14178	-2.519	-1.948
Pinfish	4	-3.2664	0.22418	-3.718	-2.815
Sand seatrout	2	-2.9329	0.31703	-3.571	-2.294
Striped mullet	9	-3.0959	0.14945	-3.397	-2.795

Std Error uses a pooled estimate of error variance

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

	pa
q*	Alpha
2.84145	0.05

Abs(Dif)-HSD	Gulf menhaden	Sand seatrout	Gulf killifish	Striped mullet	Pinfish
Gulf menhaden	-0.5697	-0.2872	0.3322	0.2774	0.2795
Sand seatrout	-0.2872	-1.2740	-0.8270	-0.8329	-0.7697
Gulf killifish	0.3322	-0.8270	-0.3603	-0.4414	-0.4617
Striped mullet	0.2774	-0.8329	-0.4414	-0.6006	-0.5950
Pinfish	0.2795	-0.7697	-0.4617	-0.5950	-0.9008


Positive values show pairs of means that are significantly different.

Level			Mean
Gulf menhaden	Α		-2.233204
Sand seatrout	Α	В	-2.932857
Gulf killifish		В	-3.042037
Striped mullet		В	-3.095909
Pinfish		В	-3.266437


Levels not connected by same letter are significantly different.

Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Gulf menhaden	Pinfish	1.033232	0.2652482	0.279542	1.786922	0.0028*
Gulf menhaden	Striped mullet	0.862704	0.2060032	0.277356	1.448052	0.0012*
Gulf menhaden	Gulf killifish	0.808832	0.1677577	0.332157	1.285508	0.0002*
Gulf menhaden	Sand seatrout	0.699652	0.3472915	-0.287160	1.686464	0.2758
Sand seatrout	Pinfish	0.333580	0.3882837	-0.769709	1.436869	0.9102
Gulf killifish	Pinfish	0.224400	0.2414446	-0.461653	0.910453	0.8840
Striped mullet	Pinfish	0.170528	0.2694256	-0.595032	0.936088	0.9689
Sand seatrout	Striped mullet	0.163052	0.3504924	-0.832855	1.158959	0.9901
Sand seatrout	Gulf killifish	0.109180	0.3294696	-0.826992	1.045352	0.9973
Gulf killifish	Striped mullet	0.053872	0.1742877	-0.441358	0.549102	0.9979


Oneway Analysis of LN Adj Result By Common_Name chemical_name=Mercury Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)




Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2), Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PAH (16) (U = 1/2), Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener (U = 1/2), Size_Class_Code=FISH (<15 CM)



Oneway Anova Summary of Fit

Rsquare	0.533399
Adj Rsquare	0.491923
Root Mean Square Error	0.392759
Mean of Response	15.03682
Observations (or Sum Wgts)	50

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	4	7.935436	1.98386	12.8605	<.0001*
Error	45	6.941687	0.15426		
C. Total	49	14.877123			

Means for Oneway Anova

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Gulf killifish	25	15.4195	0.07855	15.261	15.578
Gulf menhaden	10	14.7295	0.12420	14.479	14.980
Pinfish	4	14.9119	0.19638	14.516	15.307
Sand seatrout	2	14.4614	0.27772	13.902	15.021
Striped mullet	9	14.4988	0.13092	14.235	14.762

Std Error uses a pooled estimate of error variance

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.84145	0.05

Abs(Dif)-HSD	Gulf killifish	Pinfish	Gulf menhaden	Striped mullet	Sand seatrout
Gulf killifish	-0.3157	-0.0934	0.2724	0.4869	0.1379
Pinfish	-0.0934	-0.7891	-0.4778	-0.2575	-0.5160
Gulf menhaden	0.2724	-0.4778	-0.4991	-0.2821	-0.5964
Striped mullet	0.4869	-0.2575	-0.2821	-0.5261	-0.8351
Sand seatrout	0.1379	-0.5160	-0.5964	-0.8351	-1.1160


Positive values show pairs of means that are significantly different.

Level			Mean
Gulf killifish	Α		15.419482
Pinfish	Α	В	14.911909
Gulf menhaden		В	14.729462
Striped mullet		В	14.498772
Sand seatrout		В	14.461441


Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Gulf killifish	Sand seatrout	0.9580408	0.2886178	0.137947	1.778135	0.0147*
Gulf killifish	Striped mullet	0.9207098	0.1526773	0.486885	1.354535	<.0001*
Gulf killifish	Gulf menhaden	0.6900200	0.1469570	0.272449	1.107591	0.0002*
Gulf killifish	Pinfish	0.5075732	0.2115072	-0.093415	1.108561	0.1341
Pinfish	Sand seatrout	0.4504677	0.3401394	-0.516022	1.416957	0.6777
Pinfish	Striped mullet	0.4131366	0.2360188	-0.257500	1.083773	0.4146
Gulf menhaden	Sand seatrout	0.2680208	0.3042299	-0.596434	1.132475	0.9024
Gulf menhaden	Striped mullet	0.2306898	0.1804604	-0.282080	0.743459	0.7056
Pinfish	Gulf menhaden	0.1824468	0.2323594	-0.477791	0.842685	0.9336

Level- LevelDifferenceStd Err DifLower CLUpper CLp-Value DifferenceStriped mulletSand seatrout0.03733100.3070339-0.8350910.9097530.9999

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener (U = 1/2), Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener TEQ 1998 (Avian) (U = 1/2), Size_Class_Code=FISH (<15 CM)



Oneway Anova Summary of Fit

Rsquare	0.71961
Adj Rsquare	0.694686
Root Mean Square Error	0.402767
Mean of Response	6.129779
Observations (or Sum Wgts)	50

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	4	18.735038	4.68376	28.8726	<.0001*
Error	45	7.299966	0.16222		
C. Total	49	26.035004			

Means for Oneway Anova

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Gulf killifish	25	6.63478	0.08055	6.4725	6.7970
Gulf menhaden	10	6.02177	0.12737	5.7652	6.2783
Pinfish	4	4.77433	0.20138	4.3687	5.1799
Sand seatrout	2	4.79363	0.28480	4.2200	5.3672
Striped mullet	9	5.74635	0.13426	5.4759	6.0168

Std Error uses a pooled estimate of error variance

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.84145	0.05

Abs(Dif)-HSD	Gulf killifish	Gulf menhaden	Striped mullet	Sand seatrout	Pinfish
Gulf killifish	-0.3237	0.1848	0.4436	1.0002	1.2441
Gulf menhaden	0.1848	-0.5118	-0.2504	0.3417	0.5704
Striped mullet	0.4436	-0.2504	-0.5395	0.0581	0.2843
Sand seatrout	1.0002	0.3417	0.0581	-1.1444	-0.9718
Pinfish	1.2441	0.5704	0.2843	-0.9718	-0.8092


Positive values show pairs of means that are significantly different.

Level			Mean
Gulf killifish	Α		6.6347809
Gulf menhaden	В		6.0217726
Striped mullet	В		5.7463480
Sand seatrout		С	4.7936258
Pinfish		С	4.7743339


Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Gulf killifish	Pinfish	1.860447	0.2168968	1.24415	2.476749	<.0001*
Gulf killifish	Sand seatrout	1.841155	0.2959723	1.00016	2.682146	<.0001*
Gulf menhaden	Pinfish	1.247439	0.2382803	0.57038	1.924501	<.0001*
Gulf menhaden	Sand seatrout	1.228147	0.3119822	0.34166	2.114629	0.0025*
Striped mullet	Pinfish	0.972014	0.2420330	0.28429	1.659739	0.0020*
Striped mullet	Sand seatrout	0.952722	0.3148577	0.05807	1.847375	0.0317*
Gulf killifish	Striped mullet	0.888433	0.1565678	0.44355	1.333313	<.0001*
Gulf killifish	Gulf menhaden	0.613008	0.1507017	0.18480	1.041220	0.0017*
Gulf menhaden	Striped mullet	0.275425	0.1850588	-0.25041	0.801260	0.5752

Level- LevelDifferenceStd Err DifLower CLUpper CLp-Value DifferenceSand seatroutPinfish0.0192920.3488067-0.971831.0104091.0000

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener TEQ 1998 (Avian) (U = 1/2), Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener TEQ 2005 (Mammal) (U = 1/2), Size_Class_Code=FISH (<15 CM)



Oneway Anova Summary of Fit

Rsquare	0.576592
Adj Rsquare	0.538956
Root Mean Square Error	0.567592
Mean of Response	3.689824
Observations (or Sum Wgts)	50

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	4	19.742183	4.93555	15.3201	<.0001*
Error	45	14.497244	0.32216		
C. Total	49	34.239427			

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Gulf killifish	25	4.26055	0.11352	4.0319	4.4892

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Gulf menhaden	10	3.36342	0.17949	3.0019	3.7249
Pinfish	4	3.55015	0.28380	2.9786	4.1217
Sand seatrout	2	2.27174	0.40135	1.4634	3.0801
Striped mullet	9	2.84434	0.18920	2.4633	3.2254

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.84145	0.05


Abs(Dif)-HSD	Gulf killifish	Pinfish	Gulf menhaden	Striped mullet	Sand seatrout
Gulf killifish	-0.4562	-0.1581	0.2937	0.7893	0.8037
Pinfish	-0.1581	-1.1404	-0.7674	-0.2634	-0.1183
Gulf menhaden	0.2937	-0.7674	-0.7213	-0.2219	-0.1576
Striped mullet	0.7893	-0.2634	-0.2219	-0.7603	-0.6882
Sand seatrout	0.8037	-0.1183	-0.1576	-0.6882	-1.6128

Positive values show pairs of means that are significantly different.


Level			Mean
Gulf killifish	Α		4.2605524
Pinfish	Α	В	3.5501546
Gulf menhaden		В	3.3634207
Striped mullet		В	2.8443424
Sand seatrout		В	2.2717378

Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Gulf killifish	Sand seatrout	1.988815	0.4170934	0.803664	3.173966	0.0002*
Gulf killifish	Striped mullet	1.416210	0.2206402	0.789271	2.043149	<.0001*
Pinfish	Sand seatrout	1.278417	0.4915493	-0.118297	2.675131	0.0873
Gulf menhaden	Sand seatrout	1.091683	0.4396551	-0.157576	2.340942	0.1127
Gulf killifish	Gulf menhaden	0.897132	0.2123736	0.293682	1.500581	0.0010*
Gulf killifish	Pinfish	0.710398	0.3056578	-0.158114	1.578910	0.1562
Pinfish	Striped mullet	0.705812	0.3410805	-0.263352	1.674976	0.2511
Striped mullet	Sand seatrout	0.572605	0.4437073	-0.688168	1.833378	0.6983
Gulf menhaden	Striped mullet	0.519078	0.2607906	-0.221946	1.260102	0.2871
Pinfish	Gulf menhaden	0.186734	0.3357921	-0.767403	1.140871	0.9806

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener TEQ 2005 (Mammal) (U = 1/2), Size_Class_Code=INVERTEBRATES (2 - 7.5 CM)



Oneway Analysis of LN Adj Result By Common_Name chemical_name=Hexachlorobenzene Size_Class_Group_Code=Invertebrates (2 - 12.5 cm)



Oneway Anova Summary of Fit

Rsquare	0.205299
Adj Rsquare	0.123089
Root Mean Square Error	0.687622
Mean of Response	6.012888
Observations (or Sum Wgts)	33

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	3	3.542270	1.18076	2.4972	0.0794
Error	29	13.711898	0.47282		
C. Total	32	17 254169			

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Blue crab	21	5.78609	0.15005	5.4792	6.0930
Brown shrimp	8	6.55107	0.24311	6.0539	7.0483

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Oyster	1	5.86363	0.68762	4.4573	7.2700
White shrimp	3	6.21507	0.39700	5.4031	7.0270

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.72451	0.05


Abs(Dif)-HSD	Brown shrimp	White shrimp	Oyster	Blue crab
Brown shrimp	-0.9367	-0.9323	-1.2996	-0.0134
White shrimp	-0.9323	-1.5297	-1.8118	-0.7273
Oyster	-1.2996	-1.8118	-2.6494	-1.8400
Blue crab	-0.0134	-0.7273	-1.8400	-0.5782

Positive values show pairs of means that are significantly different.

Level		Mean
Brown shrimp	Α	6.5510750
White shrimp	Α	6.2150725
Oyster	Α	5.8636312
Blue crab	Α	5.7860888

Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Brown shrimp	Blue crab	0.7649862	0.2856894	-0.01338	1.543350	0.0554
Brown shrimp	Oyster	0.6874438	0.7293333	-1.29963	2.674520	0.7823
White shrimp	Blue crab	0.4289837	0.4244095	-0.72732	1.585292	0.7445
White shrimp	Oyster	0.3514413	0.7939975	-1.81181	2.514696	0.9705
Brown shrimp	White shrimp	0.3360024	0.4655223	-0.93232	1.604323	0.8876
Oyster	Blue crab	0.0775424	0.7038036	-1.83998	1.995062	0.9995

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Mercury Size_Class_Group_Code=Invertebrates (2 - 12.5 cm)



Oneway Anova Summary of Fit

Rsquare	0.052858
Adj Rsquare	-0.04512
Root Mean Square Error	0.42943
Mean of Response	-3.13156
Observations (or Sum Wgts)	33

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	3	0.2984544	0.099485	0.5395	0.6590
Error	29	5.3478946	0.184410		
C. Total	32	5 6463490			

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Blue crab	21	-3.0728	0.09371	-3.264	-2.881
Brown shrimp	8	-3.2277	0.15183	-3.538	-2.917

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Oyster	1	-2.9759	0.42943	-3.854	-2.098
White shrimp	3	-3.3380	0.24793	-3.845	-2.831

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.72451	0.05


Abs(Dif)-HSD	Oyster	Blue crab	Brown shrimp	White shrimp
Oyster	-1.6546	-1.1006	-0.9891	-0.9889
Blue crab	-1.1006	-0.3611	-0.3312	-0.4570
Brown shrimp	-0.9891	-0.3312	-0.5850	-0.6818
White shrimp	-0.9889	-0.4570	-0.6818	-0.9553

Positive values show pairs of means that are significantly different.

Level		Mean
Oyster	Α	-2.975930
Blue crab	Α	-3.072841
Brown shrimp	Α	-3.227740
White shrimp	Α	-3.337976

Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Oyster	White shrimp	0.3620462	0.4958631	-0.98894	1.713030	0.8842
Blue crab	White shrimp	0.2651348	0.2650500	-0.45700	0.987266	0.7504
Oyster	Brown shrimp	0.2518104	0.4554793	-0.98915	1.492768	0.9450
Blue crab	Brown shrimp	0.1548990	0.1784172	-0.33120	0.640999	0.8211
Brown shrimp	White shrimp	0.1102359	0.2907255	-0.68185	0.902320	0.9811
Oyster	Blue crab	0.0969114	0.4395357	-1.10061	1.294431	0.9961

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener (U = 1/2) Size_Class_Group_Code=Invertebrates (2 - 12.5 cm)



Oneway Anova Summary of Fit

Rsquare	0.588632
Adj Rsquare	0.546076
Root Mean Square Error	0.372565
Mean of Response	13.62859
Observations (or Sum Wgts)	33

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	3	5.7599054	1.91997	13.8321	<.0001*
Error	29	4.0253392	0.13880		
C. Total	32	9.7852446			

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Blue crab	21	13.3532	0.08130	13.187	13.519
Brown shrimp	8	14.3214	0.13172	14.052	14.591

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Oyster	1	14.1728	0.37257	13.411	14.935
White shrimp	3	13.5275	0.21510	13.088	13.967

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.72451	0.05


Abs(Dif)-HSD	Brown shrimp	Oyster	White shrimp	Blue crab
Brown shrimp	-0.5075	-0.9280	0.1068	0.5465
Oyster	-0.9280	-1.4355	-0.5268	-0.2193
White shrimp	0.1068	-0.5268	-0.8288	-0.4522
Blue crab	0.5465	-0.2193	-0.4522	-0.3133

Positive values show pairs of means that are significantly different.

Level			Mean
Brown shrimp	Α		14.321430
Oyster	Α	В	14.172793
White shrimp		В	13.527474
Blue crab		В	13.353181

Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Brown shrimp	Blue crab	0.9682494	0.1547913	0.546519	1.389980	<.0001*
Oyster	Blue crab	0.8196116	0.3813326	-0.219333	1.858556	0.1617
Brown shrimp	White shrimp	0.7939567	0.2522278	0.106759	1.481154	0.0188*
Oyster	White shrimp	0.6453189	0.4302012	-0.526769	1.817406	0.4505
White shrimp	Blue crab	0.1742927	0.2299522	-0.452214	0.800800	0.8725
Brown shrimp	Oyster	0.1486378	0.3951650	-0.927993	1.225269	0.9815

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener TEQ 1998 (Avian) (U = 1/2) Size_Class_Group_Code=Invertebrates (2 - 12.5 cm)



Oneway Anova Summary of Fit

Rsquare	0.239998
Adj Rsquare	0.161377
Root Mean Square Error	0.41256
Mean of Response	5.914452
Observations (or Sum Wgts)	33

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	3	1.5587055	0.519568	3.0526	0.0442*
Error	29	4.9359585	0.170205		
C Total	32	6.4946640			

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Blue crab	21	5.84026	0.09003	5.6561	6.0244
Brown shrimp	8	6.26186	0.14586	5.9635	6.5602

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Oyster	1	5.88946	0.41256	5.0457	6.7332
White shrimp	3	5.51572	0.23819	5.0286	6.0029

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.72451	0.05


Abs(Dif)-HSD	Brown shrimp	Oyster	Blue crab	White shrimp
Brown shrimp	-0.5620	-0.8198	-0.0454	-0.0148
Oyster	-0.8198	-1.5896	-1.1013	-0.9242
Blue crab	-0.0454	-1.1013	-0.3469	-0.3692
White shrimp	-0.0148	-0.9242	-0.3692	-0.9178

Positive values show pairs of means that are significantly different.

Level		Mean
Brown shrimp	Α	6.2618561
Oyster	Α	5.8894595
Blue crab	Α	5.8402594
White shrimp	Α	5.5157170

Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Brown shrimp	White shrimp	0.7461390	0.2793042	-0.01483	1.507106	0.0562
Brown shrimp	Blue crab	0.4215967	0.1714080	-0.04541	0.888600	0.0881
Oyster	White shrimp	0.3737425	0.4763829	-0.92417	1.671652	0.8608
Brown shrimp	Oyster	0.3723965	0.4375856	-0.81981	1.564603	0.8296
Blue crab	White shrimp	0.3245424	0.2546373	-0.36922	1.018304	0.5860
Oyster	Blue crab	0.0492001	0.4222683	-1.10127	1.199674	0.9994

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total PCB Congener TEQ 2005 (Mammal) (U = 1/2) Size_Class_Group_Code=Invertebrates (2 - 12.5 cm)



Oneway Anova Summary of Fit

Rsquare	0.109824
Adj Rsquare	0.017737
Root Mean Square Error	0.458341
Mean of Response	3.113981
Observations (or Sum Wgts)	33

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Ratio	Prob > F
Common_Name	3	0.7516149	0.250538	1.1926	0.3299
Error	29	6.0922056	0.210076		
C Total	32	6 8438204			

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Blue crab	21	3.17924	0.10002	2.9747	3.3838
Brown shrimp	8	3.12799	0.16205	2.7966	3.4594

Level	Number	Mean	Std Error	Lower 95%	Upper 95%
Oyster	1	3.03206	0.45834	2.0947	3.9695
White shrimp	3	2.64711	0.26462	2.1059	3.1883

Means Comparisons

Comparisons for all pairs using Tukey-Kramer HSD

q*	Alpha
2.72451	0.05


Abs(Dif)-HSD	Blue crab	Brown shrimp	Oyster	White shrimp
Blue crab	-0.3854	-0.4676	-1.1310	-0.2386
Brown shrimp	-0.4676	-0.6244	-1.2286	-0.3645
Oyster	-1.1310	-1.2286	-1.7660	-1.0570
White shrimp	-0.2386	-0.3645	-1.0570	-1.0196

Positive values show pairs of means that are significantly different.


Level		Mean
Blue crab	Α	3.1792398
Brown shrimp	Α	3.1279933
Oyster	Α	3.0320642
White shrimp	Α	2.6471117


Level	- Level	Difference	Std Err Dif	Lower CL	Upper CL	p-Value Difference
Blue crab	White shrimp	0.5321281	0.2828939	-0.23862	1.302875	0.2581
Brown shrimp	White shrimp	0.4808817	0.3102980	-0.36453	1.326292	0.4222
Oyster	White shrimp	0.3849525	0.5292461	-1.05698	1.826889	0.8853
Blue crab	Oyster	0.1471756	0.4691265	-1.13096	1.425316	0.9891
Brown shrimp	Oyster	0.0959291	0.4861436	-1.22857	1.420432	0.9972
Blue crab	Brown shrimp	0.0512464	0.1904288	-0.46758	0.570072	0.9930


APPENDIX C-1.2 SPATIAL ANALYSIS OF TISSUE DATA





*Non-detects graphed at 1/2 the detection limit

















*Non-detects graphed at 1/2 the detection limit





*Non-detects graphed at 1/2 the detection limit





*Non-detects graphed at 1/2 the detection limit





































*Non-detects graphed at 1/2 the detection limit























































*Non-detects graphed at 1/2 the detection limit























































*Non-detects graphed at 1/2 the detection limit



































































*Non-detects graphed at 1/2 the detection limit
































APPENDIX C-2 OUTPUTS OF 95 UCL CALCULATIONS FOR TISSUE DATA

GROUPING: BLUE CRAB, BROWN SHRIMP, WHITE SHRIMP, OYSTER

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

0	Number of Detected Data	33	Number of Valid Data
33	Number of Non-Detect Data	0	Number of Distinct Detected Data
100.00%	Percent Non-Detects		

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The dataset for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

G	eneral Statis	tics	
Number of Valid Data	33	Number of Detected Data	20
Number of Distinct Detected Data	20	Number of Non-Detect Data	13
		Percent Non-Detects	39.39%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	191	Minimum Detected	5.252
Maximum Detected	5430	Maximum Detected	8.6
Mean of Detected	821	Mean of Detected	6.401
SD of Detected	1104	SD of Detected	0.66
Minimum Non-Detect	347	Minimum Non-Detect	5.849
Maximum Non-Detect	748	Maximum Non-Detect	6.617
Note: Data have multiple DLs. Llcs of VM Method is recommended	lad	Number treated as Non-Detect	28
Note: Data have multiple DLs - Use of KM Method is recommend For all methods (except KM, DL/2, and ROS Methods),	ieu	Number treated as Non-Detect Number treated as Detected	20 5
• • • • • • • • • • • • • • • • • • • •			84.85%
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	64.65%
	UCL Statistic	s	
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values C	nly
Shapiro Wilk Test Statistic	0.407	Shapiro Wilk Test Statistic	0.839
5% Shapiro Wilk Critical Value	0.905	5% Shapiro Wilk Critical Value	0.905
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	591.1	Mean	6.013
SD	900.2	SD	0.734
95% DL/2 (t) UCL	856.5	95% H-Stat (DL/2) UCL	707.1
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE yields a negative mean	,	Mean in Log Scale	6.094
,		SD in Log Scale	0.652
		Mean in Original Scale	608.6
		SD in Original Scale	892.5
		95% t UCL	871.8
		95% Percentile Bootstrap UCL	904.4
		95% BCA Bootstrap UCL	1088
		95% H-UCL	695.2
Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	/

1.533

Data do not follow a Discernable Distribution (0.05)

k star (bias corrected)

Theta Star	535.4		
nu star	61.33		
A-D Test Statistic	2.019	Nonparametric Statistics	
5% A-D Critical Value	0.755	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.755	Mean	621
5% K-S Critical Value	0.197	SD	877.2
Data not Gamma Distributed at 5% Significance Lev	el	SE of Mean	157.7
		95% KM (t) UCL	888
Assuming Gamma Distribution		95% KM (z) UCL	880.3
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	886.5
Minimum	0.000001	95% KM (bootstrap t) UCL	1415
Maximum	5430	95% KM (BCA) UCL	975.7
Mean	546.8	95% KM (Percentile Bootstrap) UCL	922.2
Median	413.5	95% KM (Chebyshev) UCL	1308
SD	925.8	97.5% KM (Chebyshev) UCL	1606
k star	0.16	99% KM (Chebyshev) UCL	2190
Theta star	3422		
Nu star	10.55	Potential UCLs to Use	
AppChi2	4.285	95% KM (BCA) UCL	975.7
95% Gamma Approximate UCL (Use when n >= 40)	1346		
95% Adjusted Gamma UCL (Use when n < 40)	1413		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 33	Number of Distinct Observations 24
Number of Valid Observations 55	Number of Distinct Observations 2

Raw Statistics	Log-transformed Statistics
Minimum 0.009	Minimum of Log Data -4.711
Maximum 0.083	Maximum of Log Data -2.489
Mean 0.0468	Mean of log Data -3.132
Geometric Mean 0.0436	SD of log Data 0.42
Median 0.046	
SD 0.0158	
Std. Error of Mean 0.00275	

Relevant UCL Statistics

Coefficient of Variation 0.338

Skewness -0.0016

Relevant UCL Statistics	
Normal Distribution Test	Lognormal Distribution Test
Shapiro Wilk Test Statistic 0.989	Shapiro Wilk Test Statistic 0.884
Shapiro Wilk Critical Value 0.931	Shapiro Wilk Critical Value 0.931
Data appear Normal at 5% Significance Level	Data not Lognormal at 5% Significance Level
Assuming Normal Distribution	Assuming Lognormal Distribution
95% Student's-t UCL 0.0515	95% H-UCL 0.0548
95% UCLs (Adjusted for Skewness)	95% Chebyshev (MVUE) UCL 0.0632
95% Adjusted-CLT UCL (Chen-1995) 0.0514	97.5% Chebyshev (MVUE) UCL 0.07
95% Modified-t UCL (Johnson-1978) 0.0515	99% Chebyshev (MVUE) UCL 0.0833

Gamma Distribution Test	Data Distribution
k star (bias corrected) 6.595	Data appear Normal at 5% Significance Level
Theta Star 0.0071	
MLE of Mean 0.0468	
MLE of Standard Deviation 0.0182	
nu star 435.3	
Approximate Chi Square Value (.05) 387.9	Nonparametric Statistics
Adjusted Level of Significance 0.0419	95% CLT UCL 0.0514
Adjusted Chi Square Value 385.6	95% Jackknife UCL 0.0515

Anderson-Darling Test Statistic 0.478
Anderson-Darling 5% Critical Value 0.748
Kolmogorov-Smirnov Test Statistic 0.143
Kolmogorov-Smirnov 5% Critical Value 0.153
Data appear Gamma Distributed at 5% Significance Level

95% Standard Bootstrap UCL 0.0514 95% Bootstrap-t UCL 0.0515 95% Hall's Bootstrap UCL 0.0517 95% Percentile Bootstrap UCL 0.0514 95% BCA Bootstrap UCL 0.0514 95% Chebyshev(Mean, Sd) UCL 0.0589 97.5% Chebyshev(Mean, Sd) UCL 0.0641 99% Chebyshev(Mean, Sd) UCL 0.0743

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0526 95% Adjusted Gamma UCL (Use when n < 40) 0.0529

Potential UCL to Use

Use 95% Student's-t UCL 0.0515

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed datasets.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 33 Number of Distinct Observations 33

Raw Statistics

Minimum 97.57 Maximum 782.9 Mean 404.3 Geometric Mean 370.4 Median 369.9 SD 160.9 Std. Error of Mean 28 Coefficient of Variation 0.398

Skewness 0.406

Log-transformed Statistics

Minimum of Log Data 4.581 Maximum of Log Data 6.663 Mean of log Data 5.914 SD of log Data 0.451

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.974 Shapiro Wilk Critical Value 0.931 Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.95 Shapiro Wilk Critical Value 0.931

at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution 95% Student's-t UCL 451.7

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 452.5 95% Modified-t UCL (Johnson-1978) 452.1

Assuming Lognormal Distribution

95% H-UCL 476.8 95% Chebyshev (MVUE) UCL 553.4 97.5% Chebyshev (MVUE) UCL 616.1 99% Chebyshev (MVUE) UCL 739.2

Gamma Distribution Test

Theta Star 75.57 MLE of Mean 404.3 MLE of Standard Deviation 174.8 nu star 353.1

k star (bias corrected) 5.35

Data Distribution
Data appear Normal at 5% Significance Level

Approximate Chi Square Value (.05) 310.5 Adjusted Level of Significance 0.0419

djusted Level of Significance 0.0419 Adjusted Chi Square Value 308.5

Nonparametric Statistics

95% CLT UCL 450.4 95% Jackknife UCL 451.7 95% Standard Bootstrap UCL 449.6 95% Bootstrap-t UCL 454.5 95% Hall's Bootstrap UCL 455.1 95% Percentile Bootstrap UCL 449.9 95% BCA Bootstrap UCL 450.6 95% Chebyshev(Mean, Sd) UCL 526.4

Anderson-Darling Test Statistic 0.251 Anderson-Darling 5% Critical Value 0.748 Kolmogorov-Smirnov Test Statistic 0.0852 Kolmogorov-Smirnov 5% Critical Value 0.153

Data appear Gamma Distributed at 5% Significance Level

97.5% Chebyshev(Mean, Sd) UCL 579.2 99% Chebyshev(Mean, Sd) UCL 682.9

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 459.7 95% Adjusted Gamma UCL (Use when n < 40) 462.7

Potential UCL to Use

Use 95% Student's-t UCL 451.7

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 33 Number of Distinct Observations 33

Raw Statistics

Minimum 8.2 Maximum 49.2 Mean 24.77 Geometric Mean 22.51 Median 23.2 SD 10.51 Std. Error of Mean 1.83 Coefficient of Variation 0.424

Skewness 0.508

Log-transformed Statistics

Minimum of Log Data 2.104 Maximum of Log Data 3.896 Mean of log Data 3.114 SD of log Data 0.462

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.959 Shapiro Wilk Critical Value 0.931

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.955 Shapiro Wilk Critical Value 0.931

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 27.87 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 27.95

95% Modified-t UCL (Johnson-1978) 27.9

Assuming Lognormal Distribution

95% H-UCL 29.28 95% Chebyshev (MVUE) UCL 34.06 97.5% Chebyshev (MVUE) UCL 38 99% Chebyshev (MVUE) UCL 45.74

Gamma Distribution Test

k star (bias corrected) 4.919 Theta Star 5.036 MLE of Mean 24.77

MLE of Standard Deviation 11.17

nu star 324.6

Approximate Chi Square Value (.05) 283.9

Adjusted Level of Significance 0.0419

Adjusted Chi Square Value 281.9 Anderson-Darling Test Statistic 0.241

Anderson-Darling 5% Critical Value 0.748 Kolmogorov-Smirnov Test Statistic 0.0941 Kolmogorov-Smirnov 5% Critical Value 0.154

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 28.33 95% Adjusted Gamma UCL (Use when n < 40) 28.52 Data appear Normal at 5% Significance Level

Data Distribution

Nonparametric Statistics

95% CLT UCL 27.78 95% Jackknife UCL 27.87 95% Standard Bootstrap UCL 27.66 95% Bootstrap-t UCL 28.04 95% Hall's Bootstrap UCL 27.99 95% Percentile Bootstrap UCL 27.86 95% BCA Bootstrap UCL 27.82 95% Chebyshev(Mean, Sd) UCL 32.75

97.5% Chebyshev(Mean, Sd) UCL 36.2 99% Chebyshev(Mean, Sd) UCL 42.97

Potential UCL to Use

Use 95% Student's-t UCL 27.87

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: BLUE CRAB, WHITE SHRIMP, OYSTER

Blue Crab, White Shrimp, Oyster

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1_a.wst Full Precision

Confidence Coefficient 95% Number of Bootstrap Operations 2000

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 25

Raw Statistics

Minimum 228677 Maximum 1429439 Mean 717711 Geometric Mean 664560 Median 695219

SD 279856 Std. Error of Mean 55971 Coefficient of Variation 0.39 Skewness 0.731

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.958 Shapiro Wilk Test Statistic 0.969 Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918 Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Log-transformed Statistics

Assuming Lognormal Distribution

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% Chebyshev (MVUE) UCL 989827

97.5% Chebyshev (MVUE) UCL 1105919

99% Chebyshev (MVUE) UCL 1333960

95% Standard Bootstrap UCL 807051

95% Percentile Bootstrap UCL 812766

95% Chebyshev(Mean, Sd) UCL 961684 97.5% Chebyshev(Mean, Sd) UCL 1067251

99% Chebyshev (Mean, Sd) UCL 1274617

95% Bootstrap-t UCL 822860

95% Hall's Bootstrap UCL 828971

95% BCA Bootstrap UCL 811262

Minimum of Log Data 12.34

Maximum of Log Data 14.17

Mean of log Data 13.41

SD of log Data 0.415

95% H-UCL 849946

95% CLT UCL 809775 95% Jackknife UCL 813471

Assuming Normal Distribution

95% Student's-t UCL 813471 95% UCLs (Adjusted for Skewness)

> 95% Adjusted-CLT UCL (Chen-1995) 818523 95% Modified-t UCL (Johnson-1978) 814836

Gamma Distribution Test

k star (bias corrected) 5.888 Theta Star 121893 MLF of Mean 717711

MLE of Standard Deviation 295777 nu star 294.4

Approximate Chi Square Value (.05) 255.7 Adjusted Level of Significance 0.0395

Adjusted Chi Square Value 253.2 Anderson-Darling Test Statistic 0.22 Anderson-Darling 5% Critical Value 0.746

Kolmogorov-Smirnov Test Statistic 0.0918 Kolmogorov-Smirnov 5% Critical Value 0.175 Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 826477 95% Adjusted Gamma UCL (Use when n < 40) 834498

Potential UCL to Use

Use 95% Student's-t UCL 813471

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: BLUE CRAB, OYSTER, WHITE SHRIMP (SIZE CLASS A)

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File C:\Documents and Settings\sballard\My Documents\Projects\Patrick Bayou\2011\ECO RA\Wildlife Risk\ProUCL\Input

Files\OYS_BCRA_WHSA.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	20	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	20
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (1,4-dichlorobenzene_ug/kg)

General Statistics

0	Number of Detected Data	20	Number of Valid Data
20	Number of Non-Detect Data	0	Number of Distinct Detected Data
100.00%	Percent Non-Detects		

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,4-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

General	Statistics

	General 3	tatistics	
Number of Valid Data	20	Number of Detected Data	10
Number of Distinct Detected Data	10	Number of Non-Detect Data	10
		Percent Non-Detects	50.00%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	191	Minimum Detected	5.252
Maximum Detected	5430	Maximum Detected	8.6
Mean of Detected	967.5	Mean of Detected	6.315
SD of Detected	1582	SD of Detected	0.917
Minimum Non-Detect	347	Minimum Non-Detect	5.849
Maximum Non-Detect	727	Maximum Non-Detect	6.589
Note: Data have multiple DLs - Use of KM Method is recommended		Number treated as Non-Detect	18
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	2
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	90.00%

	UCL Statistics		
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.486	Shapiro Wilk Test Statistic	0.829
5% Shapiro Wilk Critical Value	0.842	5% Shapiro Wilk Critical Value	0.842
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	592.5	Mean	5.828
SD	1156	SD	0.828
95% DL/2 (t) UCL	1039	95% H-Stat (DL/2) UCL	584.9
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE method failed to converge properly		Mean in Log Scale	5.888
		SD in Log Scale	0.774
		Mean in Original Scale	602.5
		SD in Original Scale	1151
		95% Percentile Bootstrap UCL	1105
		95% BCA Bootstrap UCL	1410
Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	
k star (bias corrected)	0.786	Data do not follow a Discernable Distribution (0.05)	
Theta Star	1231		
nu star	15.72		
A-D Test Statistic	1.365	Nonparametric Statistics	
5% A-D Critical Value	0.748	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.748	Mean	625.7
5% K-S Critical Value	0.274	SD	1116
Data not Gamma Distributed at 5% Significance Level		SE of Mean	264.1
		95% KM (t) UCL	1082
Assuming Gamma Distribution		95% KM (z) UCL	1060
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	1072
Minimum	191	95% KM (bootstrap t) UCL	2788
Maximum	5430	95% KM (BCA) UCL	1119
Mean	970.2	95% KM (Percentile Bootstrap) UCL	1132
Median	967.1	95% KM (Chebyshev) UCL	1777
SD	1088	97.5% KM (Chebyshev) UCL	2275
k star	1.679	99% KM (Chebyshev) UCL	3253
Theta star	577.7		
Nu star	67.17	Potential UCLs to Use	
AppChi2	49.31	95% KM (t) UCL	1082
95% Gamma Approximate UCL	1322	95% KM (% Bootstrap) UCL	1132
95% Adjusted Gamma UCL	1355		
not a recommended method.			

Note: DL/2 is no

Result_Value (hexachlorobutadiene_ug/kg)

General Statistics

Number of Valid Data 20 Number of Detected Data 0 Number of Distinct Detected Data 0 Number of Non-Detect Data 20 Percent Non-Detects 100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs! Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (hexachlorobutadiene_ug/kg) was not processed!

Result_Value (lead_mg/kg)

General Statistics

Number of Valid Observations 20 Number of Distinct Observations 20

Raw Statistics Log-transformed Statistics

Minimum 0.092 Minimum of Log Data -2.386

Maximum 0.518 Maximum of Log Data -0.658

Mean 0.183 Mean of log Data -1.837

Median 0.126 SD of log Data 0.503

Coefficient of Variation 0.616

Approximate Chi Square Value (.05) 105.7

Skewness 1.761

SD 0.112

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.757 Shapiro Wilk Test Statistic 0.858
Shapiro Wilk Critical Value 0.905 Shapiro Wilk Critical Value 0.905

Data not Normal at 5% Significance Level

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 0.226 95% H-UCL 0.228

95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 0.271

 95% Adjusted-CLT UCL 0.235
 97.5% Chebyshev (MVUE) UCL 0.31

 95% Modified-t UCL 0.228
 99% Chebyshev (MVUE) UCL 0.388

Gamma Distribution Test Data Distribution

k star (bias corrected) 3.28 Data do not follow a Discernable Distribution (0.05)

Theta Star 0.0557
MLF of Mean 0.183

MLE of Standard Deviation 0.101 nu star 131.2

Adjusted Level of Significance 0.038 95% CLT UCL 0.224

Nonparametric Statistics

Adjusted Chi Square Value 103.9 95% Jackknife UCL 0.226
95% Standard Bootstrap UCL 0.223
Anderson-Darling Test Statistic 1.449 95% Bootstrap-t UCL 0.249
Anderson-Darling 5% Critical Value 0.746 95% Hall's Bootstrap UCL 0.237

Kolmogorov-Smirnov 7est Statistic 0.236

Kolmogorov-Smirnov 5% Critical Value 0.195

95% Percentile Bootstrap UCL 0.225

95% BCA Bootstrap UCL 0.234

Data not Gamma Distributed at 5% Significance Level95% Chebyshev(Mean, Sd) UCL 0.29297.5% Chebyshev(Mean, Sd) UCL 0.34

Assuming Gamma Distribution 99% Chebyshev(Mean, Sd) UCL 0.433
95% Approximate Gamma UCL 0.227

95% Adjusted Gamma UCL 0.23

Potential UCL to Use Use 95% Chebyshev (Mean, Sd) UCL 0.292

Result_Value (mercury_mg/kg)

Number of Valid Observations 20 Number of Distinct Observations 15

Baseline Ecological Risk AssessmentMarch 2013Patrick Bayou Superfund SitePage 3 of 8040284-01

General Statistics

Raw Statistics Log-transformed Statistics

 Minimum 0.009
 Minimum of Log Data -4.711

 Maximum 0.072
 Maximum of Log Data -2.631

 Mean 0.0474
 Mean of log Data -3.127

 Median 0.048
 SD of log Data 0.461

SD 0.0155

Coefficient of Variation 0.327

MLE of Standard Deviation 0.0198

nu star 229.1

Skewness -0.548

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.96 Shapiro Wilk Test Statistic 0.784
Shapiro Wilk Critical Value 0.905 Shapiro Wilk Critical Value 0.905

Data appear Normal at 5% Significance Level Data not Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 0.0533 95% H-UCL 0.0601

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 0.071

 95% Adjusted-CLT UCL 0.0526
 97.5% Chebyshev (MVUE) UCL 0.0807

95% Modified-t UCL 0.0533 99% Chebyshev (MVUE) UCL 0.0998

Gamma Distribution Test Data Distribution

k star (bias corrected) 5.728 Data appear Normal at 5% Significance Level
Theta Star 0.00827

MLF of Mean 0.0474

Approximate Chi Square Value (.05) 195.1 Nonparametric Statistics

Adjusted Level of Significance 0.038
Adjusted Chi Square Value 192.6
95% Jackknife UCL 0.0533
95% Standard Bootstrap UCL 0.0529
Anderson-Darling Test Statistic 0.796
95% Bootstrap-t UCL 0.0532
Anderson-Darling 5% Critical Value 0.744
95% Hall's Bootstrap UCL 0.0528

Kolmogorov-Smirnov Test Statistic 0.2

Kolmogorov-Smirnov 5% Critical Value 0.194

95% BCA Bootstrap UCL 0.0528

Data not Gamma Distributed at 5% Significance Level

95% Chebyshev(Mean, Sd) UCL 0.0624

97.5% Chebyshev(Mean, Sd) UCL 0.069

Assuming Gamma Distribution 99% Chebyshev (Mean, Sd) UCL 0.0818

95% Adjusted Gamma UCL 0.0563

Potential UCL to Use Use 95% Student's-t UCL 0.0533

95% Approximate Gamma UCL 0.0556

General Statistics

Number of Valid Observations 20

Number of Distinct Observations 20

Minimum 2.513 Minimum of Log Data 0.921

Log-transformed Statistics

Maximum 44.76 Maximum of Log Data 3.801

Result_Value (total dioxin/furan teq 1998 (avian) (u = 1/2)_ng/kg)

Raw Statistics

Mean 18 1 Mean of log Data 2.769 Median 17.57 SD of log Data 0.578 SD 8 604

Coefficient of Variation 0.475 Skewness 1.308

MLE of Standard Deviation 9.63

Relevant UCI Statistics

Normal Distribution Test Lognormal Distribution Test

> Shapiro Wilk Test Statistic 0.898 Shapiro Wilk Test Statistic 0.857 Shapiro Wilk Critical Value 0.905 Shapiro Wilk Critical Value 0.905

Data not Normal at 5% Significance Level Data not Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% H-UCL 24.89 95% Student's-t UCL 21.42

95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 29.69 95% Adjusted-CLT UCL 21.86 97.5% Chebyshev (MVUE) UCL 34.45

> 95% Modified-t UCL 21.52 99% Chebyshev (MVUE) UCL 43.82

Gamma Distribution Test Data Distribution

> k star (bias corrected) 3.532 Data appear Gamma Distributed at 5% Significance Level

Theta Star 5.124

MLE of Mean 18.1

nu star 141.3

Approximate Chi Square Value (.05) 114.8 95% CLT UCL 21.26 Adjusted Level of Significance 0.038

Adjusted Chi Square Value 112.9 95% Jackknife UCL 21.42

Nonparametric Statistics

95% Standard Bootstrap UCL 21.25

95% Bootstrap-t UCL 22.13 Anderson-Darling Test Statistic 0.634 Anderson-Darling 5% Critical Value 0.745 95% Hall's Bootstrap UCL 24.57

Kolmogorov-Smirnov Test Statistic 0.157 95% Percentile Bootstrap UCL 21.22

95% BCA Bootstrap UCL 21.86 Kolmogorov-Smirnov 5% Critical Value 0.195

Data appear Gamma Distributed at 5% Significance Level 95% Chebyshev(Mean, Sd) UCL 26.48

97.5% Chebyshev(Mean, Sd) UCL 30.11

99% Chebyshev(Mean, Sd) UCL 37.24 **Assuming Gamma Distribution**

95% Approximate Gamma UCL 22.27

Potential UCL to Use Use 95% Approximate Gamma UCL 22.27

95% Adjusted Gamma UCL 22.64

Result_Value (total pah (16) (u = 1/2)_ug/kg)

General Statistics

Number of Valid Data 20 Number of Detected Data 7 Number of Distinct Detected Data 7 Number of Non-Detect Data 13

Raw Statistics **Log-transformed Statistics**

Minimum Detected 60.91 Minimum Detected 4.109 Maximum Detected 552.1 Maximum Detected 6.314 Mean of Detected 157.2 Mean of Detected 4.706 SD of Detected 178.3 SD of Detected 0.803

65.00%

Percent Non-Detects

Minimum Non-Detect6.93Minimum Non-Detect1.936Maximum Non-Detect14.8Maximum Non-Detect2.695

Note: Data have multiple DLs - Use of KM Method is recommendedNumber treated as Non-Detect13For all methods (except KM, DL/2, and ROS Methods),Number treated as Detected7Observations < Largest ND are treated as NDs</td>Single DL Non-Detect Percentage65.00%

Warning: There are only 7 Detected Values in this data

Note: It should be noted that even though bootstrap may be performed on this data set the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

	UCL Statistic	s	
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.622	Shapiro Wilk Test Statistic	0.792
5% Shapiro Wilk Critical Value	0.803	5% Shapiro Wilk Critical Value	0.803
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	59.01	Mean	2.796
SD	124.5	SD	1.531
95% DL/2 (t) UCL	107.2	95% H-Stat (DL/2) UCL	69.93
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE yields a negative mean		Mean in Log Scale	2.988
		SD in Log Scale	1.405
		Mean in Original Scale	60.5
		SD in Original Scale	123.9
		95% Percentile Bootstrap UCL	110.2
		95% BCA Bootstrap UCL	136.1
Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	
k star (bias corrected)	0.992	Data Follow Appr. Gamma Distribution at 5% Significance Level	
Theta Star	158.5		
nu star	13.89		
A-D Test Statistic	0.892	Nonparametric Statistics	
5% A-D Critical Value	0.72	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.72	Mean	94.62
5% K-S Critical Value	0.317	SD	107.9
Data follow Appr. Gamma Distribution at 5% Significance Level		SE of Mean	26.07
		95% KM (t) UCL	139.7
Assuming Gamma Distribution		95% KM (z) UCL	137.5
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	137.4
Minimum	60.91	95% KM (bootstrap t) UCL	316.1
Maximum	552.1	95% KM (BCA) UCL	148.3
Mean	157.3	95% KM (Percentile Bootstrap) UCL	143.3
Median	157.2	95% KM (Chebyshev) UCL	208.2
SD	104.7	97.5% KM (Chebyshev) UCL	257.4
k star	3.163	99% KM (Chebyshev) UCL	354
Theta star	49.72		
Nu star	126.5	Potential UCLs to Use	

AppChi2 101.5 95% KM (t) UCL 139.7

95% Gamma Approximate UCL 196 95% Adjusted Gamma UCL 199.4

Note: DL/2 is not a recommended method.

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 20

Number of Distinct Observations 20

Raw Statistics

Minimum 228677 Maximum 1429439 Mean 703263 Median 677904 SD 304128

Coefficient of Variation 0.432 Skewness 0.866 **Log-transformed Statistics**

Minimum of Log Data 12.34
Maximum of Log Data 14.17
Mean of log Data 13.37
SD of log Data 0.448

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.938 Shapiro Wilk Critical Value 0.905

Shapiro Wilk Test Statistic 0.974 Shapiro Wilk Critical Value 0.905

Data appear Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 820853

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL 829190 95% Modified-t UCL 823047 Assuming Lognormal Distribution

Data appear Lognormal at 5% Significance Level

Lognormal Distribution Test

95% H-UCL 869904 95% Chebyshev (MVUE) UCL 1024356 97.5% Chebyshev (MVUE) UCL 1161945 99% Chebyshev (MVUE) UCL 1432211

Gamma Distribution Test

Theta Star 144750

MLE of Mean 703263

MLE of Standard Deviation 319057

nu star 194.3

k star (bias corrected) 4.858

Approximate Chi Square Value (.05) 163.1

Adjusted Level of Significance 0.038 Adjusted Chi Square Value 160.8

Anderson-Darling Test Statistic 0.236
Anderson-Darling 5% Critical Value 0.745
Kolmogorov-Smirnov Test Statistic 0.108
Kolmogorov-Smirnov 5% Critical Value 0.194

Data appear Gamma Distributed at 5% Significance Level

Nonparametric Statistics

Data Distribution

Data appear Normal at 5% Significance Level

95% CLT UCL 815122
95% Jackknife UCL 820853
95% Standard Bootstrap UCL 810356
95% Bootstrap-t UCL 838530
95% Hall's Bootstrap UCL 845932
95% Percentile Bootstrap UCL 819939
95% BCA Bootstrap UCL 829918
95% Chebyshev(Mean, Sd) UCL 999691
97.5% Chebyshev(Mean, Sd) UCL 1127955
99% Chebyshev(Mean, Sd) UCL 1379906

Assuming Gamma Distribution

95% Approximate Gamma UCL 838028 95% Adjusted Gamma UCL 849804

Potential UCL to Use

Use 95% Student's-t UCL 820853

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 20 Number of Distinct Observations 20

Raw Statistics **Log-transformed Statistics**

> Minimum 97.57 Minimum of Log Data 4.581 Maximum 782.9 Maximum of Log Data 6.663 Mean 358.2 Mean of log Data 5.784 Median 321.9 SD of log Data 0.467

SD 163.3 Coefficient of Variation 0.456

Skewness 1.225

95% Adjusted-CLT UCL 428.9

MLE of Standard Deviation 168.3

95% Adjusted Gamma UCL 435.9

Approximate Chi Square Value (.05) 151.1

Relevant UCL Statistics

Normal Distribution Test **Lognormal Distribution Test**

> Shapiro Wilk Test Statistic 0.888 Shapiro Wilk Test Statistic 0.935 Shapiro Wilk Critical Value 0.905 Shapiro Wilk Critical Value 0.905

Data appear Lognormal at 5% Significance Level Data not Normal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% H-UCL 448.6 95% Student's-t UCL 421.3 95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 529.7

> 95% Modified-t UCL 423 99% Chebyshev (MVUE) UCL 747.1

97.5% Chebyshev (MVUE) UCL 603.1

Nonparametric Statistics

Gamma Distribution Test Data Distribution

> k star (bias corrected) 4.53 Data appear Gamma Distributed at 5% Significance Level Theta Star 79.06

MLE of Mean 358.2

nu star 181 2

Adjusted Level of Significance 0.038 95% CLT UCL 418.2

Adjusted Chi Square Value 148.9 95% Jackknife UCL 421.3 95% Standard Bootstrap UCL 415.7

Anderson-Darling Test Statistic 0.549 95% Bootstrap-t UCL 438 Anderson-Darling 5% Critical Value 0.745 95% Hall's Bootstrap UCL 456.3 Kolmogorov-Smirnov Test Statistic 0.161 95% Percentile Bootstrap UCL 415.6

Kolmogorov-Smirnov 5% Critical Value 0.194 95% BCA Bootstrap UCL 427.7

Data appear Gamma Distributed at 5% Significance Level 95% Chebyshev(Mean, Sd) UCL 517.4

97.5% Chebyshev(Mean, Sd) UCL 586.3

Assuming Gamma Distribution 99% Chebyshev(Mean, Sd) UCL 721.6 95% Approximate Gamma UCL 429.6

Potential UCL to Use Use 95% Approximate Gamma UCL 429.6

GROUPING: BROWN SHRIMP

Brown Shrimp

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1_b.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 8

Number of Distinct Observations 8

Raw Statistics

Minimum 996653 Maximum 2192372

Mean 1716066 Geometric Mean 1658510

Median 1847710

SD 443301

Std. Error of Mean 156730 Coefficient of Variation 0.258

Skewness -0.73

Log-transformed Statistics

Minimum of Log Data 13.81 Maximum of Log Data 14.6 Mean of log Data 14.32 SD of log Data 0.29

Warning: There are only 8 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this dataset, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on datasets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.9 Shapiro Wilk Critical Value 0.818

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.867 Shapiro Wilk Critical Value 0.818

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 2013004

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 1930652

95% Modified-t UCL (Johnson-1978) 2006264

Assuming Lognormal Distribution

95% H-UCL 2165751 95% Chebyshev (MVUE) UCL 2493860 97.5% Chebyshev (MVUE) UCL 2828440 99% Chebyshev (MVUE) UCL 3485657

Gamma Distribution Test

k star (bias corrected) 9.347

Theta Star 183603

MLE of Mean 1716066

MLE of Standard Deviation 561315

nu star 149.5

Approximate Chi Square Value (.05) 122.3

Adjusted Level of Significance 0.0195 Adjusted Chi Square Value 116

Anderson-Darling Test Statistic 0.498 Anderson-Darling 5% Critical Value 0.716

Kolmogorov-Smirnov Test Statistic 0.232 Kolmogorov-Smirnov 5% Critical Value 0.294

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 2098698 95% Adjusted Gamma UCL (Use when n < 40) 2211517 Nonparametric Statistics

Data Distribution

Data appear Normal at 5% Significance Level

95% CLT UCL 1973865 95% Jackknife UCL 2013004

95% Standard Bootstrap UCL 1952853 95% Bootstrap-t UCL 1970574 95% Hall's Bootstrap UCL 1916436

95% Percentile Bootstrap UCL 1951384 95% BCA Bootstrap UCL 1928821 95% Chebyshev(Mean, Sd) UCL 2399238 97.5% Chebyshev(Mean, Sd) UCL 2694847

99% Chebyshev(Mean, Sd) UCL 3275514

Potential UCL to Use

Use 95% Student's-t UCL 2013004

Brown Shrimp

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed datasets.

GROUPING: GULF KILLIFISH

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1_c.wst Full Precision OFF Confidence Coefficient 95% Number of Bootstrap Operations 2000

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Distinct Observations 25 Number of Valid Observations 25

Raw Statistics

Minimum 1964622 Minimum of Log Data 14.49 Maximum of Log Data 16 Maximum 8847995 Mean 5215609 Mean of log Data 15.42 Geometric Mean 4972742 SD of log Data 0.327 Median 5144332 SD 1568411 Std. Error of Mean 313682 Coefficient of Variation 0.301

Log-transformed Statistics

Lognormal Distribution Test

Assuming Lognormal Distribution

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.99 Shapiro Wilk Test Statistic 0.962 Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918 Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 5752282 95% H-UCL 5929489 95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 6756490 95% Adjusted-CLT UCL (Chen-1995) 5749406 97.5% Chebyshev (MVUE) UCL 7414908 95% Modified-t UCL (Johnson-1978) 5755064 99% Chebyshev (MVUE) UCL 8708242

Gamma Distribution Test

Data Distribution Data appear Normal at 5% Significance Level k star (bias corrected) 9.398 Theta Star 554953

MLE of Mean 5215609 MLE of Standard Deviation 1701299 nu star 469.9

Skewness 0.266

Approximate Chi Square Value (.05) 420.7 **Nonparametric Statistics** Adjusted Level of Significance 0.0395 95% CLT UCL 5731570 Adjusted Chi Square Value 417.5 95% Jackknife UCL 5752282 95% Standard Bootstrap UCL 5709166

Anderson-Darling Test Statistic 0.164 95% Bootstrap-t UCL 5782797 Anderson-Darling 5% Critical Value 0.745 95% Hall's Bootstrap UCL 5788260 Kolmogorov-Smirnov Test Statistic 0.0718 95% Percentile Bootstrap UCL 5715854 Kolmogorov-Smirnov 5% Critical Value 0.174 95% BCA Bootstrap UCL 5731928 Data appear Gamma Distributed at 5% Significance Level 95% Chebyshev(Mean, Sd) UCL 6582918 97.5% Chebyshev(Mean, Sd) UCL 7174554 99% Chebyshev(Mean, Sd) UCL 8336708

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 5826419 95% Adjusted Gamma UCL (Use when n < 40) 5870672

> Potential UCL to Use Use 95% Student's-t UCL 5752282

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 25

Raw Statistics

Minimum 270.7 Maximum 1173 Mean 794.6 Geometric Mean 761.1 Median 792.8 SD 214.9 Std. Error of Mean 42.99 Coefficient of Variation 0.27 Skewness -0.349

Log-transformed Statistics

Minimum of Log Data 5.601 Maximum of Log Data 7.067 Mean of log Data 6.635 SD of log Data 0.32

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.982 Shapiro Wilk Critical Value 0.918 Data appear Normal at 5% Significance Level

Shapiro Wilk Test Statistic 0.898 Shapiro Wilk Critical Value 0.918

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 868.2 95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 862.1 95% Modified-t UCL (Johnson-1978) 867.7

Assuming Lognormal Distribution

Lognormal Distribution Test

95% H-UCL 902.8 95% Chebyshev (MVUE) UCL 1027 97.5% Chebyshev (MVUE) UCL 1125 99% Chebyshev (MVUE) UCL 1318

Gamma Distribution Test

k star (bias corrected) 10.38 Theta Star 76.54 MLE of Mean 794.6 MLE of Standard Deviation 246.6

nu star 519.1 Approximate Chi Square Value (.05) 467.2 Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 463.9

Anderson-Darling Test Statistic 0.406 Anderson-Darling 5% Critical Value 0.745 Kolmogorov-Smirnov Test Statistic 0.142 Kolmogorov-Smirnov 5% Critical Value 0.174 Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 882.8 95% Adjusted Gamma UCL (Use when n < 40) 889.2

Data Distribution Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 865.3 95% Jackknife UCL 868.2 95% Standard Bootstrap UCL 864.9 95% Bootstrap-t UCL 862.8 95% Hall's Bootstrap UCL 861.6 95% Percentile Bootstrap UCL 865.5 95% BCA Bootstrap UCL 862.1 95% Chebyshev(Mean, Sd) UCL 982 97.5% Chebyshev(Mean, Sd) UCL 1063 99% Chebyshev(Mean, Sd) UCL 1222

Potential UCL to Use

Use 95% Student's-t UCL 868.2

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

> Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed datasets.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 25

Raw Statistics

Minimum 15.6 Maximum 137.1 Mean 77.42 Geometric Mean 70.85 Median 80.9 SD 28.09 Std. Error of Mean 5.618 Coefficient of Variation 0.363 Skewness -0.237

Log-transformed Statistics

Minimum of Log Data 2.747 Maximum of Log Data 4.921 Mean of log Data 4.261 SD of log Data 0.482

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.98 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Assuming Normal Distribution 95% Student's-t UCL 87.03

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 86.38 95% Modified-t UCL (Johnson-1978) 86.99

Gamma Distribution Test

k star (bias corrected) 5.129
Theta Star 15.1
MLE of Mean 77.42
MLE of Standard Deviation 34.19
nu star 256.4
Approximate Chi Square Value (.05) 220.4

Adjusted Level of Significance 0.0395
Adjusted Chi Square Value 218.1

Anderson-Darling Test Statistic 0.787

Anderson-Darling lest Statistic 0.787
Anderson-Darling 5% Critical Value 0.747
Kolmogorov-Smirnov Test Statistic 0.149
Kolmogorov-Smirnov 5% Critical Value 0.175
Data follow Appr. Gamma Distribution at 5% Significance Level

Data follow Appr. Gamma Distribution at 5% Signif

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 90.1 95% Adjusted Gamma UCL (Use when n < 40) 91.04

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.852 Shapiro Wilk Critical Value 0.918 Data not Lognormal at 5% Significance Level

Assuming Lognormal Distribution

95% H-UCL 96.33 95% Chebyshev (MVUE) UCL 113.7 97.5% Chebyshev (MVUE) UCL 128.6 99% Chebyshev (MVUE) UCL 158

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 86.66
95% Jackknife UCL 87.03
95% Standard Bootstrap UCL 86.64
95% Bootstrap-t UCL 86.82
95% Hall's Bootstrap UCL 87.06
95% Percentile Bootstrap UCL 85.91
95% Chebyshev(Mean, Sd) UCL 101.9
97.5% Chebyshev(Mean, Sd) UCL 122.5
99% Chebyshev(Mean, Sd) UCL 133.3

Potential UCL to Use

Use 95% Student's-t UCL 87.03

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positively skewed datasets.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 21

Raw Statistics

Minimum 0.025
Maximum 0.109
Mean 0.0524
Geometric Mean 0.0477
Median 0.044
SD 0.0242
Std. Error of Mean 0.00485
Coefficient of Variation 0.463
Skewness 1.097

Log-transformed Statistics

Minimum of Log Data -3.689 Maximum of Log Data -2.216 Mean of log Data -3.042 SD of log Data 0.43

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.871 Shapiro Wilk Critical Value 0.918 Data not Normal at 5% Significance Level

cic vant Oct Statistics

Lognormal Distribution Test
Shapiro Wilk Test Statistic 0.944
Shapiro Wilk Critical Value 0.918
Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 0.0615 95% Modified-t UCL (Johnson-1978) 0.0608

Assuming Lognormal Distribution

95% H-UCL 0.0619 95% Chebyshev (MVUE) UCL 0.0723 97.5% Chebyshev (MVUE) UCL 0.0811 99% Chebyshev (MVUE) UCL 0.0982

Gamma Distribution Test

k star (bias corrected) 4.929
Theta Star 0.0106

95% Student's-t UCL 0.0607

MLE of Mean 0.0524 MLE of Standard Deviation 0.0236 nu star 246.5

Approximate Chi Square Value (.05) 211.1 Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 208.9

Anderson-Darling Test Statistic 0.639
Anderson-Darling 5% Critical Value 0.747
Kolmogorov-Smirnov Test Statistic 0.162
Kolmogorov-Smirnov 5% Critical Value 0.175
Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0611 95% Adjusted Gamma UCL (Use when n < 40) 0.0618

Data Distribution

Data appear Gamma Distributed at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 0.0603
95% Jackknife UCL 0.0607
95% Standard Bootstrap UCL 0.0607
95% Bootstrap UCL 0.0625
95% Hall's Bootstrap UCL 0.0612
95% Percentile Bootstrap UCL 0.0603
95% BCA Bootstrap UCL 0.0618
95% Chebyshev(Mean, Sd) UCL 0.0735
97.5% Chebyshev(Mean, Sd) UCL 0.0826

Potential UCL to Use

Use 95% Approximate Gamma UCL 0.0611

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (fish) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 23

Raw Statistics

Minimum 2.2 Maximum 9.7 Mean 6.553 Geometric Mean 6.198 Median 6.9 SD 2.013 Std. Error of Mean 0.403 Coefficient of Variation 0.307 Skewness -0.319

Log-transformed Statistics

Minimum of Log Data 0.788 Maximum of Log Data 2.272 Mean of log Data 1.824 SD of log Data 0.363

Relevant UCL Statistics

Normal Distribution Test

Assuming Normal Distribution

Shapiro Wilk Test Statistic 0.968 Shapiro Wilk Critical Value 0.918 Data appear Normal at 5% Significance Level

Lognormal Distribution Test Shapiro Wilk Test Statistic 0.907 Shapiro Wilk Critical Value 0.918 Data not Lognormal at 5% Significance Level

95% Student's-t UCL 7.242 95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 7.188

95% Modified-t UCL (Johnson-1978) 7.238

Gamma Distribution Test k star (bias corrected) 8.061

MLE of Mean 6.553 MLE of Standard Deviation 2.308 nu star 403 Approximate Chi Square Value (.05) 357.5 Adjusted Level of Significance 0.0395

Theta Star 0.813

Adjusted Chi Square Value 354.6 Anderson-Darling Test Statistic 0.5

Anderson-Darling 5% Critical Value 0.745 Kolmogorov-Smirnov Test Statistic 0.143 Kolmogorov-Smirnov 5% Critical Value 0.175 Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 7.388 95% Adjusted Gamma UCL (Use when n < 40) 7.449

Assuming Lognormal Distribution 95% H-UCL 7.598 95% Chebyshev (MVUE) UCL 8.741 97.5% Chebyshev (MVUE) UCL 9.666 99% Chebyshev (MVUE) UCL 11.48

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 7.216 95% Jackknife UCL 7.242 95% Standard Bootstrap UCL 7,201 95% Bootstrap-t UCL 7.228 95% Hall's Bootstrap UCL 7.217 95% Percentile Bootstrap UCL 7.187 95% BCA Bootstrap UCL 7.161 95% Chebyshev(Mean, Sd) UCL 8.308 97.5% Chebyshev(Mean, Sd) UCL 9.068 99% Chebyshev(Mean, Sd) UCL 10.56

Potential UCL to Use

Use 95% Student's-t UCL 7.242

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

> Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

GROUPING: GULF KILLIFISH, GULF MENHADEN, STRIPED MULLET, PINFISH, SAND SEATROUT

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1_d.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	50	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	50
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The dataset for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

	General Statis	ntion.	
Number of Valid Data	50	Number of Detected Data	28
Number of Distinct Detected Data	27	Number of Non-Detect Data	22
Number of distinct betected data	21	Percent Non-Detects	44.00%
		reitellt Noir-Detects	44.00%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	200	Minimum Detected	5.298
Maximum Detected	2360	Maximum Detected	7.766
Mean of Detected	680	Mean of Detected	6.28
SD of Detected	576.9	SD of Detected	0.66
Minimum Non-Detect	351	Minimum Non-Detect	5.861
Maximum Non-Detect	765	Maximum Non-Detect	6.64
Note: Data have multiple DLs - Use of KM Method is recommende	d	Number treated as Non-Detect	43
For all methods (except KM, DL/2, and ROS Methods),	u .	Number treated as Detected	7
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	86.00%
	UCL Statisti		
Normal Distribution Test with Detected Values Only	•	Lognormal Distribution Test with Detected Values Onl	•
Shapiro Wilk Test Statistic	0.709	Shapiro Wilk Test Statistic	0.923
5% Shapiro Wilk Critical Value	0.924	5% Shapiro Wilk Critical Value	0.924
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	519.2	Mean	6.029
SD	468.9	SD	0.603
95% DL/2 (t) UCL	630.3	95% H-Stat (DL/2) UCL	590.3
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE yields a negative mean		Mean in Log Scale	6.056
, 0		SD in Log Scale	0.573
		Mean in Original Scale	525.4
		SD in Original Scale	465.4
		95% t UCL	635.7
		95% Percentile Bootstrap UCL	643.2
		95% BCA Bootstrap UCL	676.8
		95% H-UCL	589.2
Gamma Distribution Test with Detected Values Onl		Data Distribution Test with Detected Values Only	
k star (bias corrected)	y 2.004	Data Distribution Test with Detected Values Only Data do not follow a Discernable Distribution (0.05)	
Theta Star	339.3	Data do not follow a Discernable Distribution (0.05)	
nu star	112.2		
A-D Test Statistic	1.384	Nonparametric Statistics	
5% A-D Critical Value	0.757	Kaplan-Meier (KM) Method	

K-S Test Statistic	0.757	Mean	528.3
5% K-S Critical Value	0.167	SD	463.9
Data not Gamma Distributed at 5% Significance Level		SE of Mean	68.18
		95% KM (t) UCL	642.6
Assuming Gamma Distribution		95% KM (z) UCL	640.4
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	641.9
Minimum	98.03	95% KM (bootstrap t) UCL	698.8
Maximum	2360	95% KM (BCA) UCL	645.9
Mean	546.6	95% KM (Percentile Bootstrap) UCL	646.4
Median	474.4	95% KM (Chebyshev) UCL	825.5
SD	470.9	97.5% KM (Chebyshev) UCL	954.1
k star	1.973	99% KM (Chebyshev) UCL	1207
Theta star	277		
Nu star	197.3	Potential UCLs to Use	
AppChi2	165.8	95% KM (t) UCL	642.6
95% Gamma Approximate UCL (Use when n >= 40)	650.4	95% KM (% Bootstrap) UCL	646.4
95% Adjusted Gamma UCL (Use when n < 40)	653.8		

Note: DL/2 is not a recommended method.

Number of Valid Observations 50

Std. Error of Mean 0.00575

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006). For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Raw Statistics	Log-transformed Statistics
Minimum 0.024	Minimum of Log Data -3.73
Maximum 0.199	Maximum of Log Data -1.614
Mean 0.0643	Mean of log Data -2.904
Geometric Mean 0.0548	SD of log Data 0.551
Median 0.0465	
SD 0.0406	

Number of Distinct Observations 37

Coefficient of Variation 0.632	
Skewness 1.635	
	Relevant UCL Statistics
Normal Distribution Test	Lognormal Distribution Test
Shapiro Wilk Test Statistic 0.82	Shapiro Wilk Test Statistic 0.944
Shapiro Wilk Critical Value 0.947	Shapiro Wilk Critical Value 0.947
Data not Normal at 5% Significance Level	Data not Lognormal at 5% Significance Level
Assuming Normal Distribution	Assuming Lognormal Distribution
95% Student's-t UCL 0.0739	95% H-UCL 0.0743
95% UCLs (Adjusted for Skewness)	95% Chebyshev (MVUE) UCL 0.0865
95% Adjusted-CLT UCL (Chen-1995) 0.0752	97.5% Chebyshev (MVUE) UCL 0.0964
95% Modified-t UCL (Johnson-1978) 0.0742	99% Chebyshev (MVUE) UCL 0.116
Gamma Distribution Test	Data Distribution
k star (bias corrected) 3 111	Data do not follow a Discernable Distribution (0

Gamma Distribution Test	Data Distribution
k star (bias corrected) 3.111	Data do not follow a Discernable Distribution (0.05)
Theta Star 0.0207	
MLE of Mean 0.0643	
MLE of Standard Deviation 0.0365	
nu star 311.1	

Approximate Chi Square Value (.05) 271.2	Nonparametric Statistics
Adjusted Level of Significance 0.0452	95% CLT UCL 0.0738
Adjusted Chi Square Value 270.1	95% Jackknife UCL 0.0739
	95% Standard Bootstrap UCL 0.0736
Anderson-Darling Test Statistic 1.152	95% Bootstrap-t UCL 0.0757
Anderson-Darling 5% Critical Value 0.756	95% Hall's Bootstrap UCL 0.0757
Kolmogorov-Smirnov Test Statistic 0.157	95% Percentile Bootstrap UCL 0.0749
Kolmogorov-Smirnov 5% Critical Value 0.126	95% BCA Bootstrap UCL 0.0744
Data not Gamma Distributed at 5% Significance Level	95% Chebyshev(Mean, Sd) UCL 0.0893
	97.5% Chebyshev(Mean, Sd) UCL 0.1
Assuming Gamma Distribution	99% Chebyshev(Mean, Sd) UCL 0.121

Assuming Gamma Distribution 95% Approximate Gamma UCL (Use when n >= 40) 0.0738 95% Adjusted Gamma UCL (Use when n < 40) 0.0741

Potential UCL to Use

Use 95% Chebyshev (Mean, Sd) UCL 0.0893

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (fish) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 50

Number of Distinct Observations 43

Raw Statistics

Minimum 0.6 Maximum 9.7 Mean 4.541 Geometric Mean 3.721 Median 3.95 SD 2 607 Std. Error of Mean 0.369 Coefficient of Variation 0.574 Skewness 0.414

Log-transformed Statistics

Minimum of Log Data -0.511 Maximum of Log Data 2.272 Mean of log Data 1.314 SD of log Data 0.689

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.928 Shapiro Wilk Critical Value 0.947 Data not Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.937 Shapiro Wilk Critical Value 0.947 Data not Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 5.159 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 5.17 95% Modified-t UCL (Johnson-1978) 5.163

Assuming Lognormal Distribution

95% H-UCL 5.764 95% Chebyshev (MVUE) UCL 6.869 97.5% Chebyshev (MVUE) UCL 7.811 99% Chebyshev (MVUE) UCL 9.661

Gamma Distribution Test

Theta Star 1.802 MLE of Mean 4.541 MLE of Standard Deviation 2.86 nu star 252

k star (bias corrected) 2.52

Approximate Chi Square Value (.05) 216.3 Adjusted Level of Significance 0.0452 Adjusted Chi Square Value 215.3

Anderson-Darling Test Statistic 0.489 Anderson-Darling 5% Critical Value 0.759 Kolmogorov-Smirnov Test Statistic 0.0954 Kolmogorov-Smirnov 5% Critical Value 0.126 Data appear Gamma Distributed at 5% Significance Level

Data Distribution

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 5.292 95% Adjusted Gamma UCL (Use when n < 40) 5.316

Nonparametric Statistics

95% CLT UCL 5.147 95% Jackknife UCL 5 159 95% Standard Bootstrap UCL 5.128 95% Bootstrap-t UCL 5.182 95% Hall's Bootstrap UCL 5.171 95% Percentile Bootstrap UCL 5.167 95% BCA Bootstrap UCL 5.194 95% Chebyshev(Mean, Sd) UCL 6.148 97.5% Chebyshev(Mean, Sd) UCL 6.844 99% Chebyshev(Mean, Sd) UCL 8.21

Potential UCL to Use

Use 95% Approximate Gamma UCL 5.292

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 50

Number of Distinct Observations 50

Raw Statistics

Maximum 8847995 Mean 3872225 Geometric Mean 3391635 Median 3898513

Minimum 596176

Log-transformed Statistics

Minimum of Log Data 13.3 Maximum of Log Data 16 Mean of log Data 15.04 SD of log Data 0.551

SD 1887270 Std Error of Mean 266900 Coefficient of Variation 0.487 Skewness 0.535

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.96 Shapiro Wilk Critical Value 0.947 Data appear Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 4319697 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 4332797 95% Modified-t UCL (Johnson-1978) 4323060

Gamma Distribution Test

k star (bias corrected) 3.709 Theta Star 1043886 MLF of Mean 3872225 MLE of Standard Deviation 2010513 nu star 370.9

Approximate Chi Square Value (.05) 327.3 Adjusted Level of Significance 0.0452 Adjusted Chi Square Value 326.1

Anderson-Darling Test Statistic 0.302 Anderson-Darling 5% Critical Value 0.754 Kolmogorov-Smirnov Test Statistic 0.0887 Kolmogorov-Smirnov 5% Critical Value 0.126 Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 4388468 95% Adjusted Gamma UCL (Use when n < 40) 4404777

Potential UCL to Use

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.962 Shapiro Wilk Critical Value 0.947 Data appear Lognormal at 5% Significance Level

Assuming Lognormal Distribution

95% H-UCL 4594363 95% Chebyshev (MVUE) UCL 5351061 97.5% Chebyshev (MVUE) UCL 5964232 99% Chebyshev (MVUE) UCL 7168686

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 4311237 95% Jackknife UCL 4319697 95% Standard Bootstrap UCL 4305645 95% Bootstrap-t UCL 4337908 95% Hall's Bootstrap UCL 4366124 95% Percentile Bootstrap UCL 4298829 95% BCA Bootstrap UCL 4329279 95% Chebyshev(Mean, Sd) UCL 5035616 97.5% Chebyshev(Mean, Sd) UCL 5539017 99% Chebyshev(Mean, Sd) UCL 6527849

Use 95% Student's-t UCL 4319697

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: GULF KILLIFISH, STRIPED MULLET, PINFISH, SAND SEATROUT

Gulf Killifish, Striped Mullet, Pinfish, Sand Seatrout

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1 e.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 40 Number of Distinct Observations 30

Raw Statistics

Minimum 0.024

Maximum 0.122

Mean 0.0513

Geometric Mean 0.0464

Median 0.044

SD 0.0252

Std. Error of Mean 0.00398

Coefficient of Variation 0.491

Skewness 1.247

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.853

Shapiro Wilk Critical Value 0.94

Data not Normal at 5% Significance Level

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.058

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 0.0587 95% Modified-t UCL (Johnson-1978) 0.0582

Assuming Lognormal Distribution

Lognormal Distribution Test

Log-transformed Statistics

Minimum of Log Data -3.73

Maximum of Log Data -2.104

Mean of log Data -3.071

SD of log Data 0.444

95% Chebyshev (MVUE) UCL 0.0672 97.5% Chebyshev (MVUE) UCL 0.0742

95% H-UCL 0.0585

Shapiro Wilk Test Statistic 0.937

Shapiro Wilk Critical Value 0.94

99% Chebyshev (MVUE) UCL 0.088

Gamma Distribution Test

k star (bias corrected) 4.719

Theta Star 0.0109

MLE of Mean 0.0513

MLE of Standard Deviation 0.0236

nu star 377.5

Approximate Chi Square Value (.05) 333.5

Adjusted Level of Significance 0.044

Adjusted Chi Square Value 332

Anderson-Darling Test Statistic 1.083

Anderson-Darling 5% Critical Value 0.751

Kolmogorov-Smirnov Test Statistic 0.168

Kolmogorov-Smirnov 5% Critical Value 0.14 Data not Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0581

95% Adjusted Gamma UCL (Use when n < 40) 0.0584

Potential UCL to Use

Data Distribution

Data do not follow a Discernable Distribution (0.05)

Nonparametric Statistics 95% CLT UCL 0.0579

95% Jackknife UCL 0.058

95% Standard Bootstrap UCL 0.0578 95% Bootstrap-t UCL 0.0589

95% Hall's Bootstrap UCL 0.0592

95% Percentile Bootstrap UCL 0.0581

95% BCA Bootstrap UCL 0.0586

95% Chebyshev(Mean, Sd) UCL 0.0687 97.5% Chebyshev(Mean, Sd) UCL 0.0762

99% Chebyshev(Mean, Sd) UCL 0.091

Use 95% Student's-t UCL 0.058 or 95% Modified-t UCL 0.0582

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: GULF MENHADEN

Gulf Menhaden

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1_f.wst
Full Precision OFF

Confidence Coefficient 95% Number of Bootstrap Operations 2000

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 10 Number of Distinct Observations 9

Raw Statistics

Minimum 0.068
Maximum 0.199
Mean 0.116
Geometric Mean 0.107
Median 0.097
SD 0.0499
Std. Error of Mean 0.0158
Coefficient of Variation 0.43

Skewness 0.64

Log-transformed StatisticsMinimum of Maximum of

Minimum of Log Data -2.688 Maximum of Log Data -1.614 Mean of log Data -2.233 SD of log Data 0.42

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.851 Shapiro Wilk Test Statistic 0.862
Shapiro Wilk Critical Value 0.842 Shapiro Wilk Critical Value 0.842

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.145 95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 0.146 95% Modified-t UCL (Johnson-1978) 0.146

Assuming Lognormal Distribution

95% H-UCL 0.157 95% Chebyshev (MVUE) UCL 0.184 97.5% Chebyshev (MVUE) UCL 0.213 99% Chebyshev (MVUE) UCL 0.271

Gamma Distribution Test

k star (bias corrected) 4.514
Theta Star 0.0257
MLE of Mean 0.116
MLE of Standard Deviation 0.0547

nu star 90.28

Approximate Chi Square Value (.05) 69.37
Adjusted Level of Significance 0.0267

Anderson-Darling Test Statistic 0.677 Anderson-Darling 5% Critical Value 0.728 Kolmogorov-Smirnov Test Statistic 0.29

Adjusted Chi Square Value 66.19

Kolmogorov-Smirnov 5% Critical Value 0.267

Data follow Appr. Gamma Distribution at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.151 95% Adjusted Gamma UCL (Use when n < 40) 0.158

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% Jackknife UCL 0.145
95% Standard Bootstrap UCL 0.14
95% Bootstrap-t UCL 0.153
95% Hall's Bootstrap UCL 0.142
95% Percentile Bootstrap UCL 0.141
95% BCA Bootstrap UCL 0.144
95% Chebyshev(Mean, Sd) UCL 0.185
97.5% Chebyshev(Mean, Sd) UCL 0.215
99% Chebyshev(Mean, Sd) UCL 0.273

95% CLT UCL 0.142

Potential UCL to Use

Use 95% Student's-t UCL 0.145

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: GULF MENHADEN, STRIPED MULLET, PINFISH, SAND SEATROUT

General UCL Statistics for Datasets with Non-Detects

User Selected Options

From File Sheet1_g.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 25

Raw Statistics Log-transformed Statistics

 Minimum
 596176
 Minimum of Log Data
 13.3

 Maximum
 4538571
 Maximum of Log Data
 15.33

 Mean
 2528842
 Mean of log Data
 14.65

 Geometric Mean
 2313249
 SD of log Data
 0.456

Median 2313245

Median 2313245

SD 1025826

Std. Error of Mean 205165

Coefficient of Variation 0.406 Skewness 0.429

skewness 0.429

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.954 Shapiro Wilk Test Statistic 0.939
Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 2879855
 95% H-UCL 3068891

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 3604346

95% Adjusted-CLT UCL (Chen-1995) 2885105 97.5% Chebyshev (MVUE) UCL 4058543 95% Modified-t UCL (Johnson-1978) 2882787 99% Chebyshev (MVUE) UCL 4950724

Gamma Distribution Test Data Distribution

k star (bias corrected) 5.107 Data appear Normal at 5% Significance Level

MLE of Mean 2528842

MLE of Standard Deviation 1119065

nu star 255.3

Theta Star 495210

Approximate Chi Square Value (.05) 219.3 Nonparametric Statistics

Adjusted Level of Significance 0.0395 95% CLT UCL 2866309
Adjusted Chi Square Value 217.1 95% Jackknife UCL 2879855
95% Standard Bootstrap UCL 2857615

Anderson-Darling Test Statistic 0.259 95% Bootstrap-t UCL 2908346
Anderson-Darling 5% Critical Value 0.747 95% Hall's Bootstrap UCL 2889135
Kolmogorov-Smirnov Test Statistic 0.104 95% Percentile Bootstrap UCL 2848567
Kolmogorov-Smirnov 5% Critical Value 0.175 95% BCA Bootstrap UCL 2883986

Data appear Gamma Distributed at 5% Significance Level 95% Chebyshev(Mean, Sd) UCL 3423136

97.5% Chebyshev (Mean, Sd) UCL 3810098

Assuming Gamma Distribution 99% Chebyshev (Mean, Sd) UCL 4570210

95% Approximate Gamma UCL (Use when n >= 40) 2943900 95% Adjusted Gamma UCL (Use when n < 40) 2974704

Potential UCL to Use Use 95% Student's-t UCL 2879855

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25

Number of Distinct Observations 25

Raw Statistics

Minimum 77.5 Maximum 789.3 Mean 337.9 Median 273.1

Geometric Mean 277.2 SD 207.7 Std. Error of Mean 41.55 Coefficient of Variation 0.615 Skewness 0.845

Log-transformed Statistics

Minimum of Log Data 4.35 Maximum of Log Data 6.671 Mean of log Data 5.625 SD of log Data 0.671

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.905 Shapiro Wilk Critical Value 0.918

95% Student's-t UCL 409

Data not Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.953 Shapiro Wilk Critical Value 0.918 Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 413.7

95% Modified-t UCL (Johnson-1978) 410.1

Assuming Lognormal Distribution

Data Distribution

Data appear Gamma Distributed at 5% Significance Level

95% H-UCL 464.2 95% Chebyshev (MVUE) UCL 558.8 97.5% Chebyshev (MVUE) UCL 652 99% Chebyshev (MVUE) UCL 834.9

Gamma Distribution Test

k star (bias corrected) 2.386 Theta Star 141.6

MLE of Mean 337.9 MLE of Standard Deviation 218.8

nu star 119.3

Approximate Chi Square Value (.05) 95.07

Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 93.6

Anderson-Darling Test Statistic 0.313 Anderson-Darling 5% Critical Value 0.753 Kolmogorov-Smirnov Test Statistic 0.125

Kolmogorov-Smirnov 5% Critical Value 0.176

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution 95% Approximate Gamma UCL (Use when n >= 40) 424 95% Adjusted Gamma UCL (Use when n < 40) 430.6 **Nonparametric Statistics**

95% CLT UCL 406.2 95% Jackknife UCL 409 95% Standard Bootstrap UCL 403.9 95% Bootstrap-t UCL 420.5 95% Hall's Bootstrap UCL 413.9 95% Percentile Bootstrap UCL 407.3 95% BCA Bootstrap UCL 413

95% Chebyshev(Mean, Sd) UCL 519 97.5% Chebyshev(Mean, Sd) UCL 597.4 99% Chebyshev(Mean, Sd) UCL 751.3

Potential UCL to Use

Use 95% Approximate Gamma UCL 424

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25

Number of Distinct Observations 25

Raw Statistics

Minimum 3.63 Maximum 51.5 Mean 27.24 Geometric Mean 22.63 Median 26.7 SD 13.74 Std. Error of Mean 2.748

Coefficient of Variation 0.504

Log-transformed Statistics

Minimum of Log Data 1.289 Maximum of Log Data 3.942 Mean of log Data 3.119 SD of log Data 0.718

Skewness 0.0618

Relevant UCL Statistics

Normal	Distrib	.tion	Toct
INOrmai	DISTRIDI	ıtıon	Test

Shapiro Wilk Test Statistic 0.968 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 31.94

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 31.8 95% Modified-t UCL (Johnson-1978) 31.95

Gamma Distribution Test

k star (bias corrected) 2.535 Theta Star 10.75

MLE of Mean 27.24

MLE of Standard Deviation 17.11

nu star 126.7

Approximate Chi Square Value (.05) 101.7

Adjusted Level of Significance 0.0395

Adjusted Chi Square Value 100.2

Anderson-Darling Test Statistic 0.644

Anderson-Darling 5% Critical Value 0.752 Kolmogorov-Smirnov Test Statistic 0.137

Kolmogorov-Smirnov 5% Critical Value 0.176

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 33.93 95% Adjusted Gamma UCL (Use when n < 40) 34.45

Potential UCL to Use

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.854 Shapiro Wilk Critical Value 0.918

Data not Lognormal at 5% Significance Level

Assuming Lognormal Distribution

95% H-UCL 40.22 95% Chebyshev (MVUE) UCL 48.45 97.5% Chebyshev (MVUE) UCL 56.91 99% Chebyshev (MVUE) UCL 73.52

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 31.76
95% Jackknife UCL 31.94
95% Standard Bootstrap UCL 31.66
95% Bootstrap-t UCL 31.77
95% Hall's Bootstrap UCL 31.85
95% Percentile Bootstrap UCL 31.55
95% BCA Bootstrap UCL 31.7
95% Chebyshev(Mean, Sd) UCL 39.22
97.5% Chebyshev(Mean, Sd) UCL 44.4
99% Chebyshev(Mean, Sd) UCL 54.58

Use 95% Student's-t UCL 31.94

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 23

Raw Statistics

Minimum 0 024 Maximum 0.199 Mean 0.0762 Geometric Mean 0.063 Median 0.068

SD 0.0499 Std. Error of Mean 0.00997 Coefficient of Variation 0.654 Skewness 1.162

Relevant UCL Statistics

Lognormal Distribution Test Normal Distribution Test

Shapiro Wilk Test Statistic 0.863 Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data not Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.0933 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 0.0951 95% Modified-t UCL (Johnson-1978) 0.0937

Gamma Distribution Test

k star (bias corrected) 2.465 Theta Star 0.0309

MLE of Mean 0.0762 MLE of Standard Deviation 0.0486

nu star 123.2

Approximate Chi Square Value (.05) 98.6 Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 97.1

Anderson-Darling Test Statistic 0.506 Anderson-Darling 5% Critical Value 0.752 Kolmogorov-Smirnov Test Statistic 0.144 Kolmogorov-Smirnov 5% Critical Value 0.176

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0953 95% Adjusted Gamma UCL (Use when n < 40) 0.0968

Potential UCL to Use

Log-transformed Statistics

Shapiro Wilk Test Statistic 0.954

Minimum of Log Data -3.73

Maximum of Log Data -1.614

Mean of log Data -2.765

SD of log Data 0.628

Data appear Lognormal at 5% Significance Level

Assuming Lognormal Distribution

95% H-UCL 0.1 95% Chebyshev (MVUE) UCL 0.12 97.5% Chebyshev (MVUE) UCL 0.139 99% Chebyshev (MVUE) UCL 0.177

Data Distribution

Data appear Gamma Distributed at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 0.0926 95% Jackknife UCL 0.0933 95% Standard Bootstrap UCL 0.0921 95% Bootstrap-t UCL 0.0986 95% Hall's Bootstrap UCL 0.0957 95% Percentile Bootstrap UCL 0.0938 95% BCA Bootstrap UCL 0.0946 95% Chebyshev(Mean, Sd) UCL 0.12 97.5% Chebyshev(Mean, Sd) UCL 0.139 99% Chebyshev(Mean, Sd) UCL 0.175

Use 95% Approximate Gamma UCL 0.0953

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (fish) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 22

Raw Statistics

Log-transformed Statistics
Minimum 0.6 Minimum of Log Data -0.511

Maximum 5.1 Maximum of Log Data 1.629
Mean 2.528 Mean of log Data 0.804

Geometric Mean 2.234 SD of log Data 0.544 Median 2.4

SD 1.179
Std. Error of Mean 0.236
Coefficient of Variation 0.466

Skewness 0.453

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.962 Shapiro Wilk Test Statistic 0.933
Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 2.932 95% H-UCL 3.234 **95% UCLs (Adjusted for Skewness)** 95% Chebyshev (MVUE) UCL 3.853

95% Adjusted-CLT UCL (Chen-1995) 2.939 97.5% Chebyshev (MVUE) UCL 4.406 95% Modified-t UCL (Johnson-1978) 2.935 99% Chebyshev (MVUE) UCL 5.494

Gamma Distribution Test Data Distribution

k star (bias corrected) 3.726 Data appear Normal at 5% Significance Level

Theta Star 0.679

MLE of Mean 2.528

MLE of Standard Deviation 1.31 nu star 186.3

95% Adjusted Gamma UCL (Use when n < 40) 3.062

Approximate Chi Square Value (.05) 155.7 Nonparametric Statistics

Adjusted Level of Significance 0.0395 95% CLT UCL 2.916

Adjusted Chi Square Value 153.8 95% Jackknife UCL 2.932 95% Standard Bootstrap UCL 2.911 Anderson-Darling Test Statistic 0.296 95% Bootstrap-t UCL 2.95

Anderson-Darling 5% Critical Value 0.748 95% Hall's Bootstrap UCL 2.951
Kolmogorov-Smirnov Test Statistic 0.126 95% Percentile Bootstrap UCL 2.909

Kolmogorov-Smirnov Test Statistic 0.126 95% Percentile Bootstrap UCL 2.909

Kolmogorov-Smirnov 5% Critical Value 0.175 95% BCA Bootstrap UCL 2.917 **Data appear Gamma Distributed at 5% Significance Level** 95% Chebyshev(Mean, Sd) UCL 3.556

97.5% Chebyshev(Mean, Sd) UCL 4.001

Assuming Gamma Distribution 99% Chebyshev(Mean, Sd) UCL 4.875 95% Approximate Gamma UCL (Use when n >= 40) 3.025

Potential UCL to Use Use 95% Student's-t UCL 2.932

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: GULF MENHADEN, STRIPED MULLET

Gulf Menhaden, Striped Mullet

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 19 Number of Distinct Observations 19

Raw Statistics Log-transformed Statistics

Minimum 103.7 Minimum of Log Data 4.642
Maximum 789.3 Maximum of Log Data 6.671
Mean 405.1 Mean of log Data 5.891

Geometric Mean 361.9 SD of log Data 0.506

Median 426.7 SD 192.8

Std. Error of Mean 44.24

Coefficient of Variation 0.476

Skewness 0.78

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.911 Shapiro Wilk Test Statistic 0.949
Shapiro Wilk Critical Value 0.901 Shapiro Wilk Critical Value 0.901

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 481.8
 95% H-UCL 523.2

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 622.6

 95% Adjusted-CLT UCL (Chen-1995) 486.3
 97.5% Chebyshev (MVUE) UCL 715.3

95% Modified-t UCL (Johnson-1978) 483.1 99% Chebyshev (MVUE) UCL 897.5

Gamma Distribution Test Data Distribution

k star (bias corrected) 3.902 Data appear Normal at 5% Significance Level

Theta Star 103.8

MLE of Mean 405.1

MLE of Standard Deviation 205.1

nu star 148.3

Approximate Chi Square Value (.05) 121.1 Nonparametric Statistics

Adjusted Level of Significance 0.0369 95% CLT UCL 477.9

Adjusted Chi Square Value 119 95% Jackknife UCL 481.8 95% Standard Bootstrap UCL 474.7

Anderson-Darling Test Statistic 0.379 95% Bootstrap-t UCL 497.2

Gulf Menhaden, Striped Mullet

Anderson-Darling 5% Critical Value 0.743 Kolmogorov-Smirnov Test Statistic 0.132 Kolmogorov-Smirnov 5% Critical Value 0.199

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 495.9 95% Adjusted Gamma UCL (Use when n < 40) 504.8

95% Hall's Bootstrap UCL 493.7 95% Percentile Bootstrap UCL 474.9 95% BCA Bootstrap UCL 480.9 95% Chebyshev(Mean, Sd) UCL 597.9 97.5% Chebyshev(Mean, Sd) UCL 681.3 99% Chebyshev(Mean, Sd) UCL 845.2

Potential UCL to Use

Use 95% Student's-t UCL 481.8

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GROUPING: PINFISH, SAND SEATROUT

Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 6

Number of Distinct Observations 6

Raw Statistics

Minimum 77.5 Maximum 188.1 Mean 125.1

Geometric Mean 119.2

Median 123.8 SD 42.2

Std. Error of Mean 17.23

Coefficient of Variation 0.337

Skewness 0.424

Log-transformed Statistics

Minimum of Log Data 4.35

Maximum of Log Data 5.237

Mean of log Data 4.781

SD of log Data 0.341

Warning: A sample size of 'n' = 6 may not adequate enough to compute meaningful and reliable test statistics and estimates!

It is suggested to collect at least 8 to 10 observations using these statistical methods!

If possible compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Warning: There are only 6 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.932 Shapiro Wilk Critical Value 0.788

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.943 Shapiro Wilk Critical Value 0.788

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 159.8

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 156.6

Assuming Lognormal Distribution

95% H-UCL 179.4 95% Chebyshev (MVUE) UCL 201 97.5% Chebyshev (MVUE) UCL 233.8

Pinfish, Sand Seatrout

95% Modified-t UCL (Johnson-1978) 160.3

99% Chebyshev (MVUE) UCL 298.3

Gamma Distribution Test

k star (bias corrected) 5.394

Theta Star 23.19 MLE of Mean 125.1

MLE of Standard Deviation 53.85

nu star 64.73

Approximate Chi Square Value (.05) 47.22

Adjusted Level of Significance 0.0122

Adjusted Chi Square Value 41.87

Anderson-Darling Test Statistic 0.302

Anderson-Darling 5% Critical Value 0.698

Kolmogorov-Smirnov Test Statistic 0.224

Kolmogorov-Smirnov 5% Critical Value 0.332

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n \ge 40$) 171.4 95% Adjusted Gamma UCL (Use when n < 40) 193.3

Potential UCL to Use

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 153.4
95% Jackknife UCL 159.8
95% Standard Bootstrap UCL 150.1
95% Bootstrap-t UCL 168.1
95% Hall's Bootstrap UCL 149.1
95% Percentile Bootstrap UCL 152.3
95% BCA Bootstrap UCL 153.2
95% Chebyshev(Mean, Sd) UCL 200.2

Use 95% Student's-t UCL 159.8

99% Chebyshev(Mean, Sd) UCL 296.5

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

APPENDIX C-3 ESTIMATION OF METHYLMERCURY CONCENTRATION IN SEDIMENT

1 ESTIMATION OF METHYLMERCURY CONCENTRATION IN SEDIMENT

The estimation of bulk sediment methylmercury (MeHg_{sed}) concentrations is based on the equilibrium constants (K) provided in Appendix B of O'Day et al. (2012) and the reference cited therein (Skyllberg 2008). This information is recreated in Table 1.

Table 1

Chemical Reactions for Methylmercury Partitioning

(after O'Day et al. [2012])

Reaction	Log K ¹
(DOM)H + CH3Hg+ = CH3Hg((DOM)) + H+	6.5
(SOM)H + CH3Hg+ = CH3Hg((SOM)) + H+	6.5

Note:

1. Equilibrium partitioning cited are from Skyllberg (2008).

In this case, dissolved organic matter (DOM) and sediment organic matter (SOM) are assumed to be equivalent to the dissolved organic carbon measured in the porewater (DOC $_{pw}$) and total organic carbon measured in the bulk sediment (TOC $_{sed}$). Additionally, as described in Table 1, the partitioning at equilibrium of MeHg in the porewater (MeHg $_{pw}$) to DOC $_{pw}$ and MeHg $_{sed}$ to TOC $_{sed}$ is assumed to be equivalent.

The reactions in Table 1 take the generic form of Equation 1; as a result, their respective K values can be calculated using Equation 2 (Smith 2006). Since these values are the same (Skyllberg 2008), by setting the two equilibrium constant equations for the above reactions equal to each other and cancelling terms, a relationship for estimating MeHg_{sed} can be developed (Equation 3).

$$A + B \Longleftrightarrow C + D \tag{1}$$

$$K = \frac{[C][D]}{[A][B]} \tag{2}$$

$$MeHg_{sed} = MeHg_{pw} \times \frac{TOC_{sed}}{DOC_{pw}}$$
(3)

Using the available data for MeHg_{pw}, TOC_{sed}, and DOC_{pw} (Table 2), an estimated MeHg_{sed} value was calculated for each sample location (Table 3). The MeHg_{pw} data¹ are available for depths from 0 cm to 20 cm; however, it is assumed that MeHg below a 10 cm depth is not available to receptors for ingestion. As a result, the MeHg_{pw} data from each sample location are averaged over the 0 cm to 10 cm interval and used in Equation 3. Additionally, the standard deviation for each average was calculated and used in Equation 3 to determine an estimated standard deviation for MeHg_{sed}.

Data for DOC_{pw} are representative of the sampling interval from 0 cm to 11 cm and are considered averaged values for each of the sample locations.

TOC_{sed} values for PB-023, PB-024, PB-044, PB-053, PB-059.1, and PB-059.2 are from bulk sediment samples collected over the 0 to 10 cm interval. TOC_{sed} for several sites (PB-006, PB-0046, and PB-0052) were sampled over multiple depth intervals from 0 cm to 10 cm; similar to the MeHg_{Pw} data. For these samples, TOC values are averaged and used in Equation 3. Finally, TOC in one sample (PB-0036) is only at the 2 cm depth, but it is assumed that this value is representative, as it does not fall outside the range of the average measured values from other sample locations.

⁻

 $^{^{1}}$ MeHg_{pw} and DOC_{pw} data were collected as part of the Patrick Bayou Mercury Study (August to September 2007).

Table 2
Observed Data Values

Lasakian	MeHg _{pw} (ng/L)		DOC_pw	TOC _{sed}
Location	Average	Standard Deviation	(μg/L)	(%)
PB-006A	2.47	1.48	20,000	1.48
PB-006B	4.7	1.68	19,000	1.48
PB-023	1.27	0.73	22,000	0.64
PB-024	3.46	1.62	24,000	0.92
PB-036	9.51	5.31	42,000	4.36
PB-044	0.99	0.42	12,000	1.26
PB-046	3.28	2.35	26,000	1.81
PB-052	0.32	0.27	26,000	5.12
PB-053	3.05	1.58	57,000	2.06
PB-059.1	0.23	0.07	50,000	1.09
PB-059.2	0.23	0.07	50,000	1

Notes:

- 1. Cells highlighted in blue are duplicated.
- 2. MeHg $_{\rm pw}$ sample values are averaged over the sampling interval from 0 to 10 cm.
- 3. TOC_{sed} values are averaged for samples taken over an interval (e.g., 0 to 10 cm).

Table 3 presents the results of applying Equation 3 to the available data for each sampling station. Two estimated MeHg_{sed} values are given: the average and the standard deviation. Average values are calculated as described above, and the estimated standard deviations in MeHg_{sed} are calculated using the standard deviation for the 0 cm to 10 cm interval from the non-averaged MeHg_{pw} data. The estimated MeHg_{sed} values range from 9,872 ng/kg to 46 ng/kg.

Table 3
Estimated MeHg_{sed} Concentrations

Location	MeHg _{sed} (ng/kg) Average	MeHg _{sed} (ng/kg) Standard Deviation
PB-006A	1,828	1,095
PB-006B	3,661	1,309
PB-023	369	212
PB-024	1,326	621
PB-036	9,872	5,512
PB-044	1,040	441
PB-046	2,283	1,636
PB-052	630	532
PB-053	1,102	571
PB-059.1	51	16
PB-059.2	46	14

Notes:

- Bulk sediment concentrations of MeHg are estimated using the data presented in O'Day et al. (2012) and the references cited therein.
- 2. Based on the research presented above in Note 1, the partitioning of MeHg to the solid and dissolved OC is assumed to be the same at equilibrium.

Variations in sampling location and thus the conditions for each sample are not accounted for in the above estimates (Table 3). The sample locations for the bulk sediment and porewater measurements are provided in Figure C-1.

To quantitatively assess any variations in sampling location an uncertainty analysis was performed for these MeHg_{sed} estimates. Based on Equation 3, increasing values for TOC_{sed} and MeHg_{pw} and lowering values for DOC_{pw} would provide the highest estimates for MeHg_{sed}.

Conversely, lowering values for TOC_{sed} and $MeHg_{pw}$ and increasing values for DOC_{pw} would provide the lowest estimates for $MeHg_{sed}$. This method was used to calculate an upper and lower bound to the average values estimated for $MeHg_{sed}$ (Table 3).

As described above, observed values for DOC_{pw} are considered averaged over the sampled depth interval; therefore, for the uncertainty analysis this parameter is assumed to be constant across the entire Site. The maximum and minimum values are $57,000 \,\mu\text{g/l}$ and $12,000 \,\mu\text{g/l}$, respectively. TOC_{sed} is treated similarly for this analysis, and the maximum and minimum values are 5.54 percent and 0.64 percent, respectively. The remaining parameter, MeHg_{pw}, was collected at various depths for most locations. Maximum and minimum values over the 0 to 10 cm depth interval for MeHg_{pw} at each sampling location were used to calculate the upper and lower bound estimates, respectively.

MeHg_{pw} was sampled over the 0 to 20 cm depth interval, but the 0 to 10 cm depth interval is the dataset used in this analysis for the MeHg_{sed} estimates; this rationale is described above. It should be noted that none of the maximum MeHg_{pw} concentrations occurs below the 0 to 10 cm depth interval; however, there are several instances where the lowest concentration occurs in the 10 to 20 cm depth interval. Using the lowest concentration in the 0 to 10 cm depth interval provides a more conservative estimate; therefore, only concentration values from the 0 to 10 cm depth interval are used for the uncertainty analysis. Additionally, there are two instances where a depth interval has two reported MeHg_{pw} concentrations (PB-006A and PB-044). In each case the higher (and thus more conservative) value is used for the uncertainty analysis.

The results of the uncertainty analysis are plotted in Figure C-2. As expected, the differences in the upper and lower bounds to the average MeHg_{sed} estimates are greater than the standard deviations presented in Table 3. The values estimated for the upper and lower bound concentrations are provided in Table 4. Upper bound estimates range from 67,173 ng/kg to 1,662 ng/kg, and lower bound estimates range from 376 ng/kg to 22 ng/kg.

Table 4

Upper and Lower Bound MeHg_{sed} Estimates

Location	MeHg _{sed} (ng/kg) Upper Bound	MeHg _{sed} (ng/kg) Lower Bound
PB-006A	22,714	144
PB-006B	34,717	376
PB-023	9,603	55
PB-024	22,483	77
PB-036	67,173	117
PB-044	6,648	35
PB-046	29,177	139
PB-052	3,740	22
PB-053	25,161	174
PB-059.1	1,662	22
PB-059.2	1,662	22

Notes:

- 1. Bulk sediment concentrations of MeHg are estimated using the data presented in O'Day et al. (2012) and the references cited therein.
- 2. Based on the research presented above in Note 1, the partitioning of MeHg to the solid and dissolved OC is assumed to be the same at equilibrium.

The MeHgsed results were used, along with the total mercury (HgT) observed in the bulk sediment, to estimate the proportion of MeHg to HgT (Table 5). For the analysis of the average MeHgsed concentrations, observed HgT values are averaged for each sample location; their sample depths and dates are similar to the TOCsed data discussed above. In addition to the averaged values the observed HgT value for each sampling location was used to calculate the upper and lower bounds of the proportion of MeHgsed to HgT. Based on the calculation, the lowest HgT concentrations will maximize the proportion (upper bound), and the highest HgT concentrations will minimize the proportion (lower bound). In general, HgT concentration was sampled over the 0 to 10 cm depth interval; details regarding the sampling interval for each sample location are similar to those described above for TOCsed. These values along with the respective estimated upper and lower bound proportions are provided in Table 6.

Table 5
Proportion of MeHg_{sed} to Total Observed Mercury

Location	Average Hg _T (mg/kg)	Proportion of MeHg _{sed} to Hg $_{\rm T}$
PB-006A	2.07	0.088%
PB-006B	2.07	0.177%
PB-023	2.87	0.013%
PB-024	5.44	0.024%
PB-036	45.60	0.022%
PB-044	13.70	0.008%
PB-046	5.57	0.041%
PB-052	9.64	0.007%
PB-053	843.00	<0.001%
PB-059.1	0.80	0.006%
PB-059.2	1.12	0.004%

Notes:

- 1. Cells highlighted in blue are duplicated.
- 2. Hg_T sample values are averaged over the sampling interval from 0 to 10 cm.

Table 6

Proportion of MeHg_{sed} to Total Observed Mercury – Bounding Cases

	Upper Bou	nd Estimate	Lower Bound Estimate		
Location	Minimum Hg _T (mg/kg)	Proportion of MeHg _{sed} to Hg _T	Maximum Hg _T (mg/kg)	Proportion of MeHg _{sed} to Hg _T	
PB-006A	1.49	1.524%	2.73	0.005%	
PB-006B	1.49	2.330%	2.73	0.014%	
PB-023	2.87	0.335%	2.87	0.002%	
PB-024	5.44	0.413%	5.44	0.001%	
PB-036	45.6	0.147%	45.6	<0.001%	
PB-044	13.7	0.049%	13.7	<0.001%	
PB-046	4.04	0.722%	7.58	0.002%	
PB-052	5.12	0.073%	12.1	<0.001%	
PB-053	843	0.003%	843	<0.001%	
PB-059.1	0.8	0.208%	0.8	0.003%	
PB-059.2	1.12	0.148%	1.12	0.002%	

Notes:

- 1. Cells highlighted in blue are duplicated.
- 2. Hg_T samples are for the 0 to 10 cm depth interval. Some locations only have one sample.


The results of this analysis indicate that the MeHg present in the sediments represents a very small portion of the HgT observed in the bulk sediment samples (0.177 percent to less than 0.001 percent). Based on the uncertainty analysis, upper bound estimates indicate that the proportion of MeHgsed to HgT is less than 2.5 percent. It is likely that other site-specific factors (e.g., mercury sulfide speciation) contribute to the low concentrations of MeHg in the sediments.

2 REFERENCES

- O'Day, P.A., B.A. Bessinger, D. Vlassopoulos, and S. Serrano, 2012. Reactive Transport Modeling of Subaqueous Sediment Caps and Implications for the Long-Term Fate of Arsenic, Mercury, and Methylmercury. Submitted to: *Aquatic Geochemistry*. In Publication.
- Skyllberg, U., 2008. Competition among thiols and inorganic sulfides and polysulfides for Hg and MeHg in wetland soils and sediments under suboxic conditions: Illumination of controversies and implications for MeHg net production. *J Geophys Res -Biogeosci* 113:536-554. doi:G00c0310.1029/2008jg000745

Smith, J.G., 2006. Organic Chemistry. Boston: McGraw Hill.

FIGURES










Figure C-2

APPENDIX C-4 BASELINE RISK ASSESSMENT

TABLE C-4.1 EXPOSURE VARIABLES FOR WILDLIFE TOTAL DAILY INTAKE MODELS

Table C-4.1 Exposure Variables for Wildlife Total Daily Intake Models

Parameter	Receptor Group: Receptor: Units	Piscivorous Birds Belted Kingfisher		Sediment-probing Birds Spotted Sandpiper		Carnivorous E Composit		Omnivorous Mai Raccoon	nmals
BW	kg	0.136	а	0.0425	а	0.870	а	3.99	а
NFMR	kcal/day/kg BW	6722	b	4016	b	3570	b	922.6	b
а	unitless	14.25	g	8.13	h	14.25	g	6.03	i
b	unitless	0.659	g	0.77	h	0.659	g	0.678	i
AE _{fish}	unitless	0.79	С	NA		0.79	С	0.91	С
AE _{invertebrates}	unitless	0.77	С	0.77	С	0.77	С	0.91	С
AE _{Plants}	unitless	NA		NA		NA		0.74	j
AE _{Animals}	unitless	NA		NA		0.78	0	0.84	k
GE _{fish}	kcal/kg	1200	С	NA		1200	С	1200	С
GE _{invertebrates}	kcal/kg	1070	С	1070	С	1070	С	1070	С
GE _{Plants}	kcal/kg	NA		NA		NA		1300	1
GE _{Animals}	kcal/kg	NA		NA		1700	m	1700	m
P_{fish}	unitless	0.89	n	0	n	0.33	n	0.233	n
P _{invertebrates}	unitless	0.11	n	1	n	0.33	n	0.233	n
P _{plants}	unitless	0	n	0	n	0	n	0.3	n
P _{mammals}	unitless	0	n	0	n	0.33	n	0.233	n
FS	unitless	0	f	0.18	d	0.01	f	0.094	e
P _{moisture}	unitless	0.74	q	0.72	q	0.69	q	0.69	q
AUF	unitless	0.69	р	0.34	р	0.36	р	0.021	р

Notes:

- a Average for adults, both sexes (see text for derivation).
- b Derived using Equation 6-7 in text.
- c See AE and GE derivation worksheet in Appendix A.
- d USEPA 1993. Semi-palmated sandpiper value used as surrogate.
- e Based on empirical data (Beyer et al. 1994).
- f Best professional judgment.
- g Power (b) and slope (a) terms for marine birds (Nagy 1999).
- h Power (b) and slope (a) terms for Charadriiformes (Nagy 1999).
- i Power (b) and slope (a) terms for omnivores (Nagy 1999).

- j Average AE for mammals consuming plants (USEPA 1993).
- k Average for mammals consuming small birds & mammals (USEPA 1993).
- I Average of terrestrial monocot grasses (USEPA 1993).
- m Average of reported values for mice, voles, and rabbits (USEPA 1993).
- n See text for derivation.
- o Average of birds of prey consuming birds and small mammals (USEPA 1993).
- p See text for discussion.

Table C-4.1 Exposure Variables for Wildlife Total Daily Intake Models

q Percent moisture calculated based on values in USEPA (1993). Based on a weighted average relative to the proportion of food in receptors diet. Moisture for each prey class based on the following:

Fish: Average of bony fishes in Table 4-1 of USEPA (1993) = 74%

Invertebrates: Average of aquatic invertebrates in Table 4-1 of USEPA (1993) = 72%

Mammals: Terrestrial mammal average in Table 4-1 of USEPA (1993) = 68%

Plants: Midpoint of aquatic emergent vegetation range in Table 4-2 of USEPA (1993) = 63%

NA - not applicable

TABLE C-4.2 EXPOSURE VARIABLES FOR CALCULATING MARSH RICE RAT COPC CONCENTRATION

Table C-4.2 Exposure Variables for Calculating Marsh Rice Rat COPC Concentration

		Mercury				PCB Congener	TEQ	PCB Congener	TEQ	
Variable	Definition	Units	Methyl Mero	Methyl Mercury Inorganic Mercury		(Mammal)	1	(Avian)		
Kow	Log octanol/water partition coefficient	unitless	2.54	а	log(0.61)	i	6.64	m	6.64	m
FCM _{TL3(OM)}	Food chain multiplier for trophic level 3 (marsh rice rat)	unitless	1.0	b	1.0	g	14	n	14	n
C _{SED}	COPC concentration in sediment	mg/kg dw	4.79E-02	С	4.74E+00	С	1.12E-04	С	2.08E-03	С
BCF _{PLANT-OM}	Plant-to-omnivore BCF	unitless	3.44E-04	d	2.30E-03	k	2.39E-02	0	2.39E-02	0
BCF _{SED-OM}	Sediment-to-omnivore BCF	unitless	8.34E-06	е	5.57E-05	I	1.70E-04	р	1.70E-04	р
FR _{SED}	Fraction of sediment in diet that is contaminated	unitless	1	f	1	f	1	f	1	f
$TL_{BIOTA,a}$	Trophic Level of Food Type 1	unitless	3	j	3	j	3	j	3	j
$TL_{BIOTA,b}$	Trophic Level of Food Type 2	unitless	3	j	3	j	3	j	3	j
TL _{BIOTA,c}	Trophic Level of Food Type 3	unitless	2	j	2	j	2	j	2	j
TL _{BIOTA,e}	Trophic Level of Food Type 5	unitless							3	j
FCM _{TL(BIOTA,a)}	Food Chain Multiplier for Food Type 1	unitless	1	b	1	i	14	n	14	n
FCM _{TL(BIOTA,b)}	Food Chain Multiplier for Food Type 2	unitless	1	b	1	i	14	n	14	n
FCM _{TL(BIOTA,c)}	Food Chain Multiplier for Food Type 3	unitless	1	b	1	i	1	n	1	n
FCM _{TL(BIOTA,e)}	Food Chain Multiplier for Food Type 3	unitless							14	n
FR _{BIOTA,k} (all Food Types)	Fraction of kth Food Group that is contaminated	unitless	1	f	1	f	1	f	1	f
P _{BIOTA,a}	Portion of diet consisting of Food Type 1	unitless	0.125	t	0.125	t	0.125	t	0.083	t
P _{BIOTA,b}	Portion of diet consisting of Food Type 2	unitless	0.125	t	0.125	t	0.125	t	0.083	t
P _{BIOTA,c}	Portion of diet consisting of Food Type 3	unitless	0.25	t	0.25	t	0.25	t	0.25	t
P _{BIOTA,d}	Portion of diet consisting of Food Type 4	unitless	0.5	t	0.5	t	0.5	t	0.5	t
P _{BIOTA,e}	Portion of diet consisting of Food Type 5	unitless							0.083	t
C _{BIOTA,a}	COPC concentration in Food Type 1	mg/kg ww	1.45E-01	q			8.70E-05	q	8.68E-04	q
C _{BIOTA,b}	COPC concentration in Food Type 2	mg/kg ww	5.80E-02	q			3.19E-05	q	4.24E-04	q
C _{BIOTA,c}	COPC concentration in Food Type 3	mg/kg ww	5.15E-02	q			2.79E-05	q	4.52E-04	q
C _{BIOTA,d}	COPC concentration in Food Type 4	mg/kg ww	2.43E-03	r	6.60E-02	r	2.32E-07	S	4.35E-06	h
C _{BIOTA,e}	COPC concentration in Food Type 5	mg/kg ww							1.60E-04	q

Notes:

- a Maximum value in methylmercury Kow range (Environment Canada 2002)
- b USEPA (1999) Table 5-2 based on a Kow of 3
- c Table 6-4
- d USEPA (1999) Table D-1 value for methyl mercury
- e USEPA (1999) Table D-3 value for methyl mercury
- f USEPA (1999) Table F-1-14 Default
- q USEPA (1999) Table 5-2 based on a Kow of 2 (smallest Kow value provided)
- h PCB Congener TEQ (Avian) Plant Concentration Calculation, BERA Appendix C
- USEPA (1999) Table A-2-130 value for mercuric chloride
 USEPA (1999) and Professional judgment. All fish were assumed to belong to Trophic
- j level of 3 (omnivore) and invertebrates were assumed to be belong to trophic level 2 (herbivore)

- k USEPA (1999) Table D-1 value for mercuric chloride
- / USEPA (1999) Table D-3 value for mercuric chloride
- m USEPA (1999) Table A-2-176
- n USEPA (1999) Table 5-2 based on a Kow of 6.6
- o USEPA (1999) Table D-1 value for 2,3,7,8-TCDD
- p USEPA (1999) Table D-3 value for 2,3,7,8-TCDD
- q Table 6-3
- r Mercury Plant Concentration Calculation, BERA Appendix C
- s PCB Congener TEQ (Mammal) Plant Concentration Calculation, BERA Appendix C Marsh Rice Rat diet indicated in BERA Workplan (Anchor QEA 2011). Diet was
- t assumed to be composed of 50% plant matter and 50% animal (25% invertebrates and 25% fish)

EXHIBIT C-4.1 WILDLIFE DAILY INTAKE EQUATIONS

Exhibit C-4.1

EQUATIONS	
GENERAL	
$TDI_{ALL} = TDI_{BIOTA} + TDI_{SEDIMENT}$	Equation 1
TISSUE	
$TDI_{BIOTA} = \sum_{k=1}^{m} (C_{BIOTA_k} \times FR_k \times NIR_k)$	Equation 2
$NIR_{TOTAL} = \frac{NFMR}{ME_{AVG}}$	Equation 3
$ME_{AVG} = \sum_{k=1}^{m} (P_{BIOTA,k} \times ME_k)$	Equation 4
$ME_k = GE_k \times AE_k$	Equation 5
$NIR_k = P_{BIOTA,k} \times NIR_{TOTAL}$	Equation 6
$NFMR = \frac{\left[\left(a \times BW_*^b\right) \times CF_1\right]}{BW}$	Equation 7
SEDIMENT	
$TDI_{SEDIMENT} = \sum_{j=1}^{i} \left(C_{SED,j} \times FS \times IR_{TOTAL} \times P_{SED,j} \right)$	Equation 8
$IR_{TOTAL} = (NIR_{TOTAL} \times BW) \times (1 - P_{MOISTURE})$	Equation 9

	VARIABLES				
	DESCRIPTION	UNITS			
TDI _{ALL}	Total daily intake of COPC from all sources	mg _{COPC} /kg _{BW} /day			
TDI _{BIOTA}	Total daily intake of COPC from ingestion of food items	mg _{COPC} /kg _{BW} /day			
TDI _{SEDIMENT}	Total daily intake of COPC from incidental ingestion of sediment	mg _{COPC} /kg _{BW} /day			
$C_{BIOTA,k}$	COPC concentration in kth food type, wet weight	mg _{COPC} /kg _{food}			
FR_k	Fraction of kth type of food that is contaminated	unitless			
NIR_k	Normalized food ingestion rate of kth type of food, wet weight	kg _{food} /kg _{BW} /day			
NIR _{TOTAL}	Total normalized food ingestion rate, wet weight	kg _{food} /kg _{BW} /day			
NFMR	Body weight normalized free-living metabolic rate	kcal/kg _{BW} /day			
ME_{AVG}	Average metabolizable energy of the diet, wet weight	kcal/kg _{food}			
P _{BIOTA,k}	Proportion of the diet consisting of the kth food type	unitless			
ME _k	Metabolizable energy of the kth food type, wet weight	kcal/kg _{food}			
GE_k	Gross energy content of the kth food type, wet weight	kcal/kg _{food}			
AE _k	Assimilation efficiency by receptor of kth food type	unitless			
BW	Body weight	kilograms			
BW*	Body weight	grams			
а	Free-living metabolic rate slope factor	unitless			

Exhibit C-4.1

	VARIABLES	
	DESCRIPTION	UNITS
b	Free-living metabolic rate power	unitless
CF ₁	Conversion factor	1 kcal/4.186 kjoules
$C_{SED,j}$	Sediment COPC concentration in the jth foraging area, dry weight	mg _{copc} /kg
FS	Fraction of sediment in diet; as proportion of ingestion rate	unitless
IR _{TOTAL}	Total daily food ingestion rate, dry weight	kg _{food} /day
$P_{SED,j}$	Proportion of food intake from jth foraging area	unitless
P _{MOISTURE}	Proportion of moisture in food	unitless


Definitions:

mg – milligrams kg – kilograms kcal – kilocalories kjoules – kilojoules

EXHIBIT C-4.2 HOME RANGE DISTRIBUTION ANALYSIS

Home Range Distribution Analysis (kilometers)¹

Shorebirds²



Quantiles

100.0%	maximum	9.6
99.5%		9.6
97.5%		9.6
90.0%		9.6
75.0%	quartile	7.225
50.0%	median	4
25.0%	quartile	2.2875
10.0%		1.5
2.5%		1.5
0.5%		1.5
0.0%	minimum	1.5


Moments

Mean	4.65625
Std Dev	2.8381882
Std Err Mean	1.0034511
Upper 95% Mean	7.0290348
Lower 95% Mean	2.2834652
N	8

¹ See Section 6 (Table 6-5) for list of home ranges used in calculations.

² Distributions, quantiles, and moments calculated using JMP® 9.0.2 (SAS 2010; www.jmp.com)

Herons and Egrets




Quantiles

100.0%	maximum	15
99.5%		15
97.5%		15
90.0%		14.1
75.0%	quartile	6.8875
50.0%	median	5
25.0%	quartile	2.7
10.0%		1.52
2.5%		1.4
0.5%		1.4
0.0%	minimum	1.4

Moments

Mean	5.6875
Std Dev	4.1643577
Std Err Mean	1.2021465
Upper 95% Mean	8.3334066
Lower 95% Mean	3.0415934
N	12

Belted Kingfisher



Quantiles

100.0%	maximum	4.4
99.5%		4.4
97.5%		4.4
90.0%		4.4
75.0%	quartile	3.6
50.0%	median	1.6
25.0%	quartile	1
10.0%		0.4
2.5%		0.4
0.5%		0.4
0.0%	minimum	0.4

Moments

Mean	2.0285714
Std Dev	1.4761598
Std Err Mean	0.5579359
Upper 95% Mean	3.3937915
Lower 95% Mean	0.6633514
N	7

CARNIVOROUS BIRD BASELINE RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.165	0.054	948	Fish	1	GM
2	0.165	0.054	948	Fish	1	GKF, STM, PNF, SAS
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1
1 value of loraging subareas $y = 1$ to y .	

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Baseline

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C_{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Baseline

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.018125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

Receptor: Composite
Receptor Group: Carnivorous Birds

Scenario: Baseline

COPC	C _{BIOTA,1}	$C_{BIOTA,2}$	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA},6}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
000000000000000000000000000000000000000	mg/kg (wet weight)					mg/kg _{BW} /day	g/kg _{BW} /day mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA			
Mercury (Methyl)	0.145	0.058	0.0515	3.83E-02				6.7E-03	0.0479							1.3E-05	6.7E-03	0.064	0.11
Mercury (Inorganic)				3.40E-04				1.9E-05	4.7391							1.3E-03	1.3E-03	0.9	0.0014

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Plant PCB Congener TEQ (Avian) Calculations

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

where:

 C_{PLANT} = Concentration of PCB Congener TEQ (Avian) in plant tissue C_{SED} = Concentration of PCB Congener TEQ (Avian) in sediment BAF_{PLANT} = Bioaccumulation factor for PCB (kgSED / kgPREY)

$$C_{PLANT, \ WW} = C_{PLANT,DW} \times \left(\frac{(100 - \% \ moisture)}{100}\right)$$

where:

C_{PLANT.WW} = Concentration in plant tissue as wet weight

C_{PLANT,DW} = Concentration in plant tissue as dry weight

% moisture = Moisture content of plant tissue

	C _{SED} (mg/kg)	BAF _{PLANT*}	C _{PLANT,DW} (mg/kg)	C _{PLANT,WW} (mg/kg)
PCB	2.10E-03	0.0056	1.18E-05	4.35E-06

^{*}USEPA (1999) Table C-2, BCF for 2,3,7,8-TCDD

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.11	0.036	948	Fish	1	GKF
2	0.11	0.036	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5	0.11	0.036	948	Fish	1	PNF, SAS
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Baseline

COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0021	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.21E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	3.52E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5	1.33E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.95E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.71E-03	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.083	8.68E-04	Table C-4.2
2	Fish	GM, STM	3	14	1	0.083	4.24E-04	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-4.2
4	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
5	Fish	PNF, SAS	3	14	1	0.083	1.60E-04	Table C-4.2
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)				mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{lw} /day	NA				
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	1.71E-03	0.0002			7.1E-05	0.0021							5.5E-07	7.2E-05	1.4E-04	0.51

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

BELTED KINGFISHER BASELINE RISK SHEETS

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.607	948	Fish	1	GKF, GM, STM, PNF, SAS
2	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	1		mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	(g (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Hexachlorobenzene	0.6464	0.9757						3.2E-01	2.1373							0.0E+00	3.2E-01	3.35	0.10
1,3-Dichlorobenzene	0.769	0.758						3.6E-01	0.0736							0.0E+00	3.6E-01	160.8	0.0022

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.445	0.304	948	Fish	1	GM
2	0.445	0.304	948	Fish	1	GKF, STM, PNF, SAS
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

3

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{sw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515					4.5E-02	0.0479							0.0E+00	4.5E-02	0.064	0.71
Mercury (Inorganic)								0.0E+00	4.7391				_			0.0E+00	0.0E+00	0.9	0

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.3	0.205	948	Fish	1	GKF
2	0.3	0.205	948	Fish	1	GM, STM
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.29	0.205	948	Fish	1	PNF, SAS
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA},6}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)					mg/kg _{BW} /day	g/kg _{BW} /day mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA		
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	0.0002				2.4E-04	0.0021							0.0E+00	2.4E-04	0.00014	1.69

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON BASELINE RISK SHEETS

Plant Mercury Calculations

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

where:

C_{PLANT} = Concentration of mercury in plant tissue

C_{SED} = Concentraiton of mercury in sediment

BAF_{PLANT} = Bioaccumulation factor for mercury (kgSED / kgPREY)

$$C_{PLANT, WW} = C_{PLANT,DW} \times \left(\frac{(100 - \% moisture)}{100}\right)$$

where:

C_{PLANT,WW} = Concentration in plant tissue as wet weight

C_{PLANT,DW} = Concentration in plant tissue as dry weight

% moisture = Moisture content of plant tissue

	C _{SED} (mg/kg)	BAF _{PLANT}	C _{PLANT,DW} (mg/kg)	C _{PLANT,WW} (mg/kg)
Methylmercury	0.048	0.137	0.007	0.002
Inorganic Mercury	4.74	0.0375	0.178	0.066

Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GM
2	0.116	0.010	1092	Fish	1	GKF, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Baseline

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	
and the second second			

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight ww - wet weight Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Baseline COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C_{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.81E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	7.25E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.29E-02	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant				1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight ww - wet weight Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{BIOTA,3}$	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
				mg/kg (wet weight)				mg/kg _{BW} /day			mg/k	g (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_w /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	0.002426492	3.83E-02			7.7E-05	0.0479							8.9E-06	8.6E-05	0.0177	0.0049
Mercury (Inorganic)				0.065754378	3.40E-04			3.8E-05	4.7391							8.8E-04	9.1E-04	3.54	0.00026

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Plant PCB Congener TEQ (Mammal) Calculations

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

where:

 C_{PLANT} = Concentration of PCB Congener TEQ (Mammal) in plant tissue

C_{SED} = Concentration of PCB Congener TEQ (Mammal) in sediment

BAF_{PLANT} = Bioaccumulation factor for PCB Congener TEQ (Mammal) (kgSED / kgPREY)

$$C_{PLANT, \ WW} = C_{PLANT,DW} imes \left(\frac{(100 - \% \ moisture)}{100} \right)$$

where:

C_{PLANT.WW} = Concentration in plant tissue as wet weight

C_{PLANT.DW} = Concentration in plant tissue as dry weight

% moisture = Moisture content of plant tissue

	C _{SED} (mg/kg) BAF _{PLANT*}		C _{PLANT,DW} (mg/kg)	C _{PLANT,WW} (mg/kg)	
PCB	1.12E-04	0.0056	6.27E-07	2.32E-07	

^{*}USEPA (1999) Table C-2, BCF for 2,3,7,8-TCDD

Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GKF
2	0.116	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1
---	---

		<u> </u>
Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		
	1 2 3 4 5	1 1.00 2 3 4 5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Baseline

COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.09E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	3.99E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	9.75E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	1.91E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.12E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.125	8.70E-05	Table C-4.2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.125	3.19E-05	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	2.79E-05	Table C-4.2
4	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor:

Raccoon

Receptor Group:

Omnivorous Mammal

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			m	g/kg (wet we	eight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	3.2E-05 2	2.8E-05	2.32E-07	1.12E-04			6.4E-08	0.0001							2.1E-08	8.4E-08	5.4E-07	0.16

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

SPOTTED SANDPIPER BASELINE RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: Baseline

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet we	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Lead	0.292							9.9E-02	48.944							3.5E-02	1.3E-01	3.26	0.04
Mercury (Methyl)	0.0533							1.8E-02	0.0479							3.4E-05	1.8E-02	0.064	0.28
Total PAH 16 (U=1/2)	0.1397							4.7E-02	7.3078							5.2E-03	5.2E-02	325	0.00016
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							7.5E-06	0.0002							1.5E-07	7.7E-06	0.00014	0.05
Hexachlorobenzene	1.132							3.8E-01	2.1373							1.5E-03	3.8E-01	3.35	0.11
Hexachlorobutadiene	0.3725							1.3E-01	3.8327							2.7E-03	1.3E-01	20	0.006
1,3-Dichlorobenzene	0.3725							1.3E-01	0.0736							5.2E-05	1.3E-01	160.8	0.0008
1,4-Dichlorobenzene	0.3725							1.3E-01	0.0744							5.2E-05	1.3E-01	160.8	0.0008
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.4E-04	0.0021							1.5E-06	1.5E-04	1.4E-04	1.0
Mercury (Inorganic)								0.0E+00	4.7391							3.3E-03	3.3E-03	9.0E-01	0.004

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5 WILDLIFE RISK ASSESSMENT UNCERTAINTY ANALYSIS

APPENDIX C-5.1 PREY GROUPING UNCERTAINTY ANALYSIS – DIET PROPORTIONS ADJUSTED BY PREY BIOMASS

Table C-5.1
Diet Proportions Adjusted by Prey Biomass

_		Total	Proportion	Proportion	Biomass Weighted							
Receptor	Grouping	Biomass	of Biomass	of Diet	Proportion of Diet							
	Mercury	•	•		•							
	GM	3,764	0.13	0.89	0.11							
	GKF, STM, PNF, SAS	25,398	0.87	0.89	0.78							
	BCR, BRS, WHS, OYS	9,686	1	0.11	0.11							
	Hexachlorobenzene											
	GKF, GM, STM, PNF, SAS	29,162	1	0.89	0.89							
	BCR, BRS, WHS, OYS	9,686	1	0.11	0.11							
Belted Kingfisher	1,3-Dichlorobenzene											
	GKF, GM, STM, PNF, SAS	29,162	1	0.89	0.89							
	BCR, BRS, WHS, OYS	9,686	1	0.11	0.11							
	PCB TEQ		-									
	GKF	15,235	0.52	0.89	0.46							
	GM, STM	7,700	0.26	0.89	0.24							
	PNF, SAS	6,226	0.21	0.89	0.19							
	BCR, BRS, WHS, OYS	9,686	1	0.11	0.11							
	Mercury											
	GM	3,764	0.13	0.33	0.04							
	GKF, STM, PNF, SAS	25,398	0.87	0.33	0.29							
	BCR, BRS, WHS, OYS	9,686	1	0.33	0.33							
Carnivorous Bird	PCB TEQ											
	GKF	15,235	0.52	0.33	0.17							
	GM, STM	7,700	0.26	0.33	0.09							
	PNF, SAS	6,226	0.21	0.33	0.07							
	BCR, BRS, WHS, OYS	9,686	1	0.33	0.33							
	Mercury											
	GM	3,764	0.13	0.233	0.03							
	GKF, STM, PNF, SAS	25,398	0.87	0.233	0.20							
Raccoon	BCR, BRS, WHS, OYS	9,686	1	0.234	0.23							
Naccoon	PCB TEQ											
	GKF	15,235	0.52	0.233	0.12							
	GM, STM, PNF, SAS	13,927	0.48	0.233	0.11							
	BCR, BRS, WHS, OYS	9,686	1	0.234	0.23							

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.042594	0.014	948	Fish	1	GM
2	0.287406	0.094	948	Fish	1	GKF, STM, PNF, SAS
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

4

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: **Terrestrial Mammal - Marsh Rice Rat** Scenario: **Biomass Based Proportion of Prey Groups**

COPC: **Methyl Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.0047125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0.012615	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.02E-02	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:		_	

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.0325	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.2175	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Biomass Based Proportion of Prey Groups

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.0325		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.2175		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	C _{BIOTA,3}	$C_{\text{BIOTA,4}}$	C _{BIOTA,5}	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg	r/kg (wet weig	ht)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	3.02E-02				5.3E-03	0.0479							1.3E-05	5.3E-03	0.064	0.08
Mercury (Inorganic)				3.40E-04				1.9E-05	4.7391							1.3E-03	1.3E-03	0.9	0.0014

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.17	0.056	948	Fish	1	GKF
2	0.09	0.029	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5	0.07	0.029	948	Fish	1	PNF, SAS
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: **Terrestrial Mammal - Marsh Rice Rat**

Scenario: **Biomass Based Proportion of Prey Groups**

COPC: **PCB Congener TEQ (Avian)**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C_SED	0.0021	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.13E-04	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	2.76E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5	8.79E-06	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.95E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.74E-03	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Number of food types $(k = 1 \text{ to } m)$:	4
--	---

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.13	8.68E-04	Table C-4.2
2	Fish	GM, STM	3	14	1	0.065	4.24E-04	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-4.2
4	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
5	Fish	PNF, SAS	3	14	1	0.055	1.60E-04	Table C-4.2
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	C _{SED,6}	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)							mg/kg _{BW} /day	mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0009 0.0005 0.0005 1.74E-03 0.0002							0.0021		•				·	5.5E-07	7.7E-05	1.4E-04	0.55

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.607	948	Fish	1	GKF, GM, STM, PNF, SAS
2	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	$C_{BIOTA,6}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)							mg/kg _{BW} /day	mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Hexachlorobenzene	0.6464	0.9757						3.2E-01	2.1373							0.0E+00	3.2E-01	3.35	0.10
1,3-Dichlorobenzene	0.769	0.758						3.6E-01	0.0736				<u> </u>			0.0E+00	3.6E-01	160.8	0.0022

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.114875	0.078	948	Fish	1	GM
2	0.775125	0.529	948	Fish	1	GKF, STM, PNF, SAS
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

COPC	$C_{BIOTA,1}$	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	$C_{BIOTA,6}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)							mg/kg _{BW} /day	mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg į	_{sw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515					3.2E-02	0.0479							0.0E+00	3.2E-02	0.064	0.49
Mercury (Inorganic)								0.0E+00	4.7391							0.0E+00	0.0E+00	0.9	0

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.46	0.314	948	Fish	1	GKF
2	0.24	0.164	948	Fish	1	GM, STM
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.19	0.164	948	Fish	1	PNF, SAS
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{\text{BIOTA,4}}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	C _{SED,5}	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)				mg/kg _{BW} /day mg/kg (dry weight)				mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA							
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	0.0002				2.8E-04	0.0021							0.0E+00	2.8E-04	0.00014	2.03

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.030074	0.003	1092	Fish	1	GM
2	0.202926	0.018	1092	Fish	1	GKF, STM, PNF, SAS
3	0.233	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6		<u> </u>				
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat
Scenario: Biomass Based Proportion of Prey Groups

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	4.71E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	1.26E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.29E-02	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.02E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.0325	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.2175	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant				1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight ww - wet weight Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Biomass Based Proportion of Prey Groups

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C_{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.0325		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.2175		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Raccoon

Omnivorous Mammal

Receptor Group: Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
mg/kg (wet weight)							mg/kg _{BW} /day	mg/kg (dry weight) mg/kg _E						mg/kg _{BW} /day	mg/kg	_{BW} /day	NA		
Mercury (Methyl)	0.145	0.058	0.0515	0.002426492	3.02E-02			6.1E-05	0.0479							8.9E-06	7.0E-05	0.0177	0.0040
Mercury (Inorganic)				0.065754378	3.40E-04			3.8E-05	4.7391							8.8E-04	9.1E-04	3.54	0.00026

Baseline Ecological Risk Assessment March 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME _{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.121727	0.011	1092	Fish	1	GKF
2	0.111273	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.233	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat Scenario: Biomass Based Proportion of Prey Groups

COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.13E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	3.83E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	9.75E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	1.91E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.13E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.13	8.70E-05	Table C-4.2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.12	3.19E-05	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	2.79E-05	Table C-4.2
4	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	$C_{BIOTA,5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
mg/kg (wet weight)						mg/kg _{BW} /day	/ mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA			
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	3.2E-05	2.8E-05	2.32E-07	1.13E-04			6.4E-08	0.0001							2.1E-08	8.5E-08	5.4E-07	0.16

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper

Receptor Group: Sediment-probing Birds

Scenario: Biomass Based Proportion of Prey Groups

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: Biomass Based Proportion of Prey Groups

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)						mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Lead	0.292							9.9E-02	48.944							3.5E-02	1.3E-01	3.26	0.04
Mercury (Methyl)	0.0533							1.8E-02	0.0479							3.4E-05	1.8E-02	0.064	0.28
Total PAH 16 (U=1/2)	0.1397							4.7E-02	7.3078							5.2E-03	5.2E-02	325	0.00016
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							7.5E-06	0.0002							1.5E-07	7.7E-06	0.00014	0.05
Hexachlorobenzene	1.132							3.8E-01	2.1373							1.5E-03	3.8E-01	3.35	0.11
Hexachlorobutadiene	0.3725							1.3E-01	3.8327							2.7E-03	1.3E-01	20	0.01
1,3-Dichlorobenzene	0.3725							1.3E-01	0.0736							5.2E-05	1.3E-01	160.8	0.0008
1,4-Dichlorobenzene	0.3725							1.3E-01	0.0744							5.2E-05	1.3E-01	160.8	0.0008
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.4E-04	0.0021	Ţ						1.5E-06	1.5E-04	1.4E-04	1.05
Mercury (Inorganic)								0.0E+00	4.7391							3.3E-03	3.3E-03	9.0E-01	0.004

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.2 PREY GROUPING UNCERTAINTY ANALYSIS – GROUPED BY FISH AND SHELLFISH

Table C-5.2 EPCs for All Fish Species and All Invertebrate Species Prey Groups

	Exposure Point Concentration (mg/kg)		
СОРС	Gulf Killifish, Gulf Menhaden, Pinfish, Sand Seatrout, Striped Mullet	Blue Crab, Brown Shrimp, White Shrimp, Oyster	Blue Crab, Oyster, White Shrimp
1,3-Dichlorobenzene	0.769 ^a	0.758 ^a	0.373 ^a
1,4-Dichlorobenzene			0.373 ^a
Dioxin/Furan TEQ		-1	0.0000223
Hexachlorobenzene	0.645	0.950	1.13
Hexachlorobutadiene			0.373 ^a
Lead		-1	0.292
Mercury	0.0893	0.0515	0.0533
PCB Congener TEQ (Avian)	0.00076	0.000452	0.000430
PCB Congener TEQ (Mammal	0.000062	0.0000279	
Total PAH			0.140

Notes:

a - Maximum non-detect value used

ALL FISH SPECIES COMBINED 95 UCL OUTPUTS

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	50	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	50
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

	General Stat	istics	
Number of Valid Data	50	Number of Detected Data	28
Number of Distinct Detected Data	27	Number of Non-Detect Data	22
		Percent Non-Detects	44.00%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	200	Minimum Detected	5.298
Maximum Detected	2360	Maximum Detected	7.766
Mean of Detected	680	Mean of Detected	6.28
SD of Detected	576.9	SD of Detected	0.66
Minimum Non-Detect	351	Minimum Non-Detect	5.861
Maximum Non-Detect	765	Maximum Non-Detect	6.64
Note: Data have multiple DLs - Use of KM Method is recommer	nded	Number treated as Non-Detect	43
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	7
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	86.00%
	UCL Statis	tics	
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values C	Only
Shapiro Wilk Test Statistic	0.709	Shapiro Wilk Test Statistic	0.923
5% Shapiro Wilk Critical Value	0.924	5% Shapiro Wilk Critical Value	0.924
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	519.2	Mean	6.029
SD	468.9	SD	0.603
95% DL/2 (t) UCL	630.3	95% H-Stat (DL/2) UCL	590.3
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE yields a negative mean		Mean in Log Scale	6.056
		SD in Log Scale	0.573
		Mean in Original Scale	525.4
		SD in Original Scale	465.4
		95% t UCL	635.7
		95% Percentile Bootstrap UCL	635.7
		95% BCA Bootstrap UCL	665.7
		95% H-UCL	589.2

Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	
k star (bias corrected)	2.004	Data do not follow a Discernable Distribution (0.05)	
Theta Star	339.3		
nu star	112.2		
A-D Test Statistic	1.384	Nonparametric Statistics	
5% A-D Critical Value	0.757	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.757	Mean	528.3
5% K-S Critical Value	0.167	SD	463.9
Data not Gamma Distributed at 5% Significance Level		SE of Mean	68.18
		95% KM (t) UCL	642.6
Assuming Gamma Distribution		95% KM (z) UCL	640.4
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	641.9
Minimum	98.03	95% KM (bootstrap t) UCL	682.8
Maximum	2360	95% KM (BCA) UCL	647.2
Mean	546.6	95% KM (Percentile Bootstrap) UCL	644.8
Median	474.4	95% KM (Chebyshev) UCL	825.5
SD	470.9	97.5% KM (Chebyshev) UCL	954.1
k star	1.973	99% KM (Chebyshev) UCL	1207
Theta star	277		
Nu star	197.3	Potential UCLs to Use	
AppChi2	165.8	95% KM (t) UCL	642.6
95% Gamma Approximate UCL (Use when n >= 40)	650.4	95% KM (% Bootstrap) UCL	644.8
95% Adjusted Gamma UCL (Use when n < 40)	653.8		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Raw Statistics	Log-transformed Statistics
Number of Valid Observations 50	Number of Distinct Observations 37

Kaw Statistics	Log-transformed Statistics
Minimum 0.024	Minimum of Log Data -3.73
Maximum 0.199	Maximum of Log Data -1.614
Mean 0.0643	Mean of log Data -2.904
Geometric Mean 0.0548	SD of log Data 0.551
Median 0.0465	
SD 0.0406	
Std. Error of Mean 0.00575	

Relevant UCL Statistics

Coefficient of Variation 0.632

Skewness 1.635

Normal Distribution Test	Lognormal Distribution Test
Shapiro Wilk Test Statistic 0.82	Shapiro Wilk Test Statistic 0.944
Shapiro Wilk Critical Value 0.947	Shapiro Wilk Critical Value 0.947
Data not Normal at 5% Significance Level	Data not Lognormal at 5% Significance Level

Assuming Normal Distribution	Assuming Lognormal Distribution	
95% Student's-t UCL 0.0739	95% H-UCL 0.0743	
95% UCLs (Adjusted for Skewness)	95% Chebyshev (MVUE) UCL 0.0865	
95% Adjusted-CLT UCL (Chen-1995) 0.0752	97.5% Chebyshev (MVUE) UCL 0.0964	
95% Modified-t UCL (Johnson-1978) 0.0742	99% Chebyshev (MVUE) UCL 0.116	

Gamma Distribution Test	Data Distribution
k star (bias corrected) 3.111	Data do not follow a Discernable Distribution (0.05)
Theta Star 0.0207	

MLE of Mean 0.0643 MLE of Standard Deviation 0.0365 nu star 311.1 Approximate Chi Square Value (.05) 271.2

proximate Chi Square Value (.05) 271.2 Adjusted Level of Significance 0.0452 Adjusted Chi Square Value 270.1

Anderson-Darling Test Statistic 1.152
Anderson-Darling 5% Critical Value 0.756
Kolmogorov-Smirnov Test Statistic 0.157
Kolmogorov-Smirnov 5% Critical Value 0.126
Data not Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n >= 40)\;0.0738$ 95% Adjusted Gamma UCL (Use when $n < 40)\;0.0741$

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 0.0738
95% Jackknife UCL 0.0739
95% Standard Bootstrap UCL 0.0737
95% Bootstrap-t UCL 0.0761
95% Hall's Bootstrap UCL 0.0755
95% Percentile Bootstrap UCL 0.0741
95% BCA Bootstrap UCL 0.0756
95% Chebyshev(Mean, Sd) UCL 0.0893
97.5% Chebyshev(Mean, Sd) UCL 0.1
99% Chebyshev(Mean, Sd) UCL 0.121

Use 95% Chebyshev (Mean, Sd) UCL 0.0893

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 50

Number of Distinct Observations 50

Raw Statistics

Minimum 77.5
Maximum 1173
Mean 566.3
Geometric Mean 459.3
Median 551.7
SD 311.4
Std. Error of Mean 44.04
Coefficient of Variation 0.55

Skewness 0.099

Log-transformed Statistics

Minimum of Log Data 4.35 Maximum of Log Data 7.067 Mean of log Data 6.13 SD of log Data 0.729

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.942 Shapiro Wilk Critical Value 0.947

Data not Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.897 Shapiro Wilk Critical Value 0.947

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 640.1

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 639.4 95% Modified-t UCL (Johnson-1978) 640.2 **Assuming Lognormal Distribution**

95% H-UCL 742.7 95% Chebyshev (MVUE) UCL 889.7 97.5% Chebyshev (MVUE) UCL 1017

99% Chebyshev (MVUE) UCL 1268

Gamma Distribution Test Data Distribution

k star (bias corrected) 2.404 Data do not follow a Discernable Distribution (0.05)

Theta Star 235.5 MLE of Mean 566.3

MLE of Standard Deviation 365.2

nu star 240.4

Approximate Chi Square Value (.05) 205.5 Nonparametric Statistics

Adjusted Level of Significance 0.0452 95% CLT UCL 638.7
Adjusted Chi Square Value 204.6 95% Jackknife UCL 640.1
95% Standard Bootstrap UCL 636.8

Anderson-Darling Test Statistic 1.016 95% Bootstrap-t UCL 640
Anderson-Darling 5% Critical Value 0.76 95% Hall's Bootstrap UCL 643.5

Kolmogorov-Smirnov Test Statistic 0.147 Kolmogorov-Smirnov 5% Critical Value 0.127 Data not Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 635 95% BCA Bootstrap UCL 641.1 95% Chebyshev(Mean, Sd) UCL 758.2 97.5% Chebyshev(Mean, Sd) UCL 841.3 99% Chebyshev(Mean, Sd) UCL 1004

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 662.4 95% Adjusted Gamma UCL (Use when n < 40) 665.5

Potential UCL to Use

Use 95% Chebyshev (Mean, Sd) UCL 758.2

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 50

Number of Distinct Observations 50

Raw Statistics

Minimum 3.63 Maximum 137.1 Mean 52.33 Geometric Mean 40.04 Median 45.55 SD 33.49 Std. Error of Mean 4.736 Coefficient of Variation 0.64

Skewness 0.55

Log-transformed Statistics

Minimum of Log Data 1.289 Maximum of Log Data 4.921 Mean of log Data 3.69 SD of log Data 0.836

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.937 Shapiro Wilk Critical Value 0.947

Assuming Normal Distribution

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.92 Shapiro Wilk Critical Value 0.947

Data not Normal at 5% Significance Level

Assuming Lognormal Distribution

Data not Lognormal at 5% Significance Level

95% H-UCL 73.62 95% Chebyshev (MVUE) UCL 89.01 97.5% Chebyshev (MVUE) UCL 103.2 99% Chebyshev (MVUE) UCL 131

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 60.51

95% Modified-t UCL (Johnson-1978) 60.33

Gamma Distribution Test

Data Distribution Data appear Gamma Distributed at 5% Significance Level

k star (bias corrected) 1.91 Theta Star 27.39

95% Student's-t UCL 60.27

MLE of Mean 52.33

MLE of Standard Deviation 37.86

nu star 191

Approximate Chi Square Value (.05) 160.1

Adjusted Level of Significance 0.0452

Adjusted Chi Square Value 159.2

Anderson-Darling Test Statistic 0.428 Anderson-Darling 5% Critical Value 0.763

Kolmogorov-Smirnov Test Statistic 0.0909 Kolmogorov-Smirnov 5% Critical Value 0.127

Data appear Gamma Distributed at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 60.12 95% Jackknife UCL 60.27 95% Standard Bootstrap UCL 60.14

> 95% Bootstrap-t UCL 60.72 95% Hall's Bootstrap UCL 60.55

95% Percentile Bootstrap UCL 60.45

95% BCA Bootstrap UCL 59.95

95% Chebyshev(Mean, Sd) UCL 72.97

97.5% Chebyshev(Mean, Sd) UCL 81.91

99% Chebyshev(Mean, Sd) UCL 99.45

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 62.46 95% Adjusted Gamma UCL (Use when n < 40) 62.79

Potential UCL to Use

Use 95% Approximate Gamma UCL 62.46

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

ALL INVERTEBRATE SPECIES COMBINED 95 UCL OUTPUTS

All Invertebrate Species Combined

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1 a.wst

Full Precision OFF nce Coefficient 95%

Confidence Coefficient 95% Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	33	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	33
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs! Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit! The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

	General Statistics		
Number of Valid Data	33	Number of Detected Data	20
Number of Distinct Detected Data	20	Number of Non-Detect Data	13
		Percent Non-Detects	39.39%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	191	Minimum Detected	5.252
Maximum Detected	5430	Maximum Detected	8.6
Mean of Detected	821	Mean of Detected	6.401
SD of Detected	1104	SD of Detected	0.66
Minimum Non-Detect	347	Minimum Non-Detect	5.849
Maximum Non-Detect	748	Maximum Non-Detect	6.617
Note: Data have multiple DLs - Use of KM Method is recor	mmended	Number treated as Non-Detect	28
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	5
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	84.85%

UCL Statistics

Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only		
Shapiro Wilk Test Statistic	0.407	Shapiro Wilk Test Statistic	0.839	
5% Shapiro Wilk Critical Value	0.905	5% Shapiro Wilk Critical Value	0.905	
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level		
Assuming Normal Distribution		Assuming Lognormal Distribution		
DL/2 Substitution Method		DL/2 Substitution Method		
Mean	591.1	Mean	6.013	
SD	900.2	SD	0.734	
95% DL/2 (t) UCL	856.5	95% H-Stat (DL/2) UCL	707.1	
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method		
MLE yields a negative mean		Mean in Log Scale	6.094	
		SD in Log Scale	0.652	
		Mean in Original Scale	608.6	
		SD in Original Scale	892.5	
		95% t UCL	871.8	
		95% Percentile Bootstrap UCL	910.7	
		95% BCA Bootstrap UCL	1105	
		95% H-UCL	695.2	

Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only		
k star (bias corrected)	1.533	Data do not follow a Discernable Distribution (0.05	5)	
Theta Star	535.4			
nu star	61.33			
A-D Test Statistic	2.019	Nonparametric Statistics		
5% A-D Critical Value	0.755	Kaplan-Meier (KM) Method		
K-S Test Statistic	0.755	Mean	621	
5% K-S Critical Value	0.197	SD	877.2	
Data not Gamma Distributed at 5% Significance	Level	SE of Mean	157.7	
		95% KM (t) UCL	888	
Assuming Gamma Distribution		95% KM (z) UCL	880.3	
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	886.5	
Minimum	0.000001	95% KM (bootstrap t) UCL	1413	
Maximum	5430	95% KM (BCA) UCL	949.8	
Mean	546.8	95% KM (Percentile Bootstrap) UCL	920.5	
Median	413.5	95% KM (Chebyshev) UCL	1308	
SD	925.8	97.5% KM (Chebyshev) UCL	1606	
k star	0.16	99% KM (Chebyshev) UCL	2190	
Theta star	3422			
Nu star	10.55	Potential UCLs to Use		
AppChi2	4.285	95% KM (BCA) UCL	949.8	
95% Gamma Approximate UCL (Use when n >= 40)	1346			
95% Adjusted Gamma UCL (Use when n < 40)	1413			
Note: DL/2 is not a recommended method.				

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006). For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 33	Number of Distinct Observations 24

Raw Statistics	Log-transformed Statistics
Minimum 0.009	Minimum of Log Data -4.711
Maximum 0.083	Maximum of Log Data -2.489
Mean 0.0468	Mean of log Data -3.132
Geometric Mean 0.0436	SD of log Data 0.42
Median 0.046	
SD 0.0158	
Std. Error of Mean 0.00275	
Coefficient of Variation 0.338	
Skewness -0.0016	

Relevant UCL Statistics

Normal Distribution Test	Lognormal Distribution Test			
Shapiro Wilk Test Statistic 0.989	Shapiro Wilk Test Statistic 0.884			
Shapiro Wilk Critical Value 0.931	Shapiro Wilk Critical Value 0.931			
Data appear Normal at 5% Significance Level	Data not Lognormal at 5% Significance Level			
Assuming Normal Distribution	Assuming Lognormal Distribution			
95% Student's-t UCL 0.0515	95% H-UCL 0.0548			

95% Student's-t UCL 0.0515	95% H-UCL 0.0548
95% UCLs (Adjusted for Skewness)	95% Chebyshev (MVUE) UCL 0.0632
95% Adjusted-CLT UCL (Chen-1995) 0.0514	97.5% Chebyshev (MVUE) UCL 0.07
95% Modified-t UCL (Johnson-1978) 0.0515	99% Chebyshev (MVUE) UCL 0.0833

Gamma Distribution Test	Data Distribution
k star (bias corrected) 6.595	Data appear Normal at 5% Significance Level
Theta Star 0.0071	

MLE of Mean 0.0468 MLE of Standard Deviation 0.0182 nu star 435.3 Approximate Chi Square Value (.05) 387.9

Adjusted Level of Significance 0.0419 Adjusted Chi Square Value 385.6

Anderson-Darling Test Statistic 0.478 Anderson-Darling 5% Critical Value 0.748 Kolmogorov-Smirnov Test Statistic 0.143 Kolmogorov-Smirnov 5% Critical Value 0.153 Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0526 95% Adjusted Gamma UCL (Use when n < 40) 0.0529

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 0.0514 95% Jackknife UCL 0.0515 95% Standard Bootstrap UCL 0.0514 95% Bootstrap-t UCL 0.0517 95% Hall's Bootstrap UCL 0.0516 95% Percentile Bootstrap UCL 0.0512 95% BCA Bootstrap UCL 0.0512 95% Chebyshev(Mean, Sd) UCL 0.0589 97.5% Chebyshev(Mean, Sd) UCL 0.0641 99% Chebyshev(Mean, Sd) UCL 0.0743

Use 95% Student's-t UCL 0.0515

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

> Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 33 Number of Distinct Observations 33

Raw Statistics

Minimum 97.57 Maximum 782.9 Mean 404.3 Geometric Mean 370.4 Median 369.9 SD 160.9 Std. Error of Mean 28

Coefficient of Variation 0.398 Skewness 0.406

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.974 Shapiro Wilk Critical Value 0.931

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Log-transformed Statistics

Shapiro Wilk Test Statistic 0.95 Shapiro Wilk Critical Value 0.931

Minimum of Log Data 4.581

Maximum of Log Data 6.663

Mean of log Data 5.914

SD of log Data 0.451

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 451.7

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 452.5 95% Modified-t UCL (Johnson-1978) 452.1

Assuming Lognormal Distribution

Data Distribution

Data appear Normal at 5% Significance Level

95% H-UCL 476.8 95% Chebyshev (MVUE) UCL 553.4 97.5% Chebyshev (MVUE) UCL 616.1 99% Chebyshev (MVUE) UCL 739.2

Gamma Distribution Test

k star (bias corrected) 5.35 Theta Star 75 57 MLE of Mean 404.3 MLE of Standard Deviation 174.8

nu star 353.1

Approximate Chi Square Value (.05) 310.5

Nonparametric Statistics

Adjusted Level of Significance 0.0419	95% CLT UCL 450.4
Adjusted Chi Square Value 308.5	95% Jackknife UCL 451.7
	95% Standard Bootstrap UCL 449.5
Anderson-Darling Test Statistic 0.251	95% Bootstrap-t UCL 454.4
Anderson-Darling 5% Critical Value 0.748	95% Hall's Bootstrap UCL 453.1
Kolmogorov-Smirnov Test Statistic 0.0852	95% Percentile Bootstrap UCL 450.6
Kolmogorov-Smirnov 5% Critical Value 0.153	95% BCA Bootstrap UCL 450.1
Data appear Gamma Distributed at 5% Significance Level	95% Chebyshev(Mean, Sd) UCL 526.4
	97.5% Chebyshev(Mean, Sd) UCL 579.2
Assuming Gamma Distribution	99% Chebyshev(Mean, Sd) UCL 682.9
95% Approximate Gamma UCL (Use when n >= 40) 459.7	

Potential UCL to Use

95% Adjusted Gamma UCL (Use when n < 40) 462.7

Use 95% Student's-t UCL 451.7

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 33

Number of Distinct Observations 33

Raw Statistics

Minimum 8.2 Maximum 49.2 Mean 24 77 Geometric Mean 22.51 Median 23.2 SD 10.51 Std. Error of Mean 1.83 Coefficient of Variation 0.424 Skewness 0.508

Log-transformed Statistics

Minimum of Log Data 2.104 Maximum of Log Data 3.896 Mean of log Data 3.114 SD of log Data 0.462

Relevant UCL Statistics

Normal Distribution Test	Normal	Distribution	Test
--------------------------	--------	--------------	------

Shapiro Wilk Test Statistic 0.959 Shapiro Wilk Critical Value 0.931

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.955 Shapiro Wilk Critical Value 0.931 Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution 95% Student's-t UCL 27.87

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 27.95 95% Modified-t UCL (Johnson-1978) 27.9

Assuming Lognormal Distribution

95% H-UCL 29.28 95% Chebyshev (MVUE) UCL 34.06 97.5% Chebyshev (MVUE) UCL 38 99% Chebyshev (MVUE) UCL 45.74

Gamma Distribution Test

k star (bias corrected) 4.919 Theta Star 5.036 MLE of Mean 24.77 MLE of Standard Deviation 11.17 nu star 324.6 Approximate Chi Square Value (.05) 283.9

Adjusted Chi Square Value 281.9

Adjusted Level of Significance 0.0419

Anderson-Darling Test Statistic 0.241 Anderson-Darling 5% Critical Value 0.748 Kolmogorov-Smirnov Test Statistic 0.0941 Kolmogorov-Smirnov 5% Critical Value 0.154

Data appear Gamma Distributed at 5% Significance Level

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 27.78 95% Jackknife UCL 27.87 95% Standard Bootstrap UCL 27.82 95% Bootstrap-t UCL 28.22 95% Hall's Bootstrap UCL 27.99 95% Percentile Bootstrap UCL 27.86 95% BCA Bootstrap UCL 27.79 95% Chebyshev(Mean, Sd) UCL 32.75

97.5% Chebyshev(Mean, Sd) UCL 36.2 99% Chebyshev(Mean, Sd) UCL 42.97

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 28.33 95% Adjusted Gamma UCL (Use when n < 40) 28.52

Potential UCL to Use

Use 95% Student's-t UCL 27.87

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Fish and Invertebrate Species Grouped

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.33	0.108	948	Fish	1	All fish
2	0.33	0.108	824	Invertebrate	1	All invertebrates
3	0.34	0.111	1326	Mammal	0.5	Marsh Rice Rat
4						
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Group: Terrestrial Mammal - Marsh Rice Rat
Scenario: Fish and Invertebrate Species Grouped

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C_SED	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.022325	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0.12875	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.51E-01	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	All fish	3	1	1	0.25	0.0893	Table C-4.2
2	Invertebrate	All invertebrates	2	1	1	0.25	0.515	Table C-4.2
3	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
4								
5								
6								

Definitions:

kg - kilogram mg - milligrams

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site Prey Group: Terrestrial Mammal - Marsh Rice Rat Scenario: Fish and Invertebrate Species Grouped

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	All fish	3	1	1	0.25		Table C-4.2
2	Invertebrate	All invertebrates	2	1	1	0.25		Table C-4.2
3	Plant				1	0.5	6.60E-02	Table C-4.2
4								
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Fish and Invertebrate Species Grouped

COPC	C _{BIOTA,1}	$C_{\text{BIOTA,2}}$	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA,5}}$	$C_{BIOTA,6}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	l		mg/	'kg (wet weigh	t)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
Mercury (Methyl)	0.0893	0.515	1.51E-01					2.6E-02	0.0479							1.3E-05	2.7E-02	0.064	0.41
Mercury (Inorganic)			3.40E-04					1.9E-05	4.7391							1.3E-03	1.3E-03	0.9	0.0014

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Fish and Invertebrate Species Grouped

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	0.33	0.108	948	Fish	1	All fish
2	0.33	0.108	824	Invertebrate	1	All invertebrates
3	0.34	0.111	1326	Mammal	0.5	Marsh Rice Rat
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Group: Terrestrial Mammal - Marsh Rice Rat
Scenario: Fish and Invertebrate Species Grouped

COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C_{SED}	0.0021	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.90E-04	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	1.58E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.95E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.78E-03	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	All Fish	3	14	1	0.25	7.58E-04	Table C-4.2
2	Invertebrate	All Invertebrates	2	1	1	0.25	4.52E-04	Table C-4.2
3	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
4								
5								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Fish and Invertebrate Species Grouped

COPC	C _{BIOTA,1}	$C_{\text{BIOTA,2}}$	C _{BIOTA,3}	$C_{\text{BIOTA,4}}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	C _{SED,6}	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)						mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA		
PCB Congener TEQ (Avian) (U=1/2)	0.00076	0.00045	1.78E-03					8.3E-05	0.0021							5.5E-07	8.3E-05	1.4E-04	0.59

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Fish and Invertebrate Species Grouped

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	948	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.672	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.598	948	Fish	1	All fish
2	0.11	0.074	948	Invert	1	All invert
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Fish and Invertebrate Species Grouped

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/l	kg (wet wei	ght)			mg/kg _{BW} /day			mg/k	(g (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
Mercury (Methyl)	0.0893	0.0515						4.0E-02	0.0479							0.0E+00	4.0E-02	0.064	0.62
Hexachlorobenzene	0.6448	0.9498						3.1E-01	2.1373							0.0E+00	3.1E-01	3.35	0.09
1,3-Dichlorobenzene	0.769	0.758						3.6E-01	0.0736							0.0E+00	3.6E-01	160.8	0.0022
PCB Congener TEQ (Avian) (U=1/2)	0.0008	0.0005						3.4E-04	0.0021			_				0.0E+00	3.4E-04	0.00014	2.40
Mercury (Inorganic)								0.0E+00	4.7391							0.0E+00	0.0E+00	9.0E-01	0

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Fish and Invertebrate Species Grouped

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.233	0.021	1092	Fish	1	All Fish
2	0.234	0.021	974	Invertebrate	1	All invertebrate
3	0.3	0.027	962	Plant	1	
4	0.233	0.021	1428	Mammal	0.5	Marsh Rice Rat
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

4

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Group: Terrestrial Mammal - Marsh Rice Rat
Scenario: Fish and Invertebrate Species Grouped

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	2.23E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	1.29E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.52E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	All Fish	3	1	1	0.25	8.93E-02	Table C-4.2
2	Invertebrate	All Invertebrate	2	1	1	0.25	5.15E-02	Table C-4.2
3	Plant				1	0.5	2.43E-03	Table C-4.2
4								
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight ww - wet weight Prey Group: Terrestrial Mammal - Marsh Rice Rat Scenario: Fish and Invertebrate Species Grouped

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	All Fish	3	1	1	0.25		Table C-4.2
2	Invertebrate	All Invertebrate	2	1	1	0.25		Table C-4.2
3	Plant				1	0.5	6.60E-02	Table C-4.2
4								
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams dw - dry weight

ww - wet weight

Receptor: Raccoon

Receptor Group: Scenario: **Omnivorous Mammal**

Fish and Invertebrate Species Grouped

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{BIOTA,3}$	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
				mg/kg (wet weight)				mg/kg _{BW} /day			mg/k	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Mercury (Methyl)	0.0893	0.0515	0.002426492	3.52E-02				7.2E-05	0.0479							8.9E-06	8.0E-05	0.0177	0.0045
Mercury (Inorganic)			0.065754378	3.40E-04				3.8E-05	4.7391							8.8E-04	9.1E-04	3.54	0.00026

Baseline Ecological Risk Assessment March 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Fish and Invertebrate Species Grouped

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

4

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.233	0.021	1092	Fish	1	All Fish
2	0.234	0.021	974	Invertebrate	1	All invertebrate
3	0.3	0.027	962	Plant	1	
4	0.233	0.021	1428	Mammal	0.5	Marsh Rice Rat
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Group: Terrestrial Mammal - Marsh Rice Rat
Scenario: Fish and Invertebrate Species Grouped

COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	2.18E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	1.12E-04	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3		mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	1.91E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.34E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	All Fish	3	14	1	0.25	8.70E-05	Table C-4.2
2	Invertebrate	All Invertebrate	2	1	1	0.25	3.19E-05	Table C-4.2
3	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
4								
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: Fish and Invertebrate Species Grouped

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)					mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA			
PCB Congener TEQ (Mammal) (U=1/2)	6.2E-05	2.8E-05	2.32E-07	1.34E-04				7.0E-08	0.0001							2.1E-08	9.0E-08	5.4E-07	0.17

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper

Receptor Group: Sediment-probing Birds

Scenario: Fish and Invertebrate Species Grouped

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.3		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	All invertebrates
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: Fish and Invertebrate Species Grouped

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet we	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Lead	0.292							9.9E-02	48.944							5.8E-02	1.6E-01	3.26	0.05
Mercury (Methyl)	0.0533							1.8E-02	0.0479							5.6E-05	1.8E-02	0.064	0.28
Total PAH 16 (U=1/2)	0.1397							4.7E-02	7.3078							8.6E-03	5.6E-02	325	0.00017
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							7.5E-06	0.0002							2.5E-07	7.8E-06	0.00014	0.06
Hexachlorobenzene	1.13							3.8E-01	2.1373							2.5E-03	3.8E-01	3.35	0.11
Hexachlorobutadiene	0.373							1.3E-01	3.8327							4.5E-03	1.3E-01	20	0.01
1,3-Dichlorobenzene	0.373							1.3E-01	0.0736							8.7E-05	1.3E-01	160.8	0.0008
1,4-Dichlorobenzene	0.373							1.3E-01	0.0744							8.7E-05	1.3E-01	160.8	0.0008
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.5E-04	0.0021							2.4E-06	1.5E-04	1.4E-04	1.05
Mercury (Inorganic)								0.0E+00	4.7391							5.6E-03	5.6E-03	9.0E-01	0.006

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.3 PREY GROUPING UNCERTAINTY ANALYSIS – ALL SPECIES INDEPENDENT

Table C-5.3
EPCs For All Individual Prey Species

COPC	Exposure Point Concentration (mg/kg)												
COPC	BCR	BRS	GKF	GM	OYS	PNF	SAS	STM	WHS				
1,3-Dichlorobenzene	0.748 ^a	0.709 ^a	0.763 ^a	0.765 ^a	0.704 ^a	0.765 ^a	0.37 ^a	0.769 ^a	0.758 ^a				
1,4-Dichlorobenzene	0.745 ^a				0.704 ^a		-		0.714 ^a				
Hexachlorobenzene	1.088	0.7977	0.5825	0.568	0.704 ^a	2.29 ^b	0.325 ^b	1.05	0.682 ^b				
Hexachlorobutadiene	0.745 ^a	-	-		0.704 ^a	-		-	0.714 ^a				
Lead	0.226	-	-		0.331 ^b	1	-	-	0.116 ^b				
Mercury	0.0567	0.0501	0.0611	0.145	0.051 ^b	0.083 ^b	0.063 ^b	0.0744	0.042 ^b				
Dioxin/Furan TEQ (Avian; U=1/2)		-	-		0.000020 ^b	1	-	-	0.000025 ^b				
Total PAH (U=1/2)	0.07592				0.55212 ^b				0.1623 ^b				
PCB Congener TEQ (Avian; U=1/2)	0.00044	0.00060	0.00087	0.00056	0.00036 ^b	0.00015 ^b	0.00019 ^b	0.00048	0.00031 ^b				
PCB Congener TEQ (Mammal; U=1/2)	0.000031	0.000031	0.000087	0.000038	0.000021 ^b	0.000051 ^b	0.000026 ^b	0.000030	0.000018 ^b				

Notes:

a - EPC based on maximum non-detected value

b - EPC based on maximum detected value

BCR - blue crab

BRS - brown shrimp

GKF - gulf killifish

GM - gulf menhaden

OYS - oyster

PNF - pinfish

SAS - sand seatrout

STM - striped mullet

WHS - white shrimp

BLUE CRAB 95 UCL OUTPUTS

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst Full Precision OFF Confidence Coefficient 95% Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	21	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	21
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs! Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit! The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (1,4-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	18	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	18
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs! Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit! The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,4-dichlorobenzene_ug/kg) was not processed!

Result Value (hexachlorobenzene ug/kg)

	General Statistics		
Number of Valid Data	21	Number of Detected Data	9
Number of Distinct Detected Data	9	Number of Non-Detect Data	12
		Percent Non-Detects	57.14%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	191	Minimum Detected	5.252
Maximum Detected	5430	Maximum Detected	8.6
Mean of Detected	1021	Mean of Detected	6.33
SD of Detected	1668	SD of Detected	0.971
Minimum Non-Detect	347	Minimum Non-Detect	5.849
Maximum Non-Detect	748	Maximum Non-Detect	6.617
Note: Data have multiple DLs - Use of KM Method is recommend	led	Number treated as Non-Detect	19
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	2

Warning: There are only 9 Detected Values in this data Note: It should be noted that even though bootstrap may be performed on this data set the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

Observations < Largest ND are treated as NDs

90.48%

Single DL Non-Detect Percentage

	UCL Stat	tistics	
Normal Distribution Test with Detected Values Only	,	Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.511	Shapiro Wilk Test Statistic	0.847
5% Shapiro Wilk Critical Value	0.829	5% Shapiro Wilk Critical Value	0.829
Data not Normal at 5% Significance Level		Data appear Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	568	Mean	5.786
SD	1131	SD	0.816
95% DL/2 (t) UCL	993.5	95% H-Stat (DL/2) UCL	694.3
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE method failed to converge properly		Mean in Log Scale	5.842
		SD in Log Scale	0.757
		Mean in Original Scale	575.3
		SD in Original Scale	1127
		95% t UCL	999.4
		95% Percentile Bootstrap UCL	1028
		95% BCA Bootstrap UCL	1355
		95% H-UCL	672
Gamma Distribution Test with Detected Values Only	1	Data Distribution Test with Detected Values Only	
k star (bias corrected)	0.719	Data appear Lognormal at 5% Significance Level	
Theta Star	1420		
nu star	12.94		
A-D Test Statistic	1.166	Nonparametric Statistics	
5% A-D Critical Value	0.745	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.745	Mean	600
5% K-S Critical Value	0.287	SD	1094
Data not Gamma Distributed at 5% Significance Leve	el .	SE of Mean	254.5
		95% KM (t) UCL	1039
Assuming Gamma Distribution		95% KM (z) UCL	1019
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	1026
Minimum	0.000001	95% KM (bootstrap t) UCL	2652
Maximum	5430	95% KM (BCA) UCL	1097
Mean	471.7	95% KM (Percentile Bootstrap) UCL	1088
Median	237.7	95% KM (Chebyshev) UCL	1709
SD	1165	97.5% KM (Chebyshev) UCL	2189
k star	0.119	99% KM (Chebyshev) UCL	3132
Theta star	3955		
Nu star	5.009	Potential UCLs to Use	
AppChi2	1.157	95% KM (t) UCL	1039
95% Gamma Approximate UCL (Use when n >= 40)	2043	95% KM (% Bootstrap) UCL	1088
95% Adjusted Gamma UCL (Use when n < 40)	2310		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

Result_Value (hexachlorobutadiene_ug/kg)

	General Statistics		
Number of Valid Data	18	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	18
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs! Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (hexachlorobutadiene_ug/kg) was not processed!

Result_Value (lead_mg/kg)

General Statistics

Number of Valid Observations 18 Number of Distinct Observations 18

Raw Statistics

Minimum 0.092
Maximum 0.518
Mean 0.178
Geometric Mean 0.156
Median 0.126
SD 0.112
Std. Error of Mean 0.0264
Coefficient of Variation 0.629

Skewness 2.011

Log-transformed Statistics

Minimum of Log Data -2.386 Maximum of Log Data -0.658 Mean of log Data -1.86 SD of log Data 0.494

Relevant LICI Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.735 Shapiro Wilk Critical Value 0.897 **Lognormal Distribution Test**

Shapiro Wilk Test Statistic 0.859 Shapiro Wilk Critical Value 0.897 Data not Lognormal at 5% Significance Level

Data not Normal at 5% Significance Level

Assuming Normal Distribution 95% Student's-t UCL 0.224

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 0.235 95% Modified-t UCL (Johnson-1978) 0.226 **Assuming Lognormal Distribution**

Data Distribution

Data do not follow a Discernable Distribution (0.05)

95% H-UCL 0.224 95% Chebyshev (MVUE) UCL 0.266 97.5% Chebyshev (MVUE) UCL 0.306 99% Chebyshev (MVUE) UCL 0.384

Gamma Distribution Test

k star (bias corrected) 3.27 Theta Star 0.0545

MLE of Mean 0.178 MLE of Standard Deviation 0.0985

nu star 117.7

Approximate Chi Square Value (.05) 93.67 Adjusted Level of Significance 0.0357

Adjusted Chi Square Value 91.61

Anderson-Darling Test Statistic 1.296 Anderson-Darling 5% Critical Value 0.743 Kolmogorov-Smirnov Test Statistic 0.232 Kolmogorov-Smirnov 5% Critical Value 0.205

Data not Gamma Distributed at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 0.221
95% Jackknife UCL 0.224
95% Standard Bootstrap UCL 0.219
95% Bootstrap-t UCL 0.263
95% Hall's Bootstrap UCL 0.251
95% Percentile Bootstrap UCL 0.221
95% BCA Bootstrap UCL 0.235
95% Chebyshev(Mean, Sd) UCL 0.293
97.5% Chebyshev(Mean, Sd) UCL 0.343
99% Chebyshev(Mean, Sd) UCL 0.441

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.224 95% Adjusted Gamma UCL (Use when n < 40) 0.229

Potential UCL to Use

Use 95% Student's-t UCL 0.224 or 95% Modified-t UCL 0.226

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 21

Number of Distinct Observations 16

Raw Statistics

Minimum 0.009 Maximum 0.083 Mean 0.0502 Geometric Mean 0.0463 Median 0.051 SD 0.0171 Std. Error of Mean 0.00373

Skewness -0.416

Log-transformed Statistics

Minimum of Log Data -4.711 Maximum of Log Data -2.489 Mean of log Data -3.073 SD of log Data 0.475

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.979 Shapiro Wilk Critical Value 0.908

Coefficient of Variation 0.34

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.804 Shapiro Wilk Critical Value 0.908 Data not Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.0567

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 0.056 95% Modified-t UCL (Johnson-1978) 0.0566

Assuming Lognormal Distribution

Data Distribution

Data appear Normal at 5% Significance Level

95% H-UCL 0.0639 95% Chebyshev (MVUE) UCL 0.0756 97.5% Chebyshev (MVUE) UCL 0.086 99% Chebyshev (MVUE) UCL 0.106

Gamma Distribution Test

k star (bias corrected) 5.406 Theta Star 0.00929 MLF of Mean 0.0502

MLE of Standard Deviation 0.0216

nu star 227.1

Approximate Chi Square Value (.05) 193.2 Adjusted Level of Significance 0.0383 Adjusted Chi Square Value 190.8

Anderson-Darling Test Statistic 0.712
Anderson-Darling 5% Critical Value 0.745
Kolmogorov-Smirnov Test Statistic 0.208
Kolmogorov-Smirnov 5% Critical Value 0.19
Data follow Appr. Gamma Distribution at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 0.0564
95% Jackknife UCL 0.0567
95% Standard Bootstrap UCL 0.0567
95% Bootstrap-t UCL 0.0567
95% Hall's Bootstrap UCL 0.0562
95% Percentile Bootstrap UCL 0.056
95% BCA Bootstrap UCL 0.0559
95% Chebyshev(Mean, Sd) UCL 0.0665
97.5% Chebyshev(Mean, Sd) UCL 0.0735
99% Chebyshev(Mean, Sd) UCL 0.0874

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n \ge 40$) 0.059 95% Adjusted Gamma UCL (Use when n < 40) 0.0598

Potential UCL to Use

Use 95% Student's-t UCL 0.0567

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

Result_Value (total pah (16) (u = 1/2)_ug/kg)

General Statistics

Number of Valid Data 18 Number of Detected Data

5

Number of Distinct Detected Data	5	Number of Non-Detect Data Percent Non-Detects	13 72.22%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	60.91	Minimum Detected	4.109
Maximum Detected	123.8	Maximum Detected	4.819
Mean of Detected	77.22	Mean of Detected	4.308
SD of Detected	26.45	SD of Detected	0.294
Minimum Non-Detect	6.93	Minimum Non-Detect	1.936
Maximum Non-Detect	14.8	Maximum Non-Detect	2.695
Note: Data have multiple DLs - Use of KM Method is recommend	ded	Number treated as Non-Detect	13
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	5
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	72.22%

Warning: There are only 5 Detected Values in this data

Note: It should be noted that even though bootstrap may be performed on this data set the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

	UCL Statist	ics	
Normal Distribution Test with Detected Values Onl	У	Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.694	Shapiro Wilk Test Statistic	0.741
5% Shapiro Wilk Critical Value	0.762	5% Shapiro Wilk Critical Value	0.762
Data not Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	25.88	Mean	2.473
SD	35.22	SD	1.213
95% DL/2 (t) UCL	40.32	95% H-Stat (DL/2) UCL	59.09
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE yields a negative mean		Mean in Log Scale	3.525
		SD in Log Scale	0.538
		Mean in Original Scale	39.83
		SD in Original Scale	27.32
		95% t UCL	51.03
		95% Percentile Bootstrap UCL	51.13
		95% BCA Bootstrap UCL	54.45
		95% H-UCL	51.36
Gamma Distribution Test with Detected Values On	ly	Data Distribution Test with Detected Values Only	
k star (bias corrected)	5.409	Data do not follow a Discernable Distribution (0.05)	
Theta Star	14.28		
nu star	54.09		
A-D Test Statistic	0.794	Nonparametric Statistics	
5% A-D Critical Value	0.679	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.679	Mean	65.44
5% K-S Critical Value	0.357	SD	14.45
Data not Gamma Distributed at 5% Significance Lev	el	SE of Mean	3.808
		95% KM (t) UCL	72.07
Assuming Gamma Distribution		95% KM (z) UCL	71.7
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	71.53
Minimum	0.000001	95% KM (bootstrap t) UCL	83.69
Maximum	123.8	95% KM (BCA) UCL	76.32
Mean	21.45	95% KM (Percentile Bootstrap) UCL	75.92
Median	0.000001	95% KM (Chebyshev) UCL	82.04
SD	37.83	97.5% KM (Chebyshev) UCL	89.22
k star	0.0965	99% KM (Chebyshev) UCL	103.3

Theta star	222.4		
Nu star	3.473	Potential UCLs to Use	
AppChi2	0.525	95% KM (t) UCL	72.07
95% Gamma Approximate UCL (Use when n >= 40)	141.8	95% KM (% Bootstrap) UCL	75.92
95% Adjusted Gamma UCL (Use when n < 40)	172.3		

Note: DL/2 is not a recommended method.

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 21 Number of Distinct Observations 21

Raw Statistics

Minimum 97.57

Maximum 782.9

Mean 378.7

Log-transformed Statistics

Minimum of Log Data 4.581

Maximum of Log Data 6.663

Mean of log Data 5.84

Mean 378.7 Mean of log Data 5.84
Geometric Mean 343.9 SD of log Data 0.474
Median 340.8
SD 163.7

Relevant UCL Statistics

Std. Error of Mean 35.73 Coefficient of Variation 0.432

MLE of Standard Deviation 176.4

95% Adjusted Gamma UCL (Use when n < 40) 457.6

Skewness 0.813

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.942
Shapiro Wilk Critical Value 0.908
Shapiro Wilk Critical Value 0.908
Data appear Normal at 5% Significance Level
Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution 95% Student's-t UCL 440.3 Assuming Lognormal Distribution 95% H-Ut

95% Student's-t UCL 440.3 95% H-UCL 473.9 95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 560.9 95% Adjusted-CLT UCL (Chen-1995) 444.2 97.5% Chebyshev (MVUE) UCL 638.1 95% Modified-t UCL (Johnson-1978) 441.4 99% Chebyshev (MVUE) UCL 789.7

Gamma Distribution Test Data Distribution

k star (bias corrected) 4.611 Data appear Normal at 5% Significance Level

Theta Star 82.14 MLE of Mean 378.7

nu star 193.7
Approximate Chi Square Value (.05) 162.5

Nonparametric Statistics

Adjusted Level of Significance 0.0383 95% CLT UCL 437.5
Adjusted Chi Square Value 160.3 95% Jackknife UCL 440.3
95% Standard Bootstrap UCL 436.1

Anderson-Darling Test Statistic 0.313

Anderson-Darling 5% Critical Value 0.745

Kolmogorov-Smirnov Test Statistic 0.115

Kolmogorov-Smirnov 5% Critical Value 0.19

Data appear Gamma Distributed at 5% Significance Level

Systematic Bootstrap CCL 452.3

95% Bootstrap UCL 454.1

95% Percentile Bootstrap UCL 438

95% BCA Bootstrap UCL 441.6

Posta appear Gamma Distributed at 5% Significance Level

97.5% Chebyshev (Mean, Sd) UCL 601.8

Assuming Gamma Distribution 99% Chebyshev (Mean, Sd) UCL 734.2
95% Approximate Gamma UCL (Use when n >= 40) 451.4

Potential UCL to Use Use 95% Student's-t UCL 440.3

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 21

Number of Distinct Observations 21

Raw Statistics

Minimum 8.35
Maximum 49.2
Mean 26.38
Geometric Mean 24.03
Median 24.8
SD 11.1
Std. Error of Mean 2.422
Coefficient of Variation 0.421

Skewness 0.412

Log-transformed Statistics

Minimum of Log Data 2.122 Maximum of Log Data 3.896 Mean of log Data 3.179 SD of log Data 0.46

Relevant UCL Statistics

Normal D	Distribu	tion	Test
----------	----------	------	------

Shapiro Wilk Test Statistic 0.97 Shapiro Wilk Critical Value 0.908

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.971 Shapiro Wilk Critical Value 0.908 Data appear Lognormal at 5% Significance Level

Data appear Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 30.55 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 30.59 95% Modified-t UCL (Johnson-1978) 30.59

Assuming Lognormal Distribution

95% H-UCL 32.67 95% Chebyshev (MVUE) UCL 38.58 97.5% Chebyshev (MVUE) UCL 43.77 99% Chebyshev (MVUE) UCL 53.98

Gamma Distribution Test

k star (bias corrected) 4.767 Theta Star 5.533 MLE of Mean 26.38 MLE of Standard Deviation 12.08

nu star 200.2 Approximate Chi Square Value (.05) 168.5

> Adjusted Level of Significance 0.0383 Adjusted Chi Square Value 166.3

Anderson-Darling Test Statistic 0.155 Anderson-Darling 5% Critical Value 0.745 Kolmogorov-Smirnov Test Statistic 0.107 Kolmogorov-Smirnov 5% Critical Value 0.19

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 31.35 95% Adjusted Gamma UCL (Use when n < 40) 31.77

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 30.36
95% Jackknife UCL 30.55
95% Standard Bootstrap UCL 30.27
95% Bootstrap-t UCL 30.87
95% Hall's Bootstrap UCL 30.62
95% Percentile Bootstrap UCL 30.48
95% BCA Bootstrap UCL 30.39
95% Chebyshev(Mean, Sd) UCL 36.93
97.5% Chebyshev(Mean, Sd) UCL 41.5
99% Chebyshev(Mean, Sd) UCL 50.48

Potential UCL to Use

Use 95% Student's-t UCL 30.55

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

BROWN SHRIMP 95 UCL OUTPUTS

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_a.wst

Full Precision OFF Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	8	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	8
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

General Statistics

Number of Valid Observations 8

Number of Distinct Observations 8

Raw Statistics

Minimum	537
Maximum	913
Mean	710.4
Geometric Mean	700
Median	694.5

SD 130.4 Std. Error of Mean 46.09 Coefficient of Variation 0.183 Skewness 0.292

Log-transformed Statistics

Lognormal Distribution Test

Assuming Lognormal Distribution

Minimum of Log Data 6.286 Maximum of Log Data 6.817 Mean of log Data 6.551 SD of log Data 0.183

Warning: There are only 8 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.967
Shapiro Wilk Critical Value 0.818
Shapiro Wilk Critical Value 0.818
Shapiro Wilk Critical Value 0.818

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 797.7 95% H-UCL 813.6

95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 911.6

95% Adjusted-CLT UCL (Chen-1995) 791.3 97.5% Chebyshev (MVUE) UCL 998.6

95% Adjusted-CLT UCL (Chen-1995) 791.3 97.5% Chebyshev (MVUE) UCL 998.6 95% Modified-t UCL (Johnson-1978) 798.5 99% Chebyshev (MVUE) UCL 1170

Gamma Distribution Test Data Distribution

k star (bias corrected) 21.42 Data appear Normal at 5% Significance Level
Theta Star 33.16
MLE of Mean 710.4
MLE of Standard Deviation 153.5

nu star 342.7 Approximate Chi Square Value (.05) 300.8 Adjusted Level of Significance 0.0195 Adjusted Chi Square Value 290.8

Anderson-Darling Test Statistic 0.186 Anderson-Darling 5% Critical Value 0.716 Kolmogorov-Smirnov Test Statistic 0.145 Kolmogorov-Smirnov 5% Critical Value 0.294

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 809.3 95% Adjusted Gamma UCL (Use when n < 40) 837.1

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 786.2 95% Jackknife UCL 797.7 95% Standard Bootstrap UCL 781.9 95% Bootstrap-t UCL 806.8 95% Hall's Bootstrap UCL 787.9 95% Percentile Bootstrap UCL 784.8 95% BCA Bootstrap UCL 780.1 95% Chebyshev(Mean, Sd) UCL 911.3 97.5% Chebyshev(Mean, Sd) UCL 998.2 99% Chebyshev(Mean, Sd) UCL 1169

Use 95% Student's-t UCL 797.7

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 8

Raw Statistics Minimum 0.025 Maximum 0.065 Mean 0.0414 Geometric Mean 0.0396 Median 0.0405 SD 0.013 Std. Error of Mean 0.00459

Coefficient of Variation 0.314 Skewness 0.622

Number of Distinct Observations 8

Log-transformed Statistics

Minimum of Log Data -3.689 Maximum of Log Data -2.733 Mean of log Data -3.228

SD of log Data 0.313

Warning: There are only 8 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.952 Shapiro Wilk Critical Value 0.818

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.974 Shapiro Wilk Critical Value 0.818

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.0501

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 0.05 95% Modified-t UCL (Johnson-1978) 0.0502

Gamma Distribution Test

k star (bias corrected) 7.511 Theta Star 0.00551 MLE of Mean 0.0414 MLE of Standard Deviation 0.0151 nu star 120.2

Assuming Lognormal Distribution

95% H-UCL 0.0532 95% Chebyshev (MVUE) UCL 0.0614 97.5% Chebyshev (MVUE) UCL 0.0701 99% Chebyshev (MVUE) UCL 0.0871

Data Distribution

Data appear Normal at 5% Significance Level

Approximate Chi Square Value (.05) 95.86 Adjusted Level of Significance 0.0195 Adjusted Chi Square Value 90.38

Anderson-Darling Test Statistic 0.212 Anderson-Darling 5% Critical Value 0.715 Kolmogorov-Smirnov Test Statistic 0.152 Kolmogorov-Smirnov 5% Critical Value 0.294

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n \ge 40$) 0.0519 95% Adjusted Gamma UCL (Use when n < 40) 0.055

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 0.0489
95% Jackknife UCL 0.0501
95% Standard Bootstrap UCL 0.0484
95% Bootstrap-t UCL 0.0513
95% Hall's Bootstrap UCL 0.0506
95% Percentile Bootstrap UCL 0.0489
95% BCA Bootstrap UCL 0.0493
95% Chebyshev(Mean, Sd) UCL 0.0614
97.5% Chebyshev(Mean, Sd) UCL 0.07

Use 95% Student's-t UCL 0.0501

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 8

Number of Distinct Observations 8

Raw Statistics

Minimum 369.9 Maximum 694.7 Mean 532.6 Geometric Mean 524.2 Median 552.1 SD 98.89 Std. Error of Mean 34.96 Coefficient of Variation 0.186

Skewness -0.164

Log-transformed Statistics

Minimum of Log Data 5.913 Maximum of Log Data 6.544 Mean of log Data 6.262 SD of log Data 0.194

Warning: There are only 8 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.965 Shapiro Wilk Critical Value 0.818

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.947 Shapiro Wilk Critical Value 0.818

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 598.9

95% UCLs (Adjusted for Skewness) 95% Adjusted-CLT UCL (Chen-1995) 588

95% Modified-t UCL (Johnson-1978) 598.5

Assuming Lognormal Distribution

95% H-UCL 615.7 95% Chebyshev (MVUE) UCL 692.5 97.5% Chebyshev (MVUE) UCL 761.6 99% Chebyshev (MVUE) UCL 897.2

Gamma Distribution Test

k star (bias corrected) 19.79 Theta Star 26.91

MLE of Mean 532.6

MLE of Standard Deviation 119.7 nu star 316.7

Approximate Chi Square Value (.05) 276.4

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

Adjusted Level of Significance 0.0195 Adjusted Chi Square Value 266.9

Anderson-Darling Test Statistic 0.305
Anderson-Darling 5% Critical Value 0.716
Kolmogorov-Smirnov Test Statistic 0.199
Kolmogorov-Smirnov 5% Critical Value 0.294

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 610.1 95% Adjusted Gamma UCL (Use when n < 40) 631.9

Potential UCL to Use

95% CLT UCL 590.1
95% Jackknife UCL 598.9
95% Standard Bootstrap UCL 586.4
95% Bootstrap-t UCL 595.4
95% Hall's Bootstrap UCL 594.3
95% Percentile Bootstrap UCL 582.1
95% BCA Bootstrap UCL 581.6
95% Chebyshev(Mean, Sd) UCL 685
97.5% Chebyshev(Mean, Sd) UCL 751
99% Chebyshev(Mean, Sd) UCL 880.5

Use 95% Student's-t UCL 598.9

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 8

Number of Distinct Observations 8

Raw Statistics

Minimum 8.2
Maximum 41.18
Mean 24.83
Geometric Mean 22.83
Median 22.85
SD 9.646
Std. Error of Mean 3.41
Coefficient of Variation 0.388
Skewness 0.00717

Log-transformed Statistics

Minimum of Log Data 2.104 Maximum of Log Data 3.718 Mean of log Data 3.128 SD of log Data 0.477

Warning: There are only 8 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.952 Shapiro Wilk Critical Value 0.818

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.863 Shapiro Wilk Critical Value 0.818

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 31.3

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 30.45 95% Modified-t UCL (Johnson-1978) 31.3 Assuming Lognormal Distribution 95% H-UCL 38.81

95% Chebyshev (MVUE) UCL 43.86 97.5% Chebyshev (MVUE) UCL 51.94 99% Chebyshev (MVUE) UCL 67.8

Gamma Distribution Test

k star (bias corrected) 3.895 Theta Star 6.376 Data Distribution

Data appear Normal at 5% Significance Level

MLE of Mean 24.83
MLE of Standard Deviation 12.58
nu star 62.32
Approximate Chi Square Value (.05) 45.16
Adjusted Level of Significance 0.0195
Adjusted Chi Square Value 41.49

Anderson-Darling Test Statistic 0.441 Anderson-Darling 5% Critical Value 0.718 Kolmogorov-Smirnov Test Statistic 0.234 Kolmogorov-Smirnov 5% Critical Value 0.295

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 34.27 95% Adjusted Gamma UCL (Use when n < 40) 37.3

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 30.44
95% Jackknife UCL 31.3
95% Standard Bootstrap UCL 30.06
95% Bootstrap-t UCL 31.91
95% Hall's Bootstrap UCL 32.47
95% Percentile Bootstrap UCL 30.17
95% BCA Bootstrap UCL 30.01
95% Chebyshev(Mean, Sd) UCL 39.7
97.5% Chebyshev(Mean, Sd) UCL 46.13
99% Chebyshev(Mean, Sd) UCL 58.77

Use 95% Student's-t UCL 31.3

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

GULF KILLIFISH 95 UCL OUTPUTS

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_b.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	25	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	25
		Percent Non-Detects	100 00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

	General Stati	stics	
Number of Valid Data	25	Number of Detected Data	10
Number of Distinct Detected Data	10	Number of Non-Detect Data	15
		Percent Non-Detects	60.00%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	200	Minimum Detected	5.298
Maximum Detected	2360	Maximum Detected	7.766
Mean of Detected	663.8	Mean of Detected	6.18
SD of Detected	653.3	SD of Detected	0.793
Minimum Non-Detect	351	Minimum Non-Detect	5.861
Maximum Non-Detect	763	Maximum Non-Detect	6.637
Note: Data have multiple DLs - Use of KM Method is recommer	nded	Number treated as Non-Detect	22
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	3
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	88.00%
	UCL Statist	ics	
Normal Distribution Test with Detected Values Only	,	Lognormal Distribution Test with Detected Values Or	nly
Shapiro Wilk Test Statistic	0.714	Shapiro Wilk Test Statistic	0.916
5% Shapiro Wilk Critical Value	0.842	5% Shapiro Wilk Critical Value	0.842
Data not Normal at 5% Significance Level		Data appear Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	440.5	Mean	5.846
SD	446.7	SD	0.62
95% DL/2 (t) UCL	593.4	95% H-Stat (DL/2) UCL	544.3
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE yields a negative mean		Mean in Log Scale	5.809
		SD in Log Scale	0.578
		Mean in Original Scale	422.1
		SD in Original Scale	448.1
		95% t UCL	575.4
		95% Percentile Bootstrap UCL	588.3
		95% BCA Bootstrap UCL	662.6
		95% H-UCL	500.8

Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	
k star (bias corrected)	1.272	Data appear Gamma Distributed at 5% Significance Lev	el
Theta Star	521.9		
nu star	25.44		
A-D Test Statistic	0.538	Nonparametric Statistics	
5% A-D Critical Value	0.738	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.738	Mean	420.3
5% K-S Critical Value	0.271	SD	442.9
Data appear Gamma Distributed at 5% Significance Leve	el	SE of Mean	94.76
		95% KM (t) UCL	582.5
Assuming Gamma Distribution		95% KM (z) UCL	576.2
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	578.1
Minimum	200	95% KM (bootstrap t) UCL	746.5
Maximum	2360	95% KM (BCA) UCL	613.6
Mean	450.2	95% KM (Percentile Bootstrap) UCL	578.5
Median	343.8	95% KM (Chebyshev) UCL	833.4
SD	439.8	97.5% KM (Chebyshev) UCL	1012
k star	2.262	99% KM (Chebyshev) UCL	1363
Theta star	199.1		
Nu star	113.1	Potential UCLs to Use	
AppChi2	89.53	95% KM (t) UCL	582.5
95% Gamma Approximate UCL (Use when n >= 40)	568.6		
95% Adjusted Gamma UCL (Use when n < 40)	577.8		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 25

Raw Statistics	Log-transformed Statistics
Minimum 0.025	Minimum of Log Data -3.689
Maximum 0.109	Maximum of Log Data -2.216
Mean 0.0524	Mean of log Data -3.042
Geometric Mean 0.0477	SD of log Data 0.43
Median 0.044	
SD 0.0242	
Std. Error of Mean 0.00485	
Coefficient of Variation 0.463	

Number of Distinct Observations 21

Relevant UCL Statistics

Skewness 1.097

Theta Star 0.0106

Normal Distribution Test	Lognormal Distribution Test
Shapiro Wilk Test Statistic 0.871	Shapiro Wilk Test Statistic 0.944
Shapiro Wilk Critical Value 0.918	Shapiro Wilk Critical Value 0.918
Data not Normal at 5% Significance Level	Data appear Lognormal at 5% Significance Level
Assuming Normal Distribution	Assuming Lognormal Distribution
95% Student's-t UCL 0.0607	95% H-UCL 0.0619
95% UCLs (Adjusted for Skewness)	95% Chebyshev (MVUE) UCL 0.0723
95% Adjusted-CLT UCL (Chen-1995) 0.0615	97.5% Chebyshev (MVUE) UCL 0.0811
95% Modified-t UCL (Johnson-1978) 0.0608	99% Chebyshev (MVUE) UCL 0.0982
Gamma Distribution Test	Data Distribution
k star (bias corrected) 4.929	Data appear Gamma Distributed at 5% Significance Level

MLE of Mean 0.0524 MLE of Standard Deviation 0.0236 nu star 246.5 Approximate Chi Square Value (.05) 211.1

Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 208.9

Anderson-Darling Test Statistic 0.639 Anderson-Darling 5% Critical Value 0.747 Kolmogorov-Smirnov Test Statistic 0.162 Kolmogorov-Smirnov 5% Critical Value 0.175

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0611 95% Adjusted Gamma UCL (Use when n < 40) 0.0618

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 0.0603 95% Jackknife UCL 0.0607 95% Standard Bootstrap UCL 0.0602 95% Bootstrap-t UCL 0.063 95% Hall's Bootstrap UCL 0.0611 95% Percentile Bootstrap UCL 0.0606 95% BCA Bootstrap UCL 0.0613 95% Chebyshev(Mean, Sd) UCL 0.0735 97.5% Chebyshev(Mean, Sd) UCL 0.0826 99% Chebyshev(Mean, Sd) UCL 0.101

Use 95% Approximate Gamma UCL 0.0611

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25

Raw Statistics

Minimum 270.7 Maximum 1173 Mean 794.6 Geometric Mean 761.1 Median 792.8 SD 214.9 Std. Error of Mean 42.99

Coefficient of Variation 0.27 Skewness -0.349 **Log-transformed Statistics**

Number of Distinct Observations 25

Minimum of Log Data 5.601 Maximum of Log Data 7.067 Mean of log Data 6.635 SD of log Data 0.32

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.982 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.898 Shapiro Wilk Critical Value 0.918 Data not Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 868.2 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 862.1 95% Modified-t UCL (Johnson-1978) 867.7 **Assuming Lognormal Distribution**

95% H-UCL 902.8 95% Chebyshev (MVUE) UCL 1027 97.5% Chebyshev (MVUE) UCL 1125 99% Chebyshev (MVUE) UCL 1318

Gamma Distribution Test

k star (bias corrected) 10.38 Theta Star 76.54 MLE of Mean 794.6 MLE of Standard Deviation 246.6

nu star 519.1

Approximate Chi Square Value (.05) 467.2 Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 463.9

Anderson-Darling Test Statistic 0.406

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 865.3 95% Jackknife UCL 868.2 95% Standard Bootstrap UCL 864 95% Bootstrap-t UCL 869.1

Anderson-Darling 5% Critical Value 0.745 Kolmogorov-Smirnov Test Statistic 0.142 Kolmogorov-Smirnov 5% Critical Value 0.174

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 882.8 95% Adjusted Gamma UCL (Use when n < 40) 889.2

Potential UCL to Use

95% Hall's Bootstrap UCL 865.8 95% Percentile Bootstrap UCL 868.2 95% BCA Bootstrap UCL 860.4 95% Chebyshev(Mean, Sd) UCL 982 97.5% Chebyshev(Mean, Sd) UCL 1063 99% Chebyshev(Mean, Sd) UCL 1222

Use 95% Student's-t UCL 868.2

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

> Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25

Raw Statistics

Minimum 15.6 Maximum 137.1 Mean 77.42 Geometric Mean 70.85 Median 80.9 SD 28.09

Std. Error of Mean 5.618 Coefficient of Variation 0.363 Skewness -0.237 **Log-transformed Statistics**

Number of Distinct Observations 25

Minimum of Log Data 2.747 Maximum of Log Data 4.921 Mean of log Data 4.261 SD of log Data 0.482

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.98 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Assuming Normal Distribution

95% UCLs (Adjusted for Skewness)

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.852 Shapiro Wilk Critical Value 0.918

Data not Lognormal at 5% Significance Level **Assuming Lognormal Distribution**

95% H-UCL 96.33

95% Chebyshev (MVUE) UCL 113.7 97.5% Chebyshev (MVUE) UCL 128.6 99% Chebyshev (MVUE) UCL 158

95% Modified-t UCL (Johnson-1978) 86.99

95% Student's-t UCL 87.03

Data Distribution

Data appear Normal at 5% Significance Level

Gamma Distribution Test

k star (bias corrected) 5.129 Theta Star 15.1 MLE of Mean 77.42 MLE of Standard Deviation 34.19 nu star 256.4

Approximate Chi Square Value (.05) 220.4 Adjusted Level of Significance 0.0395 Adjusted Chi Square Value 218.1

95% Adjusted-CLT UCL (Chen-1995) 86.38

Anderson-Darling Test Statistic 0.787 Anderson-Darling 5% Critical Value 0.747

Kolmogorov-Smirnov Test Statistic 0.149 Kolmogorov-Smirnov 5% Critical Value 0.175

Nonparametric Statistics

95% CLT UCL 86.66 95% Jackknife UCL 87.03 95% Standard Bootstrap UCL 86.65 95% Bootstrap-t UCL 87 95% Hall's Bootstrap UCL 86.72 95% Percentile Bootstrap UCL 85.88 95% BCA Bootstrap UCL 86.29

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site

March 2013 040284-01

Data follow Appr. Gamma Distribution at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 90.1 95% Adjusted Gamma UCL (Use when n < 40) 91.04

Potential UCL to Use

95% Chebyshev(Mean, Sd) UCL 101.9 97.5% Chebyshev(Mean, Sd) UCL 112.5 99% Chebyshev(Mean, Sd) UCL 133.3

Use 95% Student's-t UCL 87.03

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

GULF MENHADEN 95 UCL OUTPUTS

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	10	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	10
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

	General Statistics		
Number of Valid Data	10	Number of Detected Data	6
Number of Distinct Detected Data	6	Number of Non-Detect Data	4
		Percent Non-Detects	40.00%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	398	Minimum Detected	5.986
Maximum Detected	671	Maximum Detected	6.509
Mean of Detected	490.7	Mean of Detected	6.179
SD of Detected	103.4	SD of Detected	0.197
Minimum Non-Detect	699	Minimum Non-Detect	6.55
Maximum Non-Detect	765	Maximum Non-Detect	6.64
Note: Data have multiple DLs - Use of KM Method is recommen-	ded	Number treated as Non-Detect	10
For all methods (except KM, DL/2, and ROS Methods),		Number treated as Detected	0
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	100.00%

Warning: There are only 6 Detected Values in this data

Note: It should be noted that even though bootstrap may be performed on this data set the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

	UCL Statistics		
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.87	Shapiro Wilk Test Statistic	0.903
5% Shapiro Wilk Critical Value	0.788	5% Shapiro Wilk Critical Value	0.788
Data appear Normal at 5% Significance Level		Data appear Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	439.9	Mean	6.066
SD	101.5	SD	0.209
95% DL/2 (t) UCL	498.7	95% H-Stat (DL/2) UCL	502
Maximum Likelihood Estimate(MLE) Method	N/A	Log ROS Method	
MLE method failed to converge properly		Mean in Log Scale	6.179

		SD in Log Scale	0.147
		Mean in Original Scale	487.4
		SD in Original Scale	77.17
		95% t UCL	532.1
		95% Percentile Bootstrap UCL	530.9
		95% BCA Bootstrap UCL	539
		95% H-UCL	533.6
Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	
k star (bias corrected)	14.95	Data appear Normal at 5% Significance Level	
Theta Star	32.83		
nu star	179.4		
A-D Test Statistic	0.401	Nonparametric Statistics	
5% A-D Critical Value	0.697	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.697	Mean	490.7
5% K-S Critical Value	0.332	SD	94.36
Data appear Gamma Distributed at 5% Significance Leve	el	SE of Mean	42.2
		95% KM (t) UCL	568
Assuming Gamma Distribution		95% KM (z) UCL	560.1
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	571.1
Minimum	398	95% KM (bootstrap t) UCL	742.1
Maximum	671	95% KM (BCA) UCL	557.9
Mean	496.7	95% KM (Percentile Bootstrap) UCL	560.2
Median	505.8	95% KM (Chebyshev) UCL	674.6
SD	77.44	97.5% KM (Chebyshev) UCL	754.2
k star	34.23	99% KM (Chebyshev) UCL	910.6
Theta star	14.51		
Nu star	684.7	Potential UCLs to Use	
AppChi2	624.9	95% KM (t) UCL	568
95% Gamma Approximate UCL (Use when n >= 40)	544.2	95% KM (Percentile Bootstrap) UCL	560.2
95% Adjusted Gamma UCL (Use when n < 40)	553		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006). For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Raw Statistics	Log-transformed Statistics
Minimum 0.068	Minimum of Log Data -2.688
Maximum 0.199	Maximum of Log Data -1.614
Mean 0.116	Mean of log Data -2.233

Number of Distinct Observations 9

SD of log Data 0.42

Geometric Mean 0.107 Median 0.097 SD 0.0499 Std. Error of Mean 0.0158 Coefficient of Variation 0.43 Skewness 0.64

Number of Valid Observations 10

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.851 Shapiro Wilk Test Statistic 0.862 Shapiro Wilk Critical Value 0.842 Shapiro Wilk Critical Value 0.842 Data appear Lognormal at 5% Significance Level

Data appear Normal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 0.145

95% UCLs (Adjusted for Skewness)

Gamma Distribution Test

95% Adjusted-CLT UCL (Chen-1995) 0.146 95% Modified-t UCL (Johnson-1978) 0.146 95% H-UCL 0.157 95% Chebyshev (MVUE) UCL 0.184 97.5% Chebyshev (MVUE) UCL 0.213 99% Chebyshev (MVUE) UCL 0.271

Data Distribution

Data appear Normal at 5% Significance Level

Theta Star 0.0257 MLE of Mean 0.116

MLE of Standard Deviation 0.0547

k star (bias corrected) 4.514

nu star 90.28

Approximate Chi Square Value (.05) 69.37 Adjusted Level of Significance 0.0267

Adjusted Chi Square Value 66.19

Anderson-Darling Test Statistic 0.677
Anderson-Darling 5% Critical Value 0.728
Kolmogorov-Smirnov Test Statistic 0.29
Kolmogorov-Smirnov 5% Critical Value 0.267

Data follow Appr. Gamma Distribution at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.151 95% Adjusted Gamma UCL (Use when n < 40) 0.158

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 0.142
95% Jackknife UCL 0.145
95% Standard Bootstrap UCL 0.141
95% Bootstrap-t UCL 0.15
95% Hall's Bootstrap UCL 0.143
95% Percentile Bootstrap UCL 0.142
95% BCA Bootstrap UCL 0.145
95% Chebyshev(Mean, Sd) UCL 0.185
97.5% Chebyshev(Mean, Sd) UCL 0.215

99% Chebyshev(Mean, Sd) UCL 0.273

Use 95% Student's-t UCL 0.145

Minimum of Log Data 5.284

Maximum of Log Data 6.671

Shapiro Wilk Test Statistic 0.934

Shapiro Wilk Critical Value 0.842

Mean of log Data 6.022

SD of log Data 0.448

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 10 Number of Distinct Observations 10

Raw Statistics

Minimum 197.1 Maximum 789.2 Mean 449.5 Geometric Mean 412.3 Median 442.6 SD 191.3 Std. Error of Mean 60.48

Coefficient of Variation 0.426 Skewness 0.532

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.922 Shapiro Wilk Critical Value 0.842

Data appear Normal at 5% Significance Level

Data appear Lognormal at 5% Significance Level

Log-transformed Statistics

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 560.3
 95% H-UCL 628.6

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 733.8

 95% Adjusted-CLT UCL (Chen-1995) 559.8
 97.5% Chebyshev (MVUE) UCL 856.2

 95% Modified-t UCL (Johnson-1978) 562
 99% Chebyshev (MVUE) UCL 1097

Gamma Distribution Test

k star (bias corrected) 4.236 Theta Star 106.1 Data Distribution

Data appear Normal at 5% Significance Level

MLE of Mean 449.5 MLE of Standard Deviation 218.4 nu star 84.72 Approximate Chi Square Value (.05) 64.51

Adjusted Level of Significance 0.0267 Adjusted Chi Square Value 61.45

Anderson-Darling Test Statistic 0.375
Anderson-Darling 5% Critical Value 0.729
Kolmogorov-Smirnov Test Statistic 0.205
Kolmogorov-Smirnov 5% Critical Value 0.267

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 590.3 95% Adjusted Gamma UCL (Use when n < 40) 619.7

Potential UCL to Use

Nonparametric Statistics

95% CLT UCL 548.9
95% Jackknife UCL 560.3
95% Standard Bootstrap UCL 543.1
95% Bootstrap-t UCL 579.7
95% Hall's Bootstrap UCL 627.6
95% Percentile Bootstrap UCL 545.4
95% BCA Bootstrap UCL 547.1
95% Chebyshev(Mean, Sd) UCL 713.1
97.5% Chebyshev(Mean, Sd) UCL 827.2
99% Chebyshev(Mean, Sd) UCL 1051

Use 95% Student's-t UCL 560.3

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 10

Number of Distinct Observations 10

Raw Statistics

Minimum 12.88 Maximum 46.6 Mean 31.19 Geometric Mean 28.89 Median 32.35 SD 11.61

Std. Error of Mean 3.671 Coefficient of Variation 0.372 Skewness -0.311 **Log-transformed Statistics**

Minimum of Log Data 2.556 Maximum of Log Data 3.842 Mean of log Data 3.363 SD of log Data 0.436

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.931 Shapiro Wilk Critical Value 0.842

Data appear Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.889 Shapiro Wilk Critical Value 0.842

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 37.92

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 36.84 95% Modified-t UCL (Johnson-1978) 37.86 **Assuming Lognormal Distribution**

95% H-UCL 43.34 95% Chebyshev (MVUE) UCL 50.63 97.5% Chebyshev (MVUE) UCL 58.93 99% Chebyshev (MVUE) UCL 75.24

Gamma Distribution Test

k star (bias corrected) 4.749 Theta Star 6.568 MLE of Mean 31.19 MLE of Standard Deviation 14.31

nu star 94.97 Approximate Chi Square Value (.05) 73.5

Adjusted Level of Significance 0.0267 Adjusted Chi Square Value 70.22

Anderson-Darling Test Statistic 0.483

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 37.23 95% Jackknife UCL 37.92 95% Standard Bootstrap UCL 37.04 95% Bootstrap-t UCL 37.62

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site *March 2013 040284-01*

Anderson-Darling 5% Critical Value 0.728 Kolmogorov-Smirnov Test Statistic 0.265 Kolmogorov-Smirnov 5% Critical Value 0.267

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 40.3 95% Adjusted Gamma UCL (Use when n < 40) 42.18

Potential UCL to Use

95% Hall's Bootstrap UCL 36.67 95% Percentile Bootstrap UCL 36.8 95% BCA Bootstrap UCL 36.83 95% Chebyshev(Mean, Sd) UCL 47.19 97.5% Chebyshev(Mean, Sd) UCL 54.12 99% Chebyshev(Mean, Sd) UCL 67.72

Use 95% Student's-t UCL 37.92

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

PINFISH 95 UCL OUTPUTS

Pinfish

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_d.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	4	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	4
		Percent Non-Detects	100 00%

Warning: This data set only has 4 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result Value (hexachlorobenzene ug/kg)

General Statistics

Number of Valid Data	4	Number of Detected Data	3
Number of Distinct Detected Data	3	Number of Non-Detect Data	1
		Percent Non-Detects	25.00%

Warning: This data set only has 4 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (hexachlorobenzene_ug/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 4

Number of Distinct Observations 4

Warning: This data set only has 4 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (mercury_mg/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 4

Number of Distinct Observations 4

Warning: This data set only has 4 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

Pinfish

The data set for variable Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 4

Number of Distinct Observations 4

Warning: This data set only has 4 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

STRIPED MULLET 95 UCL OUTPUTS

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	9	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	9
		Percent Non-Detects	100.00%

Warning: All observations are Non-Detects (NDs), therefore all statistics and estimates should also be NDs!

Specifically, sample mean, UCLs, UPLs, and other statistics are also NDs lying below the largest detection limit!

The Project Team may decide to use alternative site specific values to estimate environmental parameters (e.g., EPC, BTV).

The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

Result_Value (hexachlorobenzene_ug/kg)

	General Stati	stics	
Number of Valid Data	9	Number of Detected Data	7
Number of Distinct Detected Data	7	Number of Non-Detect Data	2
		Percent Non-Detects	22.22%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	389	Minimum Detected	5.964
Maximum Detected	1830	Maximum Detected	7.512
Mean of Detected	822.6	Mean of Detected	6.554
SD of Detected	529.7	SD of Detected	0.593
Minimum Non-Detect	727	Minimum Non-Detect	6.589
Maximum Non-Detect	742	Maximum Non-Detect	6.609
Note: Data have multiple DLs - Use of KM Method is recommen	ded	Number treated as Non-Detect	6
For all methods (except KM, DL/2, and ROS Methods),	Number treated as Detected	3	
Observations < Largest ND are treated as NDs		Single DL Non-Detect Percentage	66.67%

Warning: There are only 7 Detected Values in this data

Note: It should be noted that even though bootstrap may be performed on this data set the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

	UCL Statistics		
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.832	Shapiro Wilk Test Statistic	0.892
5% Shapiro Wilk Critical Value	0.803	5% Shapiro Wilk Critical Value	0.803
Data appear Normal at 5% Significance Level		Data appear Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	721.4	Mean	6.41
SD	500.8	SD	0.587
95% DL/2 (t) UCL	1032	95% H-Stat (DL/2) UCL	1190
Maximum Likelihood Estimate(MLE) Method		Log ROS Method	
Mean	1360	Mean in Log Scale	6.47

SD	368.9	SD in Log Scale	0.539
95% MLE (t) UCL	1588	Mean in Original Scale	746.7
95% MLE (Tiku) UCL	1748	SD in Original Scale	482.8
		95% t UCL	1046
		95% Percentile Bootstrap UCL	1016
		95% BCA Bootstrap UCL	1103
		95% H UCL	1157
Gamma Distribution Test with Detected Values Only		Data Distribution Test with Detected Values Only	
k star (bias corrected)	1.984	Data appear Normal at 5% Significance Level	
Theta Star	414.6		
nu star	27.78		
A-D Test Statistic	0.472	Nonparametric Statistics	
5% A-D Critical Value	0.711	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.711	Mean	740.9
5% K-S Critical Value	0.313	SD	459.7
Data appear Gamma Distributed at 5% Significance Lev	el	SE of Mean	166
		95% KM (t) UCL	1050
Assuming Gamma Distribution		95% KM (z) UCL	1014
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	1045
Minimum	389	95% KM (bootstrap t) UCL	1310
Maximum	1830	95% KM (BCA) UCL	1057
Mean	758.1	95% KM (Percentile Bootstrap) UCL	1033
Median	532.6	95% KM (Chebyshev) UCL	1465
SD	476.3	97.5% KM (Chebyshev) UCL	1778
k star	2.586	99% KM (Chebyshev) UCL	2393
Theta star	293.2		
Nu star	46.54	Potential UCLs to Use	
AppChi2	31.89	95% KM (t) UCL	1050
95% Gamma Approximate UCL (Use when n >= 40)	1107	95% KM (Percentile Bootstrap) UCL	1033
95% Adjusted Gamma UCL (Use when n < 40)	1202		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 9 Number of Distinct Observations 8

Raw Statistics

Minimum 0.024
Maximum 0.122
Maximum of Log Data -3.73
Mean 0.0516
Mean of log Data -3.096
Geometric Mean 0.0452
Median 0.041
SD 0.0317
Std. Error of Mean 0.0106
Coefficient of Variation 0.614
Skewness 1.75

Warning: There are only 9 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.768 Shapiro Wilk Critical Value 0.829

Data not Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.0712

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 0.0755 95% Modified-t UCL (Johnson-1978) 0.0722

Gamma Distribution Test

k star (bias corrected) 2.728 Theta Star 0.0189

MLE of Mean 0.0516

MLE of Standard Deviation 0.0312

nu star 49.11

Approximate Chi Square Value (.05) 34.02

Adjusted Level of Significance 0.0231

Adjusted Chi Square Value 31.39

Anderson-Darling Test Statistic 0.685

Anderson-Darling 5% Critical Value 0.725

Kolmogorov-Smirnov Test Statistic 0.283

Kolmogorov-Smirnov 5% Critical Value 0.28 Data follow Appr. Gamma Distribution at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0744 95% Adjusted Gamma UCL (Use when n < 40) 0.0806

Potential UCL to Use

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.896 Shapiro Wilk Critical Value 0.829

Data appear Lognormal at 5% Significance Level

Assuming Lognormal Distribution

95% H-UCL 0.0775

95% Chebyshev (MVUE) UCL 0.0891

97.5% Chebyshev (MVUE) UCL 0.106

99% Chebyshev (MVUE) UCL 0.138

Data Distribution

Data Follow Appr. Gamma Distribution at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 0.0689

95% Jackknife UCL 0.0712

95% Standard Bootstrap UCL 0.0678

95% Bootstrap-t UCL 0.115

95% Hall's Bootstrap UCL 0.19

95% Percentile Bootstrap UCL 0.0699

95% BCA Bootstrap UCL 0.0746

95% Chebyshev(Mean, Sd) UCL 0.0975 97.5% Chebyshev(Mean, Sd) UCL 0.117

99% Chebyshev(Mean, Sd) UCL 0.157

Use 95% Approximate Gamma UCL 0.0744

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 9

Number of Distinct Observations 9

Raw Statistics

Minimum 103 7

Maximum 789 3

Mean 355.8 Geometric Mean 313

Median 321.5

SD 193.1

Std. Error of Mean 64.38

Coefficient of Variation 0.543

Skewness 1.422

Log-transformed Statistics Minimum of Log Data 4.642

Maximum of Log Data 6.671

Mean of log Data 5.746

SD of log Data 0.553

Warning: There are only 9 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

The literature suggests to use bootstrap methods on data sets having more than 10-15 observations.

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site

March 2013 040284-01

Shapiro Wilk Test Statistic 0.879
Shapiro Wilk Critical Value 0.829

Data appear Normal at 5% Significance Level

Shapiro Wilk Test Statistic 0.944
Shapiro Wilk Critical Value 0.829
Data appear Lognormal at 5% Significance Level

95% H-UCL 574.8

95% CLT UCL 461.7

95% Jackknife UCL 475.5

97.5% Chebyshev(Mean, Sd) UCL 757.9

99% Chebyshev(Mean, Sd) UCL 996.4

95% Chebyshev (MVUE) UCL 650.6

97.5% Chebyshev (MVUE) UCL 777.2

99% Chebyshev (MVUE) UCL 1026

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 475.5 95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 494.3 95% Modified-t UCL (Johnson-1978) 480.6

Gamma Distribution Test Data Distribution

k star (bias corrected) 2.784 Data appear Normal at 5% Significance Level
Theta Star 127.8
MLE of Mean 355.8

MLE of Standard Deviation 213.3 nu star 50.11

Adjusted Chi Square Value 32.19

Approximate Chi Square Value (.05) 34.85 **Nonparametric Statistics**Adjusted Level of Significance 0.0231 95% C

95% Standard Bootstrap UCL 456.5
Anderson-Darling Test Statistic 0.322
Anderson-Darling 5% Critical Value 0.725
Kolmogorov-Smirnov Test Statistic 0.172
Kolmogorov-Smirnov 5% Critical Value 0.28
95% BCA Bootstrap UCL 458.7
85% Percentile Bootstrap UCL 458.7

Data appear Gamma Distributed at 5% Significance Level 95% Chebyshev (Mean, Sd) UCL 636.4

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 511.5 95% Adjusted Gamma UCL (Use when n < 40) 553.8

Potential UCL to Use Use 95% Student's-t UCL 475.5

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 9 Number of Distinct Observations 9

Raw Statistics

Minimum 4.26 Maximum 51.5 Mean 21.63 Geometric Mean 17.19 Median 19.27

Std. Error of Mean 4.679 Coefficient of Variation 0.649

Skewness 1.024

SD 14.04

Log-transformed Statistics

Minimum of Log Data 1.449
Maximum of Log Data 3.942
Mean of log Data 2.844
SD of log Data 0.788

Warning: There are only 9 Values in this data

Note: It should be noted that even though bootstrap methods may be performed on this data set, the resulting calculations may not be reliable enough to draw conclusions

 $The \ literature \ suggests \ to \ use \ bootstrap \ methods \ on \ data \ sets \ having \ more \ than \ 10-15 \ observations.$

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.909 Shapiro Wilk Test Statistic 0.902

Shapiro Wilk Critical Value 0.829 Shapiro Wilk Critical Value 0.829 Data appear Normal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 30.33

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 31.03 95% Modified-t UCL (Johnson-1978) 30.59

Gamma Distribution Test

k star (bias corrected) 1.628 Theta Star 13.28 MLE of Mean 21.63 MLE of Standard Deviation 16.95 nu star 29.31

Approximate Chi Square Value (.05) 17.95 Adjusted Level of Significance 0.0231 Adjusted Chi Square Value 16.1

Anderson-Darling Test Statistic 0.372 Anderson-Darling 5% Critical Value 0.729 Kolmogorov-Smirnov Test Statistic 0.207 Kolmogorov-Smirnov 5% Critical Value 0.282

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 35.31 95% Adjusted Gamma UCL (Use when n < 40) 39.36

Potential UCL to Use

Data appear Lognormal at 5% Significance Level

Assuming Lognormal Distribution

95% H-UCL 50.94 95% Chebyshev (MVUE) UCL 48.94 97.5% Chebyshev (MVUE) UCL 60.38

99% Chebyshev (MVUE) UCL 82.87

Data Distribution

Data appear Normal at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 29.32 95% Jackknife UCL 30.33 95% Standard Bootstrap UCL 28.99 95% Bootstrap-t UCL 31.99 95% Hall's Bootstrap UCL 38.45 95% Percentile Bootstrap UCL 29.28 95% BCA Bootstrap UCL 30.6 95% Chebyshev(Mean, Sd) UCL 42.02 97.5% Chebyshev(Mean, Sd) UCL 50.85 99% Chebyshev(Mean, Sd) UCL 68.18

Use 95% Student's-t UCL 30.33

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

WHITE SHRIMP 95 UCL OUTPUTS

White Shrimp

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1 e.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (1,3-dichlorobenzene_ug/kg)

General Statistics

Number of Valid Data	3	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	3
		Percent Non-Detects	100.00%

Warning: This data set only has 3 observations! Data set is too small to compute reliable and meaningful statistics and estimates! The data set for variable Result_Value (1,3-dichlorobenzene_ug/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods! If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result Value (1.4-dichlorobenzene ug/kg)

General Statistics

Number of Valid Data	1	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	1
		Percent Non-Detects	100.00%

Warning: This data set only has 1 observations! Data set is too small to compute reliable and meaningful statistics and estimates! The data set for variable Result_Value (1,4-dichlorobenzene_ug/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods! If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (hexachlorobenzene_ug/kg)

General Statistics

Number of Valid Observations 3

Number of Distinct Observations 3

Warning: This data set only has 3 observations! Data set is too small to compute reliable and meaningful statistics and estimates! The data set for variable Result_Value (hexachlorobenzene_ug/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods! If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (hexachlorobutadiene_ug/kg)

General Statistics

Number of Valid Data	1	Number of Detected Data	0
Number of Distinct Detected Data	0	Number of Non-Detect Data	1
		Percent Non-Detects	100.00%

Warning: This data set only has 1 observations!

White Shrimp

Data set is too small to compute reliable and meaningful statistics and estimates! The data set for variable Result_Value (hexachlorobutadiene_ug/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (lead_mg/kg)

General Statistics

Number of Valid Observations 1

Number of Distinct Observations 1

Warning: This data set only has 1 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (lead_mg/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (mercury_mg/kg)

General Statistics

Number of Valid Observations 3

Number of Distinct Observations 3

Warning: This data set only has 3 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (mercury_mg/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (total dioxin/furan teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 3

Number of Distinct Observations 3

Warning: This data set only has 3 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (total dioxin/furan teq 1998 (avian) (u = 1/2)_ng/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (total pah (16) (u = 1/2)_ug/kg)

General Statistics

Number of Valid Observations 1

Number of Distinct Observations 1

Warning: This data set only has 1 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (total pah (16) (u = 1/2)_ug/kg) was not processed!

White Shrimp

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 3

Number of Distinct Observations 3

Warning: This data set only has 3 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (total pcb congener teq 1998 (avian) (u = 1/2)_ng/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 3

Number of Distinct Observations 3

Warning: This data set only has 3 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (total pcb congener teq 2005 (mammal) (u = 1/2)_ng/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: All Prey Species Separate

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

10

Food Type	$P_{BIOTA,k}$	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.066	0.022	948	Fish	1	GKF
2	0.066	0.022	948	Fish	1	STM
3	0.066	0.022	948	Fish	1	GM
4	0.066	0.022	948	Fish	1	PNF
5	0.066	0.022	948	Fish	1	SAS
6	0.0825	0.022	824	Invertebrate	1	BCR
7	0.0825	0.022	824	Invertebrate	1	OYS
8	0.0825	0.022	824	Invertebrate	1	BRS
9	0.0825	0.022	824	Invertebrate	1	WHS
10	0.34	0.022	1326	Mammal	0.5	Marsh Rice Rat

Number of foraging subareas (j = 1 to i):

		- 1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: All Prey Species Separate

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.003055	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0.00372	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	0.00415	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 5	0.00315	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 6	0.00354375	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 7	0.0031875	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 8	0.00313125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 9	0.002625	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.38E-02	mg/kg ww	
Number of food types ($k = 1$ to m):			

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	1	1	0.05	0.0611	Table C-4.2
2	Fish	STM	3	1	1	0.05	0.0744	Table C-4.2
3	Fish	GM	3	1	1	0.05	0.145	Table C-4.2
4	Fish	PNF	3	1	1	0.05	0.083	Table C-4.2
5	Fish	SAS	3	1	1	0.05	0.063	Table C-4.2
6	Invertebrate	BCR	2	1	1	0.0625	0.0567	Table C-4.2
7	Invertebrate	OYS	2	1	1	0.0625	0.051	Table C-4.2
8	Invertebrate	BRS	2	1	1	0.0625	0.0501	Table C-4.2
9	Invertebrate	WHS	2	1	1	0.0625	0.042	Table C-4.2
10	Plant	Plant			1	0.5	2.43E-03	Table C-4.2

Definitions:

kg - kilogram mg - milligrams Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: All Prey Species Separate

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 5	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 6	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 7	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 8	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 9	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	
Number of food types ($k = 1$ to m):			

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	1	1	0.05		Table C-4.2
2	Fish	STM	3	1	1	0.05		Table C-4.2
3	Fish	GM	3	1	1	0.05		Table C-4.2
4	Fish	PNF	3	1	1	0.05		Table C-4.2
5	Fish	SAS	3	1	1	0.05		Table C-4.2
6	Invertebrate	BCR	2	1	1	0.0625		Table C-4.2
7	Invertebrate	OYS	2	1	1	0.0625		Table C-4.2
8	Invertebrate	BRS	2	1	1	0.0625		Table C-4.2
9	Invertebrate	WHS	2	1	1	0.0625		Table C-4.2
10	Plant	Plant			1	0.5	6.60E-02	Table C-4.2

Definitions:

kg - kilogram

mg - milligrams

Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: All Prey Species Separate

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	$C_{\text{BIOTA,8}}$	$C_{\text{BIOTA},9}$	$C_{\text{BIOTA},10}$	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)							mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _{BW} /day		NA			
Mercury (Methyl)	0.0611	0.0744	0.145	0.083	0.063	0.0567	0.051	0.0501	0.042	3.38E-02	5.0E-03	0.0479							1.3E-05	5.0E-03	0.064	0.08
Mercury (Inorganic)										3.40E-04	3.7E-06	4.7391							1.3E-03	1.3E-03	0.9	0.0014

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: All Prey Species Separate

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	0.066	0.022	948	Fish	1	GKF
2	0.066	0.022	948	Fish	1	STM
3	0.066	0.022	948	Fish	1	GM
4	0.066	0.022	948	Fish	1	PNF
5	0.066	0.022	948	Fish	1	SAS
6	0.0825	0.027	824	Invertebrate	1	BCR
7	0.0825	0.027	824	Invertebrate	1	OYS
8	0.0825	0.027	824	Invertebrate	1	BRS
9	0.0825	0.027	824	Invertebrate	1	WHS
10	0.34	0.111	1326	Mammal	0.5	Marsh Rice Rat

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: All Prey Species Separate
COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0021	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	4.34E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	2.38E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	2.80E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	7.38E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 5	9.41E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 6	3.85E-04	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 7	3.16E-04	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 8	5.24E-04	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 9	2.73E-04	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.95E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.62E-03	mg/kg ww	
Number of food types ($k = 1$ to m):		-	

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.05	8.68E-04	Table C-4.2
2	Fish	STM	3	14	1	0.05	4.76E-04	Table C-4.2
3	Fish	GM	3	14	1	0.05	5.60E-04	Table C-4.2
4	Fish	PNF	3	14	1	0.05	1.48E-04	Table C-4.2
5	Fish	SAS	3	14	1	0.05	1.88E-04	Table C-4.2
6	Invertebrate	BCR	2	1	1	0.0625	4.40E-04	Table C-4.2
7	Invertebrate	OYS	2	1	1	0.0625	3.61E-04	Table C-4.2
8	Invertebrate	BRS	2	1	1	0.0625	5.99E-04	Table C-4.2
9	Invertebrate	WHS	2	1	1	0.0625	3.12E-04	Table C-4.2
10	Plant	Plant			1	0.5	4.35E-06	Table C-4.2

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: All Prey Species Separate

COPC	$C_{\text{BIOTA},1}$	$C_{BIOTA,2}$	$C_{BIOTA,3}$	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA,6}}$	$C_{BIOTA,7}$	C _{BIOTA,8}	C _{BIOTA,9}	C _{BIOTA,10}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	C _{SED,6}	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
					mg/kg (we	t weight)					mg/kg _{BW} /day			mg/k	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{sw} /day	NA
PCB Congener TEQ (Avian) (U=1/2)	0.00087	0.00048	0.00056	0.000148	0.00019	0.00044	0.00036	0.0006	0.00031	1.62E-03	6.6E-05	0.0021							5.5E-07	6.7E-05	1.4E-04	0.48

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: All Prey Species Separate

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.178	0.121	948	Fish	1	GKF
2	0.178	0.121	948	Fish	1	STM
3	0.178	0.121	948	Fish	1	GM
4	0.178	0.121	948	Fish	1	PNF
5	0.178	0.121	948	Fish	1	SAS
6	0.028	0.019	824	Invertebrate	1	BCR
7	0.028	0.019	824	Invertebrate	1	BRS
8	0.027	0.018	824	Invertebrate	1	WHS
9	0.027	0.018	824	Invertebrate	1	OYS

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

9

Definitions:

kg - kilogram mg - milligrams kcal - kilocalories

Belted Kingfisher Receptor: Receptor Group: **Piscivorous Birds** Scenario:

All Prey Species Separate

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	C _{BIOTA,8}	C _{BIOTA,9}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	C _{SED,6}	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
500000000000000000000000000000000000000			mg/l	kg (wet wei	ght)					mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Mercury (Methyl)	0.0611	0.0744	0.145	0.083	0.063	0.0567	0.0501	0.042	0.051	3.8E-02	0.0479							0.0E+00	3.8E-02	0.064	0.60
Hexachlorobenzene	0.5825	1.05	0.568	2.29	0.325	1.088	0.7977	0.682	0.704	4.5E-01	2.1373							0.0E+00	4.5E-01	3.35	0.13
1,3-Dichlorobenzene	0.763	0.769	0.765	0.765	0.37	0.748	0.709	0.758	0.704	3.3E-01	0.0736							0.0E+00	3.3E-01	160.8	0.0020
PCB Congener TEQ (Avian) (U=1/2)	0.00087	0.00048	0.00056	0.00015	0.00019	0.00044	0.0006	0.00031	0.00036	2.1E-04	0.0021							0.0E+00	2.1E-04	0.00014	1.50
Mercury (Inorganic)										0.0E+00	4.7391							0.0E+00	0.0E+00	9.0E-01	0

Baseline Ecological Risk Assessment March 2013 Patrick Bayou Superfund Site 040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal
Scenario: All Prey Species Separate

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME _{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTS}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	0.0466	0.004	1092	Fish	1	GKF
2	0.0466	0.004	1092	Fish	1	STM
3	0.0466	0.004	1092	Fish	1	GM
4	0.0466	0.004	1092	Fish	1	PNF
5	0.0466	0.004	1092	Fish	1	SAS
6	0.0585	0.005	974	Invertebrate	1	BCR
7	0.0585	0.005	974	Invertebrate	1	OYS
8	0.0585	0.005	974	Invertebrate	1	BRS
9	0.0585	0.005	974	Invertebrate	1	WHS
10	0.3	0.027	962	Plant	1	
11	0.233	0.021	1428	Mammal	0.5	Marsh Rice Rat

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

11

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: **Terrestrial Mammal - Marsh Rice Rat**

Scenario: **All Prey Species Separate**

COPC: **Methyl Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	3.06E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.72E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	7.25E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	4.15E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 5	3.15E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 6	3.54E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 7	3.19E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 8	3.13E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 9	2.63E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.38E-02	mg/kg ww	

Number of food types (k = 1 to m): 10

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	1	1	0.05	6.11E-02	Table C-4.2
2	Fish	STM	3	1	1	0.05	7.44E-02	Table C-4.2
3	Fish	GM	3	1	1	0.05	1.45E-01	Table C-4.2
4	Fish	PNF	3	1	1	0.05	8.30E-02	Table C-4.2
5	Fish	SAS	3	1	1	0.05	6.30E-02	Table C-4.2
6	Invertebrate	BCR	2	1	1	0.0625	5.67E-02	Table C-4.2
7	Invertebrate	OYS	2	1	1	0.0625	5.10E-02	Table C-4.2
8	Invertebrate	BRS	2	1	1	0.0625	5.01E-02	Table C-4.2
9	Invertebrate	WHS	2	1	1	0.0625	4.20E-02	Table C-4.2
10	Plant				1	0.5	2.43E-03	Table C-4.2

Definitions:

kg - kilogram

mg - milligrams dw - dry weight ww - wet weight

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: **All Prey Species Separate**

COPC: **Inorganic Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
P _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 5	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 6	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 7	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 8	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 9	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	

Number of food types (k = 1 to m): 10

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$P_{BIOTA,k}$	FR _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	1	1	0.05		Table C-4.2
2	Fish	STM	3	1	1	0.05		Table C-4.2
3	Fish	GM	3	1	1	0.05		Table C-4.2
4	Fish	PNF	3	1	1	0.05		Table C-4.2
5	Fish	SAS	3	1	1	0.05		Table C-4.2
6	Invertebrate	BCR	2	1	1	0.0625		Table C-4.2
7	Invertebrate	OYS	2	1	1	0.0625		Table C-4.2
8	Invertebrate	BRS	2	1	1	0.0625		Table C-4.2
9	Invertebrate	WHS	2	1	1	0.0625		Table C-4.2
10	Plant				1	0.5	6.60E-02	Table C-4.2

Definitions:

kg - kilogram

mg - milligrams dw - dry weight

ww - wet weight

Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: All Prey Species Separate

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{BIOTA,3}$	C _{BIOTA,4}	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	C _{BIOTA,8}	C _{BIOTA,9}	C _{BIOTA,10}	C _{BIOTA,11}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)											mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg	_{BW} /day	NA	
Mercury (Methyl)	0.0611	0.0744	0.145	0.083	0.063	0.0567	0.051	0.0501	0.042	0.00243	3.38E-02	6.9E-05	0.0479							8.9E-06	7.8E-05	0.0177	0.0044
Mercury (Inorganic)										0.06575	3.40E-04	3.8E-05	4.7391							8.8E-04	9.1E-04	3.54	0.00026

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Raccoon

Receptor Group: Omnivorous Mammal
Scenario: All Prey Species Separate

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME _{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTS}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	0.0466	0.004	1092	Fish	1	GKF
2	0.0466	0.004	1092	Fish	1	STM
3	0.0466	0.004	1092	Fish	1	GM
4	0.0466	0.004	1092	Fish	1	PNF
5	0.0466	0.004	1092	Fish	1	SAS
6	0.0585	0.005	974	Invertebrate	1	BCR
7	0.0585	0.005	974	Invertebrate	1	OYS
8	0.0585	0.005	974	Invertebrate	1	BRS
9	0.0585	0.005	974	Invertebrate	1	WHS
10	0.3	0.027	962	Plant	1	
11	0.233	0.021	1428	Mammal	0.5	Marsh Rice Rat

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

11

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: All Prey Species Separate
COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	4.35E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	1.52E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.90E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	2.57E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 5	1.30E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 6	2.67E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 7	1.81E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 8	2.74E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 9	1.58E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	1.91E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	9.98E-05	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:	,		

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.05	8.70E-05	Table C-4.2
2	Fish	STM	3	14	1	0.05	3.03E-05	Table C-4.2
3	Fish	GM	3	14	1	0.05	3.79E-05	Table C-4.2
4	Fish	PNF	3	14	1	0.05	5.14E-05	Table C-4.2
5	Fish	SAS	3	14	1	0.05	2.59E-05	Table C-4.2
6	Invertebrate	BCR	2	1	1	0.0625	3.06E-05	Table C-4.2
7	Invertebrate	OYS	2	1	1	0.0625	2.07E-05	Table C-4.2
8	Invertebrate	BRS	2	1	1	0.0625	3.13E-05	Table C-4.2
9	Invertebrate	WHS	2	1	1	0.0625	1.81E-05	Table C-4.2
10	Plant	Plant			1	0.5	2.32E-07	Table C-4.2

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor:

Raccoon

Receptor Group: Omnivorous Mammal
Scenario: All Prey Species Separate

COPC	C _{BIOTA,1}	$C_{\text{BIOTA,2}}$	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	C _{BIOTA,8}	C _{BIOTA,9}	C _{BIOTA,10}	C _{BIOTA,11}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)										mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA	
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	.7E-05 3E-05 3.8E-05 5.1E-05 0.0000259 3.1E-05 2.1E-05 3.1E-05 1.8E-05 2.32E-07 9.98						9.98E-05	5.4E-08	0.0001							2.1E-08	7.5E-08	5.4E-07	0.14			

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
All Prey Species Separate

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.3		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	0.33	0.328	824	Invert	1	OYS
2	0.33	0.328	824	Invert	1	BCR
3	0.34	0.337	824	Invert	1	WHS
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description						
1	1.00							
2								
3								
4								
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Scenario: All Prey Species Separate

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	C _{SED,3}	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet we	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Lead	0.331	0.226	0.116					7.5E-02	48.944							5.8E-02	1.3E-01	3.26	0.04
Mercury (Methyl)	0.051	0.0567	0.042					1.7E-02	0.0479							5.6E-05	1.7E-02	0.064	0.26
Total PAH 16 (U=1/2)	0.55212	0.07592	0.1623					8.9E-02	7.3078							8.6E-03	9.7E-02	325	0.00030
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2E-05	2.2E-05	2.5E-05					7.6E-06	0.0002							2.5E-07	7.8E-06	0.00014	0.06
Hexachlorobenzene	0.704	1.088	0.682					2.8E-01	2.1373							2.5E-03	2.8E-01	3.35	0.08
Hexachlorobutadiene	0.704	0.745	0.714					2.4E-01	3.8327							4.5E-03	2.5E-01	20	0.01
1,3-Dichlorobenzene	0.704	0.748	0.758					2.5E-01	0.0736							8.7E-05	2.5E-01	160.8	0.0015
1,4-Dichlorobenzene	0.704	0.745	0.714					2.4E-01	0.0744							8.7E-05	2.4E-01	160.8	0.0015
PCB Congener TEQ (Avian) (U=1/2)	0.00036	0.00044	0.00031					1.3E-04	0.0021							2.4E-06	1.3E-04	1.4E-04	0.91
Mercury (Inorganic)								0.0E+00	4.7391							5.6E-03	5.6E-03	9.0E-01	0.006

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.4 SEDIMENT EPC UNCERTAINTYANALYSIS SITE-WIDE SWAC AS EPC

Table C-5.4
Comparison of Baseline SWAC and SWAC Calculated With All Tidal Zones

СОРС	Units	Intertidal, and Supratidal SWAC (Baseline)	Subtidal, Intertidal, and Supratidal SWAC
1,3-Dichlorobenzene	mg/kg	0.074	0.11
1,4-Dichlorobenzene	mg/kg	0.074	0.08
Hexachlorobenzene	mg/kg	2.14	3.37
Hexachlorobutadiene	mg/kg	3.83	6.07
Lead	mg/kg	48.94	46.24
Mercury	mg/kg	4.79	10.18
Total Dioxin/Furan TEQ 1998 (Avian; U=1/2)	mg/kg	0.00022	0.00028
Total PAH (U=1/2)	mg/kg	7.31	9.36
Total PCB Congener TEQ 1998 (Avian; U=1/2)	mg/kg	0.0021	0.0023
Total PCB Congener TEQ 2005 (Mammal; U=1/2)	mg/kg	0.00011	0.00012

Notes:

SWAC - surface weighted average concentration

mg/kg - milligrams per kilogram

COPC - chemical of potential concern PAH - polycyclic aromatic hydrocarbon

PCB - polychlorinated biphenyl

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.165	0.054	948	Fish	1	GM
2	0.165	0.054	948	Fish	1	GKF, STM, PNF, SAS
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5						
6						
7						

Number of foraging subareas (j = 1 to i):

		<u> </u>
Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

4

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: **Terrestrial Mammal - Marsh Rice Rat**

Scenario: **Sediment Site-Wide SWAC**

COPC: **Methyl Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.1018	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.018125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	8.49E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Sediment Site-Wide SWAC

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	10.0815	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	5.62E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	6.37E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2		GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6					•			
7								

Definitions:

kg - kilogram

mg - milligrams

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA},6}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	C _{SED,5}	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
38.000000000000000000000000000000000000		mg/kg (wet weight)						mg/kg _{BW} /day	mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Mercury (Methyl)	0.145	.145 0.058 0.0515 3.83E-02						6.7E-03	0.1018							2.7E-05	6.7E-03	0.064	0.11
Mercury (Inorganic)				6.37E-04				3.5E-05	10.081							2.7E-03	2.7E-03	0.9	0.0030

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.11	0.036	948	Fish	1	GKF
2	0.11	0.036	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5	0.11	0.036	948	Fish	1	PNF, SAS
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Sediment Site-Wide SWAC COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C_{SED}	0.0023	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.21E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.52E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5	1.33E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	5.39E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.71E-03	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.083	8.68E-04	Table C-4.2
2	Fish	GM, STM	3	14	1	0.083	4.24E-04	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-4.2
4	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
5	Fish	PNF, SAS	3	14	1	0.083	1.60E-04	Table C-4.2
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds

Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
60.000.000.000.000.000	mg/kg (wet weight)						mg/kg _{BW} /day	/kg _{BW} /day mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA	
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	1.71E-03	0.0002			7.1E-05	0.0023							6.0E-07	7.2E-05	1.4E-04	0.51

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.607	948	Fish	1	GKF, GM, STM, PNF, SAS
2	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
3						
4						
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA,6}}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
50 - 50 - 50 - 50 - 50 - 50		mg/kg (wet weight)							mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg	_{BW} /day	NA	
Hexachlorobenzene	0.6464	0.9757						3.2E-01	3.3726							0.0E+00	3.2E-01	3.35	0.10
1,3-Dichlorobenzene	0.769	0.758						3.6E-01	0.106			•	-			0.0E+00	3.6E-01	160.8	0.0022

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.445	0.304	948	Fish	1	GM
2	0.445	0.304	948	Fish	1	GKF, STM, PNF, SAS
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4						
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1
---	---

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
8. 80. 80. 80. 80. 80.			mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515					4.5E-02	0.1018							0.0E+00	4.5E-02	0.064	0.71
Mercury (Inorganic)							_	0.0E+00	10.081				_			0.0E+00	0.0E+00	0.9	0

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.3	0.205	948	Fish	1	GKF
2	0.3	0.205	948	Fish	1	GM, STM
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.29	0.205	948	Fish	1	PNF, SAS
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	* mg/kg (wet weight)				mg/kg _{BW} /day			mg/k	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA			
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	0.0002				2.4E-04	0.0023							0.0E+00	2.4E-04	0.00014	1.69

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GM
2	0.116	0.010	1092	Fish	1	GKF, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: Sediment Site-Wide SWAC

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	1.02E-01	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.81E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	7.25E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.29E-02	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	8.49E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant				1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight ww - wet weight

Scenario: Sediment Site-Wide SWAC

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C_{SED}	10.0815	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	5.62E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	6.37E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{BIOTA,3}$	C _{BIOTA,4}	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
mg/kg (wet weight)						mg/kg _{BW} /day	y mg/kg (dry weight) mg/k						mg/kg _{BW} /day	mg/kg	_{BW} /day	NA			
Mercury (Methyl)	0.145	0.058	0.0515	0.002426492	3.83E-02			7.7E-05	0.1018							1.9E-05	9.6E-05	0.0177	0.0054
Mercury (Inorganic)				0.065754378	6.37E-04			3.8E-05	10.081							1.9E-03	1.9E-03	3.54	0.00054

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GKF
2	0.116	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1
1 value of loraging subareas $y = 1$ to y .	

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: Sediment Site-Wide SWAC
COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.09E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	3.99E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	9.75E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.06E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.12E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.125	8.70E-05	Table C-4.2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.125	3.19E-05	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	2.79E-05	Table C-4.2
4	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor: Raccoon

Receptor Group: Omnivorous Mammal
Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA},6}$	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	•	mg/kg (wet weight)					mg/kg _{BW} /day	mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA	
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	3.2E-05	2.8E-05	2.32E-07	1.12E-04			6.4E-08	0.0001							2.2E-08	8.6E-08	5.4E-07	0.16

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Scenario: Sediment Site-Wide SWAC

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Scenario: Sediment Site-Wide SWAC

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	C _{BIOTA,5}	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Lead	0.292							9.9E-02	46.238							3.3E-02	1.3E-01	3.26	0.04
Mercury (Methyl)	0.0533							1.8E-02	0.1018							7.2E-05	1.8E-02	0.064	0.28
Total PAH 16 (U=1/2)	0.1397							4.7E-02	9.3648							6.6E-03	5.4E-02	325	0.00017
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							7.5E-06	0.0003							2.0E-07	7.7E-06	0.00014	0.06
Hexachlorobenzene	1.132							3.8E-01	3.3726							2.4E-03	3.8E-01	3.35	0.11
Hexachlorobutadiene	0.3725							1.3E-01	6.07							4.3E-03	1.3E-01	20	0.006
1,3-Dichlorobenzene	0.3725							1.3E-01	0.106							7.5E-05	1.3E-01	160.8	0.0008
1,4-Dichlorobenzene	0.3725							1.3E-01	0.084							5.9E-05	1.3E-01	160.8	0.0008
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.4E-04	0.0023							1.6E-06	1.5E-04	1.4E-04	1.05
Mercury (Inorganic)								0.0E+00	10.081							7.1E-03	7.1E-03	9.0E-01	0.008

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.5 AUF UNCERTAINTY ANALYSIS — LOWER-BOUND ESTIMATE OF HOME RANGE

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.17		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

 ME_k $\overline{\mathsf{FR}}_{\mathsf{BIOTA},\mathsf{k}}$ Food Type $P_{\text{BIOTA},k}$ NIR_k Prey Group Description 0.165 0.054 948 Fish 0.165 0.054 948 Fish GKF, STM, PNF, SAS 2 1 BCR, BRS, WHS, OYS 3 0.33 0.108 824 Invertebrate 0.34 0.111 1326 Mammal March Rice Rat 0.5 5 6 7

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

4

Definitions:

kg - kilogram

 $\ mg\ \hbox{-}\ milligrams$

kcal - kilocalories

Scenario: AUF_{Lower}

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C_SED	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.018125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams

Scenario: AUF_{Lower}

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter		Value	Units	Source
FCM _{TL3(OM)}		1.0		Table C-4.2
BCF _{PLANT-OM}		2.30E-03		Table C-4.2
C _{SED}		4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}		5.57E-05		Table C-4.2
FR _{SED}		1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Typ	e 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Typ		0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Typ	e 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant		7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sedimen	t	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	-	3.40E-04	mg/kg ww	
November of food to part of the second	4			·

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1		GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6					•			
7					•			

Definitions:

kg - kilogram mg - milligrams Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	C _{BIOTA,5}	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
35.000.000.000.000.000.000.000			mg	/kg (wet weig	ht)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	3.83E-02				3.2E-03	0.0479							6.0E-06	3.2E-03	0.064	0.05
Mercury (Inorganic)				3 40F-04				1.9F-05	4 7391							6.0F-04	6 1F-04	0.9	0.0007

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.17		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.11	0.036	948	Fish	1	GKF
2	0.11	0.036	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5	0.11	0.036	948	Fish	1	PNF, SAS
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: AUF_{Lower}

COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0021	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.21E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.52E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5	1.33E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.95E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.71E-03	mg/kg ww	
Number of food types ($k = 1$ to m):		_	

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.083	8.68E-04	Table C-4.2
2	Fish	GM, STM	3	14	1	0.083	4.24E-04	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-4.2
4	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
5	Fish	PNF, SAS	3	14	1	0.083	1.60E-04	Table C-4.2
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA,6}}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
50 - 65 - 60 - 65 - 65 - 65			mg.	kg (wet weig	pht)			mg/kg _{BW} /day			mg/k	g (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	1.71E-03	0.0002			3.4E-05	0.0021							2.6E-07	3.4E-05	1.4E-04	0.24

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.57		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.607	948	Fish	1	GKF, GM, STM, PNF, SAS
2	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

2

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Hexachlorobenzene	0.6464	0.9757						2.7E-01	2.1373							0.0E+00	2.7E-01	3.35	0.08
1,3-Dichlorobenzene	0.769	0.758						3.0E-01	0.0736							0.0E+00	3.0E-01	160.8	0.0019

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.57		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.445	0.304	948	Fish	1	GM
2	0.445	0.304	948	Fish	1	GKF, STM, PNF, SAS
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
8. 80. 80. 80. 80. 80.			mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515					3.7E-02	0.0479							0.0E+00	3.7E-02	0.064	0.58
Mercury (Inorganic)							-	0.0E+00	4.7391				_			0.0E+00	0.0E+00	0.9	0

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTS}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.57		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.3	0.205	948	Fish	1	GKF
2	0.3	0.205	948	Fish	1	GM, STM
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.29	0.205	948	Fish	1	PNF, SAS
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	C _{SED,6}	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	0.0002				2.0E-04	0.0021							0.0E+00	2.0E-04	0.00014	1.40

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.019		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GM
2	0.116	0.010	1092	Fish	1	GKF, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: **AUF**_{Lower}

COPC: **Methyl Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.81E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	7.25E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.29E-02	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant				1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams dw - dry weight ww - wet weight

Scenario: AUF_{Lower}

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C_SED	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1		GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Raccoon

Receptor Group: Scenario: **Omnivorous Mammal**

AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{BIOTA,3}$	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)						mg/kg _{BW} /day	mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	0.002426492	3.83E-02			7.0E-05	0.0479							8.0E-06	7.8E-05	0.0177	0.0044
Mercury (Inorganic)				0.065754378	3.40E-04			3.4E-05	4.7391							7.9E-04	8.3E-04	3.54	0.00023

Baseline Ecological Risk Assessment March 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.019		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GKF
2	0.116	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Scenario: AUF_{Lower}

COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.09E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.99E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	9.75E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	1.91E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.12E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.125	8.70E-05	Table C-4.2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.125	3.19E-05	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	2.79E-05	Table C-4.2
4	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor:

Raccoon

Receptor Group:

Omnivorous Mammal

Scenario:

AUFLower

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{\text{SED,5}}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	ma/ka (wet welant)						mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA		
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	3.2E-05	2.8E-05	2.32E-07	1.12E-04			5.8E-08	0.0001							1.9E-08	7.6E-08	5.4E-07	0.14

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: AUF_{Lower}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.26		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

	Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
	1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
I	2						
	3						
	4						
	5						
	6						
	7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: AUFLower

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{BIOTA,3}$	$C_{BIOTA,4}$	C _{BIOTA,5}	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)						mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA	
Lead	0.292							7.5E-02	48.944							2.6E-02	1.0E-01	3.26	0.031
Mercury (Methyl)	0.0533							1.4E-02	0.0479							2.6E-05	1.4E-02	0.064	0.22
Total PAH 16 (U=1/2)	0.1397							3.6E-02	7.3078							3.9E-03	4.0E-02	325	0.00012
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							5.7E-06	0.0002							1.2E-07	5.9E-06	0.00014	0.042
Hexachlorobenzene	1.132							2.9E-01	2.1373							1.2E-03	2.9E-01	3.35	0.09
Hexachlorobutadiene	0.3725							9.6E-02	3.8327							2.1E-03	9.8E-02	20	0.0049
1,3-Dichlorobenzene	0.3725							9.6E-02	0.0736							4.0E-05	9.6E-02	160.8	0.00060
1,4-Dichlorobenzene	0.3725							9.6E-02	0.0744							4.0E-05	9.6E-02	160.8	0.00060
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.1E-04	0.0021							1.1E-06	1.1E-04	1.4E-04	0.80
Mercury (Inorganic)								0.0E+00	4.7391							2.6E-03	2.6E-03	9.0E-01	0.0028

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.6 AUF UNCERTAINTY ANALYSIS – UPPER-BOUND ESTIMATE OF HOME RANGE

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	1		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

4

Food Type	$P_{BIOTA,k}$	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.165	0.054	948	Fish	1	GM
2	0.165	0.054	948	Fish	1	GKF, STM, PNF, SAS
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5						
6						
7						

Number of foraging subareas (j = 1 to i):

		4
		7

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: AUF_{Upper}

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C_SED	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.018125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	
Number of food types ($k = 1$ to m):		•	

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams **Prey Group: Terrestrial Mammal - Marsh Rice Rat**

Scenario: AUF_{Upper}

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter		Value	Units	Source
FCM _{TL3(OM)}		1.0		Table C-4.2
BCF _{PLANT-OM}		2.30E-03		Table C-4.2
C _{SED}		4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}		5.57E-05		Table C-4.2
FR _{SED}		1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Typ	e 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Typ		0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Typ	e 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant		7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sedimen	2.64E-04	mg/kg ww		
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww		
Neverbon of food to part of the second	4			·

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: AUFUpper

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{\text{BIOTA,4}}$	C _{BIOTA,5}	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
mg/kg (wet weight)					mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA				
Mercury (Methyl)	0.145	0.058	0.0515	3.83E-02				1.9E-02	0.0479							3.5E-05	1.9E-02	0.064	0.29
Mercury (Inorganic)				3.40E-04				1.9E-05	4.7391							3.5E-03	3.5E-03	0.9	0.0039

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	1		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.11	0.036	948	Fish	1	GKF
2	0.11	0.036	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5	0.11	0.036	948	Fish	1	PNF, SAS
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: AUF_{Upper}

COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0	-	Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0021	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.21E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.52E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5	1.33E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.95E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.71E-03	mg/kg ww	
Number of food types ($k = 1$ to m): 5			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.083	8.68E-04	Table C-4.2
2	Fish	GM, STM	3	14	1	0.083	4.24E-04	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-4.2
4	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
5	Fish	PNF, SAS	3	14	1	0.083	1.60E-04	Table C-4.2
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: AUFUpper

COPC	C _{BIOTA,1}	$C_{BIOTA,2}$	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA},6}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)					mg/kg _{BW} /day			mg/k	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA		
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	1.71E-03	0.0002			2.0E-04	0.0021							1.5E-06	2.0E-04	1.4E-04	1.42

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	1		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.607	948	Fish	1	GKF, GM, STM, PNF, SAS
2	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

2

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: AUFUpper

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	C _{BIOTA,5}	$C_{BIOTA,6}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA
Hexachlorobenzene	0.6464	0.9757						4.7E-01	2.1373							0.0E+00	4.7E-01	3.35	0.14
1,3-Dichlorobenzene	0.769	0.758						5.2E-01	0.0736							0.0E+00	5.2E-01	160.8	0.0033

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME _{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTS}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	1		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.445	0.304	948	Fish	1	GM
2	0.445	0.304	948	Fish	1	GKF, STM, PNF, SAS
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram mg - milligrams kcal - kilocalories kjoule - kilojoules Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: AUFUpper

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	$C_{BIOTA,7}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	C _{SED,6}	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
5. 50. 50. 50. 50. 50. 50.			mg/l	kg (wet wei	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{lw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515					6.5E-02	0.0479							0.0E+00	6.5E-02	0.064	1.02
Mercury (Inorganic)								0.0E+00	4.7391							0.0E+00	0.0E+00	0.9	0

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	1		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.3	0.205	948	Fish	1	GKF
2	0.3	0.205	948	Fish	1	GM, STM
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.29	0.205	948	Fish	1	PNF, SAS
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: AUFUpper

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	$C_{BIOTA,4}$	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	C _{SED,3}	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)						mg/kg _{BW} /day	g/kg _{BW} /day mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA		
PCB Congener TEQ (Avian) (U=1/2)	0.0009	0.0005	0.0005	0.0002				3.4E-04	0.0021	,	·					0.0E+00	3.4E-04	0.00014	2.45

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.023		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GM
2	0.116	0.010	1092	Fish	1	GKF, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas (j = 1 to i):

		<u> </u>
Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: **Terrestrial Mammal - Marsh Rice Rat**

Scenario: AUF_{Upper}

COPC: **Methyl Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0479	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.81E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	7.25E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.29E-02	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.99E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant				1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams dw - dry weight ww - wet weight

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: AUF_{Upper}

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	4.7391	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.64E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.40E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Raccoon

Receptor Group: Scenario: **Omnivorous Mammal**

AUFUpper

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{BIOTA,3}$	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
				mg/kg (wet weight)				mg/kg _{BW} /day			mg/k	g (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_w /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	0.002426492	3.83E-02			8.5E-05	0.0479							9.7E-06	9.5E-05	0.0177	0.0053
Mercury (Inorganic)				0.065754378	3.40E-04			4.1E-05	4.7391							9.6E-04	1.0E-03	3.54	0.00028

Baseline Ecological Risk Assessment March 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Raccoon

Receptor Group: Omnivorous Mammal

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.023		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GKF
2	0.116	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: AUF_{Upper}

COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.09E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.99E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	9.75E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	1.91E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.12E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.125	8.70E-05	Table C-4.2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.125	3.19E-05	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	2.79E-05	Table C-4.2
4	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor:

Raccoon

Receptor Group:

Omnivorous Mammal

Scenario:

AUFUpper

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA},3}$	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{\text{SED,7}}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			m	g/kg (wet we	eight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg į	_{sw} /day	NA
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	3.2E-05	2.8E-05	2.32E-07	1.12E-04			7.0E-08	0.0001							2.3E-08	9.2E-08	5.4E-07	0.17

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: AUF_{Upper}

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	1		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor Group: Sediment-probing Birds

Scenario: AUFUpper

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet we	ight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Lead	0.292							2.9E-01	48.944							1.0E-01	3.9E-01	3.26	0.12
Mercury (Methyl)	0.0533							5.3E-02	0.0479							9.9E-05	5.3E-02	0.064	0.83
Total PAH 16 (U=1/2)	0.1397							1.4E-01	7.3078							1.5E-02	1.5E-01	325	0.00047
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							2.2E-05	0.0002							4.5E-07	2.3E-05	0.00014	0.16
Hexachlorobenzene	1.132							1.1E+00	2.1373							4.4E-03	1.1E+00	3.35	0.34
Hexachlorobutadiene	0.3725							3.7E-01	3.8327							7.9E-03	3.8E-01	20	0.019
1,3-Dichlorobenzene	0.3725							3.7E-01	0.0736							1.5E-04	3.7E-01	160.8	0.0023
1,4-Dichlorobenzene	0.3725							3.7E-01	0.0744							1.5E-04	3.7E-01	160.8	0.0023
PCB Congener TEQ (Avian) (U=1/2)	0.00043					,		4.3E-04	0.0021			_				4.3E-06	4.3E-04	1.4E-04	3.08
Mercury (Inorganic)								0.0E+00	4.7391							9.8E-03	9.8E-03	9.0E-01	0.011

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.7 PCB RISK UNCERTAINTY ANALYSIS – EXPOSURE AND EFFECTS BASED ON TOTAL PCB CONGENERS

Plant Total PCB Congeners Calculations

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

Where:

 C_{PLANT} = Concentration of PCB Congener TEQ (Mammal) in plant tissue

C_{SED} = Concentration of PCB Congener TEQ (Mammal) in sediment

BAF_{PLANT} = Bioaccumulation factor for PCB Congener TEQ (Mammal) (kgSED / kgPREY)

$$C_{PLANT, \ WW} = C_{PLANT,DW} \times \left(\frac{(100 - \% \ moisture)}{100}\right)$$

Where:

C_{PLANT.WW} = Concentration in plant tissue as wet weight

C_{PLANT.DW} = Concentration in plant tissue as dry weight

% moisture = Moisture content of plant tissue

	C _{SED} (mg/kg)	BAF _{PLANT*}	C _{PLANT,DW} (mg/kg)	C _{PLANT,WW} (mg/kg)
Total PCB Congeners	7.87E+00	0.01	7.87E-02	2.91E-02

^{*}USEPA (1999) Table C-2, BCF for Arcolor 1254

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds
Scenario: PCB Uncertainty

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.165	0.054	948	Fish	1	GKF
2	0.165	0.054	948	Fish	1	GM, STM, PNF, SAS
3	0.165	0.054	824	Invertebrate	1	BRS
4	0.165	0.054	824	Invertebrate	1	BCR, WHS, OYS
5	0.34	0.111	1326	Mammal	0.5	
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: PCB Uncertainty
COPC: Total PCB Congeners

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	12		USEPA (1999) Table 5-2 for log Kow of 6.2 (Aroclor 1254)
BCF _{PLANT-OM}	1.78E-02		USEPA (1999) Table D-1 for Aroclor 1254
C _{SED}	7.8689	mg/kg dw	Risk Characterization
BCF _{SED-OM}	7.42E-05		USEPA (1999) Table D-3 for Aroclor 1254
FR _{SED}	1.0		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.19E-01	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 1	3.60E-01	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	3.02E+00	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	1.22E+00	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	1.81E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	5.84E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	5.32E+00	mg/kg ww	
Number of food types ($k = 1$ to m):			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	12	1	0.125	5.75E+00	Risk Characterization
2	Fish	GM, STM, PNF, SAS	3	12	1	0.125	2.88E+00	Risk Characterization
3	Invertebrate	BRS	2	1	1	0.125	2.01E+00	Risk Characterization
4	Invertebrate	BCR, WHS, OYS	2	1	1	0.125	8.13E-01	Risk Characterization
5	Plant	Plant			1	0.5	2.91E-02	Plant Total PCB Congeners Calculations
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Receptor:CompositeReceptor Group:Carnivorous BirdsScenario:PCB Uncertainty

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)						mg/kg _{BW} /day	day mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg į	_{sw} /day	NA
Total PCB Congener (U=1/2)	5.7523	5.7523 2.8799 2.013 0.8135 5.32E+00					3.3E-01	7.8689							2.1E-03	3.3E-01	2.05	0.16	

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: PCB Uncertainty

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.445	0.304	948	Fish	1	GKF
2	0.445	0.304	948	Fish	1	GM, STM, PNF, SAS
3	0.055	0.038	824	Invertebrate	1	BRS
4	0.055	0.038	824	Invertebrate	1	BCR, WHS, OYS
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:

		<u> </u>
Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

4

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher
Receptor Group: Piscivorous Birds
Scenario: PCB Uncertainty

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	$C_{\text{BIOTA,6}}$	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	C _{SED,3}	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
50 - 50 - 50 - 50 - 50 - 50		mg/kg (wet weight)						mg/kg _{BW} /day	mg/kg _{BW} /day mg/kg (dry weight)							mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA
Total PCB Congener (U=1/2)	5.7523	5.7523 2.8799 2.013 0.8135					1.9E+00	7.8689							0.0E+00	1.9E+00	2.05	0.92	

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

March 2013
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: PCB Uncertainty

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME _{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GKF
2	0.116	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.117	0.011	974	Invertebrate	1	BRS
4	0.117	0.011	974	Invertebrate	1	BCR, WHS, OYS
5	0.3	0.027	962	Plant	1	
6	0.234	0.021	1428	Mammal	0.5	
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1
---	---

		<u> </u>
Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: PCB Uncertainty
COPC: Total PCB Congeners

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	12		USEPA (1999) Table 5-2 for log Kow of 6.2 (Aroclor 1254)
BCF _{PLANT-OM}	1.78E-02		USEPA (1999) Table D-1 for Aroclor 1254
C _{SED}	7.8689	mg/kg dw	Risk Characterization
BCF _{SED-OM}	7.42E-05		USEPA (1999) Table D-3 for Aroclor 1254
FR _{SED}	1.0		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.19E-01	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.60E-01	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	3.02E+00	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 4	1.22E+00	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	1.81E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	5.84E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	5.32E+00	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	12	1	0.125	5.75E+00	Risk Characterization
2	Fish	GM, STM, PNF, SAS	3	12	1	0.125	2.88E+00	Risk Characterization
3	Invertebrate	BRS	2	1	1	0.125	2.01E+00	Risk Characterization
4	Invertebrate	BCR, WHS, OYS	2	1	1	0.125	8.13E-01	Risk Characterization
5	Plant	Plant			1	0.5	2.91E-02	Plant Total PCB Congeners Calculations
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Raccoon

Receptor: Receptor Group: **Omnivorous Mammal** Scenario: **PCB Uncertainty**

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	$C_{\text{BIOTA,5}}$	$C_{\text{BIOTA},6}$	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			m	g/kg (wet w	eight)			mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Total PCB Congener (U=1/2)	5.75228	2.87986	2.013	0.81347	2.91E-02	5.32E+00		3.7E-03	7.8689							1.5E-03	5.2E-03	0.106	0.049

Baseline Ecological Risk Assessment March 2013 Patrick Bayou Superfund Site 040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds

Scenario: PCB Uncertainty

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Scenario: PCB Uncertainty

COPC	$C_{\text{BIOTA},1}$	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet we	ght)			mg/kg _{BW} /day			mg/k	kg (dry wei	ight)			mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA
Total PCB Congener (U=1/2)	0.82085							2.8E-01	7.8689							5.5E-03	2.8E-01	2.05	0.14

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX C-5.8 SWACS CALCULATED BY EXCLUDING GUNITE PORTION OF PATRICK BAYOU

Table C-5.8
Comparison of Baseline SWAC and Excluding Gunite Channel SWAC

		Intertidal and	Supratidal SWAC
COPC	Units	Whole Bayou (Baseline)	Excluding Gunite Channel
1,3-Dichlorobenzene	mg/kg	0.074	0.088
1,4-Dichlorobenzene	mg/kg	0.074	0.090
Hexachlorobenzene	mg/kg	2.14	2.58
Hexachlorobutadiene	mg/kg	3.83	4.69
Lead	mg/kg	48.94	45.16
Mercury	mg/kg	4.79	5.85
Total Dioxin/Furan TEQ 1998 (Avian; U=1/2)	mg/kg	0.00022	0.00026
Total PAH (U=1/2)	mg/kg	7.31	6.20
Total PCB Congener TEQ 1998 (Avian; U=1/2)	mg/kg	0.0021	0.0023
Total PCB Congener TEQ 2005 (Mammal; U=1/2)	mg/kg	0.00011	0.00013

Notes:

SWAC - surface weighted average concentration

mg/kg - milligrams per kilogram

COPC - chemical of potential concern

PAH - polycyclic aromatic hydrocarbon

PCB - polychlorinated biphenyl

CARNIVOROUS BIRD RISK SHEETS

Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Baseline

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.165	0.054	948	Fish	1	GM
2	0.165	0.054	948	Fish	1	GKF, STM, PNF, SAS
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Excluding Gunite Channel

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0585	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0.018125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.88E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	TL _{BIOTA,k}	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant	Plant			1	0.5	2.43E-03	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams **Prey Group: Terrestrial Mammal - Marsh Rice Rat**

Scenario: **Excluding Gunite Channel**

COPC: **Inorganic Mercury**

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	5.7915	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.23E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.98E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7					•			

Definitions:

kg - kilogram

mg - milligrams

Receptor: Receptor Group: Scenario: Composite Carnivorous Birds Baseline

COPC	$C_{BIOTA,1}$	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg	/kg (wet weig	ht)			mg/kg _{BW} /day			mg/k	g (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	3.83E-02				6.7E-03	0.0585							1.6E-05	6.7E-03	0.064	0.11
Mercury (Inorganic)				3.98E-04				2.2E-05	5.7915							1.5E-03	1.6E-03	0.9	0.0017

Baseline Ecological Risk Assessment Patrick Bayou Superfund Site February 2013 040284-01 Receptor: Composite

Receptor Group: Carnivorous Birds

Scenario: Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.11	0.036	948	Fish	1	GKF
2	0.11	0.036	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	March Rice Rat
5	0.11	0.036	948	Fish	1	PNF, SAS
6						
7						

1

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

5

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Excluding Gunite Channel COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0023	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	7.21E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.52E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.20E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Food Type 5	1.33E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	5.52E-06	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.71E-03	mg/kg ww	
Number of food types ($k = 1$ to m):		<u> </u>	

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.083	8.68E-04	Table C-4.2
2	Fish	GM, STM	3	14	1	0.083	4.24E-04	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-4.2
4	Plant	Plant			1	0.5	4.35E-06	Table C-4.2
5	Fish	PNF, SAS	3	14	1	0.083	1.60E-04	Table C-4.2
6								_
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor:CompositeReceptor Group:Carnivorous BirdsScenario:Excluding Gunite Channel

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},1}$ $C_{\text{BIOTA},2}$ $C_{\text{BIOTA},3}$ $C_{\text{BIOTA},4}$ $C_{\text{BIOTA},5}$ $C_{\text{BIOTA},6}$ $C_{\text{BIOTA},7}$						TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
0.000.000.000.000.000.000	mg/kg (wet weight)					mg/kg _{BW} /day	mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{BW} /day	NA			
PCB Congener TEQ (Avian) (U=1/2)	0.00087	0.00048	0.00045	1.71E-03	0.00016			7.1E-05	0.0023							6.2E-07	7.2E-05	1.4E-04	0.51

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site

640284-01

BELTED KINGFISHER RISK SHEETS

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR _k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.89	0.607	948	Fish	1	GKF, GM, STM, PNF, SAS
2	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Receptor Group: Scenario: Belted Kingfisher Piscivorous Birds

Excluding Gunite Channel

COPC	$C_{\text{BIOTA,1}}$	CBIOTA,1 CBIOTA,2 CBIOTA,3 CBIOTA,4 CBIOTA,5 CBIOTA,6 CBIOTA,7						TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	1	mg/kg (wet weight)						mg/kg _{BW} /day	/day mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA	
Hexachlorobenzene	0.6464	0.9757						3.2E-01	2.5829							0.0E+00	3.2E-01	3.35	0.10
1,3-Dichlorobenzene	0.769	0.758		,	,	,		3.6E-01	0.0885							0.0E+00	3.6E-01	160.8	0.0022

Baseline Ecological Risk Assessment February 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.445	0.304	948	Fish	1	GM
2	0.445	0.304	948	Fish	1	GKF, STM, PNF, SAS
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Receptor Group: Scenario: Belted Kingfisher Piscivorous Birds

Excluding Gunite Channel

COPC	C _{BIOTA,1}	$_{ ext{BIOTA},1}$ $C_{ ext{BIOTA},2}$ $C_{ ext{BIOTA},3}$ $C_{ ext{BIOTA},4}$ $C_{ ext{BIOTA},5}$ $C_{ ext{BIOTA},6}$ $C_{ ext{BIOTA}}$					C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
		mg/kg (wet weight)						mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515					4.5E-02	0.0585							0.0E+00	4.5E-02	0.064	0.71
Mercury (Inorganic)								0.0E+00	5.7915							0.0E+00	0.0E+00	0.9	0

Baseline Ecological Risk Assessment February 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	136	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	637	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	934	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.682	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.024	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.69		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

4

Food Type	$P_{BIOTA,k}$	NIR _k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.3	0.205	948	Fish	1	GKF
2	0.3	0.205	948	Fish	1	GM, STM
3	0.11	0.075	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.29	0.205	948	Fish	1	PNF, SAS
5						
6						
7						

Number of foraging subareas (j = 1 to i):

1

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Receptor: Belted Kingfisher Receptor Group: Piscivorous Birds

Scenario: Excluding Gunite Channel

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	C _{BIOTA,4}	$C_{\text{BIOTA},5}$	C _{BIOTA,6}	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	$C_{SED,1}$	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
	mg/kg (wet weight)						mg/kg _{BW} /day	kg _{BW} /day mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA		
PCB Congener TEQ (Avian) (U=1/2)	0.00087	0.00048	0.00045	0.00016				2.4E-04	0.0023							0.0E+00	2.4E-04	0.00014	1.69

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

RACCOON RISK SHEETS

Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME _k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GM
2	0.116	0.010	1092	Fish	1	GKF, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: **Terrestrial Mammal - Marsh Rice Rat**

Scenario: **Excluding Gunite Channel**

COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	3.44E-04		Table C-4.2
C _{SED}	0.0585	mg/kg dw	Table C-4.2
BCF _{SED-OM}	8.34E-06		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.81E-02	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	7.25E-03	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	1.29E-02	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	4.17E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.88E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	FR _{BIOTA,k}	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125	1.45E-01	Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125	5.80E-02	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	5.15E-02	Table C-4.2
4	Plant				1	0.5	2.43E-03	Table C-4.2
5								
6								
7					•			

Definitions:

kg - kilogram mg - milligrams

dw - dry weight ww - wet weight

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Excluding Gunite Channel

COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1.0		Table C-4.2
BCF _{PLANT-OM}	2.30E-03		Table C-4.2
C _{SED}	5.7915	mg/kg dw	Table C-4.2
BCF _{SED-OM}	5.57E-05		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	7.59E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.23E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.98E-04	mg/kg ww	
Number of food types $(k-1 \text{ to } m)$:			

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GM	3	1	1	0.125		Table C-4.2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-4.2
4	Plant				1	0.5	6.60E-02	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Raccoon

Receptor: Receptor Group: Scenario: **Omnivorous Mammal Excluding Gunite Channel**

COPC	C _{BIOTA,1}	$C_{\text{BIOTA,2}}$	$C_{\text{BIOTA,3}}$	$C_{BIOTA,4}$	$C_{\text{BIOTA,5}}$	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	C _{SED,1}	C _{SED,2}	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
				mg/kg (wet weight)				mg/kg _{BW} /day			mg/l	kg (dry we	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
Mercury (Methyl)	0.145	0.058	0.0515	0.002426492	3.83E-02			7.7E-05	0.0585							1.1E-05	8.8E-05	0.0177	0.0050
Mercury (Inorganic)				0.065754378	3.98E-04			3.8E-05	5.7915							1.1E-03	1.1E-03	3.54	0.00031

Baseline Ecological Risk Assessment February 2013 Patrick Bayou Superfund Site 040284-01 Receptor: Raccoon

Receptor Group: Omnivorous Mammal Scenario: Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	3990	g	Table C-4.1	AE _{FISH}	0.91		Table C-4.1
а	6.0		Table C-4.1	AE _{INVERTs}	0.91		Table C-4.1
b	0.678		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	100	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.84		Table C-4.1
ME_{AVG}	1104	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.090	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.094	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.094		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.021		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.116	0.010	1092	Fish	1	GKF
2	0.116	0.010	1092	Fish	1	GM, STM, PNF, SAS
3	0.234	0.021	974	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.3	0.027	962	Plant	1	
5	0.234	0.021	1428	Mammal	0.5	Marsh Rice Rat
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Excluding Gunite Channel
COPC: PCB Congener TEQ (Mammal)

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	14.0		Table C-4.2
BCF _{PLANT-OM}	2.39E-02		Table C-4.2
C _{SED}	0.0001	mg/kg dw	Table C-4.2
BCF _{SED-OM}	1.70E-04		Table C-4.2
FR _{SED}	1.000		Table C-4.2
COPC concentration in Marsh Rice Rat from Food Type 1	1.09E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	3.99E-06	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	9.75E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	2.77E-09	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	2.21E-08	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	1.12E-04	mg/kg ww	

Number of food types (k = 1 to m):

Food Type	Prey Group	Description	TL _{BIOTA,k}	FCM _{TL(BIOTA,k)}	$FR_{BIOTA,k}$	$P_{BIOTA,k}$	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.125	8.70E-05	Table C-4.2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.125	3.19E-05	Table C-4.2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	2.79E-05	Table C-4.2
4	Plant	Plant			1	0.5	2.32E-07	Table C-4.2
5								
6								
7								

Definitions:

kg - kilogram mg - milligrams dw - dry weight Raccoon

Receptor: Receptor Group: Scenario: Omnivorous Mammal Excluding Gunite Channel

COPC	C _{BIOTA,1}	C _{BIOTA,2}	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	$C_{\text{BIOTA},5}$	$C_{BIOTA,6}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
50.0000.0000.0000.0000.0000.0000.0000	mg/kg (wet weight)						mg/kg _{BW} /day			mg/l	kg (dry wei	ight)			mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA	
PCB Congener TEQ (Mammal) (U=1/2)	8.7E-05	3.7E-05 3.2E-05 2.8E-05 2.32E-07 1.12E-04					6.4E-08	0.0001							2.4E-08	8.8E-08	5.4E-07	0.16	

Baseline Ecological Risk Assessment February 2013 Patrick Bayou Superfund Site 040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Excluding Gunite Channel

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR_{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories


Receptor:Spotted SandpiperReceptor Group:Sediment-probing BirdsScenario:Excluding Gunite Channel

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$C_{SED,1}$	C _{SED,2}	$C_{SED,3}$	C _{SED,4}	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	HQ
			mg/	kg (wet we	ight)			mg/kg _{BW} /day	mg/kg (dry weight)					mg/kg _{BW} /day	mg/kg į	_{sw} /day	NA		
Lead	0.292							9.9E-02	45.16							3.2E-02	1.3E-01	3.26	0.04
Mercury (Methyl)	0.0533							1.8E-02	0.0585							4.1E-05	1.8E-02	0.064	0.28
Total PAH 16 (U=1/2)	0.1397							4.7E-02	6.198							4.4E-03	5.2E-02	325	0.00016
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							7.5E-06	0.0003							1.8E-07	7.7E-06	0.00014	0.05
Hexachlorobenzene	1.132							3.8E-01	2.5829							1.8E-03	3.8E-01	3.35	0.11
Hexachlorobutadiene	0.3725							1.3E-01	4.687							3.3E-03	1.3E-01	20	0.006
1,3-Dichlorobenzene	0.3725							1.3E-01	0.0885							6.2E-05	1.3E-01	160.8	0.0008
1,4-Dichlorobenzene	0.3725							1.3E-01	0.0897							6.3E-05	1.3E-01	160.8	0.0008
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.4E-04	0.0023							1.6E-06	1.5E-04	1.4E-04	1.0
Mercury (Inorganic)								0.0E+00	5.7915							4.1E-03	4.1E-03	9.0E-01	0.005

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01


APPENDIX C-5.9 SEDIMENT HOT SPOT UNCERTAINTY ANALYSIS

FIGURES















Feet

550






Baseline Ecological Risk Assessment















Feet


550

825 1,100





























TABLES

Table C-5.9 1
Comparison of Site-wide and Hot Spot Intertidal and Supratidal Weighted Averages and HQs

		Interti	dal and Supratidal Weighted Ave	rages ^a	HQs						
СОРС	Units	Site-wide ^b	Hot Spot Area Based on the Spotted Sandpiper Minimum Home-range (1.5 km)	Hot Spot Area Based on the Carnivorous Bird Minimum Home-range (1.4 km)	Site-wide Spotted Sandpiper	Hot Spot Spotted Sandpiper	Site-wide Carnivorous Bird	Hot Spot Carnivorous Bird			
Lead	mg/kg	48.9	51.1		0.04	0.04					
Mercury (Methyl) ^c	mg/kg	0.05	0.06	0.06	0.28	0.28	0.11	0.11			
Mercury (Inorganic) ^d	mg/kg	4.74	5.77	5.80	0.004	0.005	0.0014	0.0017			
Total Mercury	mg/kg	4.79	5.83	5.86	0.28	0.29	0.11	0.11			
Total PAH	μg/kg	7310	7420		0.00016	0.00016					
1,3-Dichlorobenzene	μg/kg	73.6	109		0.0008	0.0008					
1,4-Dichlorobenzene	μg/kg	74.4	112		0.0008	0.0008					
Hexachlorobenzene	μg/kg	2140	3180		0.11	0.11					
Hexachlorobutadiene	μg/kg	3830	5820		0.006	0.006					
PCB Congener TEQ (Avian)	ng/kg	2080	2424	2451	1.0	1.0	0.76	0.76			
Dioxin/Furan TEQ (Avian)	ng/kg	108	260		0.05	0.05					

^a - Intertidal and Supratidal Weighted Average is the sediment exposure point concentration.

Notes:

COPC - Chemical of potential concern

SWAC - surface weighted average concentration

PAH - polycyclic aromatic hydrocarbon

PCB - polychlorinated biphenyl

TEQ - Toxic equivalent

mg/kg - milligram per kilogram

ng/kg - nanogram per kilogram

μg/kg - microgram per kilogram

b - From Table 6-4

^c- Based on the Total Mercury sediment concentration multiplied by 0.01

^d - Based on the Total Mercury sediment concentration multiplied 0.99

Table C-5.9 2
Hot Spot Analysis Exposure Variables for Calculating Marsh Rice Rat COPC Concentration

				Me	ercury		PCB Congener TEQ		
Variable	Definition	Units	Methyl Merc	ury	Inorganic Mer	cury	(Avian)		
Kow	Log octanol/water partition coefficient	unitless	2.54	а	log(0.61)	i	6.64	m	
FCM _{TL3(OM)}	Food chain multiplier for trophic level 3 (marsh rice rat)	unitless	1.0	b	1.0	g	14	n	
Hot_Spot_C _{SED}	Hot spot COPC concentration in sediment	mg/kg dw	5.86E-02	С	5.80E+00	С	2.45E-03	С	
BCF _{PLANT-OM}	Plant-to-omnivore BCF	unitless	3.44E-04	d	2.30E-03	k	2.39E-02	0	
BCF _{SED-OM}	Sediment-to-omnivore BCF	unitless	8.34E-06	е	5.57E-05	I	1.70E-04	р	
FR _{SED}	Fraction of sediment in diet that is contaminated	unitless	1	f	1	f	1	f	
$TL_{BIOTA,a}$	Trophic Level of Food Type 1	unitless	3	j	3	j	3	j	
$TL_{BIOTA,b}$	Trophic Level of Food Type 2	unitless	3	j	3	j	3	j	
TL _{BIOTA,c}	Trophic Level of Food Type 3	unitless	2	j	2	j	2	j	
TL _{BIOTA,e}	Trophic Level of Food Type 5	unitless					3	j	
FCM _{TL(BIOTA,a)}	Food Chain Multiplier for Food Type 1	unitless	1	b	1	g	14	n	
FCM _{TL(BIOTA,b)}	Food Chain Multiplier for Food Type 2	unitless	1	b	1	g	14	n	
FCM _{TL(BIOTA,c)}	Food Chain Multiplier for Food Type 3	unitless	1	b	1	g	1	n	
FCM _{TL(BIOTA,e)}	Food Chain Multiplier for Food Type 3	unitless					14	n	
FR _{BIOT,k} (all Food Types)	Portion of kth Food Group that is contaminated	unitless	1	f	1	f	1	f	
P _{BIOTA,a}	Portion of diet composed of Food Type 1	unitless	0.125	S	0.125	S	0.083	S	
P _{BIOTA,b}	Portion of diet composed of Food Type 2	unitless	0.125	S	0.125	S	0.083	S	
P _{BIOTA,c}	Portion of diet composed of Food Type 3	unitless	0.25	S	0.25	S	0.25	S	
P _{BIOTA,d}	Portion of diet composed of Food Type 4	unitless	0.5	S	0.5	S	0.5	S	
P _{BIOTA,e}	Portion of diet composed of Food Type 5	unitless					0.083	S	
C _{BIOTA,a}	COPC concentration in Food Type 1	mg/kg ww	1.45E-01	q			8.68E-04	q	
C _{BIOTA,b}	COPC concentration in Food Type 2	mg/kg ww	5.80E-02	q			4.24E-04	q	
C _{BIOTA,c}	COPC concentration in Food Type 3	mg/kg ww	5.15E-02	q			4.52E-04	q	
$C_{BIOTA,d}$	COPC concentration in Food Type 4	mg/kg ww	3.00E-03	r	8.10E-02	r	5.18E-06	h	
$C_{BIOTA,e}$	COPC concentration in Food Type 5	mg/kg ww					1.60E-04	q	

Notes:

- a Maximum value in methyl mercury Kow range (Environment Canada 2002)
- b USEPA (1999) Table 5-2 based on a Kow of 3
- c Table C-5.9 1
- d USEPA (1999) Table D-1 value for methyl mercury
- e USEPA (1999) Table D-3 value for methyl mercury
- f USEPA (1999) Table F-1-14 Default
- g USEPA (1999) Table 5-2 based on a Kow of 2 (smallest Kow value provided)
- h Hot Spot PCB Congener TEQ (Avian) Plant Concentration Calculation, BERA Appendix
- USEPA (1999) Table A-2-130 value for mercuric chloride
 USEPA (1999) and Professional judgment. All fish were assumed to belong to
- j Trophic level of 3 (omnivore) and invertebrates were assumed to be belong to trophic level 2 (herbivore)

- $k\;\;$ USEPA (1999) Table D-1 value for mercuric chloride
- / USEPA (1999) Table D-3 value for mercuric chloride
- m USEPA (1999) Table A-2-176
- n USEPA (1999) Table 5-2 based on a Kow of 6.6
- o USEPA (1999) Table D-1 value for 2,3,7,8-TCDD
- p USEPA (1999) Table D-3 value for 2,3,7,8-TCDD
- q Table 6-3
- r Hot Spot Mercury Plant Concentration Calculation
- s Marsh Rice Rat diet indicated in BERA Workplan (Anchor QEA 2011). Diet was assumed to be composed of 50% plant matter and 50% animal (25% invertebrates and 25% fish)

CARNIVOROUS BIRD RISK SHEETS

Hot Spot Plant Mercury Calculations

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

Where:

C_{PLANT} = Concentration of mercury in plant tissue

C_{SED} = Concentration of mercury in sediment

BAF_{PLANT} = Bioaccumulation factor for mercury (kgSED / kgPREY)

$$C_{PLANT, WW} = C_{PLANT,DW} \times \left(\frac{(100 - \% moisture)}{100}\right)$$

Where:

C_{PLANT,WW} = Concentration in plant tissue as wet weight

C_{PLANT,DW} = Concentration in plant tissue as dry weight

% moisture = Moisture content of plant tissue

	Hot_Spot_C _{SED} (mg/kg)	BAF _{PLANT}	Hot_Spot_C _{PLANT,DW} (mg/kg)	Hot_Spot_C _{PLANT,WW} (mg/kg)
Methylmercury	0.059	0.137	0.008	0.003
Inorganic Mercury	5.80	0.0375	0.218	0.081

Hot Spot Plant PCB Congener TEQ (Avian) Calculations

$$C_{PLANT} = C_{SED} \times BAF_{PLANT}$$

Where:

C_{PLANT} = Concentration of PCB Congener TEQ (Avian) in plant tissue

C_{SED} = Concentration of PCB Congener TEQ (Avian) in sediment

BAF_{PLANT} = Bioaccumulation factor for PCB (kgSED / kgPREY)

$$C_{PLANT, \ WW} = C_{PLANT,DW} imes \left(rac{(100 - \% \ moisture)}{100}
ight)$$

Where:

C_{PLANT,WW} = Concentration in plant tissue as wet weight

C_{PLANT,DW} = Concentration in plant tissue as dry weight

% moisture = Moisture content of plant tissue

	Hot_Spot_C _{SED} (mg/kg)	BAF _{PLANT*}	Hot_Spot_C _{PLANT,DW} (mg/kg)	Hot_Spot_C _{PLANT,WW} (mg/kg)		
PCB Congener TEQ (Avian)	2.50E-03	0.0056	1.40E-05	5.18E-06		

^{*}USEPA (1999) Table C-2, BCF for 2,3,7,8-TCDD

Receptor: Composite

Receptor Group: Carnivorous Birds
Scenario: Hot Spot Analysis

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	$P_{BIOTA,k}$	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.165	0.054	948	Fish	1	GM
2	0.165	0.054	948	Fish	1	GKF, STM, PNF, SAS
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	0.5	
5						
6						
7						

Number of foraging subareas $(j = 1 \text{ to } i)$:	1
1 value of loraging subareas $y = 1$ to y .	

	Subarea	$P_{SED,j}$	Description
	1	1.00	
	2		
	3		
	4		
	5		
	6		
L	7		

4

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Hot Spot Analysis COPC: Methyl Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1		Table C-5.9 2
BCF _{PLANT-OM}	3.44E-04		Table C-5.9 2
Hot_Spot_C _{SED}	0.0586	mg/kg dw	Table C-5.9 2
BCF _{SED-OM}	8.34E-06		Table C-5.9 2
FR _{SED}	1		Table C-5.9 2
COPC concentration in Marsh Rice Rat from Food Type 1	0.018125	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0.00725	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0.012875	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	5.16E-07	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	4.89E-07	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	3.83E-02	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

C_{BIOTA,k} mg/kg Description $\mathsf{FR}_{\mathsf{BIOTA},k}$ $\mathsf{P}_{\mathsf{BIOTA},k}$ Food Type Prey Group TL_{BIOTA,k} FCM_{TL(BIOTA,k)} Source ww GM 0.125 1.45E-01 Table C-5.9 2 Fish 3 GKF, STM, PNF, SAS 2 Fish 3 0.125 5.80E-02 Table C-5.9 2

BCR, BRS, WHS, OYS 3 Invertebrate 2 0.25 5.15E-02 Table C-5.9 2 4 Plant Plant 1 0.5 3.00E-03 Table C-5.9 2 5 6

Definitions:

kg - kilogram

mg - milligrams

Prey Group: Terrestrial Mammal - Marsh Rice Rat

Scenario: Hot Spot Analysis COPC: Inorganic Mercury

Model: USEPA (1999) Equation 5-12

Parameter	Value	Units	Source
FCM _{TL3(OM)}	1		Table C-5.9 2
BCF _{PLANT-OM}	2.30E-03		Table C-5.9 2
Hot_Spot_C _{SED}	5.8042	mg/kg dw	Table C-5.9 2
BCF _{SED-OM}	5.57E-05		Table C-5.9 2
FR _{SED}	1		Table C-5.9 2
COPC concentration in Marsh Rice Rat from Food Type 1	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 2	0	mg/kg ww	
COPC concentration in Marsh Rice Rat from Food Type 3	0	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Plant	9.32E-05	mg/kg ww	
COPC Concentration in Marsh Rice Rat from Sediment	3.23E-04	mg/kg ww	
Total COPC concentration in Marsh Rice Rat	4.16E-04	mg/kg ww	
Number of food types $(k = 1 \text{ to } m)$:			

Food Type	Prey Group	Description	Description TL _{BIOTA,k} FCM _{TL(BIOTA,k)} FR _{BIOTA,k} P _{BIOTA,k}		C _{BIOTA,k} mg/kg ww	Source		
1	Fish	GM	3	1	1	0.125		Table C-5.9 2
2	Fish	GKF, STM, PNF, SAS	3	1	1	0.125		Table C-5.9 2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25		Table C-5.9 2
4	Plant				1	0.5	8.10E-02	Table C-5.9 2
5								
6					•			
7								

Definitions:

kg - kilogram

mg - milligrams

Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: Hot Spot Analysis

COPC	C _{BIOTA,1}	$C_{\text{BIOTA},2}$	$C_{\text{BIOTA,3}}$	$C_{\text{BIOTA,4}}$	C _{BIOTA,5}	$C_{\text{BIOTA},6}$	C _{BIOTA,7}	TDI _{BIOTA}	Hot_Spot_C _{SED,1}	$C_{SED,2}$	$C_{SED,3}$	$C_{SED,4}$	$C_{SED,5}$	$C_{SED,6}$	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	Hot_Spot_HQ
mg/kg (wet weight)					mg/kg _{BW} /day	n _{BW} /day mg/kg (dry weight)						mg/kg _{BW} /day	mg/kg į	_{BW} /day	NA				
Mercury (Methyl)	0.145	0.058	0.0515	3.83E-02				6.7E-03	0.058629							1.6E-05	6.7E-03	0.064	0.11
Mercury (Inorganic)				4.16E-04				2.3E-05	5.804239							1.5E-03	1.6E-03	0.9	0.0017

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

Receptor: Composite

Receptor Group: Carnivorous Birds
Scenario: Hot Spot Analysis

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	870	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	14.3		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.659		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	339	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	1036	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.327	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.074	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.74		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.01		Table C-4.1	CF ₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.36		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR_k	ME_k	Prey Group	$FR_{BIOTA,k}$	Description
1	0.11	0.036	948	Fish	1	GKF
2	0.11	0.036	948	Fish	1	GM, STM
3	0.33	0.108	824	Invertebrate	1	BCR, BRS, WHS, OYS
4	0.34	0.111	1326	Mammal	1	
5	0.11	0.036	948	Fish	1	PNF, SAS
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Prey Grou, Terrestrial Mammal - Marsh Rice Rat

Scenario: Hot Spot Analysis

COPC: PCB Congener TEQ (Avian)

Model: USEPA (1999) Equation 5-12

Parameter		Value	Units	Source
FCM _{TL3(OM)}		14	-	Table C-5.9 2
BCF _{PLANT-OM}		2.39E-02		Table C-5.9 2
Hot_Spot_C _{SED}		0.0025	mg/kg dw	Table C-5.9 2
BCF _{SED-OM}		1.70E-04		Table C-5.9 2
FR _{SED}		1		Table C-5.9 2
COPC concentration in Marsh Rice Rat from	COPC concentration in Marsh Rice Rat from Food Type 1			
COPC concentration in Marsh Rice Rat from	Food Type 2	3.52E-05	mg/kg ww	
COPC concentration in Marsh Rice Rat from	Food Type 3	1.58E-03	mg/kg ww	
COPC Concentration in Marsh Rice Rat from	n Plant	6.19E-08	mg/kg ww	
COPC Concentration in Marsh Rice Rat from	1.33E-05	mg/kg ww		
COPC Concentration in Marsh Rice Rat from	5.83E-06	mg/kg ww		
Total COPC concentration in Marsh Rice I	1.71E-03	mg/kg ww		
Number of food types $(k = 1 \text{ to } m)$:	5			

Food Type	Prey Group	Description	$TL_{BIOTA,k}$	$FCM_{TL(BIOTA,k)}$	$FR_{BIOTA,k}$	P _{BIOTA,k}	C _{BIOTA,k} mg/kg ww	Source
1	Fish	GKF	3	14	1	0.083	8.68E-04	Table C-5.9 2
2	Fish	GM, STM, PNF, SAS	3	14	1	0.083	4.24E-04	Table C-5.9 2
3	Invertebrate	BCR, BRS, WHS, OYS	2	1	1	0.25	4.52E-04	Table C-5.9 2
4	Plant	Plant			1	0.5	5.18E-06	Table C-5.9 2
5	Fish	PNF, SAS	3	14	1	0.083	1.60E-04	Table C-5.9 2
6	·							
7								

Definitions:

kg - kilogram

mg - milligrams

dw - dry weight

ww - wet weight

Receptor: Composite
Receptor Group: Carnivorous Birds
Scenario: Hot Spot Analysis

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	C _{BIOTA,5}	C _{BIOTA,6}	C _{BIOTA,7}	TDI _{BIOTA}	$Hot_Spot_C_{SED,1}$	C _{SED,2}	C _{SED,3}	C _{SED,4}	$C_{SED,5}$	C _{SED,6}	C _{SED,7}	TDI _{SEDIMENT}	TDI _{ALL}	TRV	Hot_Spot_HQ
			mg	/kg (wet weig	ght)			mg/kg _{BW} /day			mg/kg (c	lry weight)				mg/kg _{BW} /day	mg/kg _E	_{sw} /day	NA
PCB Congener TEQ (Avian) (U=1/2)	0.00087	0.00048	0.00045	1.71E-03	0.00016			1.1E-04	0.002451							6.5E-07	1.1E-04	1.4E-04	0.76

Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

SPOTTED SANDPIPER RISK SHEETS

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Scenario: Hot Spot Analysis

Model Parameters

Parameter	Value	Units	Source	Parameter	Value	Units	Source
BW	43	g	Table C-4.1	AE _{FISH}	0.79		Table C-4.1
а	8.1		Table C-4.1	AE _{INVERTs}	0.77		Table C-4.1
b	0.77		Table C-4.1	AE _{PLANTS}	0.74		Table C-4.1
NFMR	818	kcal/kg _{BW} /day	Equation 7 (Exhibit C-4.1)	AE _{MAMMALS}	0.78		Table C-4.1
ME_{AVG}	824	kcal/kg	Equation 4 (Exhibit C-4.1)	GE _{FISH}	1200	kcal/kg _{FISH}	Table C-4.1
NIR _{TOTAL}	0.992	kg/kg _{BW} /day	Equation 3 (Exhibit C-4.1)	GE _{INVERTs}	1070	kcal/kg _{INVERTS}	Table C-4.1
IR _{TOTAL}	0.012	kg _{food} /day	Equation 9 (Exhibit C-4.1)	GE _{PLANTS}	1300	kcal/kg _{PLANTS}	Table C-4.1
P _{MOISTURE}	0.73		Table C-4.1	GE _{MAMMALS}	1700	kcal/kg _{MAMMALS}	Table C-4.1
FS	0.18		Table C-4.1	CF₁	0.239	kcal/kjoule	Equation 7 (Exhibit C-4.1)
AUF	0.34		Table C-4.1	CF ₂	0.001	kg/g	Equation 7 (Exhibit C-4.1)

Number of food types (k = 1 to m):

Food Type	P _{BIOTA,k}	NIR _k	ME _k	Prey Group	FR _{BIOTA,k}	Description
1	1	0.992	824	Invert	1	OYS,BCRA,WHSA
2						
3						
4						
5						
6						
7						

Number of foraging subareas (j = 1 to i):

Subarea	$P_{SED,j}$	Description
1	1.00	
2		
3		
4		
5		
6		
7		

Definitions:

kg - kilogram

mg - milligrams

kcal - kilocalories

kjoule - kilojoules

Receptor: Spotted Sandpiper
Receptor Group: Sediment-probing Birds
Scenario: Hot Spot Analysis

COPC	C _{BIOTA,1}	C _{BIOTA,2}	C _{BIOTA,3}	C _{BIOTA,4}	C _{BIOTA,5}	$C_{\text{BIOTA},6}$	$C_{\text{BIOTA,7}}$	TDI _{BIOTA}	Hot_Spot_C _{SED,1}	C _{SED,2}	$C_{SED,3}$	C _{SED,4}	C _{SED,5}	C _{SED,6}	$C_{SED,7}$	TDI _{SEDIMENT}	TDI _{ALL}	TRV	Hot_Spot_HQ
800.000.000.000.000.000.000.000.000			mg/	kg (wet we	ight)			mg/kg _{BW} /day			mg/kg (d	ry weight))			mg/kg _{BW} /day	mg/kg	_{BW} /day	NA
Lead	0.292							9.9E-02	51.082824							3.6E-02	1.3E-01	3.26	0.04
Mercury (Methyl)	0.0533							1.8E-02	0.058257							4.1E-05	1.8E-02	0.064	0.28
Total PAH 16 (U=1/2)	0.1397							4.7E-02	7.420385							5.2E-03	5.2E-02	325	0.00016
Dioxin/Furan Congener TEQ (Avian) (U=1/2)	2.2E-05							7.5E-06	0.000259718							1.8E-07	7.7E-06	0.00014	0.05
Hexachlorobenzene	1.132							3.8E-01	3.179603							2.2E-03	3.8E-01	3.35	0.11
Hexachlorobutadiene	0.3725							1.3E-01	5.820095							4.1E-03	1.3E-01	20	0.006
1,3-Dichlorobenzene	0.3725							1.3E-01	0.109273							7.7E-05	1.3E-01	160.8	0.0008
1,4-Dichlorobenzene	0.3725							1.3E-01	0.111573							7.9E-05	1.3E-01	160.8	0.0008
PCB Congener TEQ (Avian) (U=1/2)	0.00043							1.4E-04	0.002424							1.7E-06	1.5E-04	1.4E-04	1.0
Mercury (Inorganic)								0.0E+00	5.767408							4.1E-03	4.1E-03	9.0E-01	0.005


Baseline Ecological Risk Assessment
Patrick Bayou Superfund Site
040284-01

APPENDIX D FISH RISK ASSESSMENT SUPPORTING INFORMATION

APPENDIX D-1.1 PREY GROUP ANALYSIS

Dioxin/Furan TEQ

Oneway Analysis of LN Adj Result By Common_Name chemical_name=Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2), Size_Class_Code=F01



Oneway Anova Summary of Fit

Rsquare	0.629744
Adj Rsquare	0.596833
Root Mean Square Error	0.372283
Mean of Response	2.050219
Observations (or Sum Wgts)	50

Analysis of Variance

DF	Sum of Squares	Mean Square	F Ratio	Prob > F
4	10.607699	2.65192	19.1344	<.0001*
45	6.236757	0.13859		
49	16.844456			
	4 45	4 10.607699 45 6.236757	4 10.607699 2.65192 45 6.236757 0.13859	4 10.607699 2.65192 19.1344 45 6.236757 0.13859

APPENDIX D-1.2 FISH TISSUE EPC CALCULATION

SELENIUM

Gulf Killifish, Gulf Menhaden, Striped Mullet, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF
Confidence Coefficient 95%
Number of Bootstrap Operations 2000

result_value (selenium_d_ug/l)

		General Statistics	
Number of Valid Data	14	Number of Detected Data	8
Number of Distinct Detected Data	6	Number of Non-Detect Data	6
		Percent Non-Detects	42.86%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	3.8	Minimum Detected	1.335
Maximum Detected	15	Maximum Detected	2.708
Mean of Detected	11.1	Mean of Detected	2.335
SD of Detected	3.699	SD of Detected	0.45
Minimum Non-Detect	5	Minimum Non-Detect	1.609
Maximum Non-Detect	5	Maximum Non-Detect	1.609

Warning: There are only 8 Detected Values in this data

Note: It should be noted that even though bootstrap may be performed on this data set
the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

UCL Statistics

Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only		
Shapiro Wilk Test Statistic	0.895	Shapiro Wilk Test Statistic	0.784	
5% Shapiro Wilk Critical Value	0.818	5% Shapiro Wilk Critical Value	0.818	
Data appear Normal at 5% Significance Level		Data not Lognormal at 5% Significance Level		
Assuming Normal Distribution		Assuming Lognormal Distribution		
DL/2 Substitution Method		DL/2 Substitution Method		
Mean	7.414	Mean	1.727	
SD	5.184	SD	8.0	
95% DL/2 (t) UCL	9.868	95% H-Stat (DL/2) UCL	13.43	
Maximum Likelihood Estimate(MLE) Method		Log ROS Method		
Mean	12.15	Mean in Log Scale	1.986	
SD	2.231	SD in Log Scale	0.564	
95% MLE (t) UCL	13.2	Mean in Original Scale	8.371	
95% MLE (Tiku) UCL	13.64	SD in Original Scale	4.328	
		95% t UCL	10.42	
		95% Percentile Bootstrap UCL	10.21	
		95% BCA Bootstrap UCL	10.13	
		95% H UCL	11.89	
Gamma Distribution Test with Detected Values O	•	Data Distribution Test with Detected Values Only		
k star (bias corrected)	4.538	Data appear Normal at 5% Significance Level		
k star (bias corrected) Theta Star	2.446	Data appear Normal at 5% Significance Level		
,		Data appear Normal at 5% Significance Level		
Theta Star	2.446	Data appear Normal at 5% Significance Level Nonparametric Statistics		
Theta Star nu star	2.446 72.61 0.649 0.717	,		
Theta Star nu star A-D Test Statistic	2.446 72.61 0.649 0.717 0.717	Nonparametric Statistics Kaplan-Meier (KM) Method Mean	7.971	
Theta Star nu star A-D Test Statistic 5% A-D Critical Value	2.446 72.61 0.649 0.717	Nonparametric Statistics Kaplan-Meier (KM) Method	7.971 4.46	
Theta Star nu star A-D Test Statistic 5% A-D Critical Value K-S Test Statistic	2.446 72.61 0.649 0.717 0.717 0.295	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean	4.46 1.274	
Theta Star nu star A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value	2.446 72.61 0.649 0.717 0.717 0.295	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD	4.46 1.274 10.23	
Theta Star nu star A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Data appear Gamma Distributed at 5% Significance Assuming Gamma Distribution	2.446 72.61 0.649 0.717 0.717 0.295	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL	4.46 1.274 10.23 10.07	
A-D Test Statistic A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Data appear Gamma Distributed at 5% Significance Assuming Gamma Distribution Gamma ROS Statistics using Extrapolated Data	2.446 72.61 0.649 0.717 0.717 0.295 Level	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL 95% KM (jackknife) UCL	4.46 1.274 10.23 10.07 10.48	
Theta Star nu star A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Data appear Gamma Distributed at 5% Significance Assuming Gamma Distribution	2.446 72.61 0.649 0.717 0.717 0.295 Level	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL 95% KM (jackknife) UCL 95% KM (bootstrap t) UCL	4.46 1.274 10.23 10.07 10.48 10.09	
A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Compared to the statistic of the statis	2.446 72.61 0.649 0.717 0.717 0.295 Level 1E-06 15	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL 95% KM (jackknife) UCL 95% KM (bootstrap t) UCL 95% KM (BCA) UCL	4.46 1.274 10.23 10.07 10.48 10.09 12.43	
A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Data appear Gamma Distributed at 5% Significance Assuming Gamma Distribution Gamma ROS Statistics using Extrapolated Data Minimum Maximum Maximum Mean	2.446 72.61 0.649 0.717 0.717 0.295 Level 1E-06 15 7.722	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL 95% KM (jackknife) UCL 95% KM (bootstrap t) UCL 95% KM (BCA) UCL 95% KM (Percentile Bootstrap) UCL	4.46 1.274 10.23 10.07 10.48 10.09 12.43	
A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Data appear Gamma Distributed at 5% Significance Assuming Gamma Distribution Gamma ROS Statistics using Extrapolated Data Minimum Maximum Mean Median	2.446 72.61 0.649 0.717 0.717 0.295 Level 1E-06 15 7.722 7.598	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL 95% KM (jackknife) UCL 95% KM (bootstrap t) UCL 95% KM (Percentile Bootstrap) UCL 95% KM (Percentile Sootstrap) UCL	4.46 1.274 10.23 10.07 10.48 10.09 12.43 12 13.53	
A-D Test Statistic 5% A-D Critical Value K-S Test Statistic 5% K-S Critical Value Data appear Gamma Distributed at 5% Significance Assuming Gamma Distribution Gamma ROS Statistics using Extrapolated Data Minimum Maximum Maximum Mean	2.446 72.61 0.649 0.717 0.717 0.295 Level 1E-06 15 7.722	Nonparametric Statistics Kaplan-Meier (KM) Method Mean SD SE of Mean 95% KM (t) UCL 95% KM (z) UCL 95% KM (jackknife) UCL 95% KM (bootstrap t) UCL 95% KM (BCA) UCL 95% KM (Percentile Bootstrap) UCL	4.46 1.274 10.23 10.07 10.48 10.09 12.43	

Gulf Killifish, Gulf Menhaden, Striped Mullet, Pinfish, Sand Seatrout

Theta star	26.92		
Nu star	8.033	Potential UCLs to Use	
AppChi2	2.754	95% KM (t) UCL	10.23
95% Gamma Approximate UCL (Use when n >= 40)	22.52	95% KM (Percentile Bootstrap) UCL	12
95% Adjusted Gamma UCL (Use when n < 40)	26.28		
Note: DL/2 is not a recommended method.			

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

result_value (selenium_t_ug/l)

		General Statistics	
Number of Valid Data	14	Number of Detected Data	8
Number of Distinct Detected Data	7	Number of Non-Detect Data	6
		Percent Non-Detects	42.86%
Raw Statistics		Log-transformed Statistics	
Minimum Detected	4.5	Minimum Detected	1.504
Maximum Detected	16	Maximum Detected	2.773
Mean of Detected	10.94	Mean of Detected	2.331
SD of Detected	3.669	SD of Detected	0.4
Minimum Non-Detect	5	Minimum Non-Detect	1.609
Maximum Non-Detect	5	Maximum Non-Detect	1.609

Warning: There are only 8 Detected Values in this data

Note: It should be noted that even though bootstrap may be performed on this data set the resulting calculations may not be reliable enough to draw conclusions

It is recommended to have 10-15 or more distinct observations for accurate and meaningful results.

		UCL Statistics	
Normal Distribution Test with Detected Values Only		Lognormal Distribution Test with Detected Values Only	
Shapiro Wilk Test Statistic	0.963	Shapiro Wilk Test Statistic	0.892
5% Shapiro Wilk Critical Value	0.818	5% Shapiro Wilk Critical Value	0.818
Data appear Normal at 5% Significance Level		Data appear Lognormal at 5% Significance Level	
Assuming Normal Distribution		Assuming Lognormal Distribution	
DL/2 Substitution Method		DL/2 Substitution Method	
Mean	7.321	Mean	1.725
SD	5.101	SD	0.783
95% DL/2 (t) UCL	9.736	95% H-Stat (DL/2) UCL	13
Maximum Likelihood Estimate(MLE) Method		Log ROS Method	
Mean	11.88	Mean in Log Scale	2.006
SD	2.587	SD in Log Scale	0.516
95% MLE (t) UCL	13.11	Mean in Original Scale	8.381
95% MLE (Tiku) UCL	13.61	SD in Original Scale	4.156
		95% t UCL	10.35
		95% Percentile Bootstrap UCL	10.33
		95% BCA Bootstrap UCL	10.35
		95% H UCL	11.35
Gamma Distribution Test with Detected Values O	nly	Data Distribution Test with Detected Values Only	
k star (bias corrected)	5.286	Data appear Normal at 5% Significance Level	
Theta Star	2.069		
nu star	84.57		
A-D Test Statistic	0.34	Nonparametric Statistics	
5% A-D Critical Value	0.716	Kaplan-Meier (KM) Method	
K-S Test Statistic	0.716	Mean	8.179
5% K-S Critical Value	0.295	SD	4.108
Data appear Gamma Distributed at 5% Significance	Level	SE of Mean	1.174
		95% KM (t) UCL	10.26
Assuming Gamma Distribution		95% KM (z) UCL	10.11
Gamma ROS Statistics using Extrapolated Data		95% KM (jackknife) UCL	10.39
Minimum	1E-06	95% KM (bootstrap t) UCL	10.4
Maximum	16	95% KM (BCA) UCL	12.07

Gulf Killifish, Gulf Menhaden, Striped Mullet, Pinfish, Sand Seatrout

Mean	7.661	95% KM (Percentile Bootstrap) UCL	11.64
Median	7.609	95% KM (Chebyshev) UCL	13.3
SD	5.111	97.5% KM (Chebyshev) UCL	15.51
k star	0.393	99% KM (Chebyshev) UCL	19.86
Theta star	19.5		
Nu star	11	Potential UCLs to Use	
AppChi2	4.578	95% KM (t) UCL	10.26
95% Gamma Approximate UCL (Use when n >= 40)	18.41	95% KM (Percentile Bootstrap) UCL	11.64
95% Adjusted Gamma UCL (Use when n < 40)	20.85		

Note: DL/2 is not a recommended method.

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Maichle, and Lee (2006).

For additional insight, the user may want to consult a statistician.

MERCURY

Gulf Killifish

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF

Confidence Coefficient 95% Number of Bootstrap Operations 2000

Result Value (mercury mg/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 21

Raw Statistics Log-transformed Statistics

Minimum 0.025 Minimum of Log Data -3.689

Maximum 0.109 Maximum of Log Data -2.216

Mean 0.0524 Mean of log Data -3.042

netric Mean 0.0477 SD of log Data 0.43

Geometric Mean 0.0477

Median 0.044

SD 0.0242

Std. Error of Mean 0.0049 Coefficient of Variation 0.463 Skewness 1.097

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.871 Shapiro Wilk Test Statistic 0.944 Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data not Normal at 5% Significance Level

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 0.0607
 95% H-UCL 0.0619

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 0.0723

 95% Adjusted-CLT UCL (Chen-1995) 0.0615
 97.5% Chebyshev (MVUE) UCL 0.0811

 95% Modified-t UCL (Johnson-1978) 0.0608
 99% Chebyshev (MVUE) UCL 0.0982

Gamma Distribution Test Data Distribution

k star (bias corrected) 4.929 Data appear Gamma Distributed at 5% Significance Level

Theta Star 0.0106 MLE of Mean 0.0524

MLE of Standard Deviation 0.0236

nu star 246.5

Approximate Chi Square Value (.05) 211.1 Nonparametric Statistics

Adjusted Level of Significance 0.0395 95% CLT UCL 0.0603

Adjusted Chi Square Value 208.9 95% Jackknife UCL 0.0607

95% Standard Bootstrap UCL 0.0602

Anderson-Darling Test Statistic 0.639 95% Bootstrap-t UCL 0.0625 Anderson-Darling 5% Critical Value 0.747 95% Hall's Bootstrap UCL 0.0612

Gulf Killifish

Kolmogorov-Smirnov Test Statistic 0.162 Kolmogorov-Smirnov 5% Critical Value 0.175

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0611 95% Adjusted Gamma UCL (Use when n < 40) 0.0618

Potential UCL to Use

95% Percentile Bootstrap UCL 0.0603 95% BCA Bootstrap UCL 0.0618 95% Chebyshev(Mean, Sd) UCL 0.0735 97.5% Chebyshev(Mean, Sd) UCL 0.0826 99% Chebyshev(Mean, Sd) UCL 0.101

Use 95% Approximate Gamma UCL 0.0611

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Gulf Killifish, Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_a.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result Value (mercury mg/kg)

General Statistics

Number of Valid Observations 50 Number of Distinct Observations 37

Raw Statistics Log-transformed Statistics

Minimum 0.024 Minimum of Log Data -3.73

Maximum 0.199 Maximum of Log Data -1.614

Mean 0.0643 Mean of log Data -2.904

Geometric Mean 0.0548 SD of log Data 0.551

Median 0.0465 SD 0.0406

Std. Error of Mean 0.0058 Coefficient of Variation 0.632 Skewness 1.635

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.82 Shapiro Wilk Test Statistic 0.944 Shapiro Wilk Critical Value 0.947 Shapiro Wilk Critical Value 0.947

Data not Normal at 5% Significance Level

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 0.0739
95% H-UCL 0.0743
95% UCLs (Adjusted for Skewness)
95% Chebyshev (MVUE) UCL 0.0865
95% Adjusted-CLT UCL (Chen-1995) 0.0752
97.5% Chebyshev (MVUE) UCL 0.0964
95% Modified-t UCL (Johnson-1978) 0.0742
99% Chebyshev (MVUE) UCL 0.116

Gamma Distribution Test Data Distribution

k star (bias corrected) 3.111 Data do not follow a Discernable Distribution (0.05)

Theta Star 0.0207

MLE of Mean 0.0643

MLE of Standard Deviation 0.0365

nu star 311.1

Approximate Chi Square Value (.05) 271.2 Nonparametric Statistics

Adjusted Level of Significance 0.0452 95% CLT UCL 0.0738
Adjusted Chi Square Value 270.1 95% Jackknife UCL 0.0739

Anderson-Darling Test Statistic 1.152

Anderson-Darling 5% Critical Value 0.756

95% Standard Bootstrap UCL 0.0736

95% Bootstrap-t UCL 0.0757

95% Hall's Bootstrap UCL 0.0757

Gulf Killifish, Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

Kolmogorov-Smirnov Test Statistic 0.157 Kolmogorov-Smirnov 5% Critical Value 0.126

Data not Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 0.0749 95% BCA Bootstrap UCL 0.0744 95% Chebyshev(Mean, Sd) UCL 0.0893 97.5% Chebyshev(Mean, Sd) UCL 0.1 99% Chebyshev(Mean, Sd) UCL 0.121

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0738 95% Adjusted Gamma UCL (Use when n < 40) 0.0741

Potential UCL to Use

Use 95% Chebyshev (Mean, Sd) UCL 0.0893

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_b.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result Value (mercury mg/kg)

General Statistics

Number of Valid Observations 25

Number of Distinct Observations 23

Minimum of Log Data -3.73

Maximum of Log Data -1.614

Mean of log Data -2.765

SD of log Data 0.628

Log-transformed Statistics

Raw Statistics

Minimum 0.024

Maximum 0.199

Mean 0.0762

Geometric Mean 0.063

Median 0.068

SD 0.0499

Std. Error of Mean 0.01

Coefficient of Variation 0.654

Skewness 1.162

Relevant UCL Statistics

Normal Distribution Test

Shapiro Wilk Test Statistic 0.863

Shapiro Wilk Critical Value 0.918

Data not Normal at 5% Significance Level

Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.954

Shapiro Wilk Critical Value 0.918

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution

95% Student's-t UCL 0.0933

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 0.0951

95% Modified-t UCL (Johnson-1978) 0.0937

Assuming Lognormal Distribution

95% H-UCL 0.1

95% Chebyshev (MVUE) UCL 0.12

97.5% Chebyshev (MVUE) UCL 0.139

99% Chebyshev (MVUE) UCL 0.177

Gamma Distribution Test

Data Distribution

k star (bias corrected) 2.465

Theta Star 0.0309

MLE of Mean 0.0762

MLE of Standard Deviation 0.0486

nu star 123.2

Approximate Chi Square Value (.05) 98.6

Adjusted Level of Significance 0.0395

Adjusted Chi Square Value 97.1

Anderson-Darling Test Statistic 0.506 Anderson-Darling 5% Critical Value 0.752

Data appear Gamma Distributed at 5% Significance Level

Nonparametric Statistics

95% CLT UCL 0.0926

95% Jackknife UCL 0.0933

95% Standard Bootstrap UCL 0.0921

95% Bootstrap-t UCL 0.0986

95% Hall's Bootstrap UCL 0.0957

Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

Kolmogorov-Smirnov Test Statistic 0.144

Kolmogorov-Smirnov 5% Critical Value 0.176

Data appear Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 0.0938 95% BCA Bootstrap UCL 0.0946 95% Chebyshev(Mean, Sd) UCL 0.12 97.5% Chebyshev(Mean, Sd) UCL 0.139 99% Chebyshev(Mean, Sd) UCL 0.175

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 0.0953 95% Adjusted Gamma UCL (Use when n < 40) 0.0968

Potential UCL to Use

Use 95% Approximate Gamma UCL 0.0953

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and laci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

TOTAL PCB CONGENER

Gulf Killifish

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_c.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result Value (total pcb congener (u = 1/2) ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 25

Raw Statistics Log-transformed Statistics

Minimum 2E+06 Minimum of Log Data 14.49

Maximum 9E+06 Maximum of Log Data 16

Mean 5E+06 Mean of log Data 15.42
Geometric Mean 5E+06 SD of log Data 0.327

Median 5E+06 SD 2E+06

Std. Error of Mean 313682 Coefficient of Variation 0.301

Skewness 0.266

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.99 Shapiro Wilk Test Statistic 0.962 Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 6E+06
 95% H-UCL 5929489

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 6756490

 95% Adjusted-CLT UCL (Chen-1995) 6E+06
 97.5% Chebyshev (MVUE) UCL 7414908

95% Modified-t UCL (Johnson-1978) 6E+06 99% Chebyshev (MVUE) UCL 8708242

Gamma Distribution Test Data Distribution

k star (bias corrected) 9.398 Data appear Normal at 5% Significance Level

Theta Star 554953

MLE of Mean 5E+06

MLE of Standard Deviation 2E+06

nu star 469.9

Approximate Chi Square Value (.05) 420.7 Nonparametric Statistics

Adjusted Level of Significance 0.0395 95% CLT UCL 5731570
Adjusted Chi Square Value 417.5 95% Jackknife UCL 5752282

Anderson-Darling Test Statistic 0.164

Anderson-Darling 5% Critical Value 0.745

95% Standard Bootstrap UCL 5709166

95% Bootstrap-t UCL 5782797

95% Hall's Bootstrap UCL 5788260

Gulf Killifish

Kolmogorov-Smirnov Test Statistic 0.0718 Kolmogorov-Smirnov 5% Critical Value 0.174

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 6E+0695% Adjusted Gamma UCL (Use when n < 40) 6E+06

Potential UCL to Use

95% Percentile Bootstrap UCL 5715854 95% BCA Bootstrap UCL 5731928 95% Chebyshev(Mean, Sd) UCL 6582918 97.5% Chebyshev(Mean, Sd) UCL 7174554 99% Chebyshev(Mean, Sd) UCL 8336708

Use 95% Student's-t UCL 5752282

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Gulf Menhaden, Striped Mullet, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_g.wst

Full Precision OFF

Confidence Coefficient

Number of Bootstrap Operations 2000

Result Value (total pcb congener (u = 1/2) ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 25

Raw Statistics Log-transformed Statistics

> Minimum 596176 Maximum 5E+06

> > Mean 3E+06

Geometric Mean 2E+06

Median 2E+06

SD 1E+06

Std. Error of Mean 205165

Coefficient of Variation 0.406

Skewness 0.429

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.954

Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Assuming Normal Distribution

Gamma Distribution Test

Assuming Lognormal Distribution

Data appear Lognormal at 5% Significance Level

95% Student's-t UCL 3E+06

95% UCLs (Adjusted for Skewness)

95% Adjusted-CLT UCL (Chen-1995) 3E+06

95% Modified-t UCL (Johnson-1978) 3E+06

Data Distribution

Data appear Normal at 5% Significance Level

Theta Star 495210

MLE of Mean 3E+06

MLE of Standard Deviation 1E+06

k star (bias corrected) 5.107

nu star 255.3

Approximate Chi Square Value (.05) 219.3

Adjusted Level of Significance 0.0395

Adjusted Chi Square Value 217.1

Anderson-Darling Test Statistic 0.259 Anderson-Darling 5% Critical Value 0.747

Nonparametric Statistics

95% CLT UCL 2866309

95% H-UCL 3068891

95% Jackknife UCL 2879855

95% Standard Bootstrap UCL 2857615

Minimum of Log Data 13.3

Maximum of Log Data 15.33

Shapiro Wilk Test Statistic 0.939

Shapiro Wilk Critical Value 0.918

95% Chebyshev (MVUE) UCL 3604346

99% Chebyshev (MVUE) UCL 4950724

97.5% Chebyshev (MVUE) UCL 4058543

Mean of log Data 14.65

SD of log Data 0.456

95% Bootstrap-t UCL 2908346

95% Hall's Bootstrap UCL 2889135

Gulf Menhaden, Striped Mullet, Pinfish, Sand Seatrout

Kolmogorov-Smirnov Test Statistic 0.104
Kolmogorov-Smirnov 5% Critical Value 0.175

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 3E+06 95% Adjusted Gamma UCL (Use when n < 40) 3E+06

Potential UCL to Use

95% Percentile Bootstrap UCL 2848567 95% BCA Bootstrap UCL 2883986 95% Chebyshev(Mean, Sd) UCL 3423136 97.5% Chebyshev(Mean, Sd) UCL 3810098 99% Chebyshev(Mean, Sd) UCL 4570210

Use 95% Student's-t UCL 2879855

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Gulf Killifish, Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total pcb congener (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 50 Number of Distinct Observations 50

Raw Statistics Log-transformed Statistics

Minimum 596176 Minimum of Log Data 13.3

Maximum 9E+06 Maximum of Log Data 16

Mean 4E+06 Mean of log Data 15.04

Geometric Mean 3E+06 SD of log Data 0.551

Median 4E+06 SD 2E+06

Std. Error of Mean 266900

Coefficient of Variation 0.487

Skewness 0.535

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.96 Shapiro Wilk Test Statistic 0.962 Shapiro Wilk Critical Value 0.947 Shapiro Wilk Critical Value 0.947

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 4E+06
 95% H-UCL 4594363

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 5351061

 95% Adjusted-CLT UCL (Chen-1995) 4E+06
 97.5% Chebyshev (MVUE) UCL 5964232

95% Modified-t UCL (Johnson-1978) 4E+06 99% Chebyshev (MVUE) UCL 7168686

Gamma Distribution Test Data Distribution

k star (bias corrected) 3.709 Data appear Normal at 5% Significance Level

Theta Star 1E+06

MLE of Mean 4E+06

MLE of Standard Deviation 2E+06

nu star 370.9

Approximate Chi Square Value (.05) 327.3 Nonparametric Statistics

Adjusted Level of Significance 0.0452 95% CLT UCL 4311237

Adjusted Chi Square Value 326.1 95% Jackknife UCL 4319697

95% Standard Bootstrap UCL 4305645

Anderson-Darling Test Statistic 0.302 95% Bootstrap-t UCL 4337908

Gulf Killifish, Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

Anderson-Darling 5% Critical Value 0.754 Kolmogorov-Smirnov Test Statistic 0.0887 Kolmogorov-Smirnov 5% Critical Value 0.126

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 4E+06 95% Adjusted Gamma UCL (Use when n < 40) 4E+06

Potential UCL to Use

95% Hall's Bootstrap UCL 4366124 95% Percentile Bootstrap UCL 4298829 95% BCA Bootstrap UCL 4329279 95% Chebyshev(Mean, Sd) UCL 5035616 97.5% Chebyshev(Mean, Sd) UCL 5539017 99% Chebyshev(Mean, Sd) UCL 6527849

Use 95% Student's-t UCL 4319697

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

PCB CONGENER TEQ (FISH)

Gulf Killifish

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result Value (total pcb congener teq 1998 (fish) (u = 1/2) ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 23

Raw Statistics Log-transformed Statistics

Minimum 2.2 Minimum of Log Data 0.788

Maximum 9.7 Maximum of Log Data 2.272

Mean 6.553 Mean of log Data 1.824

Geometric Mean 6.198 SD of log Data 0.363

Median 6.9

SD 2.013

Coefficient of Variation 0.307

Std. Error of Mean 0.403

Skewness -0.319

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.968 Shapiro Wilk Test Statistic 0.907
Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 7.242
 95% H-UCL 7.598

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 8.741

 95% Adjusted-CLT UCL (Chen-1995) 7.188
 97.5% Chebyshev (MVUE) UCL 9.666

95% Modified-t UCL (Johnson-1978) 7.238 99% Chebyshev (MVUE) UCL 11.48

Gamma Distribution Test Data Distribution

k star (bias corrected) 8.061 Data appear Normal at 5% Significance Level

Theta Star 0.813

MLE of Mean 6.553

MLE of Standard Deviation 2.308

nu star 403

Approximate Chi Square Value (.05) 357.5 Nonparametric Statistics

Adjusted Level of Significance 0.0395 95% CLT UCL 7.216
Adjusted Chi Square Value 354.6 95% Jackknife UCL 7.242

95% Standard Bootstrap UCL 7.201

Anderson-Darling Test Statistic 0.5 95% Bootstrap-t UCL 7.228

Anderson-Darling 5% Critical Value 0.745 95% Hall's Bootstrap UCL 7.217

Gulf Killifish

Kolmogorov-Smirnov Test Statistic 0.143 Kolmogorov-Smirnov 5% Critical Value 0.175

Data appear Gamma Distributed at 5% Significance Level

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n \ge 40$) 7.388 95% Adjusted Gamma UCL (Use when n < 40) 7.449

Potential UCL to Use

95% Percentile Bootstrap UCL 7.187 95% BCA Bootstrap UCL 7.161 95% Chebyshev(Mean, Sd) UCL 8.308 97.5% Chebyshev(Mean, Sd) UCL 9.068 99% Chebyshev(Mean, Sd) UCL 10.56

Use 95% Student's-t UCL 7.242

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

Gulf Killifish, Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_a.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result Value (total pcb congener teq 1998 (fish) (u = 1/2) ng/kg)

General Statistics

Number of Valid Observations 50 Number of Distinct Observations 43

Raw Statistics Log-transformed Statistics

Minimum 0.6 Maximum 9.7

Mean 4.541 Mean of log Data 1.314

Minimum of Log Data -0.511

Maximum of Log Data 2.272

99% Chebyshev (MVUE) UCL 9.661

Geometric Mean 3.721 SD of log Data 0.689

Median 3.95

SD 2.607

Std. Error of Mean 0.369

Coefficient of Variation 0.574

Skewness 0.414

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.928 Shapiro Wilk Test Statistic 0.937
Shapiro Wilk Critical Value 0.947 Shapiro Wilk Critical Value 0.947

Data not Normal at 5% Significance Level

Data not Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 5.159
 95% H-UCL 5.764

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 6.869

 95% Adjusted-CLT UCL (Chen-1995) 5.17
 97.5% Chebyshev (MVUE) UCL 7.811

Gamma Distribution Test Data Distribution

k star (bias corrected) 2.52 Data appear Gamma Distributed at 5% Significance Level

Theta Star 1.802

MLE of Mean 4.541

MLE of Standard Deviation 2.86

95% Modified-t UCL (Johnson-1978) 5.163

nu star 252

Approximate Chi Square Value (.05) 216.3 Nonparametric Statistics

Adjusted Level of Significance 0.0452 95% CLT UCL 5.147

Adjusted Chi Square Value 215.3 95% Jackknife UCL 5.159 95% Standard Bootstrap UCL 5.128

Anderson-Darling Test Statistic 0.489 95% Bootstrap-t UCL 5.182

Anderson-Danning rest statistic 0.469

Anderson-Darling 5% Critical Value 0.759 95% Hall's Bootstrap UCL 5.171

Gulf Killifish, Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

Kolmogorov-Smirnov Test Statistic 0.0954

Kolmogorov-Smirnov 5% Critical Value 0.126

Data appear Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 5.167 95% BCA Bootstrap UCL 5.194 95% Chebyshev(Mean, Sd) UCL 6.148 97.5% Chebyshev(Mean, Sd) UCL 6.844 99% Chebyshev(Mean, Sd) UCL 8.21

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n \ge 40$) 5.292 95% Adjusted Gamma UCL (Use when n < 40) 5.316

Potential UCL to Use

Use 95% Approximate Gamma UCL 5.292

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1 b.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total pcb congener teq 1998 (fish) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 25 Number of Distinct Observations 22

Raw Statistics Log-transformed Statistics

Minimum 0.6 Minimum of Log Data -0.511

Maximum 5.1 Maximum of Log Data 1.629

Mean 2.528 Mean of log Data 0.804

etric Mean 2.234 SD of log Data 0.544

Geometric Mean 2.234 Median 2.4

SD 1.179

Std. Error of Mean 0.236 Coefficient of Variation 0.466

Skewness 0.453

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.962 Shapiro Wilk Test Statistic 0.933 Shapiro Wilk Critical Value 0.918 Shapiro Wilk Critical Value 0.918

Data appear Normal at 5% Significance Level Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 2.932
 95% H-UCL 3.234

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 3.853

 95% Adjusted-CLT UCL (Chen-1995) 2.939
 97.5% Chebyshev (MVUE) UCL 4.406

 95% Modified-t UCL (Johnson-1978) 2.935
 99% Chebyshev (MVUE) UCL 5.494

Gamma Distribution Test Data Distribution

k star (bias corrected) 3.726 Data appear Normal at 5% Significance Level

Theta Star 0.679

MLE of Mean 2.528

MLE of Standard Deviation 1.31

nu star 186.3

Approximate Chi Square Value (.05) 155.7 Nonparametric Statistics

Adjusted Level of Significance 0.0395 95% CLT UCL 2.916

Adjusted Chi Square Value 153.8 95% Jackknife UCL 2.932

95% Standard Bootstrap UCL 2.911

Anderson-Darling Test Statistic 0.296 95% Bootstrap-t UCL 2.95

Striped Mullet, Gulf Menhaden, Pinfish, Sand Seatrout

Anderson-Darling 5% Critical Value 0.748 Kolmogorov-Smirnov Test Statistic 0.126 Kolmogorov-Smirnov 5% Critical Value 0.175

Data appear Gamma Distributed at 5% Significance Level

95% Approximate Gamma UCL (Use when n >= 40) 3.025 95% Adjusted Gamma UCL (Use when n < 40) 3.062

Assuming Gamma Distribution

Use 95% Student's-t UCL 2.932

95% Hall's Bootstrap UCL 2.951

95% BCA Bootstrap UCL 2.917

95% Percentile Bootstrap UCL 2.909

95% Chebyshev(Mean, Sd) UCL 3.556 97.5% Chebyshev(Mean, Sd) UCL 4.001

99% Chebyshev(Mean, Sd) UCL 4.875

Potential UCL to Use

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL. These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

DIOXIN/FURAN TEQ (FISH)

Gulf Menhaden

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_a.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result Value (total dioxin/furan teq 1998 (fish) (u = 1/2) ng/kg)

General Statistics

Number of Valid Observations 10 Number of Distinct Observations 10

Raw Statistics Log-transformed Statistics

Minimum 9.495 Minimum of Log Data 2.251 Maximum 18.13 Maximum of Log Data 2.898

Mean of log Data 2.619

Geometric Mean 13.73 SD of log Data 0.224

Median 14.65

SD 2.942

Std. Error of Mean 0.93

95% Student's-t UCL 15.73

Coefficient of Variation 0.21

Skewness -0.456

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.92 Shapiro Wilk Test Statistic 0.893

Shapiro Wilk Critical Value 0.842 Shapiro Wilk Critical Value 0.842

Data appear Normal at 5% Significance Level

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 18.4

 95% Adjusted-CLT UCL (Chen-1995) 15.41
 97.5% Chebyshev (MVUE) UCL 20.29

95% Modified-t UCL (Johnson-1978) 15.71 99% Chebyshev (MVUE) UCL 23.99

Gamma Distribution Test Data Distribution

k star (bias corrected) 16.35 Data appear Normal at 5% Significance Level

Theta Star 0.858

MLE of Mean 14.03

MLE of Standard Deviation 3.469

nu star 327.1

Approximate Chi Square Value (.05) 286.2 Nonparametric Statistics

Adjusted Level of Significance 0.0267 95% CLT UCL 15.56

Adjusted Chi Square Value 279.5 95% Jackknife UCL 15.73

95% Standard Bootstrap UCL 15.48

Anderson-Darling Test Statistic 0.518 95% Bootstrap-t UCL 15.61

Anderson-Darling 5% Critical Value 0.725 95% Hall's Bootstrap UCL 15.39

95% H-UCL 16.23

Gulf Menhaden

Kolmogorov-Smirnov Test Statistic 0.206
Kolmogorov-Smirnov 5% Critical Value 0.266

Data appear Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 15.52 95% BCA Bootstrap UCL 15.33 95% Chebyshev(Mean, Sd) UCL 18.08 97.5% Chebyshev(Mean, Sd) UCL 19.84 99% Chebyshev(Mean, Sd) UCL 23.28

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when $n \ge 40$) 16.03 95% Adjusted Gamma UCL (Use when n < 40) 16.41

Potential UCL to Use

Use 95% Student's-t UCL 15.73

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

Note: For highly negative-skewed data, confidence limits (e.g., Chen, Johnson, Lognormal, and Gamma) may not be reliable. Chen's and Johnson's methods provide adjustments for positvely skewed data sets.

Sand Seatrout

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1_b.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

Result_Value (total dioxin/furan teq 1998 (fish) (u = 1/2)_ng/kg)

General Statistics

Number of Valid Observations 2

Number of Distinct Observations 2

Warning: This data set only has 2 observations!

Data set is too small to compute reliable and meaningful statistics and estimates!

The data set for variable Result_Value (total dioxin/furan teq 1998 (fish) (u = 1/2)_ng/kg) was not processed!

It is suggested to collect at least 8 to 10 observations before using these statistical methods!

If possible, compute and collect Data Quality Objectives (DQO) based sample size and analytical results.

Gulf Killifish, Striped Mullet, Pinfish

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst
Full Precision OFF

Confidence Coefficient 95%
Number of Bootstrap Operations 2000

Result Value (total dioxin/furan teg 1998 (fish) (u = 1/2) ng/kg)

General Statistics

Number of Valid Observations 38 Number of Distinct Observations 38

Raw Statistics Log-transformed Statistics

Minimum 2.897 Minimum of Log Data 1.064
Maximum 20.44 Maximum of Log Data 3.017
Mean 8.009 Mean of log Data 1.997

Geometric Mean 7.37 SD of log Data 0.411

Median 7.121 SD 3.481

Std. Error of Mean 0.565 Coefficient of Variation 0.435 Skewness 1.395

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.908 Shapiro Wilk Test Statistic 0.991
Shapiro Wilk Critical Value 0.938 Shapiro Wilk Critical Value 0.938

Data not Normal at 5% Significance Level

Data appear Lognormal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

 95% Student's-t UCL 8.962
 95% H-UCL 9.092

 95% UCLs (Adjusted for Skewness)
 95% Chebyshev (MVUE) UCL 10.41

 95% Adjusted-CLT UCL (Chen-1995) 9.075
 97.5% Chebyshev (MVUE) UCL 11.45

 95% Modified-t UCL (Johnson-1978) 8.983
 99% Chebyshev (MVUE) UCL 13.49

Gamma Distribution Test Data Distribution

k star (bias corrected) 5.699 Data appear Gamma Distributed at 5% Significance Level

Theta Star 1.405

MLE of Mean 8.009

MLE of Standard Deviation 3.355

nu star 433.1

Approximate Chi Square Value (.05) 385.9 Nonparametric Statistics

Adjusted Level of Significance 0.0434 95% CLT UCL 8.938
Adjusted Chi Square Value 384 95% Jackknife UCL 8.962

95% Standard Bootstrap UCL 8.911

Anderson-Darling Test Statistic 0.286 95% Bootstrap-t UCL 9.121

Anderson-Darling 5% Critical Value 0.75 95% Hall's Bootstrap UCL 9.201

Gulf Killifish, Striped Mullet, Pinfish

Kolmogorov-Smirnov Test Statistic 0.103 Kolmogorov-Smirnov 5% Critical Value 0.143

Data appear Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 8.916 95% BCA Bootstrap UCL 9.065 95% Chebyshev(Mean, Sd) UCL 10.47 97.5% Chebyshev(Mean, Sd) UCL 11.54 99% Chebyshev(Mean, Sd) UCL 13.63

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 8.99 95% Adjusted Gamma UCL (Use when n < 40) 9.033

Potential UCL to Use

Use 95% Approximate Gamma UCL 8.99

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

APPENDIX D-2 FISH TISSUE LOEDS FOR MERCURY

Appendix D-2 Fish Tissue LOEDs for Mercury

Ref ID	Year	Author	Publication Source	Species Scientific Name	Species Common Name	Species TSN Code	Analyte Name	CAS No.	Conc_Wet	Conc_Units	No. Reps	QA Stats Method	QA Chem Method	QA Chem DL
SEQ97-31	1982	Snarski, V.M., Olson, G.F.	Aquat Toxicol	Pimephales promelas	Fathead minnow	163517	Mercury	7439-97-6	1.31	MG/KG	6	Υ	Υ	N
SEQ97-31	1982	Snarski, V.M., Olson, G.F.	Aquat Toxicol	Pimephales promelas	Fathead minnow	163517	Mercury	7439-97-6	1.36	MG/KG	30	Υ	Υ	N
SEQ97-31	1982	Snarski, V.M., Olson, G.F.	Aquat Toxicol	Pimephales promelas	Fathead minnow	163517	Mercury	7439-97-6	1.36	MG/KG	7	Υ	Υ	N
URS21	1996	Friedmann, A.S., M.C. Watzin, T. Brinck-Johnsen and J.C. Leiter	Aquat Toxicol 35:265-278	Stizostedion vitreum	Walleye	168506	Mercury	7439-97-6	2.37	MG/KG	22	Υ	Y	Y
SEQ97-31	1982	Snarski, V.M., Olson, G.F.	Aquat Toxicol	Pimephales promelas	Fathead minnow	163517	Mercury	7439-97-6	4.76	MG/KG	6	Υ	Υ	N
JA376	1982	Rodgers DW, FWH Beamish	Aquat Toxicol 02:271-290	Oncorhynchus mykiss	Trout - Rainbow	161989	Methyl Mercury	22967-92-6	8	MG/KG	1	Υ	Υ	Υ
JA376	1982	Rodgers DW, FWH Beamish	Aquat Toxicol 02:271-290	Oncorhynchus mykiss	Trout - Rainbow	161989	Methyl Mercury	22967-92-6	10	MG/KG	1	Υ	Υ	Υ
SEQ97-31	1982	Snarski, V.M., Olson, G.F.	Aquat Toxicol	Pimephales promelas	Fathead minnow	163517	Mercury	7439-97-6	18.8	MG/KG	4	Υ	Υ	N
SEQ97-31	_	Snarski, V.M., Olson, G.F.	Aquat Toxicol	Pimephales promelas	Fathead minnow	163517	Mercury	7439-97-6	4.18	MG/KG	6	Υ	Υ	N
JA488	1975	Wobeser, G.	J Fish Res Bd Can 32(11):2005-2013	Oncorhynchus mykiss	Trout - Rainbow	161989	Mercury	7439-97-6	6.9	MG/KG	3	Υ	Υ	N
JA489	1975	Wobeser, G.	J Fish Res Bd Can 32(11), 2015-2023	Oncorhynchus mykiss	Trout - Rainbow	161989	Mercury	7439-97-6	10	MG/KG	5	N	Υ	Υ
MEC03-134	2002	Hammerschmidt, Chad, Mark Sandheinrich, James Wiener, and Ronald Rada	Environ Sci Tech 36: 877-883	Pimephales promelas	Fathead minnow	163517	Methyl Mercury	22967-92-6	0.7	MG/KG	8	Υ	Y	Υ
MEC04-120	1998	Vollestad LA, E Fjeld, T Haugen, SA Oxnevad	Environ Pollut 101:349-354	Thymallus thymallus	Grayling	162022	Mercury	7439-97-6	3.8	MG/KG	1	Υ	Υ	N

Appendix D-2 Fish Tissue LOEDs for Mercury

Ref ID	QA Exp Method	QA Effects Method	QA Reps	Effect Class	Toxicity Measure	Exposure Route	Species Body Part	Species Start Lifestage	Species Habitat	Species Feeding Behavior	Comments
SEQ97-31	Υ	Υ	Υ	Growth	LOED	Combined	Whole Body	Adult	Lakes, ponds, streams	Not Specified	
SEQ97-31	Υ	Υ	Υ	Growth	LOED	Combined	Whole Body	Adult	Lakes, ponds, streams	Not Specified	30-day exposure: Progeny of exposed fish.
SEQ97-31	Υ	Υ	Υ	Growth	LOED	Combined	Whole Body	Adult	Lakes, ponds, streams	Not Specified	Weight of females.
URS21	Y	Υ	Υ	Growth	LOED	Ingestion	Whole Body		Great Lakes region, upper Mississippi north to Assiniboia, east to VT, PA, GA, AB; in clear water with rocks, gravel, sand bottom	Carnivorous; largely on various minnows, occasionally young whitefish	Significant reduction in length and weight of males, but not females.
SEQ97-31	Υ	Υ	Υ	Growth	LOED	Combined	Whole Body	Adult	Lakes, ponds, streams	Not Specified	
JA376	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Juvenile	Cool streams	Carnivore-juv fish, inverts	Growth significantly reduced. Fish fed the 25/2% diet and 75/2% diet were not significantly different in their wet weights.
JA376	Υ	Y	Υ	Growth	LOED	Ingestion	Whole Body	Juvenile	Cool streams	Carnivore-juv fish, inverts	Growth significantly reduced. Fish fed the 25/ad lib diet and 45/ad lib diet were not significantly different in their wet weights.
SEQ97-31	Υ	Υ	Υ	Growth	LOED	Combined	Whole Body	Adult	Lakes, ponds, streams	Not Specified	Weight of males.
SEQ97-31	Υ	Υ	Υ	Mortality	LOED	Combined	Whole Body	Adult	Lakes, ponds, streams	Not Specified	
JA488	Υ	Υ	Υ	Mortality	LOED	Absorption	Carcass	Immature	Cool streams	Carnivore-juv fish, inverts	Increase in mortality.
JA489	Υ	Υ	Υ	Mortality	LOED	Ingestion	Whole Body	Juvenile	Cool streams	Carnivore-juv fish, inverts	Increase in mortality. Residue is injected dose (i.p.).
MEC03-134	Υ	Υ	Υ	Reproduction	LOED	Ingestion	Whole Body	Egg	Lakes, ponds, streams	Not Specified	Daily number of eggs produced relative to carcass concentration of methylmercury in female fish fed the same diet throughout exposure periods.
MEC04-120	Υ	Υ	N	Reproduction	LOED	Water	Whole Body	Egg	NS	Not Specified	Hatching success.

APPENDIX D-3 FISH TISSUE LOEDS FOR 2,3,7,8-TCDD

Appendix D-3 Fish Tissue LOEDs for 2,3,7,8-TCDD

Ref ID	Year	Author	Publication Source	Species Scientific Name	Species Common Name	Species TSN Code	Analyte Name	CAS No.	Conc_Wet	Conc_Units	No. Reps	QA Stats Method	QA Chem Method	QA Chem DL
JA470	1994	Walker MK, PM Cook, AR Batterman, BC Butterworth, C Berini, JJ Libal, LC Hufnagle, RE Peterson	Can J Fish Aquat Sci 51:1410-1419	Salvelinus namaycush	Trout -Lake	162002	2,3,7,8-TCDD	1746-01-6	4.00E-05	MG/KG	3	Υ	Υ	Y
JA470	1994	Walker MK, PM Cook, AR Batterman, BC Butterworth, C Berini, JJ Libal, LC Hufnagle, RE Peterson	Can J Fish Aquat Sci 51:1410-1419	Salvelinus namaycush	Trout -Lake	162002	2,3,7,8-TCDD	1746-01-6	5.00E-05	MG/KG	3	Y	Υ	Y
JA474	1991	Walker MK, JM Spitsbergen, JR Olson, RE Peterson	Can J Fish Aquat Sci 48:875-883	Salvelinus namaycush	Trout -Lake	162002	2,3,7,8-TCDD	1746-01-6	5.50E-05	MG/KG	4	Υ	Υ	Υ
JA474	1991	Walker MK, JM Spitsbergen, JR Olson, RE Peterson	Can J Fish Aquat Sci 48:875-883	Salvelinus namaycush	Trout -Lake	162002	2,3,7,8-TCDD	1746-01-6	5.50E-05	MG/KG	4	Υ	Υ	Υ
JA470	1994	Walker MK, PM Cook, AR Batterman, BC Butterworth, C Berini, JJ Libal, LC Hufnagle, RE Peterson	Can J Fish Aquat Sci 51:1410-1419	Salvelinus namaycush	Trout -Lake	162002	2,3,7,8-TCDD	1746-01-6	5.50E-05	MG/KG	3	Υ	Υ	Y
JA474	1991	Walker MK, JM Spitsbergen, JR Olson, RE Peterson	Can J Fish Aquat Sci 48:875-883	Salvelinus namaycush	Trout -Lake	162002	2,3,7,8-TCDD	1746-01-6	0.000226	MG/KG	4	Υ	Υ	Υ
JA368	1995	Prince, R. and K.R. Cooper	Environ Tox & Chem 14(4):579-587	Fundulus heteroclitus	Killfish	165647	2,3,7,8-TCDD	1746-01-6	0.000635	MG/KG	5	Υ	Υ	Υ
URS34	1985	Branson, D.R., L.T. Takahashi, W.M. Parker and G.E. Blau	Environ Tox & Chem 04:779-788	Oncorhynchus mykiss	Trout - Rainbow	161989	2,3,7,8-TCDD	1746-01-6	0.00065	MG/KG	3	Υ	Υ	Y
URS34	1985	Branson, D.R., L.T. Takahashi, W.M. Parker and G.E. Blau	Environ Tox & Chem 04:779-788	Oncorhynchus mykiss	Trout - Rainbow	161989	2,3,7,8-TCDD	1746-01-6	0.00065	MG/KG	3	Υ	Υ	Y
URS212	1994	Tietge, J.E.	Society of Environmental Toxicology and Chemistry 15th Annual Meeting Abstract	Salvelinus fontinalis	Trout - Brook	162003	2,3,7,8-TCDD	1746-01-6	0.0012	MG/KG	1	N	N	N
URS199	1979	Miller, R.A., L.A. Norris and B.R. Loper	Trans Am Fish Soc 108:401-407	Oncorhynchus kisutch	Salmon-coho	161977	2,3,7,8-TCDD	1746-01-6	0.00217	MG/KG	1	Υ	Υ	Y
URS199	1979	Miller, R.A., L.A. Norris and B.R. Loper	Trans Am Fish Soc 108:401-407	Oncorhynchus kisutch	Salmon-coho	161977	2,3,7,8-TCDD	1746-01-6	0.00217	MG/KG	1	Υ	Υ	Υ
JA185	1977	Hawkes, C.L. and L.A. Norris	Trans Am Fish Soc 106:641-645	Oncorhynchus mykiss	Trout - Rainbow	161989	2,3,7,8-TCDD	1746-01-6	1.38	MG/KG	1	Υ	Υ	Υ
JA185	1977	Hawkes, C.L. and L.A. Norris	Trans Am Fish Soc 106:641-645	Oncorhynchus mykiss	Trout - Rainbow	161989	2,3,7,8-TCDD	1746-01-6	1.38	MG/KG	1	Υ	Υ	Υ

Appendix D-3 Fish Tissue LOEDs for 2,3,7,8-TCDD

									FISH TISSUE LOLDS for 2,5,7,8-1CI		
Ref ID	QA Exp Method	QA Effects Method	QA Reps	Effect Class	Toxicity Measure	Exposure Route	Species Body Part	Species Start Lifestage	Species Habitat	Species Feeding Behavior	Comments
JA470	Υ	Y	Υ	Mortality	LOED	Ingestion	Whole Body	Egg	Deep waters of cold northern lakes	Carnivore-adults piscivorous	Eggs groups recieved one injection out of 7 graded doses of TCDD.
JA470	Υ	Υ	Υ	Mortality	LOED	Ingestion	Whole Body	Egg	Deep waters of cold northern lakes	Carnivore-adults piscivorous	TCDD exposure resulted from feeding adult female trout for up to 149d prior to spawning. No significant differences were found between exp. dose and egg bioaccumulation.
JA474	Υ	Υ	Υ	Growth	LOED	Water	Whole Body	Egg	Deep waters of cold northern lakes	Carnivore-adults piscivorous	Significant increase in sac fry weight at this TCDD concentration.
JA474	Υ	Υ	Υ	Mortality	LOED	Water	Whole Body		Deep waters of cold northern lakes	Carnivore-adults piscivorous	Significant increase in mortality at this concentration - the lowest concentration to do so. (Called LOAEL in the paper.)
JA470	Υ	Υ	Υ	Mortality	LOED	Ingestion	Whole Body	Egg	Deep waters of cold northern lakes	Carnivore-adults piscivorous	Eggs were treated with waterborne TCDD.
JA474	Υ	Υ	Υ	Growth	LOED	Water	Whole Body	Egg	Deep waters of cold northern lakes	Carnivore-adults piscivorous	Significant sac fry wet weight increase at this TCDD concentration. Significant decrease in sac fry length at this TCDD concentration.
JA368	Υ	Υ	Υ	Mortality	LOED	Absorption	Whole Body	Embryo	Not Specified	Not Specified	28% decrease in hatchability.
URS34	Υ	Υ	Υ	Growth	LOED	Absorption	Whole Body	Immature	Cool streams	Carnivore-juv fish, inverts	Reduced growth, exposed fish weighed 50 G vs. 130 G for controls. And, liver/body weight ratio nearly twice that of controls.
URS34	Υ	Υ	Υ	Mortality	LOED	Absorption	Whole Body	Immature	Cool streams	Carnivore-juv fish, inverts	Lethal to 7 of 90 fish over 139 days.
URS212	Υ	N	N	Reproduction	LOED	Ingestion	Whole Body	Immature	Cool streams, gravel bottoms	Carnivore-aquatic insects, fish	Significant delay in spawning; TCDD-spiked diet to produce desired body burden; abstract with minimal information.
URS199	Υ	Υ	Υ	Growth	LOED	Absorption	Whole Body	Immature	NW America and Asian Coast. Streams in Spring, Fall	Carnivore-aquatic insects, fish, inverts	Reduced weight gain. Residue measured after 114 days in clean water.
URS199	Υ	Υ	Υ	Mortality	LOED	Absorption	Whole Body	Immature	NW America and Asian Coast. Streams in Spring, Fall	Carnivore-aquatic insects, fish, inverts	Reduced survival. Residue measured after 114 days in clean water.
JA185	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Juvenile	Cool streams	Carnivore-juv fish, inverts	Weight loss.
JA185	Υ	Υ	Υ	Mortality	LOED	Ingestion	Whole Body	Juvenile	Cool streams	Carnivore-juv fish, inverts	Over 90% mortality.

APPENDIX D-4 FISH TISSUE LOEDS FOR TOTAL PCBS

Appendix D-4 Fish Tissue LOEDs for Total PCBs

Ref ID	Year	Author	Publication Source	Species Scientific Name	Species Common Name	Species TSN Code	Analyte Name	CAS No	Conc_Wet	Conc_Units	No. Reps	QA Stats Method	QA Chem Method	QA Chem DL
MEC04-046	1998	Orn, S., P.L. Anderson, L. Forlin, M. Tysklind, L. Norrgren	Arch Environ Contam Toxicol 35:53-57	Danio rerio	Zebra Danio	163699	PCBs	1336-36-3	1.40E-01	MG/KG	1	Υ	Υ	N
MEC04-046	1998	Orn, S., P.L. Anderson, L. Forlin, M. Tysklind, L. Norrgren	Arch Environ Contam Toxicol 35:53-57	Danio rerio	Zebra Danio	163699	PCBs	1336-36-3	1.10E+00	MG/KG	1	Υ	Υ	N
SEQ99J_11	1973	Hansen, D.J., S.C. Schimmel and J. Forester	Proceedings of the Twenty-Seventh Annual Conference Southeastern Association of Game and Fish Commissioners, Hot Springs, Arkansas	Cyprinodon variegatus	Sheepshead minnow	165631	PCBs	1336-36-3	8.80E-01	MG/KG	24	N	Υ	N
MEC04-046	1998	Orn, S., P.L. Anderson, L. Forlin, M. Tysklind, L. Norrgren	Arch Environ Contam Toxicol 35:53-57	Danio rerio	Zebra Danio	163699	PCBs	1336-36-3	1.10E+00	MG/KG	1	Υ	Υ	N
URS14	1980	Bengtsson, B.E.	Water Res. 14:681-687	Phoxinus phoxinus	Minnow	163596	PCBs	1336-36-3	7.00E+01	MG/KG	4	Υ	Υ	N
URS14	1980	Bengtsson, B.E.	Water Res. 14:681-687	Phoxinus phoxinus	Minnow	163596	PCBs	1336-36-3	15	MG/KG	4	Y	Υ	N
URS14	1980	Bengtsson, B.E.	Water Res. 14:681-687	Phoxinus phoxinus	Minnow	163596	PCBs	1336-36-3	180	MG/KG	4	Υ	Υ	N
URS103	1975	Hansen, D.J., S.C. Schimmel and J. Forester	Trans Am Fish Soc 104:584-588	Cyprinodon variegatus	Sheepshead minnow	165631	PCBs	1336-36-3	200	MG/KG	1	Y	Υ	Υ
JA278	1978	Mauck, W.L., P.M. Mehrle, and F.L. Mayer.	J Fish Res Bd Can 35:1084-1088	Salvelinus fontinalis	Trout - Brook	162003	Aroclor 1254	11097-69-1	125	MG/KG	1	Υ	Υ	Υ
JA28	1981	Berlin, W.H., R.J. Hesselberg, and M.J. Mac	In Chlorinated Hydrocarbons as a Factor in the Reproduction and Survival of Lake Trout (Salvelinus namaycush) in Lake Michigan. Technical Paper 105. U.S. Fish and Wildlife Service	Salvelinus namaycush	Trout -Lake	162002	Aroclor 1254	11097-69-1	1.53	MG/KG	2	Υ	Υ	N
URS157	1979	Leatherland, J.F., B.A. Sonstegard and N.V. Holdrient	Comp Biochem Physiol 063c:243-246	Oncorhynchus kisutch	Salmon-coho	161977	PCBs	1336-36-3	9.8	MG/KG	6	N	Υ	N
URS104	1976	Hansen, L.G., W.B. Wiekhorst and J. Simon	J Fish Res Bd Can 33:1343-1352	Ictalurus punctatus	Catfish-Channel	163998	PCBs	1336-36-3	14.3	MG/KG	3	N	Υ	N
URS157	1979	Leatherland, J.F., B.A. Sonstegard and N.V. Holdrient	Comp Biochem Physiol 063c:243-246	Oncorhynchus kisutch	Salmon-coho	161977	PCBs	1336-36-3	43	MG/KG	6	N	Y	N
URS102	1974	Hansen, D.J., P.R. Parrish and J. Forester	Environ Res 7:363-373	Lagodon rhomboides	Pinfish	169187	PCBs	1336-36-3	65	MG/KG	1	N	Υ	Υ
URS94	1976	Gruger, E.H., T. Hurley and N.L. Karrick	Environ Sci Tech 10:1033-1037	Oncorhynchus kisutch	Salmon-coho	161977	PCBs	1336-36-3	2.30E+00	MG/KG	3	Υ	Υ	N
URS103	1975	Hansen, D.J., S.C. Schimmel and J. Forester	Trans Am Fish Soc 104:584-588	Cyprinodon variegatus	Sheepshead minnow	165631	PCBs	1336-36-3	1.100E+03	MG/KG	1	Υ	Υ	Y

Appendix D-4 Fish Tissue LOEDs for Total PCBs

									Fish Tissue LOEDs for Total PC	.03	
Ref ID	QA Exp Method	QA Effects Method	QA Reps	Effect Class	Toxicity Measure	Exposure Route	Species Body Part	Species Start Lifestage	Species Habitat	Species Feeding Behavior	Comments
MEC04-046	Υ	Υ	N	Growth	LOED	Ingestion	Whole Body	Adult	Not Specified	Not Specified	PCBs 41, 51, 60, 68, 91, 99, 104, 112, 115, 126, 143, 153, 169, 184, 193 - weight.
MEC04-046	Υ	Υ	N	Mortality	LOED	Ingestion	Whole Body	Adult	Not Specified	Not Specified	PCBs 41, 51, 60, 68, 91, 99, 104, 112, 115, 126, 143, 153, 169, 184, 193 - no significance calculated.
SEQ99J_11	Υ	Υ	Y	Reproduction	LOED	Absorption	Whole Body	Adult	Shallow brackish waters, Cape Cod to Mexico	Wide variety of small aquatic animals and plants	Increase in fecundity.
MEC04-046	Υ	Υ	N	Reproduction	LOED	Ingestion	Whole Body	Adult	Not Specified	Not Specified	PCBs 41, 51, 60, 68, 91, 99, 104, 112, 115, 126, 143, 153, 169, 184, 193 - ovary weight.
URS14	Υ	Υ	Υ	Mortality	LOED	Ingestion	Whole Body	Adult	Not Specified	Not Specified	Doubling of mortality rate compared to controls after 300 days
URS14	Y	Υ	Υ	Reproduction	LOED	Ingestion	Whole Body	Adult	Not Specified	Not Specified	Reduction in time to hatch. Fry death within 1 week of hatching. Spawning induced at higher water temperatures. Residues measured 2 months before spawning.
URS14	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Adult	Not Specified	Not Specified	Significantly increased growth at 79 days. Residue measured at 82 days.
URS103	Υ	Υ	Υ	Mortality	LOED	Absorption	Whole Body	Egg-embryo	Shallow brackish waters, Cape Cod to Mexico	Wide variety of small aquatic animals and plants	Lethal to 86% of fry in 28 days.
JA278	Υ	Υ	Υ	Mortality	LOED	Absorption	Whole Body	Embryo	Cool streams, gravel bottoms	Carnivore-aquatic insects, fish	18% increase in mortality.
JA28	Υ	Υ	Y	Mortality	LOED	Combined	Whole Body	Fry	Deep waters of cold northern lakes	Carnivore-adults piscivorous	Significant increase in mortality. Exposure to 20.8 ng/L in water and 1.05 ug/g in diet.
URS157	Y	Υ	Y	Growth	LOED	Ingestion	Carcass	Immature	NW America and Asian Coast. Streams in Spring, Fall	Carnivore-aquatic insects, fish, inverts	Significant increase in hepatosomatic index. Concurrent exposure to 5 mg/kg Mirex in food.
URS104	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Immature	Rapid water streams	Omnivore	40% Reduction in body weight. Increased liver/body weight ratio.
URS157	Υ	Υ	Υ	Growth	LOED	Ingestion	Carcass	Immature	NW America and Asian Coast. Streams in Spring, Fall	Carnivore-aquatic insects, fish, inverts	Significant increase in hepatosomatic index.
URS102	Υ	Y	Y	Mortality	LOED	Absorption	Whole Body	Immature	Lower portions of estuaries, Massachusetts to Texas, most common Virginia south	Fish, shellfish, worms, crustaceans, seaweed	18% mortality in 96 hours.
URS94	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Juvenile	NW America and Asian Coast. Streams in Spring, Fall	Carnivore-aquatic insects, fish, inverts	Significant increased rate of growth (weight gain).
URS103	Y	Υ	Υ	Mortality	LOED	Absorption	Whole Body	Juvenile	Shallow brackish waters, Cape Cod to Mexico	Wide variety of small aquatic animals and plants	88% juvenile mortality in 28 days.

APPENDIX D-5 FISH TISSUE LOEDS FOR SELENIUM

Appendix D-5
Fish Tissue LOEDs for Selenium

Pof ID Voc				Species Scientific	Species Common	Chasias	Analista		l	I	I	OA State	QA Chem	QA Chem
Ref ID	Year	Author	Publication Source	Name	Species Common Name	Species TSN Code	Analyte Name	CAS No	Conc_Wet	Conc Units	No. Reps	QA Stats Method	Method	DL QA Chem
JA29	1986	Bertram, P.E. and A.S. Brooks.	Wat. Res. 20(7):877-884.	Pimephales promelas	Fathead minnow	163517	Selenium	7782-49-2	0.18	mg/kg	1	Υ	Υ	N
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	0.66	mg/kg	10	Υ	Υ	Υ
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	0.68	mg/kg	20	Y	Y	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	0.8	mg/kg	4	Υ	Υ	Υ
MEC04-272	2003	Hamilton SJ	Ecotoxicol Environ Saf 56:201-210	Lepomis macrochirus	Bluegill	168141	Selenium	7782-49-2	0.92	mg/kg				
JA200	1982	Hilton, J.W., P.V. Hodson, and S.J. Slinger.	Comp Biochem Physiol 071C:49-55.	Oncorhynchus mykiss	Trout - Rainbow	161989	Selenium	7782-49-2	0.92	mg/kg	1	N	Υ	N
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	1.08	mg/kg	4	Y	Y	Y
URS159	1993	Lemly, A.D.	Aquat Toxicol 27:133-158	Lepomis macrochirus	Bluegill	168141	Selenium	7782-49-2	1.2	mg/kg	3	Υ	Y	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	1.3	mg/kg	4	Υ	Y	Y
JA336	1989	Ogle, R.S. and A.W. Knight	Arch Environ Contam Toxicol 18:795-803	Pimephales promelas	Fathead minnow	163517	Selenium	7782-49-2	1.4	mg/kg	4	Υ	Υ	N
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	1.6	mg/kg	6	Υ	Υ	Y
URS159	1993	Lemly, A.D.	Aquat Toxicol 27:133-158	Lepomis macrochirus	Bluegill	168141	Selenium	7782-49-2	1.6	mg/kg	3	Y	Y	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	1.7	mg/kg	20	Υ	Y	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	2.01	mg/kg	10	Y	Υ	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	2.16	mg/kg	4	Υ	Υ	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	2.88	mg/kg	6	Y	Y	Y
JB6	1992	Hermanutz, R.O., Allen, K.N., Roush, T.H., and S.F. Hedtke	Environ Tox & Chem 11: 217-224	Lepomis macrochirus	Bluegill	168141	Selenium	7782-49-2	4.6	mg/kg	6	Υ	Υ	N

Appendix D-5
Fish Tissue LOEDs for Selenium

Ref ID	QA Exp Method	QA Effects Method	QA Reps	Effect Class	Toxicity Measure	Exposure Route	Species Body Part	Species Start Lifestage	Species Habitat	Species Feeding Behavior	Comments
JA29	Υ	Υ	N	Growth	LOED	Absorption	Whole Body	Adult	Lakes, ponds, streams	Not Specified	Decrease weight gain.
URS98	Y	Y	Υ	Growth	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	invertebrates	Reduced Weight And Length Gain In 60 Days
URS98	Y	Y	Υ	Growth	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Weight And Length Gain In 30 Days
URS98	Y	Y	Y	Growth	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Weight And Length Gain In 60 Days
MEC04-272				Growth	LOED	NS	Whole Body	NS	Vegetation in lakes, ponds, streams	Carnivore-aquatic insects, fish, inverts	Review paper Cleveland et al 1993
JA200	Υ	Υ	Υ	Mortality	LOED	Ingestion	Carcass	Early Juvenile	Cool streams	Carnivore-juv fish, inverts	Significant increase in mortality. Residue measured at 12 weeks.
URS98	Y	Y	Υ	Mortality	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Survival In 90 Days
URS159	Υ	Υ	Υ	Growth	LOED	Combined	Whole Body	Immature	Vegetation in lakes, ponds, streams	Carnivore-aquatic insects, fish, inverts	Reduced weight to length ratio. Exposure at 4 degrees C. Concurrent epxosure to 5 ug inorganic Se/L in water and 5.1 ug Seleno-L-Methionine/g in food.
URS98	Y	Y	Υ	Mortality	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Survival In 90 Days
JA336	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Adult	Lakes, ponds, streams	Not Specified	Significant decrease in body weight gain.
URS98	Y	Y	Υ	Growth	LOED	Ingestion	Whole Body	Immature	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Weight Gain After 120 Days
URS159	Υ	Υ	Y	Mortality	LOED	Combined	Whole Body	Immature	Vegetation in lakes, ponds, streams	Carnivore-aquatic insects, fish,	35% Increase in mortality. Exposure at 4 degrees C. Concurrent epxosure to 5 ug inorganic Se/L in water and 5.1 ug Seleno-L-Methionine/g in food.
URS98	Y	Y	Υ	Growth	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Weight And Length Gain In 30 Days
URS98	Y	Y	Υ	Growth	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	invertebrates	Reduced Weight And Length Gain In 60 Days
URS98	Y	Υ	Y	Growth	LOED	Ingestion	Whole Body	Larval	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	invertebrates	Reduced Weight And Length Gain
URS98	Y	Υ	Y	Growth	LOED	Ingestion	Whole Body	Immature	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	invertebrates	Reduced Length After 120 Days
JB6	Υ	Y	Υ	Mortality	LOED	Combined	Whole Body	Adult	Vegetation in lakes, ponds, streams	Carnivore-aquatic insects, fish, inverts	

Appendix D-5
Fish Tissue LOEDs for Selenium

Ref ID	Year	Author	Publication Source	Species Scientific Name	Species Common Name	Species TSN Code	Analyte Name	CAS No	Conc_Wet	Conc Units	No. Reps	QA Stats Method	QA Chem Method	QA Chem DL
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	4.64	mg/kg	6	Υ	Υ	Y
URS98	1990	Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer and F.A. Bullard	Environ Tox & Chem 09:347-358	Oncorhynchus tshawytscha	Salmon - Chinook	161980	Selenium	7782-49-2	4.68	mg/kg	10	Υ	Υ	Y
URS15	1986	Bennett, W.N., A.S. Brooks and M.E. Boraas	Arch Environ Contam Toxicol 15:513-517	Pimephales promelas	Fathead minnow	163517	Selenium	7782-49-2	8.6	mg/kg	3	Υ	Υ	N
URS71	1996	Dobbs, M.G., D.S. Cherry and J. Cairns, Jr.	Environ Tox & Chem 15:340-347	Pimephales promelas	Fathead minnow	163517	Selenium	7782-49-2	12	mg/kg	2	Υ	Υ	N
URS71	1996	Dobbs, M.G., D.S. Cherry and J. Cairns, Jr.	Environ Tox & Chem 15:340-347	Pimephales promelas	Fathead minnow	163517	Selenium	7782-49-2	17.8	mg/kg	2	Υ	Y	N

Appendix D-5
Fish Tissue LOEDs for Selenium

Ref ID	QA Exp Method	QA Effects Method	QA Reps	Effect Class	Toxicity Measure	Exposure Route	Species Body Part	Species Start Lifestage	Species Habitat Species Feeding Behavior		Comments
URS98	Y	Y	Υ	Growth	LOED	Ingestion	Whole Body	Immature	Both coasts of Pacific; Monterey Bay and China north to Bering Straits, ascending all large streams, e.g., Columbia River	Feeds on variety of forms, e.g., larval, adult insects; small fish, invertebrates	Reduced Weight And Length Gain In 120 Days
URS98	Y	Y	Y	Mortality	LOED	Ingestion	Whole Body	Larval	<u>u</u> ,	Feeds on variety of forms, e.g., larval,	37% Reduced Survival In 60 Days
URS15	Υ	Υ	Υ	Growth	LOED	Ingestion	Whole Body	Larval	Lakes, ponds, streams	Not Specified	46% Reduction in weight gain. Larval uptake was 1.46 ng selenium per day.
URS71	Y	Y	Y	Growth	LOED	Combined	Whole Body	Larval	Lakes, ponds, streams	Not Specified	Significant reduction in weight.
URS71	Y	Y	Υ	Mortality	LOED	Combined	Whole Body	Larval	Lakes, ponds, streams	Not Specified	Mortality

APPENDIX E BENTHIC INVERTEBRATE RISK ASSESSMENT SUPPORTING INFORMATION

APPENDIX E-1 BENTHIC MODEL SUPPORTING INFORMATION

TABLE OF CONTENTS

1 BENTHI	C PREDICTIVE MODEL
1.1 Mod	lel Development Summary
	Assessing Model Performance
	Model Optimization4
	ernative Model Evaluation and Interspecies Comparisons
1.2.1	Interspecies Comparisons
	Alternative Model Evaluations
1.3 Re-	evaluation of COPC without SQGs
	lictive Model Uncertainty10
1.4.1	Non-Chemical Stressors1
1.4.2	Confounding Factors12
1.4.3	Analytical Uncertainty13
2 REFEREI	NCES15
List of Table	
Table E-1.1	Initial Reliability Analysis Results for Mean ERM-Q, Mean PEC-Q, Mean
	PEL-Q, PAH ESTBU, NOC ESTBU, and Metals ESBTU
Table E-1.2	Reliability Analysis Results for Initial and Optimized Mean PEL-Q and PEC-Q
Table E-1.3	Performance Assessment of Optimized Mean PEL-Q Model Applied to Test
	Species Other Than L. plumulosus
Table E-1.4	Performance Assessment of Initial SQG and Mechanistic Models for Test
	Species Other Than <i>L. plumulosus</i>

List of Exhibits

Exhibit E-1.1 Graphic Analysis of Benthic Risk Models

Exhibit E-1.2 Hypothesis Testing COPC Without SQGs

1 BENTHIC PREDICTIVE MODEL

In the BERA Work Plan, several predictive toxicity models were evaluated in order to define the relationship between surface sediment chemistry and toxic responses in the benthic community observed at the Site. The models identify the predictive relationship from different perspectives; although the goals of the models are similar: to develop a predictive relationship based on empirical data (i.e., sediment chemistry and toxicity test data) and to identify the principal chemical(s) that appear to best predict the relationship between sediment quality and toxicity. The models generally fall into two groups: 1) generic sediment quality guideline quotient (SQG-Q) models (Long et al. 1998; Ingersoll et al. 2001; MacDonald et al. 1996); and 2) mechanistic models (USEPA 2003d, 2008b).

The generic SQG-Q models rely on the use of threshold effect concentration (e.g., effects range median [ERM], probable effects concentration [PEC], and probable effect level [PEL]) quotients of mixtures of COPCs. Briefly, the quotient is the ratio of each chemical concentration to its threshold effect concentration (i.e., ERM, PEC, and PEL). Individual quotients can then be summed or averaged to evaluate the likelihood that a sediment sample would be expected to be toxic or non-toxic based on the narrative intent of the sediment quality guideline (SQG). Equilibrium partitioning (EqP; i.e., mechanistic) models are used to estimate equilibrium partitioning sediment benchmarks (ESB) to determine if a sample would be toxic based on partitioning between sediment and porewater, which represents the bioavailable fraction. These models rely on a toxic units (TU) approach whereby the toxicity of a class of related chemicals is considered additive. Sediment EqP models have been developed to estimate ESBs for PAH Equilibrium Partitioning Sediment Benchmark Toxic Unit (PAH ESBTU), non-ionic organics (NOC ESBTU), and divalent metals (metals ESBTU). Further details regarding the basis and application of the generic SQG-Q and EqP models used in the development of a predictive model can be found in Section 4.3 of the BERA Work Plan.

Each of these approaches are considered 'generalized' models in the sense that they rely on data from a compilation of sites and/or studies performed under a wide range of locations and conditions. Thus, they are intended to be representative of a broad range of conditions but may not account for potentially unique or confounding conditions that may be present at a

site to which the model is extrapolated. Thus, it is important to evaluate a potential model's performance using Site-specific data (i.e., bioassay data) in order to determine if a model provides an adequate degree of confidence in its interpretation of sediment toxicity at the Site.

The generalized models were evaluated using 51 samples with co-located sediment chemistry and toxicity test data. Samples were collected between September 2000 and August 2006. Treatment and development of the chemistry and toxicity database for model development are detailed in Section 4.3 of the BERA Work Plan. Datasets used in the benthic risk assessment are also summarized in Section 4 above.

In the BERA Work Plan, predictive model development focused on survival of *Leptocheirus plumulosus* in 10-day whole sediment bioassay tests as the effect endpoint for the model. This decision was based on several factors evaluated in detail in the BERA Work Plan. Briefly, they are:

- 1. This species was the most frequently tested with Site sediments (n = 43),
- 2. This species showed equal or greater sensitivity and good concordance with other species tested (see exception noted below), and
- 3. This species has been shown to be equally sensitive under acute, chronic, and subchronic test conditions and endpoints.

1.1 Model Development Summary

This section summarizes the steps taken in the BERA Work Plan to develop a predictive model for the benthic invertebrate risk characterization. Further details on the model development and optimization can be found there.

1.1.1 Assessing Model Performance

The predictive ability of SQG-Q and ESBTU models were initially evaluated by comparing the observed toxicity in the *L. plumulosus* bioassay test and the predicted toxicity based on chemistry data for each sample. Initially, the mean effects range median-quotient (ERM-Q), mean probable effect concentration-quotient (PEC-Q), mean probable effect level-quotient (PEL-Q), PAH ESBTU, NOC ESBTU, and metals ESBTU were included in the predictive ability analysis. Initially, all COPCs with SQGs available were included in the respective models. To evaluate the predictive ability of each, the following statistics were calculated initially assuming the effects threshold was equal to 1.0:

- False negative rate: number of samples incorrectly predicted as non-toxic/number of toxic samples.
- **False positive rate**: number of samples incorrectly predicted as toxic/number of non-toxic samples.
- Reliability: number of samples correctly predicted/total number of samples.

An SQG-Q or ESBTU model was deemed acceptable for use in the benthic invertebrate risk characterization based on the following acceptability criteria applied in similar evaluations of model performance (Long et al. 1998; MacDonald et al. 2000):

- False negative and false positive rates were less than 25 percent
- Overall reliability greater than or equal to 75 percent

Minimizing the false negative frequency in the model is important to ensure that the model is conservative and protective of the endpoint being evaluated. That is, the consequence of identifying a sample as non-toxic when in fact it is toxic results in a situation where adverse ecological effects may occur but are not identified in the risk characterization. Minimization of the false positive rate is important to avoid unnecessary potential remediation of locations that are not toxic; resulting in an unnecessary use of resources and possibly causing a greater impact (e.g., 'risk of remediation') to the Site than is warranted. Finally, maximizing the overall reliability ensures that a reasonable level of confidence can be placed in the predictive power of the model.

¹ Toxicity was defined as control adjusted survival less than 80 percent.

Table E-1.1 presents the acceptability criteria results for ERM-Q, PEC-Q, PEL-Q, PAH ESBTU, non-ionic COPCs ESBTU, and metals ESBTU models (graphical analysis results are provided in Exhibit E-1.1). All SQG-Q and ESBTU models had false negative rates below or near the acceptability criteria of 25 percent (0 to 28 percent) but only one, the PEC-Q model, had a false positive rate (16 percent) within the acceptability criteria limit. In addition, the PEC-Q model was the only quotient or ESBTU model with a reliability exceeding 75 percent.

Because three of the four SQG-Q or ESBTU models using the default threshold of 1.0 did not meet the acceptability criterion of less than 25 percent false positives, the effect of adjusting the threshold from 1.0 was evaluated. Similar to the approach used in other studies (Long et al. 2006), the thresholds were adjusted in an attempt to achieve acceptable error rates in comparison to toxicity test results for the Site. Effect thresholds were incrementally raised until acceptable error rates were achieved, if possible.

In summary, the mean PEL-Q model could meet the initial performance criteria and the model's overall performance was maximized by adjusting the effects threshold to 2.44. For the mean PEC-Q model, the performance criteria were nearly met at a threshold of 1.0 and further adjustment to the initial 1.0 threshold did not result in any increase in overall model performance. For all other models, the performance criteria were not met at a threshold of 1.0 or at any other threshold value; indicating they are not reliable predictors of toxicity at this Site. As a result, the mean PEL-Q and PEC-Q models were carried forward through a series of model optimization and calibration steps in an attempt to further simplify and improve the model performance and to identify the most appropriate model for characterizing risk to the benthic community.

1.1.2 Model Optimization

To optimize the mean quotient models that met the acceptability criteria (i.e., the PEC-Q and PEL-Q models), steps were taken to identify the COPCs that contributed significantly to the predictive ability of the model and to remove COPCs that did not, with the ultimate goal of improving the model's performance compared to the initial mean quotient model.

Several steps were taken to identify COPCs that contribute significantly to the model's performance. First, COPCs that did not show a significant difference in chemical concentrations between toxic and non-toxic sample groups were removed from the model. The reasoning for this step is that only COPCs that show some positive relationship with observed toxicity are likely to be significant contributors to toxicity at the Site. Next, mean quotients for PEC-Q and PEL-Q were recalculated using only those COPCs identified as having a significant difference between toxic and non-toxic datasets. More specifically, the PEC-Q was recalculated as the mean of the individual PEC-Q for total PAH, total PCB, and lead. The PEL-Q was recalculated using the mean of the individual PEL-Q for total PAH, total PCB, lead, and bis(2-ethylhexyl)phthalate (BEHP).

Finally, graphical analysis of the optimized models was performed similar to the original models. The false negative, false positive, and overall reliability of the optimized models were calculated for a range of quotient values and graphed. Graphs showing false negative, false positive and reliability of the optimized mean PEC-Q and PEL-Q are provided in Exhibit E-1.1. Both optimized models have lower error rates and higher reliability than their respective full mean quotient models (Table E-1.2). Again, the models were initially evaluated at a threshold of 1.0. At a threshold of 1.0 the optimized mean PEL-Q model still had an unacceptably high false positive and low reliability rate. Thus, an incremental evaluation of different thresholds was performed to identify the optimum threshold (e.g., highest reliability and lowest false positives and negatives). At a threshold of 1.6 the optimized mean PEL-Q had a higher overall reliability (81 percent vs. 79 percent), lower false negative (17 percent vs. 22 percent), and same false positive (20 percent) rate as the full mean PEL-Q model at threshold of 2.44. The optimized mean PEC-Q at a threshold of 1.21 had a higher overall reliability (88 percent vs. 79 percent) rate as the full model, a lower false negative rate (22 percent vs. 28 percent), and a lower false positive rate (4 percent vs. 16 percent).

Thus, the optimization step of removing COPCs from the quotient that did not contribute significantly to the predictive ability of the model did improve the performance of both models. In both cases, use of a toxicity threshold greater than 1.0 also significantly improved the predictive ability of both models.

1.2 Alternative Model Evaluation and Interspecies Comparisons

In the BERA Work Plan, an analysis of potential uncertainty associated with the model development was performed. Several sources of uncertainty were addressed, including:

- 1. Potential contribution of non-ionic organic COPCs to toxicity,
- 2. Metals bioavailability,
- 3. COPCs exceeding SQG that are not model variables,
- 4. COPCs without SQGs, and
- 5. Reliance on a single species and endpoint to assess community effects.

Nonetheless, USEPA and the Trustees requested that additional analysis of the bioassay and chemistry data be conducted during the BERA to further evaluate potential models that may incorporate additional Site COPCs and species into this LOE. To further assess the representativeness of a benthic toxicity model based on *L. plumulosus* to characterize risks to the benthic community, two types of additional analysis were performed. For the first analysis, the performance of the optimized mean PEL-Q model was assessed when applied to other test species (both individually and pooled) included in the Site toxicity database. For the second analysis, other SQG-Q and mechanistic models were evaluated using *A. bahia*, *N. arenaceodentata*, and *A. abdita*² individually and combined to determine if another model may more accurately predict the presence or absence of toxicity across all species tested in Site sediments. Results of this analysis are evaluated below.

1.2.1 Interspecies Comparisons

The purpose of this analysis was to evaluate whether the model could predict the presence or absence of toxicity for other species as determined by the empirical toxicity and sediment chemistry data for the Site. To address this issue, the rate of false negatives, false positives, and overall reliability of the optimized mean PEL-Q model was calculated using a threshold of 1.6 for *A. bahia*, *N. arenaceodentata*, *A. abdita*, and all species combined. Results are provided in Table E-1.3 (graphical analysis provided in Exhibit E-1.1).

At a threshold of 1.6 the rates of false negatives for *M. Americamysis* and *A. abdita* were less than or equal to the acceptability threshold of 25 percent while *N. arenaceodentata* had a

² See the BERA Work Plan for a discussion on the performance of other models in relation to *L. plumulosus*.

false negative rate of 67 percent. All species combined had a false negative rate of 27 percent which slightly exceeded the acceptability threshold of 25 percent. Thus, with the exception of *N. arenaceodentata*, a threshold of 1.6 would not result in an unacceptable number of locations (i.e., > 25 percent) being incorrectly identified as non-toxic to other test species based on the optimized mean PEL-Q. However, as can be seen from the rate of false positives (48 – 75 percent) for other test species, a threshold of 1.6 is likely to result in samples frequently being incorrectly identified as toxic based on the optimized mean PEL-Q.

Thus, the optimized mean PEL-Q threshold, or 1.6, is highly conservative with regard to predicting toxicity to other test species, with the possible exception of *N. arenaceodentata* (for the reasons described above) and likely result in frequent false positives. Based on the conservative nature of the threshold, the narrative intent of this threshold should be considered a threshold below which adverse effects to the benthic community would rarely be expected to occur.

If the observed responses from other test species are considered, there exists a range of optimized mean PEL-Q values above 1.6 where the response varies among the different species tested (i.e., adverse effects will be observed more frequently for all species but at different rates). At some threshold, adverse effects would consistently be expected for all species. This threshold would narratively be defined as a threshold above which adverse effects to the benthic community would be frequently expected to occur. Table E-1.3 provides the threshold(s) with the best predictive power for each species and all species combined (excepting *N. arenaceodentata*). If the acceptability criteria for false negatives, false positives, and reliability could not be met at any threshold, then the highest threshold that met the false negative acceptability criteria was chosen.

One species, A. bahia, met the acceptability criteria at a threshold range of 3.0 to 14.9. A. abdita did not meet all three criteria at any threshold but had an acceptable false negative rate up to a threshold of 2.3^3 . Similarly, for all species combined (excepting N. arenaceodentata), all three criteria were not met at any threshold but an acceptable false

_

³ Interestingly, at the next highest threshold of 2.4 had a false negative rate of 31 percent but the false positive rate decreased from 75 percent to 25 percent.

negative rate was observed up to a threshold of 2.3⁴. Finally, *N. arenaceodentata* had an acceptable false negative rate up to a threshold of 0.57. Based on this analysis, the threshold of 2.3 was selected as the upper bound threshold above which adverse effects to the benthic community would frequently be expected to occur.

1.2.2 Alternative Model Evaluations

For the second analysis, other SQG-Q and mechanistic models were evaluated using *A. bahia, N. arenaceodentata,* and *A. abdita*⁵ individually and combined to determine if another model may more accurately predict the presence or absence of toxicity across all species tested in Site sediments. The same methodology was applied that was used for development of the *L. plumulosus* model. The reliability, false positives, and false negatives were calculated at a range of thresholds for each test species. The threshold (or range of thresholds) that met the acceptability criteria for all three metrics was then determined and recorded in Table E-1.4. If all three acceptability criteria could not be met, the highest threshold meeting the false negative criteria was recorded. Exhibit E-1.1 provides the results of the graphical analysis used to evaluate these alternative models.

None of the other SQG-Q or mechanistic models met the acceptability criteria for *N. arenaceodentata* or *A. abdita*. The PEL-Q, PEC-Q, and ERM-Q models (full models with all available COPCs) met the acceptability criteria for *A. bahia* at a range of thresholds.

1.3 Re-evaluation of COPC without SQGs

Section 4.4.3.4 of the BERA Work Plan evaluated the potential risk associated with benthic invertebrate COPCs that were not included in the model development due to their lack of SQGs or inclusion in mechanistic models. USEPA and the Trustees requested additional analyses be considered during the BERA to provide a more robust assessment of these COPCs. They include:

- Ammonia
- Barium

⁴ As with *A. abdita*, the next highest threshold of 2.35 for the combined species model had a false negative rate of 26 percent but the false negative rate decreased from 37 percent to 22 percent.

⁵ See the BERA Work Plan for a discussion on the performance of other models in relation to *L. plumulosus*.

- Hexachlorobutadiene
- Benzidine
- Bis(2-chloroisopropyl)ether
- Hexachlorobenzene
- Total xylene
- Isopropylbenzene
- n-Butylbenzene
- n-Propylbenzene
- sec-Butylbenzene
- tert-Butylbenzene

In the BERA Work Plan, concentrations of these COPCs in sediment were compared to an arbitrary threshold of ten times their Site average (based on the exposure assessment dataset).

To support this initial analysis, two evaluations were performed in the BERA. The first used the sediment chemistry data from the toxicity model dataset to statistically evaluate differences in the concentration of these COPCs in the toxic versus non-toxic dataset based on empirical observations of toxicity to L. plumulosus. This analysis followed the same approach as the model optimization step for COPCs included in the optimized mean PEL-Q development (Section 4.4.2 of the BERA Work Plan). In this analysis, COPCs with greater than 40 percent frequency of detection were tested using the non-parametric Wilcoxon-Mann-Whitney (WMW) while Gehan's test was used for COPCs with less than 40 percent frequency of detection as recommended by USEPA's ProUCL Version 4.0 Technical Guide (2007). Both tests used ranks of the data to test whether the measurements from one sample group tend to be consistently larger than those in the second group. Samples with less than 8 detected observations in either the toxic or non-toxic dataset could not be evaluated. For each test, α was set at 0.05 and the null and alternative hypotheses are:

$$H_o: \widetilde{\mu}_{TOXIC} \leq \widetilde{\mu}_{NONTOXIC}$$

$$H_A: \widetilde{\mu}_{TOXIC} > \widetilde{\mu}_{NONTOXIC}$$

where:

 H_0 = null hypothesis

H_A = alternative hypothesis

 $\tilde{\mu}_{TOXIC}$ = Toxic sample group median

 $\tilde{\mu}_{NONTOXIC}$ = Non-toxic sample group median

Results are provided in Exhibit E-1.2. Based on the low frequency of detection or low number of results, ammonia, total xylene, n-butylbenzene, n-propylbenzene, secbutylbenzene, and tert-butylbenzene were not evaluated.

With one exception, none of the remaining COPCs (hexachlorobutadiene, hexachlorobenzene, benzidine, and bis(2-chloroisopropyl)ether) demonstrated a significant difference between toxic and non-toxic samples. This analysis confirms the previous assessment of these COPCs which concludes that they show no significant association with the observed toxicity in Site sediments and that exclusion from the benthic model does not significantly compromise its predictive power.

Barium was the only COPC to demonstrate a significant difference between toxic and non-toxic sample groups. This may be due to simple covariance of barium with other COPCs that may be causing toxicity. A review of the available literature indicate that relevant bulk sediment toxicity SQG have not been established for barium. However, concentrations in porewater were below the corresponding water quality screening benchmark (Anchor 2008b).

1.4 Predictive Model Uncertainty

The following subsections describe three important potential sources of uncertainty: 1) the potential contribution of non-chemical stressors that may erroneously imply causality of toxicity by the COPC used in the current model; 2) the uncertainty related to confounding factors in the historical sediment bioassay investigations; and 3) analytical uncertainty related to PCBs in bulk sediment. These three sources of uncertainty may compromise the applicability of the current benthic toxicity model for risk characterization and risk management applications. If non-chemical stressors and confounding factors are significant

but unaccounted for, the result is likely to be an overly conservative characterization of risk due to Site COPCs that could potentially result in significant remedial actions that may ultimately not be warranted and that would not otherwise be considered. Analytical uncertainty in the toxicity dataset affects any extrapolation related to the concentration of COPC in sediments without empirical toxicity data.

1.4.1 Non-Chemical Stressors

Within a mesohaline benthic community there are a variety of non-chemical stressors present. These non-chemical stressors may influence the apparent response of organisms or communities observed in typical measurement endpoints selected for ecological risk assessment. These non-chemical stressors can increase the apparent response by test organisms to COPC and may often be the primary cause of adverse effects in toxicity tests. They may also play a significant role in shaping the benthic community and influence many of the benthic community properties (i.e., abundance, diversity, etc.) that are evaluated in ecological risk assessments. These non-chemical stressors may include, but are not limited to, grain size, organic carbon, salinity, and temperature. These stressors may be natural or anthropogenic (e.g., nutrient enrichment) but unrelated to historical or ongoing releases of COPC to a system. However, if non-chemical stressors are not accounted for when evaluating benthic invertebrate measurement endpoints, their effects may be erroneously attributed to chemical stressors (e.g., COPCs) that are measured.

Controlling or quantifying effects of non-chemical stressors when evaluating benthic invertebrate measurement endpoints can be challenging. Nonetheless, there are effective methods to address these issues. Arguably the most important is to incorporate an appropriate reference area into any testing program. An effective reference site would duplicate all the conditions, including non-chemical stressors, of a study site and would provide a point of reference to account for the potential effect of these stressors in the absence of Site COPCs. The incremental effect of these non-chemical stressors can be quantitatively evaluated and incorporated into the observed responses of Site data (e.g., reference envelope approach for bioassay endpoints [Hunt et al. 1998; Smith 2002]).

Unfortunately, the historical Site bioassay and community dataset does not include reference area data⁶. This is a significant data gap. Without appropriate reference area observations for comparison, overly conservative conclusions may be reached based on the responses observed in sediment toxicity tests when compared to levels of chemical stressors in sediment at the Site. Moreover, assessing the incremental contribution of non-chemical stressors to effects observed in Site sediments cannot be performed retroactively. Only additional, concurrent testing and observations of Site and reference area sediments are appropriate to adequately assess the significance of non-chemical stressors on benthic invertebrate bioassay measurement endpoints.

1.4.2 Confounding Factors

Confounding factors (CFs) are characteristics or features that may be present in laboratory sediment bioassays that are not related to the chemical stressors of interest but contribute to observed biological effects (e.g., mortality). These confounding factors can be divided into matrix features and non-matrix features:

- Matrix features vary due to laboratory conditions; including such characteristics as ammonia or sulfide concentrations, the quality of organic carbon in sediment, and acclimation of sediments to test conditions.
- Non-matrix features include selection of appropriate test organisms, test organism health, and test organism acclimation and handling techniques.

Toxicity testing may not provide effective data for assessing and selecting risk management strategies unless it recognizes the potential for CFs and incorporates methods to explain or remove the effects of CF's during testing. Failure to account for CFs may lead to misinterpretation of the causal factors of toxicity; leading to risk characterization focusing on irrelevant or marginally relevant chemical stressors (i.e., COPC) or overestimating the magnitude of risk associated with Site COPC. Awareness of the potential for CFs to impact

⁶ Benthic community analysis performed by Parsons Engineering (2002, 2004) of the historical data did compare Site results to 'non-degraded sites in Galveston Bay'. However, these sites were sampled under a completely different program separated in time from the Patrick Bayou sampling efforts and should not be considered a true reference area dataset.

toxicity data interpretation allows the inclusion of those factors in the assessment which, if properly managed, could provide a more robust characterization of risk⁷.

Typical CFs that may be present during sediment toxicity tests include:

- 1. Test organism acclimation to test conditions
- 2. Test organism health prior to test initiation
- 3. Sediment acclimation to test conditions
- 4. Water quality in test chambers

1.4.3 Analytical Uncertainty

Total PCBs are included in the model developed using the paired toxicity and chemistry database. These data were generated outside of the Patrick Bayou Superfund RI and were reviewed for use in the RI during the historical data review process (Anchor 2007b). Analysis of PCBs for samples in the toxicity dataset was performed on an Aroclor basis (i.e., USEPA Method 1656). In each case, only PCBs consistent with Aroclor 1248 standards were identified and reported. More specifically, one tri-chlorinated congener peak and four tetrachlorinated congener peaks were chosen to quantify the Aroclor 1248 results for samples collected in September 2000 and April 2001 (Parsons Engineering 2002)⁸. Thus, PCB congeners outside of the profile of an Aroclor 1248 standard are not reported.

For the RI surface sediment dataset being used to characterize risk in the BERA (Section 4), PCBs were analyzed on a congener specific basis (i.e., USEPA Method 1668). As a result, the full range of congeners (n = 209) were quantified and reported. Thus total PCB congener values based on all congeners were not consistent with the toxicity dataset and would result in an overestimation of PCBs relative to the methods used in the toxicity dataset.

The uncertainty related to the differences in the two methods is biased conservative. That is, estimates of PCBs based on the Aroclor analytical methods used are likely to underestimate

⁷ It should be noted that the real level of risk in the environment may not be accurately predicted from laboratory tests. These standardized tests expose organisms in a controlled environment, where temperature, oxygen, and salinity stay constant.

⁸ Commercial Aroclor 1248 mixtures typically consist of 70 percent tri- and tetra-chlorinated homologs (USEPA 1996) (USEPA n.d.)

the concentration of congeners in an environmental sample. Aroclor analysis is based on a pattern-recognition technique and requires significant professional judgment on the part of the analyst to identify these patterns and match them to an expected standard Aroclor mixture profile. These judgments are made using only a subset of the information (e.g., less than 28 congeners). In contrast, congener-specific analysis is much less subjective and can reliably identify all 209 congeners. As a result, comparison between data based on Aroclor versus congener specific analysis is problematic. It is expected that this uncertainty may overestimate benthic risks associated with PCBs.

2 REFERENCES

- Anchor, 2007b. Work Package 2 Vertical Profiling Hydrodynamic Field Data Collection and Contaminant Source Evaluation. Patrick Bayou Superfund Site. Deer Park, Texas.

 Prepared for USEPA and Patrick Bayou Joint Defense Group. April 2007.
- Anchor, 2008b. Amendment to the Selection of Contaminants of Potential Concern for Ecological Risk Assessment Technical Memorandum. Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. October 2008.
- Anchor QEA, 2011a. *Final Baseline Ecological Risk Assessment Work Plan.* Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. May 2011.
- Anchor QEA, 2011b. *Final Upstream Patrick Bayou Characterization Sampling and Analysis Plan.* Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. June 2011.
- Hunt J.W., B.S. Anderson, B.M. Phillips, J. Newman, R.S. Tjeerdema, K. Taberski, C.J. Wilson, M. Stephenson, H.M. Puckett, R. Fairey, J. Oakden, 1998. Sediment Quality and Biological Effects in San Francisco Bay. Bay protection and toxic cleanup program. Final technical report. California State Water Resources Control Board, San Francisco Bay Regional Water Quality Control Board, California Dept. of Fish and Game, California State University and University of California, Santa Cruz.
- Ingersoll C.G., D.D. MacDonald, N. Wang, J.L. Crane, L.J. Field, P.S. Haverland, N.E.Kemble, R.A. Lindskoog, C.G. Severn, and D.E. Smorong, 2001. Predictions ofSediment Toxicity Using Consensus-Based Freshwater Sediment Quality Guidelines.Archives of Environmental Contamination and Toxicology 41:8-21.
- Long, E.R., L.J. Field, and D.D. MacDonald, 1998. Predicting Toxicity in Marine Sediments with Numerical Sediment Quality Guidelines. *Environmental Toxicology and Chemistry* 17(4): 714 727.
- MacDonald, D.D., R.S. Carr, F.D. Calder, E.R. Long, and C.G. Ingersoll, 1996. Development and Evaluation of Sediment Quality Guidelines for Florida Coastal Waters. *Ecotoxicology* 5:253–278.

- MacDonald D.D., C.G. Ingersoll, T.A. Berger, 2000. Development and Evaluation of consensus based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* 39(5):20-31.
- Parsons et al., 2002. Assessment of Sediment Toxicity and Quality in Patrick Bayou, Segment 1006, Harris County, Texas. Prepared for Patrick Bayou TMDL Lead Organization.
- Parsons et al., 2004. Assessment of Sediment Toxicity and Quality in Patrick Bayou, Segment 1006, Harris County, Texas. Prepared for Patrick Bayou TMDL Lead Organization.
- Smith R.W., 2002. The use of random-model tolerance intervals in environmental monitoring and regulation. *J Agric Biol Environ Statist* 7(1):74-94.
- USEPA, 1996. PCBs: Cancer dose-response assessment and application to environmental mixtures. Washington, DC: Office of Research and Development.
- USEPA, 2003d. *Technical Basis for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms: Non-Ionic Organics.*Draft. USEPA-600-R-02-014. Office of Research and Development, Washington, DC.
- USEPA, 2008b. *Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms.* Compendium of Tier 2 values for nonionic organics. USEPA/600/R-02/016. Office of Research and Development, Washington, DC.
- USEPA, (nod). Plots of Aroclor composition. Retrieved March 21, 2012, from http://www.epa.gov/epawaste/hazard/tsd/pcbs/pubs/aroclorplots.pdf

TABLES

Table E-1.1
Initial Reliability Analysis Results for Mean ERM-Q, Mean PEC-Q, Mean PEL-Q, PAH ESBTU, NOC ESBTU, and Metals ESBTU

	Effect Threshold = 1.0							
	ERM-Q							
False Negatives	6%							
False Positives	64%							
Reliability	60%							
	PEC-Q							
False Negatives	28%							
False Positives	16%							
Reliability	79%							
	PEL-Q							
False Negatives	0%							
False Positives	80%							
Reliability	53%							
	PAH ESBTU							
False Negatives	17%							
False Positives	68%							
Reliability	53%							
	NOC ESBTU							
False Negatives	100%							
False Positives	0%							
Reliability	52%							
Metals ESBTU								
False Negatives	16%							
False Positives	100%							
Reliability	49%							
M = 4 = = .								

Notes:

Shaded cells meet initial acceptability criteria

ERM-Q - Effect Range Median quotient

PEC-Q - probable effects concentration quotient

PEL-Q = probable effect limit quotient

PAH ESBTU - polynuclear aromatic hydrocarbon equilibrium screening benchmark toxic unit

NOC ESBTU - non-ionic COPC equilibrium screening benchmark toxic unit

Table E-1.2
Reliability Analysis Results for Initial and Optimized Mean PEL-Q and PEC-Q

SQG	M	lean PEC-Q	Mean PEL-Q			
Model	Initial	Optimized	Initial	Optimized		
COPC	All	PAH, PCB, Lead	All	PAH, PCB, BEHP, Lead		
Threshold	1.00	1.21	2.44	1.56		
False Negatives	28%	22%	22%	17%		
False Positives	16%	4%	20%	20%		
Reliability	79%	88%	79%	81%		

Notes:

PEC-Q - probable effects concentation quotient

PEL-Q - probable effect level quotient

Table E-1.3
Performance Assessment of Optimzed Mean PEL-Q Model Applied to Test Species Other Than *L. plumulosus*

Toxicity Metric	Test Group	Predictive Performance ^a	Optimized Mean PEL-Q @ 1.6	Optimi	zed Mea	an PEL-Q ^b
Control Adjusted Survival < 80%	M. americamysis	Reliability	55%	85%		90%
	n=20	False Negative	13%	13%		13%
		False Positive	67%	17%		8%
		Threshold Number		3.0		15
	A. abdita	Reliability	65%	65%		70% ^c
	n=20	False Negative	25%	25%		31%с
		False Positive	75%	75%		25%c
		Threshold Number		2.3		2.4
	N. arenaceodentata	Reliability	47%		39%	
	n=36	False Negative	67%		22%	
		False Positive	48%		74%	
		Threshold Number			0.57	
	All species ≠ N. arenaceodentata	Reliability	64%	70%		76%c
	n=83	False Negative	27%	24%		26%c
		False Positive	43%	37%		22%c
		Threshold Number		2.3		2.4

Notes:

a Acceptability criteria for predictive performance:

Reliability ≥ 75% False negatives ≤ 25%

False positives ≤ 25%

b If the acceptability criteria could not be met at any threshold, the highest threshold that met the false negative acceptability criteria is provided c Although these thresholds did not meet the performance crteria they are provided for context. See text for discussion.

Table E-1.4 Performance Assessment of Initial SQG and Mechanistic Models for Test Species Other Than L. plumulosus

Toxicity Metric	Test Group	Predictive Performance ^a	Mean PEL-Q ^b	Mean PEC-Q ^b	Mean ERM-Q ^b	PAH-ESBTU	NOC-ESBTU	PAH ESBTU + NOC ESBTU	
Control Adjusted Survival < 80%	M. americamysis	Reliability	90% 85%	85% 90%	85% 85%	55%	40%	55%	
	n=20	False Negative	13% 25%	13% 13%	13% 25%	0%	0%	0%	
		False Positive	8% 8%	17% 8%	17% 8%	75%	100%	75%	
		Threshold Number	5.15 30.48	1.54 6.07	5.56 41.41	0.45	0	0.45	
	A. abdita	Reliability	65%	65%	65%	75%	80%	75%	
	n=20	False Negative	25%	25%	25%	13%	0%	13%	
		False Positive	75%	75%	75%	75%	100%	75%	
		Threshold Number	3.7	1.2	4.67	0.45	0	0.45	
	N. arenaceodentata	Reliability	28%	31%	28%	50%	50%	44%	
	n=36	False Negative	22%	22%	22%	22%	22%	22%	
		False Positive	89%	85%	89%	59%	59%	67%	
		Threshold Number	1.09	0.47	0.63	1.24	0.007	1.26	
	All species ≠ N. arenaceodentata	Reliability	69%	70%	66%	51%	51%	51%	
	n=83	False Negative	24%	24%	19%	21%	0%	21%	
		False Positive	39%	37%	49%	78%	100%	78%	
1		Threshold Number	2.94	1.12	2.05	0.69	0	0.71	


a Acceptability criteria for predictive performance: Reliability ≥ 75%


False negatives ≤ 25% False positives ≤ 25%


b If the acceptability criteria could not be met at any threshold, the highest threshold that met the false negative acceptability criteria is provided.

EXHIBIT E-1.1 GRAPHIC ANALYSIS OF BENTHIC RISK MODELS

Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models









Exhibit E-1.1
Graphical Analysis of Benthic Risk Models











Exhibit E-1.1
Graphical Analysis of Benthic Risk Models






EXHIBIT E-1.2 HYPOTHESIS TESTING COPC WITHOUT SQGS

From File: C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02 BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model worksheet for ProUCL.wst

Summary Statistics for Raw Data Sets With NDs Using Detected Data Only

						Raw S	Statistics Usin	g Detected O	bservations		
Variable	Num Ds	NumNDs	% NDs	Minimum	Maximum	Mean	Median	SD	MAD/0.675	Skewness	CV
Ammonia (0)	2	0	0.00%	17.5	39.7	28.6	28.6	15.7	16.46	N/A	0.549
Ammonia (1)	2	0	0.00%	134	362	248	248	161.2	169	N/A	0.65
Barium (0)	25	0	0.00%	60.8	447	277.3	270	100.3	105.3	-0.101	0.362
Barium (1)	18	0	0.00%	218	23500	1700	365.5	5443	155.7	4.235	3.201
Hexachlorobenzene (0)	15	10	40.00%	13.8	201000	32934	3400	58847	5020	2.201	1.787
Hexachlorobenzene (1)	12	6	33.33%	39.9	8790	2176	579.4	2953	798.3	1.407	1.357
Hexachlorobutadiene (0)	13	12	48.00%	57.8	222000	32940	7260	64051	10678	2.629	1.944
Hexachlorobutadiene (1)	10	8	44.44%	55.7	23300	5416	425.2	8026	522.5	1.549	1.482
Benzidine (0)	6	19	76.00%	103	503	261.5	187	169.7	94.89	0.834	0.649
Benzidine (1)	10	8	44.44%	119	27000	3148	464	8387	480.4	3.154	2.664
bis(2-Chloroisopropyl) ether (0)	10	15	60.00%	0.257	410	88.07	36.1	126.4	29.68	2.233	1.435
bis(2-Chloroisopropyl) ether (1)	13	5	27.78%	12.4	3320	661.4	417	957.8	511.5	2.214	1.448
Total Xylene (0)	0	21	100.00%	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Total Xylene (1)	4	12	75.00%	11	1900	771.3	587	841.6	664.2	0.976	1.091
Isopropylbenzene (Cumene) (0)	3	18	85.71%	15	22.9	18.3	17	4.107	2.965	1.282	0.224
Isopropylbenzene (Cumene) (1)	8	8	50.00%	25.4	29000	7282	1620	11730	2248	1.489	1.611
n-Butylbenzene (0)	0	21	100.00%	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
n-Butylbenzene (1)	2	14	87.50%	176	699	437.5	437.5	369.8	387.7	N/A	0.845
n-Propylbenzene (0)	0	21	100.00%	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
n-Propylbenzene (1)	4	12	75.00%	66	1400	529	325	605.1	299.5	1.561	1.144
sec-Butylbenzene (0)	0	21	100.00%	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
sec-Butylbenzene (1)	4	12	75.00%	10	383	147.9	99.35	162.5	73.39	1.572	1.098
tert-Butylbenzene (0)	0	21	100.00%	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
tert-Butylbenzene (1)	3	13	81.25%	40.9	1160	436.3	108	627.6	99.48	1.71	1.439

NOTES:

Number in parenthesis after chemical name denotes toxic and non-toxic datasets based on L. plumulosus survival; (0) = nontoxic, (1) = toxic

Wilcoxon-Mann-Whitney Site vs Background Comparison Test for Full Data Sets without NDs

User Selected Options

From File C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02

BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model

worksheet for ProUCL.wst

Full Precision OFF

Confidence Coefficient 95% Substantial Difference 0

Selected Null Hypothesis Site or AOC Mean/Median Less Than or Equal to Background Mean/Median (Form 1)

Alternative Hypothesis Site or AOC Mean/Median Greater Than Background Mean/Median

Area of Concern Data: Barium(1)
Background Data: Barium(0)

Raw Statistics

	Site	Background
Number of Valid Observations	18	25
Number of Distinct Observations	18	25
Minimum	218	60.8
Maximum	23500	447
Mean	1700	277.3
Median	365.5	270
SD	5443	100.3
SE of Mean	1283	20.07

Wilcoxon-Mann-Whitney (WMW) Test

H0: Mean/Median of Site or AOC <= Mean/Median of Background

Site Rank Sum W-Stat 503.5 WMW Test U-Stat 2.634 WMW Critical Value (0.050) 1.645

P-Value 0.00422

Wilcoxon-Mann-Whitney Site vs Background Comparison Test for Data Sets with Non-Detects

User Selected Options

From File C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02

BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model

worksheet for ProUCL.wst

Full Precision OFF

Confidence Coefficient 95% Substantial Difference (S) 0

Selected Null Hypothesis Site or AOC Mean/Median Less Than or Equal to Background Mean/Median (Form 1)

Alternative Hypothesis Site or AOC Mean/Median Greater Than Background Mean/Median

Area of Concern Data: Benzidine(1)
Background Data: Benzidine(0)

Raw Statistics

	Site	Background
Number of Valid Data	18	25
Number of Non-Detect Data	8	19
Number of Detect Data	10	6
Minimum Non-Detect	41.9	31
Maximum Non-Detect	417	408
Percent Non detects	44.44%	76.00%
Minimum Detected	119	103
Maximum Detected	27000	503
Mean of Detected Data	3148	261.5
Median of Detected Data	464	187
SD of Detected Data	8387	169.7

Wilcoxon-Mann-Whitney Site vs Background Test

All observations <= 417 (Max DL) are ranked the same

Wilcoxon-Mann-Whitney (WMW) Test

H0: Mean/Median of Site or AOC <= Mean/Median of Background

Site Rank Sum W-Stat 456
WMW Test U-Stat 1.465
WMW Critical Value (0.050) 1.645
P-Value 0.0715

Wilcoxon-Mann-Whitney Site vs Background Comparison Test for Data Sets with Non-Detects

User Selected Options

From File C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02

BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model

worksheet for ProUCL.wst

Full Precision OFF

Confidence Coefficient 95% Substantial Difference (S) 0

Selected Null Hypothesis Site or AOC Mean/Median Less Than or Equal to Background Mean/Median (Form 1)

Alternative Hypothesis Site or AOC Mean/Median Greater Than Background Mean/Median

Area of Concern Data: bis(2-Chloroisopropyl) ether(1)
Background Data: bis(2-Chloroisopropyl) ether(0)

Raw Statistics

	Site	Background
Number of Valid Data	18	25
Number of Non-Detect Data	5	15
Number of Detect Data	13	10
Minimum Non-Detect	16.4	12.1
Maximum Non-Detect	417	660
Percent Non detects	27.78%	60.00%
Minimum Detected	12.4	0.257
Maximum Detected	3320	410
Mean of Detected Data	661.4	88.07
Median of Detected Data	417	36.1
SD of Detected Data	957.8	126.4

Wilcoxon-Mann-Whitney Site vs Background Test

All observations <= 660 (Max DL) are ranked the same

Wilcoxon-Mann-Whitney (WMW) Test

H0: Mean/Median of Site or AOC <= Mean/Median of Background

Site Rank Sum W-Stat 446

WMW Test U-Stat 1.219

WMW Critical Value (0.050) 1.645

P-Value 0.111

Wilcoxon-Mann-Whitney Site vs Background Comparison Test for Data Sets with Non-Detects

User Selected Options

HCB C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02

BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model worksheet

for ProUCL.wst

Full Precision OFF

Confidence Coefficient 95%

Substantial Difference (S) 0

Selected Null Hypothesis Site or AOC Mean/Median Less Than or Equal to Background Mean/Median (Form 1)

Alternative Hypothesis Site or AOC Mean/Median Greater Than Background Mean/Median

Area of Concern Data: Hexachlorobenzene(1)
Background Data: Hexachlorobenzene(0)

Raw Statistics

	Site	Background
Number of Valid Data	18	25
Number of Non-Detect Data	6	10
Number of Detect Data	12	15
Minimum Non-Detect	377	24.9
Maximum Non-Detect	417	400
Percent Non detects	33.33%	40.00%
Minimum Detected	39.9	13.8
Maximum Detected	8790	201000
Mean of Detected Data	2176	32934
Median of Detected Data	579.4	3400
SD of Detected Data	2953	58847

Wilcoxon-Mann-Whitney Site vs Background Test

All observations <= 417 (Max DL) are ranked the same

Wilcoxon-Mann-Whitney (WMW) Test

H0: Mean/Median of Site or AOC <= Mean/Median of Background

Site Rank Sum W-Stat 364 WMW Test U-Stat -0.8 WMW Critical Value (0.050) 1.645

P-Value 0.212

Wilcoxon-Mann-Whitney Site vs Background Comparison Test for Data Sets with Non-Detects

User Selected Options

From File C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02

BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model

worksheet for ProUCL.wst

Full Precision OFF

Confidence Coefficient 95% Substantial Difference (S) 0

Selected Null Hypothesis Site or AOC Mean/Median Less Than or Equal to Background Mean/Median (Form 1)

Alternative Hypothesis Site or AOC Mean/Median Greater Than Background Mean/Median

Area of Concern Data: Hexachlorobutadiene(1)
Background Data: Hexachlorobutadiene(0)

Raw Statistics

	Site	Background
Number of Valid Data	18	25
Number of Non-Detect Data	8	12
Number of Detect Data	10	13
Minimum Non-Detect	27.6	20.5
Maximum Non-Detect	417	408
Percent Non detects	44.44%	48.00%
Minimum Detected	55.7	57.8
Maximum Detected	23300	222000
Mean of Detected Data	5416	32940
Median of Detected Data	425.2	7260
SD of Detected Data	8026	64051

Wilcoxon-Mann-Whitney Site vs Background Test

All observations <= 417 (Max DL) are ranked the same

Wilcoxon-Mann-Whitney (WMW) Test

H0: Mean/Median of Site or AOC <= Mean/Median of Background

Site Rank Sum W-Stat 345.5 WMW Test U-Stat -1.256 WMW Critical Value (0.050) 1.645 P-Value 0.105

Wilcoxon-Mann-Whitney Site vs Background Comparison Test for Data Sets with Non-Detects

User Selected Options

From File C:\Users\jkase\Documents\AnchorQEA\Projects_Active\Patrick Bayou\Active\Task 01_02

BERA\Analysis\Benthic\Sediment models\Feb 2012 Evaluation\Individual COPC results for model

worksheet for ProUCL.wst

Full Precision OFF

Confidence Coefficient 95% Substantial Difference (S) 0

Selected Null Hypothesis Site or AOC Mean/Median Less Than or Equal to Background Mean/Median (Form 1)

Alternative Hypothesis Site or AOC Mean/Median Greater Than Background Mean/Median

Area of Concern Data: Silver(1)
Background Data: Silver(0)

Raw Statistics

	Site	Background
Number of Valid Data	18	25
Number of Non-Detect Data	0	2
Number of Detect Data	18	23
Minimum Non-Detect	N/A	0.0695
Maximum Non-Detect	N/A	0.0745
Percent Non detects	0.00%	8.00%
Minimum Detected	0.173	0.0739
Maximum Detected	1.53	1.57
Mean of Detected Data	0.612	0.402
Median of Detected Data	0.579	0.269
SD of Detected Data	0.349	0.358

Wilcoxon-Mann-Whitney Site vs Background Test

All observations <= 0.0745 (Max DL) are ranked the same

Wilcoxon-Mann-Whitney (WMW) Test

H0: Mean/Median of Site or AOC <= Mean/Median of Background

Site Rank Sum W-Stat 506.5 WMW Test U-Stat 2.708 WMW Critical Value (0.050) 1.645 P-Value 0.00338

APPENDIX E-2 DATA AND CALCULATIONS FOR THE BENTHIC TOXICITY LINE OF EVIDENCE

Appendix E-2.1
Sediment Toxicity Data Evaluated in Benthic Risk Assessment

	Data			Percent Cont	rol Adjusted Sur	<i>i</i> ival	Number of	Number of	Proportion
Location	Source ^a	Sample Date	Americamysis bahia	Ampelisca abdita	Leptocheirus plumulosus	Neanthes arenaceodentata	Toxic Results ^b	Bioassay Tests	Toxic
2.5	TMDL	Sep-2000			90	96	0	2	0.00
2.5	TMDL	Apr-2001	-		87	94	0	2	0.00
3	EPA/TCEQ	Aug-2003	93	87		-	0	2	0.00
3	EPA/TCEQ	Aug-2006	10	0			2	2	1.00
3	TMDL	Sep-2000			76	109	0	2	0.00
3	TMDL	Apr-2001			0	86	1	2	0.50
3	TMDL	Aug-2003	98	68	95		0	3	0.00
4A	EPA/TCEQ	Aug-2003	79	0			1	2	0.50
4A	EPA/TCEQ	Aug-2006	93	0			1	2	0.50
4A	TMDL	Sep-2000			89	96	0	2	0.00
4A	TMDL	Apr-2001			6	69	1	2	0.50
4A	TMDL	Aug-2003	60	1	30		2	3	0.67
5	TMDL	Sep-2000			103	83	0	2	0.00
5	TMDL	Apr-2001			0	94	1	2	0.50
6A	EPA/TCEQ	Aug-2003	21	0			2	2	1.00
6A	EPA/TCEQ	Aug-2006	100	67			0	2	0.00
6A	TMDL	Sep-2000			25	109	1	2	0.50
6A	TMDL	Apr-2001			0	61	1	2	0.50
6A	TMDL	Aug-2003	32	0	31		3	3	1.00
E	TMDL	Sep-2000			104	87	0	2	0.00
Е	TMDL	Apr-2001			57	98	1	2	0.50
G	TMDL	Sep-2000			104	91	0	2	0.00
G	TMDL	Apr-2001			93	37	1	2	0.50
Q	TMDL	Sep-2000			80	100	0	2	0.00
Q	TMDL	Apr-2001			4	82	1	2	0.50
S	TMDL	Sep-2000			100	91	0	2	0.00
S	TMDL	Apr-2001			66	86	0	2	0.00
Т	TMDL	Sep-2000			91	104	0	2	0.00
T	TMDL	Apr-2001			72	78	0	2	0.00
U	TMDL	Sep-2000			98	91	0	2	0.00
U	TMDL	Apr-2001			83	69	0	2	0.00
V	TMDL	Sep-2000			102	70	0	2	0.00
V	TMDL	Apr-2001			79	94	0	2	0.00

Appendix E-2.1
Sediment Toxicity Data Evaluated in Benthic Risk Assessment

ſ		Data			Percent Cont	rol Adjusted Surv	/ival	Number of	Number of	Proportion
	Location	Source	Sample Date	Americamysis	Ampelisca	Leptocheirus	Neanthes	Toxic Results ^b	Bioassay Tests	Toxic ^c
L		Source		bahia	abdita	plumulosus	arenaceodentata	TOXIC RESUITS	bloassay rests	TOXIC

^a Data Source References:


TMDL - Parsons et al. (2002, 2004)

EPA/TCEQ - L. Broach (no date)

^b Control adjusted survival < 60%

^c Proportion Toxic = (Number of toxic results)/(Number of bioassay tests)

Appendix E-2.2
Flowchart Describing Calculation of Average Relative % Difference from Median Optimized Mean PEL-Q



PEL-Q - probable effect level quotient $%\Delta$ - percent difference

Appendix E-2.3
Sediment Chemistry and PEL-Q Values by Sample

Г	Comple Tatal DCD Tatal DCD Tatal DAU Tatal DAU DEUD DEUD DEUD DEUD DEUD DEUD DEUD																					
Location	Sample ID	Sample	Source ^a	Total PCB	Units	Total PCB	Total PAH	Units	Total PAH	BEHP	Units	BEHP	Lead	Units	Lead	Total PCB	Total PAH	BEHP	Lead	Optimized	Median Optimized	Relative % Δ from
2.5	2 5 4/22/2004	Date	TMD	Result	/!	PEL	Result	/1	PEL 1.C. 770	Result	/!	PEL	Result	/	PEL	PEL-Q	PEL-Q		PEL-Q	`	Mean PEL-Q	Median
2.5 2.5	2.5-4/23/2001 2.5-4/25/2001	4/1/2001 4/1/2001	TMDL TMDL	525 525	μg/kg	189 189	43,235 43,235	μg/kg	16,770 16,770	2,800 2,800	μg/kg	2,647 2,647	34.8 34.8	mg/kg	112 112	2.78 2.78	2.58 2.58	1.06	0.31	1.68 1.68	2.35 2.35	-29% -29%
2.5	2.5-9/5/2000	9/1/2000	TMDL	328	μg/kg	189	30,858	μg/kg	16,770	1,060	μg/kg	2,647	34.6	mg/kg	112	1.74	1.84	0.40	0.31	1.06	2.35	-55%
2.5	2.5-9/8/2000	9/1/2000	TMDL	328	μg/kg	189	30,858	μg/kg	16,770	1,060	μg/kg	2,647	30	mg/kg mg/kg	112	1.74	1.84	0.40	0.27	1.06	2.35	-55%
2.3	3-4/24/2001	4/1/2001	TMDL	6,720	μg/kg	189	230,683	μg/kg	16,770	3,940	μg/kg	2,647	308		112	35.6	13.8	1.49	2.75	13.4	2.35	469%
2	3-4/25/2001		TMDL	6,720	μg/kg	189	230,683	μg/kg	16,770		μg/kg	_	 	mg/kg	112	35.6	13.8		2.75	13.4	2.35	469%
2	· · ·	4/1/2001			μg/kg			μg/kg		3,940	μg/kg	2,647	308	mg/kg				1.49				
3	3-8/12/2003	8/1/2003	TMDL	1,360	μg/kg	189	18,236	μg/kg	16,770	1,120		2,647	43.4	mg/kg	112	7.20	1.09	0.42	0.39	2.27	2.35	-3% -3%
3	3-8/12/2003	8/1/2003	TMDL	1,360	μg/kg	189	18,236	μg/kg	16,770	1,120		2,647	43.4	mg/kg	112	7.20	1.09	0.42	0.39		2.35	
3	3-8/12/2003	8/1/2003	TMDL	1,360	μg/kg	189	18,236	μg/kg	16,770	1,120	μg/kg	2,647	43.4	mg/kg	112	7.20	1.09	0.42	0.39	2.27	2.35	-3%
3	3-8/12/2003	8/1/2003	TMDL	1,360	μg/kg	189	18,236	μg/kg	16,770	1,120	1.0	2,647	43.4	mg/kg	112	7.20	1.09	0.42	0.39	2.27	2.35	-3%
3	3-8/12/2003	8/1/2003	TMDL	1,360	μg/kg	189	18,236	μg/kg	16,770	1,120	μg/kg	2,647	43.4	mg/kg	112	7.20	1.09	0.42	0.39	2.27	2.35	-3%
3	3-8/12/2003-EPA/TCEQ	8/1/2003	EPA/TCEQ	1,360	μg/kg	189	18,505	μg/kg	16,770	1,340	μg/kg	2,647	43.4	mg/kg	112	7.20	1.10	0.51	0.39	2.30	2.35	-2%
3	3-8/12/2003-EPA/TCEQ	8/1/2003	EPA/TCEQ	1,360	μg/kg	189	18,505	μg/kg	16,770	1,340	μg/kg	2,647	43.4	mg/kg	112	7.20	1.10	0.51	0.39	2.30	2.35	-2%
3	3-8/31/2006-EPA/TCEQ	8/1/2006	EPA/TCEQ	17,000	μg/kg	189	111,530	μg/kg	16,770	2,060	μg/kg	2,647	498	mg/kg	112	89.9	6.65	0.78	4.45	25.5	2.35	982%
3	3-8/31/2006-EPA/TCEQ	8/1/2006	EPA/TCEQ	17,000	μg/kg	189	111,530	μg/kg	16,770	2,060	μg/kg	2,647	498	mg/kg	112	89.9	6.65	0.78	4.45	25.5	2.35	982%
3	3-9/5/2000	9/1/2000	TMDL	1,080	μg/kg	189	42,091	μg/kg	16,770	1,540		2,647	67.7	mg/kg	112	5.71	2.51	0.58	0.60	2.35	2.35	0%
3	3-9/6/2000	9/1/2000	TMDL	1,080	μg/kg	189	42,091	μg/kg	16,770	1,540	μg/kg	2,647	67.7	mg/kg	112	5.71	2.51	0.58	0.60	2.35	2.35	0%
4A	4A-4/24/2001	4/1/2001	TMDL	398	μg/kg	189	16,133	μg/kg	16,770	4,630	μg/kg	2,647	39.7	mg/kg	112	2.11	0.96	1.75	0.35	1.29	2.35	-45%
4A	4A-4/25/2001	4/1/2001	TMDL	398	μg/kg	189	16,133	μg/kg	16,770	4,630	μg/kg	2,647	39.7	mg/kg	112	2.11	0.96	1.75	0.35	1.29	2.35	-45%
4A	4a-8/12/2003	8/1/2003	TMDL	34,600	μg/kg	189	14,201	μg/kg	16,770	1,135	μg/kg	2,647	55.9	mg/kg	112	183.1	0.85	0.43	0.50	46.2	2.35	1864%
4A	4a-8/12/2003	8/1/2003	TMDL	34,600	μg/kg	189	14,201	μg/kg	16,770	1,135	μg/kg	2,647	55.9	mg/kg	112	183.1	0.85	0.43	0.50	46.2	2.35	1864%
4A	4A-8/12/2003	8/1/2003	TMDL	34,600	μg/kg	189	14,201	μg/kg	16,770	1,135	μg/kg	2,647	55.9	mg/kg	112	183.1	0.85	0.43	0.50	46.2	2.35	1864%
4A	4a-8/12/2003	8/1/2003	TMDL	34,600	μg/kg	189	14,201	μg/kg	16,770	1,135	μg/kg	2,647	55.9	mg/kg	112	183.1	0.85	0.43	0.50	46.2	2.35	1864%
4A	4a-8/12/2003	8/1/2003	TMDL	34,600	μg/kg	189	14,201	μg/kg	16,770	1,135	μg/kg	2,647	55.9	mg/kg	112	183.1	0.85	0.43	0.50	46.2	2.35	1864%
4A	4A-8/12/2003-EPA/TCEQ	8/1/2003	EPA/TCEQ	34,600	μg/kg	189	16,192	μg/kg	16,770	1,500	μg/kg	2,647	55.9	mg/kg	112	183.1	0.97	0.57	0.50	46.3	2.35	1867%
4A	4A-8/12/2003-EPA/TCEQ	8/1/2003	EPA/TCEQ	34,600	μg/kg	189	16,192	μg/kg	16,770	1,500	μg/kg	2,647	55.9	mg/kg	112	183.1	0.97	0.57	0.50	46.3	2.35	1867%
4A	4A-8/31/2006-EPA/TCEQ	8/1/2006	EPA/TCEQ	150,000	μg/kg	189	18,430	μg/kg	16,770	2,305	μg/kg	2,647	58.8	mg/kg	112	793.7	1.10	0.87	0.53	199.0	2.35	8360%
4A	4A-8/31/2006-EPA/TCEQ	8/1/2006	EPA/TCEQ	150,000	μg/kg	189	18,430	μg/kg	16,770	2,305	μg/kg	2,647	58.8	mg/kg	112	793.7	1.10	0.87	0.53	199.0	2.35	8360%
4A	4A-9/5/2000	9/1/2000	TMDL	1,690	μg/kg	189	14,655	μg/kg	16,770	1,780	μg/kg	2,647	36	mg/kg	112	8.94	0.87	0.67	0.32	2.70	2.35	15%
4A	4A-9/5/2000	9/1/2000	TMDL	1,690	μg/kg	189	14,655	μg/kg	16,770	1,780	μg/kg	2,647	36	mg/kg	112	8.94	0.87	0.67	0.32	2.70	2.35	15%
5	5-4/24/2001	4/1/2001	TMDL	1,910	μg/kg	189	78,263	μg/kg	16,770	7,670	μg/kg	2,647	125	mg/kg	112	10.1	4.67	2.90	1.12	4.70	2.35	100%
5	5-4/25/2001	4/1/2001	TMDL	1,910	μg/kg	189	78,263	μg/kg	16,770	7,670	μg/kg	2,647	125	mg/kg	112	10.1	4.67	2.90	1.12	4.70	2.35	100%
5	5-9/5/2000	9/1/2000	TMDL	780	μg/kg	189	11,337	μg/kg	16,770	1,700		2,647		mg/kg	112	4.13	0.68	0.64	0.35	1.45	2.35	-38%
5	5-9/5/2000	9/1/2000	TMDL	780	μg/kg	189	11,337	μg/kg	16,770	1,700	μg/kg	2,647	39.5	mg/kg	112	4.13	0.68	0.64	0.35	1.45	2.35	-38%
6A	6A-4/25/2001	4/1/2001	TMDL	2,060	μg/kg	189	5,886	μg/kg	16,770	2,330	μg/kg	2,647	50.6	mg/kg	112	10.9	0.35	0.88	0.45	3.15	2.35	34%
6A	6A-4/25/2001	4/1/2001	TMDL	2,060	μg/kg	189	5,886	μg/kg	16,770	2,330	μg/kg	2,647	50.6	mg/kg	112	10.9	0.35	0.88	0.45	3.15	2.35	34%
6A	6a-8/12/2003	8/1/2003	TMDL	10,900	μg/kg	189	11,974	μg/kg	16,770	1,068	μg/kg	2,647	76.6	mg/kg	112	57.7	0.71	0.40	0.68	14.9	2.35	532%
6A	6a-8/12/2003	8/1/2003	TMDL	10,900	μg/kg	189	11,974	μg/kg	16,770				76.6	mg/kg	112	57.7	0.71	0.40	0.68	14.9	2.35	532%
6A	6A-8/12/2003	8/1/2003	TMDL	10,900	μg/kg	189	11,974	μg/kg	16,770	1,068	μg/kg	2,647	76.6	mg/kg	112	57.7	0.71	0.40	0.68	14.9	2.35	532%
6A	6a-8/12/2003	8/1/2003	TMDL	10,900	μg/kg	189	11,974	μg/kg	16,770	1,068	μg/kg	2,647	76.6	mg/kg	112	57.7	0.71	0.40	0.68	14.9	2.35	532%
6A	6a-8/12/2003	8/1/2003	TMDL	10,900	μg/kg	189	11,974	μg/kg	16,770	1,068	μg/kg	2,647	76.6	mg/kg	112	57.7	0.71	0.40	0.68	14.9	2.35	532%
6A	6A-8/12/2003-EPA/TCEQ	8/1/2003	EPA/TCEQ	10,900	μg/kg	189	27,538	μg/kg	16,770	1,860		2,647		mg/kg	112	57.7	1.64	0.70	0.68	15.2	2.35	545%
6A	6A-8/12/2003-EPA/TCEQ	8/1/2003	EPA/TCEQ	10,900	μg/kg	189	27,538	μg/kg	16,770	1,860	μg/kg	2,647	76.6	mg/kg	112	57.7	1.64	0.70	0.68	15.2	2.35	545%
6A	6A-8/31/2006-EPA/TCEQ	8/1/2006	EPA/TCEQ	7,700	μg/kg	189	3,642	μg/kg	16,770	1,195	μg/kg	2,647	23.3	mg/kg	112	40.7	0.22	0.45	0.21	10.4	2.35	342%
6A	6A-8/31/2006-EPA/TCEQ	8/1/2006	EPA/TCEQ	7,700	μg/kg	189	3,642	μg/kg	16,770	1,195	μg/kg	2,647	23.3	mg/kg	112	40.7	0.22	0.45	0.21	10.4	2.35	342%
6A	6A-9/5/2000	9/1/2000	TMDL	9,370	μg/kg	189	60,074	μg/kg	16,770	1,610		2,647	34.6	mg/kg	112	49.6	3.58	0.61	0.31	13.5	2.35	475%
6A	6A-9/5/2000	9/1/2000	TMDL	9,370	μg/kg	189	60,074	μg/kg	16,770	1,610	μg/kg	2,647	34.6	mg/kg	112	49.6	3.58	0.61	0.31	13.5	2.35	475%
E	E-4/23/2001	4/1/2001	TMDL	1,580	μg/kg	189	45,108	μg/kg	16,770	2,550		2,647	47.5	mg/kg	112	8.36	2.69	0.96	0.42	3.11	2.35	32%
E	E-4/25/2001	4/1/2001	TMDL	1,580	μg/kg	189	45,108	μg/kg	16,770	2,550	μg/kg	2,647	47.5	mg/kg	112	8.36	2.69	0.96	0.42	3.11	2.35	32%
E	E-9/5/2000	9/1/2000	TMDL	370	μg/kg	189	13,475	μg/kg	16,770	1,170	_	2,647	25	mg/kg	112	1.96	0.80	0.44	0.22	0.86	2.35	-64%
E	E-9/6/2000	9/1/2000	TMDL	370	μg/kg	189	13,475	μg/kg	16,770	1,170			25	mg/kg	112	1.96	0.80	0.44	0.22	0.86	2.35	-64%

Appendix E-2.3
Sediment Chemistry and PEL-Q Values by Sample

		Sample		Total PCB		Total PCB	Total PAH		Total PAH	ВЕНР		ВЕНР	Lead		Lead	Total PCB	Total PAH	BEHP	Lead	Optimized	Median Optimized	Relative % Δ from
Location	Sample ID	Date	Source ^a	Result	Units	PEL	Result	Units	PEL	Result	Units	PEL	Result	Units	PEL	PEL-Q	PEL-Q	PEL-Q	PEL-Q	Mean PEL-Q	Mean PEL-Q	Median
G	G-4/24/2001	4/1/2001	TMDL	197	μg/kg	189	9,737	μg/kg	16,770	507	μg/kg	2,647	52	mg/kg	112	1.04	0.58	0.19	0.46	0.57	2.35	-76%
G	G-4/25/2001	4/1/2001	TMDL	197	μg/kg	189	9,737	μg/kg	16,770	507	μg/kg	2,647	52	mg/kg	112	1.04	0.58	0.19	0.46	0.57	2.35	-76%
G	G-9/5/2000	9/1/2000	TMDL	399	μg/kg	189	14,235	μg/kg	16,770	809	μg/kg	2,647	39.8	mg/kg	112	2.11	0.85	0.31	0.36	0.91	2.35	-62%
G	G-9/6/2000	9/1/2000	TMDL	399	μg/kg	189	14,235	μg/kg	16,770	809	μg/kg	2,647	39.8	mg/kg	112	2.11	0.85	0.31	0.36	0.91	2.35	-62%
Q	Q-4/25/2001	4/1/2001	TMDL	1,070	μg/kg	189	16,350	μg/kg	16,770	12,000	μg/kg	2,647	193	mg/kg	112	5.66	0.97	4.53	1.72	3.22	2.35	37%
Q	Q-4/25/2001	4/1/2001	TMDL	1,070	μg/kg	189	16,350	μg/kg	16,770	12,000	μg/kg	2,647	193	mg/kg	112	5.66	0.97	4.53	1.72	3.22	2.35	37%
Q	Q-9/5/2000	9/1/2000	TMDL	3,780	μg/kg	189	30,926	μg/kg	16,770	1,960	μg/kg	2,647	27.3	mg/kg	112	20.0	1.84	0.74	0.24	5.71	2.35	143%
Q	Q-9/5/2000	9/1/2000	TMDL	3,780	μg/kg	189	30,926	μg/kg	16,770	1,960	μg/kg	2,647	27.3	mg/kg	112	20.0	1.84	0.74	0.24	5.71	2.35	143%
S	S-4/24/2001	4/1/2001	TMDL	2,090	μg/kg	189	116,399	μg/kg	16,770	14,400	μg/kg	2,647	124	mg/kg	112	11.1	6.94	5.44	1.11	6.14	2.35	161%
S	S-4/25/2001	4/1/2001	TMDL	2,090	μg/kg	189	116,399	μg/kg	16,770	14,400	μg/kg	2,647	124	mg/kg	112	11.1	6.94	5.44	1.11	6.14	2.35	161%
S	S-9/5/2000	9/1/2000	TMDL	113	μg/kg	189	6,812	μg/kg	16,770	196	μg/kg	2,647	47.8	mg/kg	112	0.60	0.41	0.07	0.43	0.38	2.35	-84%
S	S-9/5/2000	9/1/2000	TMDL	113	μg/kg	189	6,812	μg/kg	16,770	196	μg/kg	2,647	47.8	mg/kg	112	0.60	0.41	0.07	0.43	0.38	2.35	-84%
Т	T-4/25/2001	4/1/2001	TMDL	626	μg/kg	189	17,373	μg/kg	16,770	2,550	μg/kg	2,647	127	mg/kg	112	3.31	1.04	0.96	1.13	1.61	2.35	-32%
Т	T-4/25/2001	4/1/2001	TMDL	626	μg/kg	189	17,373	μg/kg	16,770	2,550	μg/kg	2,647	127	mg/kg	112	3.31	1.04	0.96	1.13	1.61	2.35	-32%
Т	T-9/5/2000	9/1/2000	TMDL	837	μg/kg	189	10,712	μg/kg	16,770	1,180	μg/kg	2,647	28.9	mg/kg	112	4.43	0.64	0.45	0.26	1.44	2.35	-39%
Т	T-9/5/2000	9/1/2000	TMDL	837	μg/kg	189	10,712	μg/kg	16,770	1,180	μg/kg	2,647	28.9	mg/kg	112	4.43	0.64	0.45	0.26	1.44	2.35	-39%
U	U-4/24/2001	4/1/2001	TMDL	13	μg/kg	189	444	μg/kg	16,770		μg/kg	2,647	16.8	mg/kg	112	0.07	0.03		0.15	0.08	2.35	-97%
U	U-4/25/2001	4/1/2001	TMDL	13	μg/kg	189	444	μg/kg	16,770		μg/kg	2,647	16.8	mg/kg	112	0.07	0.03		0.15	0.08	2.35	-97%
U	U-9/5/2000	9/1/2000	TMDL	79	μg/kg	189	2,602	μg/kg	16,770	200	μg/kg	2,647	15	mg/kg	112	0.42	0.16	0.08	0.13	0.20	2.35	-92%
U	U-9/6/2000	9/1/2000	TMDL	79	μg/kg	189	2,602	μg/kg	16,770	200	μg/kg	2,647	15	mg/kg	112	0.42	0.16	0.08	0.13	0.20	2.35	-92%
V	V-4/23/2001	4/1/2001	TMDL	442	μg/kg	189	24,025	μg/kg	16,770	2,250	μg/kg	2,647	50.9	mg/kg	112	2.34	1.43	0.85	0.45	1.27	2.35	-46%
V	V-4/25/2001	4/1/2001	TMDL	442	μg/kg	189	24,025	μg/kg	16,770	2,250	μg/kg	2,647	50.9	mg/kg	112	2.34	1.43	0.85	0.45	1.27	2.35	-46%
V	V-9/5/2000	9/1/2000	TMDL	447	μg/kg	189	26,994	μg/kg	16,770	1,390	μg/kg	2,647	48.2	mg/kg	112	2.37	1.61	0.53	0.43	1.23	2.35	-48%
V	V-9/5/2000	9/1/2000	TMDL	447	μg/kg	189	26,994	μg/kg	16,770	1,390	μg/kg	2,647	48.2	mg/kg	112	2.37	1.61	0.53	0.43	1.23	2.35	-48%

^a Data Source References:

TMDL - Parsons et al. (2002, 2004) EPA/TCEQ - L. Broach (no date)

μg/kg - microgram per kilogram mg/kg - milligram per kilogram

PCB - polychorinated biphenyl
PAH - polycyclic aromatic hydrocarbon

BEHP - bis(2-ethylhexyl)phthalate

PEL - probable effect level

PEL-Q - probable effect level quotient

 $\%\Delta$ - percent difference

Appendix E-2.4
Average Optimized Mean PEL-Q Values by Location

Location	Sample Date	Data Source ^a	Average Total PCB PEL-Q	Average Total PAH PEL-Q	Average BEHP PEL-Q	Average Lead PEL-Q	Average Optimized Mean PEL-Q
2.5	9/1/2000	TMDL	1.74	1.84	0.400	0.268	1.06
2.5	4/1/2001	TMDL	2.78	2.58	1.06	0.311	1.68
3	9/1/2000	TMDL	5.71	2.51	0.582	0.604	2.35
3	4/1/2001	TMDL	35.6	13.8	1.49	2.75	13.4
3	8/1/2003	EPA/TCEQ	7.20	1.10	0.506	0.388	2.30
3	8/1/2003	TMDL	7.20	1.09	0.423	0.388	2.27
3	8/1/2006	EPA/TCEQ	89.9	6.65	0.778	4.45	25.5
4A	9/1/2000	TMDL	8.94	0.874	0.672	0.321	2.70
4A	4/1/2001	TMDL	2.11	0.962	1.75	0.354	1.29
4A	8/1/2003	EPA/TCEQ	183	0.966	0.567	0.499	46.3
4A	8/1/2003	TMDL	183	0.847	0.429	0.499	46.2
4A	8/1/2006	EPA/TCEQ	794	1.10	0.871	0.525	199
5	9/1/2000	TMDL	4.13	0.676	0.642	0.353	1.45
5	4/1/2001	TMDL	10.1	4.67	2.90	1.12	4.70
6A	9/1/2000	TMDL	49.6	3.58	0.608	0.309	13.5
6A	4/1/2001	TMDL	10.9	0.351	0.880	0.452	3.15
6A	8/1/2003	EPA/TCEQ	57.7	1.64	0.703	0.684	15.2
6A	8/1/2003	TMDL	57.7	0.714	0.403	0.684	14.9
6A	8/1/2006	EPA/TCEQ	40.7	0.217	0.451	0.208	10.4
E	9/1/2000	TMDL	1.96	0.804	0.442	0.223	0.857
E	4/1/2001	TMDL	8.36	2.69	0.963	0.424	3.11
G	9/1/2000	TMDL	2.11	0.849	0.306	0.355	0.905
G	4/1/2001	TMDL	1.04	0.581	0.192	0.464	0.570
Q	9/1/2000	TMDL	20.0	1.84	0.740	0.244	5.71
Q	4/1/2001	TMDL	5.66	0.975	4.53	1.72	3.22
S	9/1/2000	TMDL	0.598	0.406	0.0740	0.427	0.376
S	4/1/2001	TMDL	11.1	6.94	5.44	1.11	6.14
Т	9/1/2000	TMDL	4.43	0.639	0.446	0.258	1.44
Т	4/1/2001	TMDL	3.31	1.04	0.963	1.13	1.61
U	9/1/2000	TMDL	0.417	0.155	0.0756	0.134	0.195
U	4/1/2001	TMDL	0.0677	0.0265		0.150	0.0814
V	9/1/2000	TMDL	2.37	1.61	0.525	0.430	1.23
V	4/1/2001	TMDL	2.34	1.43	0.850	0.454	1.27

Appendix E-2.4
Average Optimized Mean PEL-Q Values by Location

Location	Sample Date	Data Source ^a	Average Total PCB PEL-Q	Average Total PAH PEL-Q	Average BEHP PEL-Q	Average Lead PEL-Q	Average Optimized Mean PEL-Q	
		MEDIAN:	7.20	1.04	0.625	0.427	2.35	

^a Data Source References:

TMDL - Parsons et al. (2002, 2004) EPA/TCEQ - L. Broach (no date)

PCB - polychorinated biphenyl

PAH - polycyclic aromatic hydrocarbon

BEHP - bis(2-ethylhexyl)phthalate

PEL-Q - probable effect level quotient

APPENDIX E-3 BENTHIC HABITAT EVALUATION

TABLE OF CONTENTS

1	BEN	VTHIC HABITAT	1
		General Conditions	
	1.2	Salinity	2
		Grain Size	
		Dissolved Oxygen	
		Summary of Non-Chemical Stressor Effects	
2	REF	'ERENCES	5

List of Tables

Table E-3.1 Salinity

Table E-3.2 Dissolved Oxygen

List of Figures

- Figure E-3.1 Visual Model Showing Relationships Among Macrobenthos Metrics,
 Contaminants, and Natural Environmental Factors
- Figure E-3.2 Hydrodynamic Continuous Monitoring Locations
- Figure E-3.3 Salinity Time Series Plot
- Figure E-3.4 Surface Water Salinity Box Plots
- Figure E-3.5 Dissolved Oxygen Time Series Plot
- Figure E-3.6 Surface Water Dissolved Oxygen Box Plots

1 BENTHIC HABITAT

In estuaries, structure and composition of the benthic assemblage is strongly related to variation in salinity, sediment type, DO, temperature, and water depth (e.g., Boesch 1973, 1977; Dauer et al. 1987; Holland et al. 1987; Schaffner et al. 1987; Chester et al. 1983; Engle et al. 1994; Ferraro et al. 1991). These non-chemical stressors impose physical limitations on the overall health of the benthic community, even in the absence of anthropogenic stressors (Wilson and Jeffrey 1994). A graphical model from Gaston et al. (1998) is useful to help visualize the interrelationships among biological and physical conditions and broad benthic patterns throughout the northern Gulf of Mexico (GOM) (Figure E-3.1). The figure illustrates that as salinity and DO increase, increases in dominance of crustaceans, trophic diversity, and benthic organism abundance occur; likewise, as silt/clay components and contaminant amounts increase, annelids increase in dominance, trophic simplification occurs, and benthos are less abundant. The following sections describe observed patterns of these non-chemical stressors within the Site and how they are expected to affect the expected condition of the benthic communities present.

1.1 General Conditions

The upstream and southern portion of the Site has a significantly higher gradient (about 10 feet of elevation change over 5,000 linear feet) when compared to the middle and lower portions of Patrick Bayou (less than 1 foot of elevation change over 8,000 linear feet) (Anchor 2006). The water depth in the upstream portions of Patrick Bayou to the downstream limit of the gunite-lined channel generally does not exceed -3 feet North American Vertical Datum 1988 (NAVD88). Below this, Patrick Bayou exhibits water depths ranging to -6 feet NAVD88. The downstream-most portion of the channel widens prior to its intersection with the HSC and areas of deposition and channel flow are more clearly defined. Depths in this area range between -3 and -6 feet (NAVD88) in depth. Water depths at the transition between the Site and HSC range between -11 and -17 feet NAVD88.

Due to the shallow nature of the Site, many areas of intertidal and even subtidal habitat can be dewatered for significant periods of time due to wind driven water level fluctuations. These prolonged periods of exposure can have significant effects on the benthic community. In addition, areas with high hydraulic gradients and shallow soft sediments are indicative of

areas where sediment stability is low. Even in areas that are considered net depositional, high bed shear stress during stormwater runoff events results in periodic erosional events. Sediment resuspension and transport may have significant effects on the benthic community at the Site, since benthic communities in the GOM tend to be concentrated in the shallowest portion of the sediment column (< 2 cm) compared to more northern estuaries (e.g., Chesapeake Bay; Brown et al. 2000; Gaston et al. 1998). For the Site, sediment scouring deeper than 2 cm would not be expected due to a 2-year storm event (Anchor QEA 2011). However, up to 10 percent of the sediment surface area at the Site would be expected to be scoured deeper than 2 cm after a 5-year storm event and 12 percent of the Site would be scoured after a 10-year storm event.

1.2 Salinity

Salinity has been shown to have a significant influence on benthic community structure in GOM estuaries (Brown et al. 2000; Dobberstine et al. 2007; TCEQ 2003). Benthic communities are generally more diverse and more abundant at high salinity ranges (>25 percent salinity) (Brown et al. 2000; Gaston et al. 1998; TCEQ 2003), and contain less subsurface deposit feeders (Gaston et al. 1998). TCEQ's sampling effort in Galveston Bay (TCEQ 2003) labeled salinity a "key determinant" in the distribution of estuarine fauna. In addition, Engle et al. (1994) observed strong salinity effects on species richness in estuaries. Stability in the salinity range is also a factor; communities in Galveston Bay were more diverse in areas with stable salinity and less diverse where salinity fluctuations are more frequent and over a greater range, such as the upper HSC (Lester and Gonzales 2002). Montague and Ley (1993) observed that the standard deviation of salinity observations was the best environmental correlate with benthic community abundance in a North Florida estuary.

Patrick Bayou is generally a mesohaline system with a salinity gradient between the downstream and upstream limits of the Site as a result of storm water runoff, upstream flow, and tidal effects. Salinity ranged from freshwater (<0.5 ppt) up to 12 ppt during continuous measurements collected at four locations monitored between October 2006 and 2007 (Figure E-3.2). Median daily salinity values ranged from 4.0 to 2.3 ppt from downstream to

upstream¹. Rapid changes in salinity associated with rainfall were observed (Figure E-3.3). Box plots and a table of salinity levels for the four continuous monitoring locations show the high variance in salinity levels at the Site (Figure E-3-4 and Table E-3.1). Salinity changes occurred over a period of hours and were often transient (returning to previous levels over a period of hours) but were commonly associated with lasting (i.e., weeks) change in salinity regime. Longer-term variability (e.g., months) was also observed. This short- and long-term variability may have a significant effect on the condition of the benthic community at the Site.

1.3 Grain Size

The average grain size distribution for portions of Patrick Bayou ranges from sand to clay (Anchor 2006). Sediment in upstream portions of the Site and upstream of the Site itself contain a higher proportion of sand, including some gravel, while downstream areas tend to have a higher proportion of silt and clay and little to no gravel. This change in the distribution of grain size is consistent with transition from a higher water velocity and steep slope environment in the upstream portion of the Site, to the lower slope lower water velocity environment in downstream areas.

Grain size is another key driver of benthic community structure, both for type of species present, as well as species diversity and organism abundance. Smaller grain size fractions (silt/clay) have been observed to contain more surface deposit feeding benthos, as opposed to suspension feeders (Rhoads and Young 1970; Levinton 1972; Day et al. 1989; Peterson 1991). Diversity and organism density have both been shown to be lower at muddy versus sandy sites in northern GOM estuaries (Brown 2000). Changes in grain size, therefore, would be expected to affect benthic community characteristics.

1.4 Dissolved Oxygen

DO conditions are known to affect species diversity, abundance, and type of organisms present in GOM estuaries. Some studies have indicated stronger relationships than others.

¹ The East Fork tributary (EF005) was distinctly lower in salinity than Patrick Bayou itself. Median daily salinity was 0.4 ppt.

Brown et al. (2000) found more species diversity, greater biota abundance, and higher proportions of certain feeding groups at sites with higher DO (>5 ppm) than at lower DO (i.e., hypoxic) sites (<2 ppm), but DO conditions were not statistically correlated with the overall structure of benthic communities. Dominant species did differ, crustaceans and mollusks were more dominant at high DO sites, while worms dominated in low DO environments (Brown et al. 2000). Gaston et al. (1998) data did not show statistically significant relationships between trophic groups and level of DO. Changes in DO conditions could thus have some effects on benthic organisms and community. One example of these effects was reported from modeling work in Chesapeake Bay, where Dauer (2003) found that modeled stations exposed to low DO events resulted in reduced biomass and species richness, and more opportunistic species overall. Intermittent hypoxia was also shown in an experimental study in the Chesapeake Bay to result in significant decreases in macrobenthic abundance and the local extinction of several worm taxa (Llanso 1992).

At the Site, continuous measurements of DO from October 2006 to June 2007 (Figure E-3.5) demonstrated that hypoxic conditions (24-hour DO average less than 2.0 milligrams per liter [mg/L]) did exist in upstream areas (i.e., PB-075) nearly 40 percent of the time (100 of 257 daily records). There was an apparent seasonality to DO levels, as well as short-term (days) fluctuations linked to runoff events. Box plots and a table of DO levels for the four continuous monitoring locations also show the high variance in DO levels at the Site (Figure E-3-6 and Table E-3.2). Thus, DO likely plays a significant role in benthic community structure in upstream areas of the Site, acting to limit diversity and abundance in upstream areas.

1.5 Summary of Non-Chemical Stressor Effects

Common events like flushing flows from storms would have implications for multiple physical variables at once, such as DO, temperature, sediment resuspension, and salinity. Flushing flows in shallow tributaries or bayous with increased impervious surface area would be expected to create multiple, frequent high peak flow events which are observed at the Site.

2 REFERENCES

- Anchor Environmental, LLC (Anchor), 2006. *Preliminary Site Characterization Report.*Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. May 2006.
- Anchor QEA, 2011. *Sediment Transport Modeling Report, Patrick Bayou*. Prepared for Patrick Bayou Joint Defense Group. September 2011.
- Boesch, D.F., 1973. Classification and community structure of macrobenthos in Hampton Roads area. Virginia. *Mar. Biol.* 21: 226-244.
- Boesch, D.F., N.E. Armstrong, J.C. Cloern, L.A. Deegan, R.D. Perkins, and S.L. Williams, 1997. Annual Report of the Florida Bay Science Oversight Panel: Perspectives from the 1996 Florida Bay Science Conference and Review of the Strategic Plan. Report to the Program Management Committee, Interagency Florida Bay Research Program.
- Brown, S.S., G.R. Gaston, C.F. Rakocinski, and R.W. Heard, 2000. Effects of sediment contaminants and environmental gradients on macrobenthic community trophic structure in Gulf of Mexico estuaries. *Estuaries, Vol 23: No 3* (June 2000), pp. 411-424.
- Chester, A.J., L. Ferguson, and G.W. Thayer, 1983. Environmental gradients and benthic macroinvertebrate distributions in a shallow North Carolina estuary. *Bulletin of Marine Science* 33:282-295.
- Dauer, D.M., 2003. Biological criteria, environmental health and estuarine macrobenthic community structure. *Marine Pollution Bulletin*, 26:5, pp. 249-257.
- Dauer, D.M., R.M. Ewing, and J.A. Ranasinghe, 1987. *Macrobenthic Communities of the Lower Chesapeake Bay*: Chesapeake Bay Program, March 1985-June 1986. Prepared for the Virginia Water Control Board. 414-S-86-47. Old Dominion University Research Foundation. Norfolk, VA.
- Day, J.R., C.A.S. Hall, W.M. Kemp, and A. Yanez-Arancibia, 1989. *Estuarine Ecology.* John Wiley and Sons, Inc., New York.

- Dobberstine, J., J. Horne, L. Brzuzy, and C. Howard, 2007. Identifying suitable reference sites for impated sites along the Houston Ship Channel. Proceedings of the Eighth Biennial State of the Bay Symposium, Galveston Bay Estuary Program.
- Engle, V.D., J.K. Summers, and G.R. Gaston, 1994. A Benthic Index of Environmental Condition of Gulf of Mexico Estuaries. *Estuaries*. Vol. 17, No. 2, pp. 372-84. June 1994.
- Ferraro, S.P., R.C. Swartz, F.A. Cole, and D.W. Schults, 1991. Temporal changes in the benthos along a pollution gradient: discriminating the effects of natural phenomena from sewage-industrial wastewater effects. *Estuarine, Coastal and Shelf Science* 33:383-407.
- Gaston, G.R., C.F. Rakocinski, S.S. Brown, and C.M. Cleveland, 1998. Trophic function in estuaries: response of macrobenthos to natural and contaminant gradients. *Marine Freshwater Research.* 49:833-846.
- Holland, A.F., A. Shaughnessey, and M.H. Heigel, 1987. Long-term variation in mesohaline Chesapeake Bay benthos: spatial and temporal patterns. *Estuaries* 10:227-245.
- Lester, J., L. Gonzalez, (eds). 2002. *The State of the Bay.* A Characterization of the Galveston Bay Ecosystem (2nd ed). Houston: Galveston Bay Estuary Program.
- Levinton, J.S., 1972. Stability and trophic structure in deposit feeding and suspension-feeding communities. *American Naturalist 106:472-486.*
- Llanso, R.J., 1992. Effects of hypoxia on estuarine benthos: the lower Rappahannock River (Chesapeake Bay), a case study. *Estuarine, Coastal, and Shelf Science*. Volume 36, Issue 5, November 1992. Pp. 491-515.
- Montague, C.L., and J.A. Ley, 1993. A possible effect of salinity fluctuation on abundance of benthic vegetation and associated fauna in Northeastern Florida Bay. *Estuaries*, 16, 703–717.
- Peterson, C.H., 1991. Intertidal zonation or marine invertebrates in sand and mud. *American Scientist* 79:236-249.
- Rhoads, D.C. and D.K. Young, 1970. The influence of deposit feeding organisms on sediment stability and community trophic structure. *Journal of Marine Research* 28: 150-178.

- Schaffner, L.C., R.J. Diaz, C.R. Olson, and I.L. Larsen, 1987. Faunal characteristics and sediment accumulation processes in the James River Estuary, Virginia. *Estuarine, Coastal and Shelf Science* 25:211-226.
- Texas Commission on Environmental Quality (TCEQ), 2003. *An evaluation of macrobenthic community structure in the Galveston Bay System from 1997 to 2000.* Texas Commission on Environmental Quality, Field Operations Division. AS-191. May 2003.
- Wilson, J.G., and D.W. Jeffrey, 1994. *Benthic biological pollution indices in estuaries*. Pages 311-327 *in* K.J.M. Kramer. Editor. Biomonitoring of coastal waters and estuaries. CRC Press. Boca Raton. Florida. USA.

TABLES


Table E-3.1 Salinity (ppt)

Station	N	Minimum	Maximum	Median	Arithmetic Mean	Standard Deviation	Coefficient of Variation	5th Percentile	20th Percentile	80th Percentile	95th Percentile
EF005	12,401	0.04	7.31	0.4	1.014	1.219	1.202	0.12	0.19	1.83	3.64
PB020	28,257	0.107	12.354	3.965	4.569	2.921	0.639	0.796	1.818	7.156	10.348
PB045	24,783	0.04	11.92	3.67	4.001	2.56	0.64	0.52	1.61	6.37	8.38
PB075	26,309	0.04	10.58	2.27	2.407	1.424	0.592	0.28	1.26	3.42	4.84

Table E-3.2 Dissolved Oxygen (mg/L)

Station	N	Minimum	Maximum	Median	Arithmetic Mean	Standard Deviation	Coefficient of Variation	5th Percentile	20th Percentile	80th Percentile	95th Percentile
EF005	12,401	0.31	32.65	5.34	5.613	2.716	0.484	1.66	3.56	7.26	10.06
PB020	28,257	0.09	15.76	6.663	6.75	2.125	0.315	3.33	4.89	8.63	10.071
PB045	24,783	0.79	30.93	6.75	6.888	2.42	0.351	3.16	4.69	8.98	10.64
PB075	26,309	0	12.56	2.78	2.978	2.234	0.75	0.11	0.68	4.89	6.95


FIGURES














































APPENDIX E-4 BENTHIC INDEX EVALUATION

TABLE OF CONTENTS

1	ENGLE AND SUMMERS	1
2	TCEQ BIOTIC INDEX	4
3	INCREMENTAL RISK CALCULATIONS	6
4	REFERENCES	7

List of Tables

Table E-4.1	ES-BI September 2000 Calculations
-------------	-----------------------------------

Table E-4.2 ES-BI April 2001 Calculations

Table E-4.3 Summary of Engle and Summers Benthic Index

Table E-4.4 TCEQ Tidal Stream-Tolerant/Opportunistic/Pollution Indicative Taxa

Table E-4.5 TCEQ Tidal Stream-Intolerant/Pollution Sensitive Taxa

Table E-4.6 TCEQ Tidal Stream Metric Score Calculations

Table E-4.7A Patrick Bayou September 2000 Benthic Community Data and TCEQ BI Scores

Table E-4.7B Patrick Bayou April 2001 Benthic Community Data and TCEQ BI Scores

Table E-4.7C Patrick Bayou August 2003 Benthic Community Data and TCEQ BI Scores

Table E-4.8 Summary of TCEQ Benthic Index

List of Attachments

Attachment E-4.1 ProUCL Statistics

1 ENGLE AND SUMMERS

Engle and Summers (1999) developed an indicator of benthic condition, Engle and Summers Biotic Index (ES-BI), for northern Gulf of Mexico estuaries. The ES-BI was developed based on data from USEPA's Environmental Monitoring and Assessment Program (EMAP) using samples collected from estuaries in the Louisiana Province from 1991 to 1994.

The ES-BI developed by Engle and Summers (1999), is based on a combination of five metrics: the percentage of expected species diversity, the mean abundance of tubificid oligochaetes, the percent total abundance represented by capitellid polychaetes (relative abundance of capitellid polychaetes), the relative abundance of bivalve mollusks, and the relative abundance of amphipods. The percentage of expected species diversity is based on a salinity adjustment (Engle and Summers 1999). The percentage of expected species diversity = the mean diversity (H')/expected diversity. Where the expected diversity is calculated based on the salinity at each site when the sample was collected (Engle and Summers 1999) using the following equation:

Expected diversity = 2.618426-(0.044795 x salinity) + (0.007278 x salinity 2) + (-0.000119 x salinity)

The ES-BI used for this evaluation is the same benthic index used by Parsons Engineering (2002) during the third party TMDL investigations performed at the Site. However, the index for this assessment was recalculated to evaluate the September 2000 and April 2001 data separately without any reference site data. Typically, the ES-BI is calculated using site of interest data and nearby reference area data. However, for this evaluation, appropriate reference area data are not available. Thus, the ES-BI was calculated using only Site data. Thus, the score is a relative index of community condition within the Site and is not related to absolute scoring to determine condition of the community. The relative index is appropriate for the analysis because only Site data are of interest in understanding the condition of the benthic community on a by station basis. Use of appropriate reference data would be of benefit if the goal was to assess the Site benthic community to determine the potential for a gradient from a stressed to a reference condition at the Site.

Samples included in this LOE were collected from Patrick Bayou in 2000 and 2001¹. Sample locations are identified in Figure E-4.1. For the recalculation, transformed data for each metric were derived directly from the Parsons Engineering (2002) report. As noted above, the ES-BI scores were calculated separating the data into two seasonal groups. The first set is the "summer" data group that included data for samples collected in September 2000 from the Site (Parsons 2002). The second data group was for spring collected samples, April 2001. No attempt was made to verify the calculation of the metrics. The recalculation of the ES-BI is presented in Tables E-4.1 and E-4.2. The ES-BI is calculated following four steps (Engle and Summers 1999).

Step 1: Calculate arithmetic mean and standard deviation of the benthic variables using the Patrick Bayou data (Table E-4.1).

Step 2: Normalize (standardize) the data using this mean and standard deviation calculated in Step 1. The normalization is conducted using the following equation:

$$x_1' = \frac{(S \times (x_1 - \bar{x}) + M)}{S_x}$$

where

 x_1' = new standardized value

S = desired standard deviation (one)

M = desired mean (zero)

 x_1 = observation's original value

x = variable's mean from Step 1

 S_x = variable's standard deviation from Step 1

With Step 3: Each new standardized value is then scored using the following relationships developed by Engle and Summers (1999) for the ES-BI:

Discriminant Score = (1.571 x Percentage of expected species diversity) +

¹ Benthic invertebrate data from 2003 were not used, as salinity data were not available. Salinity data are needed to calculate the ES-BI. Furthermore, only four stations were evaluated in October 2001, so the data were not used.

```
(-1.0335 x mean abundance of tubificid oligochaetes) + (-0.5607 x relative abundance of capitellid polychaetes) + (-0.447 x relative abundance of bivalves) + (0.5023 x relative abundance of amphipods)
```

Finally, with Step 4: The benthic index is calculated from the discriminant scores as follows:

$$ES-BI = [((Discriminant Score - (-3.21)) / 7.50)]*10$$

Patrick Bayou stations 2, Y, R, 7, 8, 9, and 10 are not being evaluated in this report as they are not within the Site area or are in the gunite channel. A summary of E-S BI Scores are presented in Table E-4.3.

2 TCEQ BIOTIC INDEX

A second benthic invertebrate community index was selected for a sensitivity analysis to evaluate Site benthic invertebrate community data. A biotic index specific to tidal streams in Texas was used. The TCEQ Biotic Index is being developed for evaluating tidal streams in Texas (TCEQ 2012). It is in draft form and has not been peer reviewed. However, this index was used here as requested by USEPA. This index has six different metrics:

- Taxa Richness
- Diversity (Shannon-Weiner)
- Percent Tolerant Taxa
- Percent Intolerant Taxa
- Presence of Five Key Taxa Groups, (Polychaetes, Crustacea, Nemertea, Bivalvia, Gastropoda)
- Three Most Dominant Taxa

The macroinvertebrate taxa that are assumed to be tolerant are presented in Table E-4.4 and intolerant in Table E-4.5. The selection of the taxa that are tolerant or intolerant in tidal streams of Texas has not been peer reviewed.

The benthic community data were scored using the approach described by TCEQ (Table E-4.6). The TCEQ BI scoring system is based on 94 samples from tidal streams and side bays in the Galveston Bay system (TCEQ 2012). Samples were included in the TCEQ evaluation if they were collected in summer (June through September) and where salinity was less than 20 ppt at time of collection (TCEQ 2012). Samples were collected between 1992 and 2008 for various special studies or for routine monitoring. The 94 samples were collected from 64 stations.

In this analysis the evaluation of a score for a station was not in comparison to a reference condition (e.g., stressed or not stressed) but in the context of the other scores at the Site (a relative score). The TCEQ BIs for this assessment were calculated to evaluate the September 2000, April 2001 and August 2003 monitoring events. The Site benthic community data, TCEQ metric scores and TCEQ BI Scores are presented in Tables E-4.7A, E-4.7B, and E-4.7C. A summary of TCEQ BI Scores is presented in Table E-4.8.

3 INCREMENTAL RISK CALCULATIONS

Next, overall incremental risk to the benthic community as measured by the overall composition and health of the benthic community risk was categorized for each location according to the following criteria:

- Probable Risk relative ES-BI score was less than the 15th percentile (lowest 15 percent) of locations within the Site
- Indeterminate Risk relative ES-BI score was greater than the 15th percentile but was equal to or less than the Site-wide average
- Low Risk relative ES-BI score was greater than the Site-wide average

These statistics were calculated using ProUCL Version 4.1 software (USEPA 2010). The statistics for the E-S BI and TCEQ BI scores are presented in Attachment E-4.1. The results of this evaluation are presented in Table 8-3.

4 REFERENCES

- Brown, S.S., G.R. Gaston, C. F. Rakocinski, and R.W. Heard. 2000. Effects of sediment contaminants and environmental gradients on macrobenthic community trophic structure in Gulf of Mexico estuaries. Estuaries 23(3):411-424.
- Dauer, D.M, A.J. Rodi, Jr., and J.A. Ranasinghe. 1992. Effects of low dissolved oxygen events on the macrobenthos of the lower Chesapeake Bay. Estuaries 15(3):384-391.
- Engle, V.D., and J.K. Summers, 1999. Refinement, Validation, and Application of a Benthic Condition Index for Northern Gulf of Mexico Estuaries. *Estuaries* 22: 624-635.
- Lerberg, S.B., A.F. Holland, and D.M. Sanger. 2000. Responses of tidal creek macrobenthic communities to the effects of watershed development. Estuaries 23(6):838-853.
- Parsons et al., 2002. Assessment of Sediment Toxicity and Quality in Patrick Bayou, Segment 1006, Harris County, Texas. Prepared for Patrick Bayou TMDL Lead Organization.
- Rakocinski, C.F., S.S. Brown, G.R. Gaston, R.W. Heard, W.W. Walker, and J.K. Summers. 1997. Macrobenthic responses to natural and contaminant-related gradients in northern Gulf of Mexico estuaries. Ecological Applications 7(4):1278-1298.
- Ritter, C. and P.A. Montagna. 1999. Seasonal hypoxia and models of benthic response in a Texas Bay. Estuaries 22(1):7-20.
- Roach, R.W., R.S. Carr, C.L. Howard, and B.W. Cain. 1992. An assessment of produced water impacts in the Galveston Bay system. Draft report prepared by U.S. Fish and Wildlife Service report. 61 pages.
- TCEQ, 2012. Draft, Development of a Tidal Stream Benthic Index in Galveston Bay, Texas. Linda Broach, May 2012.
- Weisberg, S.B., J.A. Ranasinghe, D.M. Dauer, L.C. Schaffner, R.J. Diaz, J.B. Frithsen. 1997.

 An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. Estuaries 20(1):149-158

TABLES

Table E-4.1 ES-BI September 2000 Calculations

									P10										
	Site Infor	mation		Trar		No	ata		Discriminant Scores						Benthic Index, "summer"				
Site	Station	Date	Mean Diversity Proportion	Log Turbellaria Abundunce		1	Arc-sin, % Amphipods	Mean Diversity Proportion	Log Turbellaria Abundunce		Arc-sin, % Capitillids	Arc-sin, % Amphipods	Mean Diversity Proportion	Log Turbellaria Abundunce			Arc-sin, % Amphipods	Sum Discriminant Score	EŞ-BI
Patrick Bayou	V	September 2000	0.078	0	0	0	0	-0.7954	-0.5756	0	-0.6879	-0.3015	-1.2495	0.5949	0	0.3857	-0.1514	-0.4204	3.719
Patrick Bayou	2.5	September 2000	0.138	0	0	0	0	-0.3207	-0.5756	0	-0.6879	-0.3015	-0.5039	0.5949	0	0.3857	-0.1514	0.3253	4.714
Patrick Bayou	S	September 2000	0.287	0	0	0.021	0	0.8579	-0.5756	0	1.9605	-0.3015	1.3478	0.5949	0	-1.0993	-0.1514	0.6920	5.203
Patrick Bayou	U	September 2000	0.002	0	0	0	0	-1.3966	-0.5756	0	-0.6879	-0.3015	-2.1940	0.5949	0	0.3857	-0.1514	-1.3649	2.460
Patrick Bayou	3	September 2000	0.127	0.097	0	0	0.041	-0.4078	-0.4325	0	-0.6879	3.0151	-0.6406	0.4470	0	0.3857	1.5145	1.7066	6.555
Patrick Bayou	G	September 2000	0.218	0.000	0	0	0	0.3121	-0.5756	0	-0.6879	-0.3015	0.4903	0.5949	0	0.3857	-0.1514	1.3195	6.039
Patrick Bayou	4A	September 2000	0.060	0.176	0	0.015	0	-0.9378	-0.3159	0	1.2038	-0.3015	-1.4732	0.3265	0	-0.6750	-0.1514	-1.9732	1.649
Patrick Bayou	5	September 2000	0.060	0.352	0	0.013	0	-0.9378	-0.0562	0	0.9516	-0.3015	-1.4732	0.0581	0	-0.5336	-0.1514	-2.1002	1.480
Patrick Bayou	Т	September 2000	0.362	0.966	0	0	0	1.4512	0.8498	0	-0.6879	-0.3015	2.2799	-0.8783	0	0.3857	-0.1514	1.6359	6.461
Patrick Bayou	6A	September 2000	0.290	2.223	0	0.011	0	0.8817	2.7046	0	0.6994	-0.3015	1.3851	-2.7952	0	-0.3921	-0.1514	-1.9537	1.675
Patrick Bayou	Q	September 2000	0.342	0.477	0	0	0	1.2930	0.1282	0	-0.6879	-0.3015	2.0314	-0.1325	0	0.3857	-0.1514	2.1331	7.124
		Mean	0.179	0.390	0	0.00545	0.00373												
		Standard Deviation	0.126	0.678	0	0.00793	0.0124												

Table E-4.2 ES-BI April 2001 Calculations

Site	Information	on		Tran	nsformed Da	ita			No	rmalized Da	ita		Discriminant Scores						
Site	Station	Date	Mean Diversity Proportion	Log Turbellaria Abundunce			Arc-sin, % Amphipods	Mean Diversity Proportion	Log Turbellaria Abundunce		Arc-sin, % Capitillids	Arc-sin, % Amphipods	Mean Diversity Proportion	Log Turbellaria Abundunce	Arc-sin, % Bivalves	Arc-sin, % Capitillids	Arc-sin, % Amphipods	Sum Discriminant Score	ES-BI
Patrick Bayou	V	April 2001	0.503	0.352	0.008	0.284	0	0.326	-0.754	0.449	2.81	-0.489	0.512	0.780	-0.201	-1.58	-0.246	-0.730	3.31
Patrick Bayou	2.5	April 2001	0.488	0.243	0.002	0.141	0	0.240	-0.932	-0.286	1.12	-0.489	0.378	0.963	0.128	-0.626	-0.246	0.597	5.08
Patrick Bayou	S	April 2001	0.482	0.903	0	0.035	0.005	0.206	0.14	-0.531	-0.140	0.522	0.324	-0.145	0.237	0.079	0.262	0.757	5.29
Patrick Bayou	Е	April 2001	0.644	0.978	0.023	0.041	0.017	1.13	0.262	2.29	-0.0691	2.95	1.78	-0.270	-1.02	0.039	1.48	2.00	6.95
Patrick Bayou	U	April 2001	0.367	0.954	0	0.008	0.004	-0.451	0.224	-0.531	-0.460	0.320	-0.708	-0.231	0.237	0.258	0.161	-0.283	3.90
Patrick Bayou	3	April 2001	0.68	0.978	0	0.005	0	1.34	0.262	-0.531	-0.496	-0.489	2.10	-0.270	0.237	0.278	-0.246	2.10	7.08
Patrick Bayou	G	April 2001	0.437	0.176	0	0.025	0.003	-0.0509	-1.04	-0.531	-0.259	0.118	-0.0800	1.08	0.237	0.145	0.0593	1.44	6.20
Patrick Bayou	4A	April 2001	0.506	0.301	0.019	0	0	0.343	-0.837	1.80	-0.555	-0.489	0.539	0.865	-0.803	0.311	-0.246	0.667	5.17
Patrick Bayou	5	April 2001	0.484	0.176	0	0	0	0.218	-1.04	-0.531	-0.555	-0.489	0.342	1.08	0.237	0.311	-0.246	1.72	6.57
Patrick Bayou	T	April 2001	0.527	0.889	0	0.023	0	0.463	0.118	-0.531	-0.282	-0.489	0.728	-0.122	0.237	0.158	-0.246	0.756	5.29
Patrick Bayou	6A	April 2001	0.147	1.90	0	0	0	-1.71	1.76	-0.531	-0.555	-0.489	-2.68	-1.82	0.237	0.311	-0.246	-4.20	-1.32
Patrick Bayou	Q	April 2001	0.086	1.95	0	0	0	-2.06	1.84	-0.531	-0.555	-0.489	-3.23	-1.90	0.237	0.311	-0.246	-4.82	-2.15
		Mean	0.446	0.817	0.00433	0.0468	0.00242												
	Standa	rd Deviation	0.175	0.616	0.00816	0.0844	0.00494												

Table E-4.3
Summary of Engle and Summers Benthic Index

Summary of Engle and Summers Benefice mack											
Station	Date	ES-BI Score ¹	Mean % Sand ²								
V	9/5/2000	3.72	21								
V	4/24/2001	3.31									
2.5	9/5/2000	4.71	52								
2.5	4/23/2001	5.08									
S	9/5/2000	5.20	46								
S	4/24/2001	5.29									
E	4/23/2001	6.95	56								
U	9/5/2000	2.46	22								
U	4/24/2001	3.90									
3	9/5/2000	6.56	43								
3	4/24/2001	7.08									
G	9/5/2000	6.04	78								
G	4/24/2001	6.20									
4A	9/5/2000	1.65	30								
4A	4/24/2001	5.17									
5	9/5/2000	1.48	42								
5	4/24/2001	6.57									
Т	9/5/2000	6.46	77								
Т	4/25/2001	5.29									
6A	9/5/2000	1.68	82								
6A	4/25/2001	-1.32									
Q	9/5/2000	7.12	67								
Q	4/25/2001	-2.15									

 $^{^{1}}$ Engle and Summers (1999) Benthic Index.

² Mean based on all sample events for each Patrick Bayou station.

Table E-4.4
TCEQ Tidal Stream-Tolerant/Opportunistic/Pollution Indicative Taxa

Class/Phylum	Organism	Reference							
Oligochaeta	Oligochaeta	Brown et al., 2000; Ritter and Montagna, 1999; Weisberg et al., 1997							
	Capitella spp	Rakocinski et al., 1997; Weisberg et al., 1997							
	Laeonereis culveri	Lerberg et al., 2000							
Polychaeta	Paraprionospio pinnata	Dauer et al., 1992; Weisberg et al., 1997							
,	Streblospio benedicti	Dauer et al., 1992; Rakocinski et al., 1997; Ritter and Montagna, 1999; Roach et al., 1992; Weisberg et al., 1997							
Insecta	Chironomidae	Weisberg et al., 1997							
Mollusca	Macoma mitchilli	Rakocinski et al., 1997							
IVIUIIUSCa	Mulinia lateralis	Dauer et al., 1992; Weisberg et al., 1997							

Table derived from draft TCEQ white paper, "Development of a Tidal Stream Benthic Index in Galveston Bay, Texas" (TCEQ, May 2012)

Table E-4.5
TCEQ Tidal Stream-Intolerant/Pollution Sensitive Taxa

Class/Phylum	Organism	Reference
	Aricidea philbinae	Rakocinski et al., 1997
	Asychis elongata	Dauer et al., 1992; Weisberg et al., 1997
	Clymenella torquata	Dauer et al., 1992; Weisberg et al., 1997
	Diopatra cuprea	Dauer et al., 1992; Weisberg et al., 1997
Polychaeta	Glycera americana	Weisberg et al., 1997
Polychaeta	Glycinde solitaria	Weisberg et al., 1997
	Heteromastus filiformis	Lerberg et al., 2000
	Mediomastus	Brown et al., 2000; Weisberg et al., 1997
	Spiochaetopterus costarum	Weisberg et al., 1997
	Spiophanes bombyx	Weisberg et al., 1997
Crustacea	Listriella clymenellae	Weisberg et al., 1997
	Cyrtopleura costata	Dauer et al., 1992; Weisberg et al., 1997
Mollusca	Rangia cuneata	Dauer et al., 1992; Weisberg et al., 1997
ivioliusca	Tagalus divisus	Dauer et al., 1992; Rakocinski et al., 1997;
	Tagelus divisus	Weisberg et al., 1997
Nemertea	Nemertea	Lerberg et al., 2000
Echinodermata	Micropholis atra	Dauer et al., 1992; Weisberg et al., 1997

Table derived from draft TCEQ white paper, "Development of a Tidal Stream Benthic Index in Galveston Bay, Texas" (TCEQ, May 2012)

Table E-4.6
TCEQ Tidal Stream Metric Score Calculations

Metric	Metric Explanation	Tidal Stream Reference Group 90th Percentile (or 10th)	Computation of Metric Score	Range ^b
Taxa	# of taxa in sample	13	(Taxa/13)*10	0 to 10
Diversity	H' = Shannon-Weiner diversity	1.85	(Div/1.85)*10	0 to 10
Major Taxa	Polychaeta, Crustacea, Nemertea, Bivalvia, Gastropoda	5	#MajorTaxa*2	0 to 10
% Intolerant	% of individuals from intolerant taxa	0.7	(%Intol/0.70)*10	0 to 10
% Tolerant ^a	% of individuals from tolerant taxa	0.16 (10 th %ile)	10*(1-%Toler)/(1-0.16)	0 to 10
% Dominant ^a	% of the sample from the top 3 species	0.71 (10 th %ile)	10*(1-%Dom)/(1-0.71)	0 to 10

- a If no organisms present in sample, value of metric is 0.
- b If a metric score exceeds 10, it is set equal to 10.

Table derived from draft TCEQ white paper, "Development of a Tidal Stream Benthic Index in Galveston Bay, Texas" (TCEQ, May 2012)

Table E-4.7A
Patrick Bayou September 2000 Benthic Community Data and TCEQ BI Scores

П	STATION*												TCEQ,	TCEQ,							
Taxon	9	8	7	R	Υ	Q	6A	т	5	4A	G	3	U	E	S	2.5	v	2	Intolerant	Tolerant	Taxa group
Nemertea	_				1					1					2			5	Υ		Nemertea
Nematoda							4											5			Nematoda
Hirudinea		17																			Annelida
Oligochaeta		20	14	10	80	9	715	36	5	2		1						5		Υ	Annelida
Streblospio benedicti						1	4	5	11	6					32	14	170	264	İ	Υ	Polychaeta
Laeonereis culveri								85	682	573	3	46	16	101	58	99	1	2	İ	Υ	Polychaeta
Amphicteis floridus ¹					23	29	168	179		3					2						Polychaeta
Parandalia americana																2		10			Polychaeta
Neanthes succinea							40											1			Polychaeta
Capitella capitata									10	9					3					Υ	Polychaeta
Mediomastus californiensis							6												Y		Polychaeta
Gastropoda (U)												1									Gastropoda
Ostracoda					1																Crustacea
Acartia tonsa											1	1									Crustacea
Gammarus sp.												1									Crustacea
Ericthonius sp.												1									Crustacea
Macrobrachium sp.											2										Crustacea
Callinectes sapidus			1																		Crustacea
Americamysis bahia																	11	22			Crustacea
Chironomidae	9465	46		1	1	3	10													Υ	Insecta
Total Abundance	9465	83	15	11	106	42	947	305	708	594	6	51	16	101	97	115	182	314			
Richness	1	3	2	2	5	4	7	4	4	6	3	6	1	1	5	3	3	8			
Diversity not salinity corrected. From Parsons 2002	NC	0.997	0.244	0.324	0.682	0.871	0.782	0.991	0.196	0.206	0.713	0.421	0.005	NC	0.956	0.46	0.26	0.701			
Diversity, calculation	0.000	0.995	0.245	0.305	0.676	0.863	0.779	0.988	0.196	0.201	1.011	0.479	0.000	0.000	0.941	0.456	0.262	0.690			
TCEQ, Taxa, Score	0.769	2.308	1.538	1.538	3.846	3.077	5.385	3.077	3.077	4.615	2.308	4.615	0.769	0.769	3.846	2.308	2.308	6.154			
TCEQ, Diversity Score, New	0.000	5.377	1.324	1.647	3.654	4.667	4.210	5.343	1.059	1.088	5.467	2.587	0.000	0.000	5.086	2.464	1.416	3.730			
TCEQ, Diversity Score, Parsons	0.000	5.389	1.319	1.751	3.686	4.708	4.227	5.357	1.059	1.114	3.854	2.276	0.027	0.000	5.168	2.486	1.405	3.789			
TCEQ, Intolerant, %; Score	0.000	0.000	0.000	0.000	0.135	0.000	0.091	0.000	0.000	0.024	0.000	0.000	0.000	0.000	0.295	0.000	0.000	0.227			
TCEQ, Tolerant, %; Score	0.000	2.438	0.794	0.000	2.808	8.220	2.740	6.987	0.000	0.080	5.952	0.934	0.000	0.000	0.491	0.207	0.720	1.630			
TCEQ Five Taxa, Score	0.000	0.000	2.000	0.000	6.000	2.000	2.000	2.000	2.000	4.000	4.000	6.000	2.000	2.000	4.000	2.000	4.000	6.000			
Top Three Taxa Score	0.000	0.000	0.000	0.000	0.651	0.821	0.874	0.565	0.244	0.348	0.000	2.028	0.000	0.000	1.422	0.000	0.000	1.977			
Total of Metric Scores, Parsons Diversity	0.769	10.135	5.651	3.290	17.126	18.826	15.317	17.986	6.380	10.181	16.114	15.853	2.796	2.769	15.221	7.001	8.433	19.778			
Total of Metric Scores	0.769	10.123	5.656	3.185	17.093	18.785	15.300	17.972	6.379	10.156	17.727	16.164	2.769	2.769	15.140	6.979	8.443	19.719			

Values are mean number of individuals per square meter

NC = No Calculation

^{*}Data from Parsons et al. 2002; samples collected September 2000

¹Formerly *Hobsonia (Hypaniola) florida*

Table E-4.7B
Patrick Bayou April 2001 Benthic Community Data and TCEQ BI Scores

	STATION*									TCEQ,	TCEQ,								
Taxon	7	R	Υ	Q	6A	Т	5	4A	G	3	U	E	s	2.5	V	2	Intolerant	Tolerant	Taxa group
Turbellaria	1.1	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	1.1	0.0	0.0	2.2	0.0	0.0	0.0			Turbellaria
Nemertea	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	1.1	0.0	0.0	0.0	Y		Nemertea
Nematoda	0.0	0.0	2.2	3.2	0.0	1.1	0.0	0.0	0.0	8.6	0.0	20.5	7.5	0.0	0.0	0.0			Nematoda
Hirudinea	3.2	24.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	11.9	0.0	0.0	0.0	0.0	0.0	0.0			Annelida
Oligochaeta	116.9	285.5	1186.3	377.1	339.4	29.1	2.2	4.3	2.2	36.6	34.5	36.6	30.2	3.2	5.4	14.0		Υ	Annelida
Streblospio benedicti	1.1	0.0	0.0	0.0	1.1	2.2	0.0	0.0	1.1	1.1	0.0	0.0	0.0	0.0	23.7	0.0		Υ	Polychaeta
Laeonereis culveri	0.0	0.0	0.0	0.0	1.1	1.1	25.9	2.2	133.6	40.9	141.2	6.5	167.0	208.0	55.0	0.0		Υ	Polychaeta
Amphicteis floridus ¹	0.0	1.1	1.1	1.1	18.3	18.3	14.0	9.7	55.0	93.7	33.4	90.5	38.8	86.2	200.4	0.0			Polychaeta
Parandalia americana	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			Polychaeta
Neanthes succinea	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	1.1	1.1	6.5	8.6	55.0	145.5	0.0			Polychaeta
Capitella capitata	0.0	1.1	0.0	0.0	0.0	1.1	0.0	0.0	1.1	1.1	1.1	6.5	8.6	55.0	145.5	0.0		Υ	Polychaeta
Mediomastus californiensis	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	Υ		Polychaeta
Polydora sp.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0			Polychaeta
Littoridina sp.	5.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			Gastropoda
Macoma mitchelli	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	0.0	5.4	0.0	1.1	1.1	0.0		Υ	Mollusca
Ostracoda	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	0.0	0.0	0.0	0.0			Crustacea
Calanoida	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	1.1	0.0	0.0	2.2	0.0			Crustacea
Acartia tonsa	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	1.1	0.0	0.0	0.0			Crustacea
Cyclopoida	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	1.1	9.7	0.0	2.2	2.2	0.0	0.0	1.1			Crustacea
Harpacticoida	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			Crustacea
Gammarus sp.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0			Crustacea
Ericthonius sp.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	2.2	1.1	0.0	0.0	0.0			Crustacea
Idoteidae	0.0	0.0	2.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	0.0	0.0	0.0	4.3			Crustacea
Cladocera	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4.3	0.0	0.0	0.0	0.0	0.0	0.0			Crustacea
Chironomidae (larvae)	78.7	176.7	19.4	14.0	11.9	25.9	14.0	35.6	8.6	50.6	0.0	42.0	9.7	0.0	1.1	2.2		Υ	Insecta
Odonata (Larvae)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0			Insecta
Total Abundance	206.4	490.3	1211.2	395.4	371.8	79.9	56.1	56.2	204.9	262.9	213.5	225.5	278.1	409.6	581.0	21.6			
Richness	6	6	5	4	5	8	4	8	9	14	7	14	12	7	10	4			
Diversity, calculation	0.905	0.875	0.116	0.219	0.376	1.406	1.177	1.224	0.954	1.827	0.967	1.821	1.394	1.281	1.515	0.987			
,																			
TCEQ, Taxa, Score	4.615	4.615	3.846	3.077	3.846	6.154	3.077	6.154	6.923	10.000	5.385	10.000	9.231	5.385	7.692	3.077			
TCEQ, Diversity Score, New	4.894	4.728	0.626	1.183	2.033	7.597	6.360	6.616	5.157	9.877	5.226	9.844	7.534	6.924	8.186	5.333			
TCEQ, Intolerant, %; Score	0.000	0.032	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.060	0.000	0.000	0.057	0.000	0.027	0.000			
TCEQ, Tolerant, %; Score	0.559	0.656	0.054	0.129	0.586	3.054	2.971	2.754	3.387	6.004	2.046	6.784	2.680	4.136	7.155	2.976			
TCEQ Five Taxa, Score	4.000	4.000	4.000	2.000	2.000	4.000	2.000	6.000	4.000	6.000	4.000	6.000	6.000	4.000	6.000	2.000			
Top Three Taxa Score	0.902	0.232	0.094	0.096	0.204	2.848	1.352	4.050	1.296	10.000	0.711	8.625	5.220	5.085	5.318	1.756			
Total of Metric Scores	14.971	14.263	8.620	6.485	8.669	23.654	15.760	25.573	20.763	41.942	17.367	41.253	30.721	25.530	34.379	15.142			

Values are mean number of individuals per square meter

^{*}Data from Parsons et al. 2002; samples collected April 2001

¹Formerly *Hobsonia (Hypaniola) florida*

Table E-4.7C
Patrick Bayou August 2003 Benthic Community Data and TCEQ BI Scores

_			STAT	ION*	TCEQ,		_		
Taxon	Е	3	4A	6A	R	7	Intolerant	TCEQ, Tolerant	Taxa group
Empididae (Diptera larva)	0	0	0	2.75	0	0			Insecta
Hirudinea	0	5.5	0	0	0	5.5			Annelida
Oligochaeta	427.75 1692 72.75 998 266.25 8		86		Υ	Annelida			
Capitella capitata	0	5.5	2.75	0	0	0		Υ	Polychaeta
Amphicteis floridus ¹	616	449.25	53.75	8	0	0			Polychaeta
Laeonereis culveri	390	1331.5	395.5	8	0	0		Υ	Polychaeta
Mediomastus ambiseta	0	2.75	0	0	0	0	Υ		Polychaeta
Polydora ligni	5.5	0	0	0	0	0			Polychaeta
Streblospio benedicti	118.25	56.5	161.5	0	0	0		Υ	Polychaeta
Mulinia lateralis	2.75	0	0	0	0	0			Bivalvia
Rangia cuneata	2.75	0	0	0	0	0	Υ		Bivalvia
Texadina sp.	13.5	2.75	0	0	0	0			Gastropoda
Chironomidae	10.75	8	0	5.5	13.5	13.5		Υ	Insecta
Gnathiidae	0	0	0	0	0	0			Crustacea
Edotia sp.²	0	8	2.75	0	0	0			Crustacea
Amphipoda	5.5	0	0	0	0	0			Crustacea
Ampelisca sp.	0	0	0	2.75	0	0			Crustacea
Total Abundance	1592.75	3561.75	689	1025	279.75	105			
Richness	10	10	6	6	2	3			
Diversity, Calculation	1.393	1.107	1.139	0.162	0.193	0.582			
TCEQ, Taxa, Score	7.692	7.692	4.615	4.615	1.538	2.308			
TCEQ, Diversity Score	7.532	5.982	6.158	0.873	1.045	3.144			
TCEQ, Intolerant, %; Score	0.025	0.011	0.000	0.000	0.000	0.000		†	
TCEQ, Tolerant, %; Score	4.828	1.565	0.976	0.157	0.000	0.624		†	
TCEQ Five Taxa, Score	8.000	6.000	4.000	4.000	0.000	0.000		† †	
Top Three Taxa Score	3.442	0.862	2.965	0.370	0.000	0.000			
Mean of Metric Scores	31.519	22.113	18.715	10.016	2.584	6.076			

Values are mean number of individuals per square meter

^{*}Data from Parsons et al. 2004; samples collected August 2003

¹Formerly *Hobsonia (Hypaniola) florida*

²Originally *Edotea* (invalid; literature misspelling)

Table E-4.8
Summary of TCEQ Benthic Index

Julillary of TCLQ Bentine Index											
Station	Date	TCEQ Total Scores									
V	9/5/2000	8.44									
V	4/24/2001	34.38									
2.5	9/5/2000	6.98									
2.5	4/23/2001	25.53									
S	9/5/2000	15.14									
S	4/24/2001	30.72									
Е	9/5/2000	2.77									
E	4/23/2001	41.25									
Е	8/12/2003	31.52									
U	9/5/2000	2.77									
U	4/24/2001	17.37									
3	9/5/2000	16.16									
3	4/24/2001	41.94									
3	8/12/2003	22.11									
G	9/5/2000	17.73									
G	4/24/2001	20.76									
4A	9/5/2000	10.16									
4A	4/24/2001	25.57									
4A	8/12/2003	18.71									
5	9/5/2000	6.38									
5	4/24/2001	15.76									
Т	9/5/2000	17.97									
Т	4/25/2001	23.65									
6A	9/5/2000	15.30									
6A	4/25/2001	8.67									
6A	8/12/2003	10.02									
Q	9/5/2000	18.78									
Q	4/25/2001	6.49									

ATTACHMENT E-4.1 PROUCL STATISTICS

Attachment E-4.1 ProUCL Statistics

From File: C:\Patrick Bayou\BERA 2012\Comments\Benthic_Community\Analysis\E_S_TCEQ_Total_2.xls.wst

Summary Statistics for Raw Full Data Sets

Variable	NumObs	Minimum	Maximum	Mean	Median	Variance	SD	MAD/0.675	Skewness	Kurtosis	CV
TCEQ Total Score	12	10.07	26.74	17.99	18.7	32.79	5.726	7.632	-0.0297	-1.341	0.318
ES-BI Relative Benthic Indexc	12	0.177	6.953	4.391	4.461	3.94	1.985	1.996	-0.627	0.275	0.452
				Percentiles	for Raw Full D	ata Sets					
Variable	NumObs	5%ile	10%ile	20%ile	25%ile(Q1)	50%ile(Q2)	75%ile(Q3)	80%ile	90%ile	95%ile	99%ile
TCEQ Total Score	12	10.62	11.1	11.59	12.31	18.7	21.79	22.63	24.96	25.88	26.57
ES-BI Relative Benthic Indexc	12	1.447	2.556	3.227	3.352	4.461	5.935	6.069	6.747	6.878	6.938

APPENDIX E-5 PROUCL CALCULATIONS FOR SURFACE WATER TOTAL PCB EXPOSURE POINT CONCENTRATION

Appendix E-5 95 UCL Results for PCB Congeners (U=1/2)

General UCL Statistics for Data Sets with Non-Detects

User Selected Options

From File Sheet1.wst

Full Precision OFF

Confidence Coefficient 95%

Number of Bootstrap Operations 2000

TPCB

General Statistics

Number of Valid Observations 18 Number of Distinct Observations 18

Raw Statistics Log-transformed Statistics

> Minimum 5.65 Minimum of Log Data 1.732 Maximum 431 Maximum of Log Data 6.066 Mean 141.6 Mean of log Data 4.708

Geometric Mean 110.8 SD of log Data 0.882 Median 114.6

Std. Error of Mean 21.85

Coefficient of Variation 0.655

Skewness 1.83

SD 92.7

Relevant UCL Statistics

Normal Distribution Test Lognormal Distribution Test

Shapiro Wilk Test Statistic 0.845 Shapiro Wilk Test Statistic 0.782 Shapiro Wilk Critical Value 0.897 Shapiro Wilk Critical Value 0.897

Data not Lognormal at 5% Significance Level

99% Chebyshev (MVUE) UCL 517.3

Data not Normal at 5% Significance Level

Assuming Normal Distribution Assuming Lognormal Distribution

95% Student's-t UCL 179.6 95% H-UCL 277.2 95% UCLs (Adjusted for Skewness) 95% Chebyshev (MVUE) UCL 315.9 95% Adjusted-CLT UCL (Chen-1995) 187.6 97.5% Chebyshev (MVUE) UCL 383.8 95% Modified-t UCL (Johnson-1978) 181.2

Gamma Distribution Test Data Distribution

> k star (bias corrected) 1.866 **Data appear Gamma Distributed at 5% Significance Level**

> > Theta Star 75.89

MLE of Mean 141.6

MLE of Standard Deviation 103.6

nu star 67.16

Approximate Chi Square Value (.05) 49.3 **Nonparametric Statistics**

Adjusted Level of Significance 0.0357 95% CLT UCL 177.5 Adjusted Chi Square Value 47.83 95% Jackknife UCL 179.6

95% Standard Bootstrap UCL 176.6

Anderson-Darling Test Statistic 0.631 95% Bootstrap-t UCL 196.8 Anderson-Darling 5% Critical Value 0.751 95% Hall's Bootstrap UCL 328.7

Appendix E-5 95 UCL Results for PCB Congeners (U=1/2)

Kolmogorov-Smirnov Test Statistic 0.184

Kolmogorov-Smirnov 5% Critical Value 0.206

Data appear Gamma Distributed at 5% Significance Level

95% Percentile Bootstrap UCL 178.5 95% BCA Bootstrap UCL 188.4 95% Chebyshev(Mean, Sd) UCL 236.8 97.5% Chebyshev(Mean, Sd) UCL 278 99% Chebyshev(Mean, Sd) UCL 359

Assuming Gamma Distribution

95% Approximate Gamma UCL (Use when n >= 40) 192.9 95% Adjusted Gamma UCL (Use when n < 40) 198.8

Potential UCL to Use

Use 95% Approximate Gamma UCL 192.9

Note: Suggestions regarding the selection of a 95% UCL are provided to help the user to select the most appropriate 95% UCL.

These recommendations are based upon the results of the simulation studies summarized in Singh, Singh, and Iaci (2002) and Singh and Singh (2003). For additional insight, the user may want to consult a statistician.

APPENDIX F COMMENTS AND RESPONSES MATRIX FOR TCEQ, NOAA, AND USEPA DRAFT BASELINE ECOLOGICAL RISK ASSESSMENT

Comments and Responses Matrix for TCEQ, NOAA, and USEPA Draft Baseline Ecological Risk Assessment

Comment No.	Section	Comment	Response to Comment - Proposed Revision			
Overall Comment - TCEQ		Using historical benthic community data from the site and nearby tidal bayous, the BERA concludes that the condition of the benthic community at the site does not differ significantly from local comparison sites, that there is no significant relationship between chemicals of potential concern (COPCs) and the condition of the benthic community, and that COPCs are not identified that represent an unacceptable adverse risk to the benthic community at the site. No risk management recommendations were deemed necessary for this receptor group. We disagree with this assessment. First, we believe the benthic community evaluation was flawed for reasons outlined in the comments that follow. In fact, our analysis indicates that the benthic community in Patrick Bayou is degraded compared to the comparison sites presented in the BERA. This conclusion withstands even when data for the gunite-lined portion of the bayou is removed from the evaluation. Secondly, Patrick Bayou appears impacted by site COPCs based on the other two legs of the sediment quality triad (i.e., Chapman 1990). Surficial sediment concentrations for mercury, PCBs (polychlorinated biphenyls), and Total PAHs (polycyclic aromatic hydrocarbons) consistently exceed the second effects level which indicate a concentration threshold above which adverse effects can frequently occur (i.e., Long et al. 1995; TCEQ 2006). Additionally, sediment toxicity tests conducted as part of the TMDL (Total Maximum Daily Load) effort and by U.S. EPA/TCEQ indicated that site sediments were frequently toxic on multiple occasions at multiple locations (e.g., BERA Work Plan, Table 17). As this was a draft document, we believe continued discussions with regulators should occur before the final BERA is submitted. We can provide details of our benthic evaluation, and we can discuss potential responses to comments along with corresponding changes to the BERA. However, we believe the ultimate risk management decision for the benthic invertebrate exposure pathway should reflect a concentra	have attempted to respond to specific agency comments and concerns regarding the benthic risk assessment in the responses that follow.			
1	4	BERA Dataset - The information for notes a and b in Table 4-1 should be added.	The requested information will be provided in the revised BERA.			
2	4.1.2.3	Toxic Equivalency Quotients - The table reference for the source of the toxic equivalency factors (TEFs) provided by U.S. EPA (2008) should be Table 4 rather than Table 4-2.	The text is referring to Table 4-2 in the BERA Report, which was taken from USEPA (2008).			
3	4.7.2	Comparison Site Data - The discussion should include additional information on the benthic sample collection effort in Dobberstine (2007). In the Dobberstine study, five replicate 4-inch cores were collected at each sample location. This methodology results in a sediment sample volume of about 405 cm2 compared to a volume of roughly 929 cm2 for samples collected in Patrick Bayou. Normally in estuarine benthic communities, a few species are relatively abundant whereas other species are less abundant or relatively rare. Amphipods, gastropods, bivalves, and nemerteans are usually among these less common species. When smaller sample sizes are used, the rare species are often missed. This affects estimates of species richness and diversity because fewer species are found. This also affects any metrics that use these rarer species, such as percent bivalves, amphipods, or intolerant organisms. Please revise the BERA text.	The use of different sample devices may result in differences in estimates of abundance and diversity due to differences in area (i.e., cm2) sampled and sample depth retrieval. Without side by side sample collection at the Site and representative reference sites it is not known if the differences in sampling devices between the Site and reference area are significant. However, the significance of these differences are not relevant as we are no longer using the Dobberstine (2007) data in the BERA.			
4	6.1.2.1.1	Prey Groupings - Analysis by Prey Species - Looking at Appendix C and Table 6-2, it seems appropriate that Gulf killifish as prey were separated out for assessing exposure to PCB toxic equivalent quotient (TEQ, avian) concentrations. However, it appears that PCB TEQ (avian) concentrations in Gulf menhaden and striped mullet were statistically higher than those for pinfish and sand sea trout. Why were these fish not separated out as well?	Gulf menhaden and striped mullet will be separated from sand seatrout and pinfish in the prey grouping and EPC calculations for PCB TEQ (avian). Hazard Quotients (HQ) for belted kingfisher and carnivorous bird will be recalculated using the revised groupings and associated EPCs. Revised HQs will be presented in the BERA.			

Comments and Responses Matrix for TCEQ, NOAA, and USEPA

Draft Baseline Ecological Risk Assessment

		Draft Baseline Ecological Risk Assessi	
5	6.1.2.1.1	Prey Groupings - Spatial Trend Analysis - Appendix C-1.2 displays plots of tissue concentrations by distance from the mouth of Patrick Bayou. The intent was to evaluate potential spatial trends in prey item concentrations along the bayou. Overall, the discussion indicates that significant spatial patterns were not apparent for most COPCs and species, and no prey subgroups were identified based on spatial differences in COPC concentrations in prey tissue. The text in this section provides a few r ² and p-values. No similar information was provided with the figures in Appendix C-	revised BERA.
		1.2. The appendix should be revised to include this information, as appropriate, for each COPC/prey item pair.	
6	6.1.2.1.2	95UCL - The exposure point concentrations (EPCs) for each receptor and their respective prey groups are provided in Table 6-3. For the most part, the various ProUCL summary pages indicating the 95% Upper Confidence Limits (UCLs) for each prey grouping were provided in Appendix C. However, we could not locate the 95% UCL information for the blue crab/oyster/white shrimp prey grouping. Please provide this information. Additionally, if a maximum or non-detect proxy value was used, please indicate this in Table 6-3.	The 95UCL output / calculation sheets for the blue crab/white shrimp/oyster prey subgroup will be provided in Appendix C-2 of the revised BERA. Table 6-3 will be revised to identify any prey item exposure point concentrations (EPCs) based on a maximum detected value or a proxy value (i.e., non-detect).
7	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Section 1.1 (Method Selection) of Appendix B compares inverse distance weighting (IDW) and Kriging interpolation methods using lead and hexachlorobenzene. Please present a similar comparison (graphics, calculated values) for PCBs (congeners, TEQ) since this was the primary risk driver for wildlife.	Appendix B will be updated to include a comparison of IDW and Kriging for PCB Congener TEQ (mammalian and avian TEQs).
8	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Was the gunite-lined portion of Patrick Bayou included in the SWAC calculations for wildlife? Either way, what was the basis for the decision?	Yes, the gunite channel was included in the SWAC calculation for estimating the $C_{k,SD}$ term, as some intertidal habitat is available in this portion of the Site. However, the available intertidal habitat $(0.006~km^2)$ in this portion of the Site is minimal and has no significant incremental effect on the Ck,SD term. To support this statement, the uncertainty section will evaluate the differences in wildlife COPC SWACs with and without the gunite channel included and the incremental effect on the risk characterization for these scenarios (Appendix C-5.8). In addition, as discussed in the December 10, 2012 meeting among USEPA, TCEQ, trustees, and the JDG, this portion of the Site has limited and poor quality benthic habitat that does not support a benthic community. Thus, feeding and foraging by wildlife receptors would be limited.
9	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Section 6.1.2.1 of the BERA Work Plan states that a hot spot analysis will be performed as part of the uncertainty assessment for the concentration term (i.e., Ck,SD) to evaluate if areas of elevated sediment COPC may occur within a foraging area that would lead to potentially higher exposure than indicated by use of a SWAC for this term. Was this done? If not, this evaluation should be added to the BERA.	Areas of elevated COPCs will be defined as a station where concentrations of a COPC exceed the 90th percentile of Site-wide surface weighted average concentration. The minimum reported home range estimate for a given receptor will be applied to define the distance from the elevated COPC location that will be used in the calculation of the sediment concentration term. Centered on the location, the exposure concentration will be calculated as the mean of the raster surface within the home range areas as appropriately defined by the area the receptor is able to forage within. For example, the sediment concentration term for raccoon would be calculated from the raster surface defined by the supratidal and intertidal zones within a 1.33 km distance centered or the elevated location. This analysis and results will be provided in the wildlife risk assessment uncertainty discussion.
10	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Methylmercury Estimates in Sediment - Available Site data were evaluated to estimate the proportion of methylmercury (MeHg) to total mercury in site sediments. These proportions were then used to estimate bulk sediment concentrations of MeHg and inorganic mercury for the incidental sediment ingestion pathway exposure assessment (for wildlife). In summary, estimates of MeHg in sediments were developed using simple equilibrium partitioning models and site-specific measurements of porewater MeHg, dissolved organic carbon (DOC), and bulk sediment total organic carbon (TOC). Details of this analysis are provided in Appendix C-3. Please provide a general discussion indicating why MeHg was not analyzed in site (bulk) sediment samples.	The requested discussion will be provided. The discussion will focus on the advantages of porewater analysis of MeHg over bulk sediment.

Comments and Responses Matrix for TCEQ, NOAA, and USEPA Draft Baseline Ecological Risk Assessment

		Diait baselile Ecological Nisk Assess	
11	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Methylmercury Estimates in Sediment - After a quick on-line search, it appears that the Skyllberg (2008) reference in Appendix C-3 is actually an article in Journal of Geophysical Research – Biogeosciences. Please update the reference as appropriate.	The reference will be updated in the revised BERA as: Skyllberg, U. (2008). Competition among thiols and inorganic sulfides and polysulfides for Hg and MeHg in wetland soils and sediments under suboxic conditions: illumination of controversies and implications for MeHg net production. J Geophys Res Biogeosci 113:536–554. doi:G00c0310.1029/2008jg000745.
12A	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Methylmercury Estimates in Sediment - Using the formulas indicated in Appendix C-3, an estimated MeHg _{sed} value was calculated for each sample location (Table 2 and 3) using bulk sediment TOC data for the nearest surface grab. It is unclear how the nearest surface grab was selected to pair with the porewater (DOC and MeHg) data. Also, since these analyses were not based on contemporaneous sampling events (2007 and 2009, footnotes page C-2), there is some uncertainty in the application of the assumptions. Please provide more information on both of these points.	A figure will be included to show the location and distances between bulk sediment TOC samples and MeHg porewater samples (Appendix C-3, Figure C-1). Sensitivity analysis will be performed using the range of observed bulk sediment TOC values to evaluate the uncertainty in the lack of synoptic sampling for this analysis.
12B	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Methylmercury Estimates in Sediment - Using the formulas indicated in Appendix C-3, an estimated MeHg _{sed} value was calculated for each sample location (Table 2 and 3) using bulk sediment TOC data for the nearest surface grab. It is unclear how the nearest surface grab was selected to pair with the porewater (DOC and MeHg) data. Also, since these analyses were not based on contemporaneous sampling events (2007 and 2009, footnotes page C-2), there is some uncertainty in the application of the assumptions. Please provide more information on both of these points.	A figure will be included to show the location and distances between bulk sediment TOC samples and MeHg porewater samples. Sensitivity analysis will be performed using the range of observed bulk sediment TOC values to evaluate the uncertainty in the lack of synoptic sampling for this analysis.
13	6.1.2.2.1	Sediment Surface Weighted Average Concentration (SWAC) - Methylmercury Estimates in Sediment - Please revise Appendix C-3 (page 2) to better explain the various depth intervals over which TOC concentrations (from any one sample location) were obtained and used in this approach. As it is, the draft language does not pair up with the information in Table 2 of the appendix. Anchor QEA representatives have proposed revised language in an e-mail to TCEQ.	Appendix C-3 will be revised to clarify the depth intervals over which TOC were obtained and included in the analysis. This revised language will reflect the previous proposed language shared with TCEQ.
14	6.1.2.3	Terrestrial Animals and Plant Matter - Regarding terrestrial animals as prey for carnivorous birds and raccoons, the discussion states that none of these prey items would be expected to have significant exposure to bioaccumulative COPCs in site sediments and surface water; so, the terrestrial animal diet component for the raccoon and carnivorous birds was set to zero for the risk characterization. This statement is in conflict with a response to comment on the BERA Work Plan. In a comment regarding Section 6.2 of the Work Plan, we had asked (see page 20 of the response to comments table) if the intent was to model hypothetical exposure to terrestrial prey that have foraged within Patrick Bayou. The response was: "Yes, our intent is to model hypothetical exposure to terrestrial prey that may forage in Patrick Bayou (see Sections 6.2.2 and 6.2.4 in the BERA [Work Plan] for references to the equations we will use)." Please justify this apparent shift in the approach.	While exposure to sediment-associated COPCs to carnivorous birds and raccoon from ingestion of terrestrial animals and plants in the riparian zone are likely to be negligible, the baseline risk estimates will be revised to include these exposure pathways. Estimation of exposure through ingestion of terrestrial plant and animal matter will follow the procedures presented in the BERA Work Plan (2011). Revised baseline HQ reflecting these exposure pathways will be presented in the baseline risk characterization discussion. The relative contribution of terrestrial animal and plants to exposure and risk will be compared to the contributions from consumption of fish and invertebrates and incidental ingestion of sediment and summarized as part of the uncertainty analysis.
15	6.1.3	Area Use Factors - We were not able to duplicate the average (belted kingfisher) and upper 75 th percentile (belted kingfisher and spotted sandpiper) home range values provided in Table 6-5. Please verify these calculations.	The average belted kingfisher home range will be revised to 2.0 in the revised BERA. Statistical documentation of the upper quartiles for belted kingfisher and spotted sandpiper will be included in Appendix C-4 (Exhibit C-4.2) of the revised BERA.
16	6.2.1	Development of Inorganic Mercury TRV (Toxicity Reference Value) - The avian LOAEL (lowest observed adverse effect level) TRV for Total PAH (325 mg/kg/day, Table 6-6) differs from the value of 40 mg/kg/day set out in the BERA Work Plan (Table 29). Please provide the justification for this change and present the derivation of the final value based on the LOAEL value in the cited study.	The difference in the TRV in the Work Plan and that used in the BERA is due to assumptions for food uptake rate and body weight used to calculate the LOAEL as presented in the BERA Work Plan and that used in the BERA itself. The same study was used in the derivation of the TRV for both the Work Plan and the BERA. The effect level at the LOAEL was 4,000 mg/kg total PAHs (Patton and Dieter 1980). In the Work Plan we assumed that the food uptake was 1% of the body weight and the body was assumed to 1.0 kg. Therefore, we assumed a low food uptake rate for a duck. For the BERA itself, the assumptions were updated so that food uptake was 100 mg/day (0.1 kg/day) and the weight was derived from the original study, 1.23 kg. These two new assumptions allowed the calculation of 325 mg/kg/day as the LOAEL.

Comments and Responses Matrix for TCEQ, NOAA, and USEPA

Draft Baseline Ecological Risk Assessment

	Diate baseline Ecological risk Assessment					
17	6.3.2.4	Spotted Sandpiper - For the spotted sandpiper, incidental sediment ingestion was modeled as 18% of the total daily intake of food (Table 6-1). This conflicts with the value of 30% that was presented in the approved BERA Work Plan (Table 28 and Section 6.2.1). Please provide a justification for this move from the value presented in the BERA Work Plan.	The incidental sediment ingestion rate of 30% was intended to be used during the selection of wildlife COPC but was not intended for use in the baseline risk assessment. During a review of Beyer et al. (1994) it was noted that sediment ingestion by four sandpiper species (i.e., stilt sandpiper, semipalmated sandpiper, least sandpiper, and western sandpiper) ranged from 7.3 to 30% of the dry diet, with an average incidental sediment ingestion of 18%. An incidental sediment ingestion rate of 18% was selected for the baseline risk assessment consistent with the selection of more realistic exposure parameters for wildlife in the baseline risk assessment and consistent with another CERCLA Site (i.e., Portland Harbor). The BERA will be revised to discuss this as a change from the Work Plan and the basis for the change (described above) will be included.			
18	6.3.3.1.1	Prey Item Grouping - It does not appear that the uncertainty associated with Total PCB congeners was part of this analysis. Please add this COPC to the uncertainty evaluation or explain why it was excluded.	The uncertainty assessment associated with prey item grouping is intended to evaluate uncertainty in a particular element of the exposure assessment. As discussed in Section 6.3.1 of the BERA, risk characterization was performed using TEQ for dioxin-like PCB congeners only due to the uncertainty in the effects assessment for commercial Aroclor mixtures applied to weathered environmental samples. Thus, combining the prey item grouping uncertainty assessment (exposure) with the uncertainty in the effects assessment for PCB congeners (effects) would in effect 'mix' the different sources of uncertainty in the outcome (HQs); a situation that should be avoided. Uncertainty in the exposure assessment (e.g., mg/kg-bw/day) for Total PCB congeners associated with prey item grouping could be performed but should stop short of characterizing the associated risk (HQ) to avoid this situation. However, we feel that this type of assessment would not provide the context to meaningfully assess the effect of this uncertainty in a way that could be applied to risk management decision making.			
19	6.3.3.1.1	Prey Item Grouping - Why were oysters excluded as food in the kingfisher evaluation (all prey species separate)?	Appendix C-5.3 will be revised to include oysters in the uncertainty assessment of prey item grouping for belted kingfisher. Oysters will be included in the diet as an individual prey item and HQs recalculated for this uncertainty scenario. Table 6-9 (Uncertainty Analysis Scenario 1) HQs for belted kingfisher will be updated based on revised Appendix C-5.3			

Comments and Responses Matrix for TCEQ, NOAA, and USEPA Draft Baseline Ecological Risk Assessment

		Diati Daseille Ecological Nisk Assessi	
20	6.3.3.1.3	Area Use Factor (AUF) - To evaluate the potential uncertainty associated with the AUF, the AUF was varied for each receptor using the upper bound and lower bound home range estimates, and hazard quotients for each scenario were then determined. For the sandpiper, the hazard quotient for PCB Congener TEQ was 0.80 based on a lower-bound estimate of its home range and 3.1 based on an upper-bound estimate of its home range. Based on this evaluation, the BERA concludes that the uncertainty associated with the range of AUF values used in the baseline risk characterization is considered low. Given this outcome for the sandpiper evaluation, please discuss the conservatism and confidence in the AUF selected for the sandpiper.	Several (n = 8) studies were evaluated for the shorebird guild (sandpiper) to determine expected home ranges. All studies included in this evaluation were conducted at relatively undisturbed habitats that represented generally high quality, continuous habitat conditions. Under such conditions, species are likely to have shorter foraging distances to meet their energetic requirements than lower quality, fragmented habitats. As such, the home ranges used in development of the AUF are all considered conservative representatives of home ranges for species that may forage at the Site. This conservatism was addressed through the use of a 75 th percentile estimate of the home ranges to determine the AUF. However, even this assumption may not fully account for the quality of habitat at the Site relative to home ranges identified in the literature. Overall, the AUF used in the baseline is considered a conservative estimate for this guild. Another consideration to include in evaluating the conservatism / uncertainty in the AUF guild is the relative amount of intertidal habitat along the Sites length. Figure 6-1 of the draft BERA indicates that much of the Site has little areal intertidal habitat. This is in contrast to many of the study sites used in assessing home ranges that are located in areas with significant intertidal habitat (mudflats, sand bars) along an estuary gradient. Thus, all stream distances may not be equal in terms of intertidal foraging area and quality; unfortunately most shorebird foraging studies are expressed in terms of distance metrics. Thus, a kilometer of habitat in many of the studies is likely to have a greater amount of foraging distance necessary for shorebirds at the Site; the result being an underestimate of the foraging distance necessary for shorebirds at the Site; the result being an underestimate of the foraging distance necessary for shorebirds at the Site; the result being an underestimate of the foraging distance necessary for shorebirds at the Site. Notwithstanding the factors di
21	6.4	Risk Conclusions - The discussion repeats that for the spotted sandpiper, the uncertainty analysis indicates that the hazard quotient varies around one depending on the exposure assumptions, and that the hazard quotients range between 3.1 and 0.8 in the uncertainty analysis indicating that the probability of adverse effects above a LOAEL threshold are expected to be relatively low. The discussion adds that this generally translates into a low probability of adverse effects at the population level for this receptor group. Please provide more discussion regarding the potential for risks to the population level for this guild.	More discussion will be provided in the revised BERA. Consistent with the response to USEPA comments #1 and #2 (below), language in the BERA will be removed that states HQs above 1.0 are not associated with potential risk to population levels for this receptor. Rather, this type of evaluation will be limited to the uncertainty discussion and will focus on the uncertainty of risk characterization using a deterministic approach based on a single TRV to represent risk to a population or guild.
22	7.1	Bioaccumulative COPCs Risk Assessment - The discussion at the beginning of this section states that based on the previous screening, the potentially bioaccumulative COPCs included in the BERA for fish are mercury, PCBs, and selenium. This conflicts with the BERA Work Plan which included dioxins/furans (as TEQ) as bioaccumulative COPCs for fish (from the surface water to fish pathway). In fact, the March 2011 response to comment document (page 9) states the following: "Based on the results of the surface water screen for COPCs, dioxins/furans were identified as COPCs. We did state that due to the lack of a surface water screening value for dioxins/furans and the uncertainty with evaluating the bioavailability of dioxins/furans in the water column, an approach that used body burden data would be the best way to evaluate this COPC. As such, risks from dioxin/furan congeners in surface water will be assessed as a bioaccumulative COPC using a body burden approach for fish. Therefore, dioxins/furans will be included as an analyte in the fish and invertebrate SAP for the BERA." Please clarify if dioxins/furans were evaluated as bioaccumulative COPCs for fish as we do not see them in the summary tables for this section. If dioxins/furans were not evaluated in the BERA as a COPC for fish, please provide a robust rationale for this omission.	
23	7.1.1	Exposure Assessment - Looking at Appendix C and Appendix D, we could not find the presentation of the 95% UCL determination for the following combinations: Mercury: all species PCB Congener TEQ (Fish): GKF; all species Total PCBs: GM, STM, PNF, SAS; all species Please provide this information.	Appendix D will be updated to include UCL95 calculation sheets for the indicated COPC / species groups in the revised BERA.

Comments and Responses Matrix for TCEQ, NOAA, and USEPA

Draft Baseline Ecological Risk Assessment

		Draft Baseline Ecological Risk Assessr	ment
24	7.1.2	Tissue Based Effects Assessment - Details of the data used to derive the fish tissue-based selenium TRV were not provided in Appendix D. Please provide this information.	We will update Appendix D to include the data used to derive the fish tissue based selenium TRV.
25	7.3	Risk Conclusions - Based on the assessment of bioaccumulation-based and surface water exposures for fish, the BERA concludes there are no COPCs for fish. With the exception of selenium, this was primarily based on comparison of tissue residue-based TRVs with empirical tissue data for smaller (< 15 cm) fish collected from Patrick Bayou. In our comments on a draft of the BERA Work Plan, we had suggested collection of larger fish to more adequately assess the risks associated with bioaccumulative COPCs in Patrick Bayou. The final Work Plan stated (page 72) that "exploratory data analysis (e.g., coefficient of variation within and between species) will also be performed to assess the representativeness of larger size class fish (i.e., greater than 30 cm) as a measurement endpoint for the fish risk analysis and risk characterization." Was this analysis performed? If not, why not?	This analysis was not performed in the draft BERA due to the very low body burdens and associated HQ for whole body fish collected as part of the BERA tissue investigation. In addition, there is continued uncertainty that the body burdens in larger size class fish, whether measured or estimated, are representative of exposure to Site COPC or reflect exposure from offsite sources (due to larger expected foraging range of larger fish). Notwithstanding this uncertainty, to evaluate potential exposure of larger, higher trophic level fish, the following approach will be evaluated for assessing potential risk to larger size class fish. Food chain multipliers (FCM) will be developed from the literature to assess potential biomagnification of COPC to larger size class fish using the empirical fish tissue data for lower trophic level fish. These estimated tissue concentrations in higher trophic level fish will be compared to whole-body TRVs to calculate HQ for higher trophic level fish. Results will be presented in the uncertainty assessment for the fish risk assessment.
26	8	Benthic Invertebrate Risk Assessment - The introductory discussion indicates that the BERA problem formulation identified three lines of evidence (LOEs) for benthic invertebrate risk. These included an assessment of predicted sediment toxicity based on sediment chemistry and bioassay tests (i.e., the chemical and toxic LOE of the sediment quality triad); benthic indices that describe the condition of the benthic community; and a surface water exposure pathway risk assessment in which chemical concentrations in water are compared to TRVs derived for the protection of aquatic organisms. This conflicts with the problem formulation discussion (Section 3.3.1) of the BERA Work Plan which limited the benthic evaluation measurement endpoints to the use of a site-specific predictive sediment toxicity model and an evaluation of surface water exposure concentrations compared to water quality criteria. In fact, the Work Plan stated that the "predictive sediment toxicity model will be the primary measurement endpoint used to assess risks to the benthic invertebrate community on a site-wide basis." Please provide a clear explanation why the toxicity model was not presented in the BERA. Was the analysis performed?	The toxicity model initially developed during the Work Plan was refined and reevaluated during the BERA (this refinement and additional evaluation can be found in Appendix E). During the development of the BERA, it became apparent that the toxicity model had several weaknesses that we were unable to resolve as initially conceived, and that led to a significant amount of uncertainty in its utility and relevance as a primary line of evidence. Specifically, and in no particular order of priority, they are; the lack of a reference area toxicity tests to account for naturally present confounding factors (salinity acclimation, porewater ammonia, etc.) results in an unknown but potentially significant amount of conservative bias in the toxicity attributed to Site COPCs; the model, when applied to other test species, results in a high percentage of false positives, leading to a potentially significant amount of Site sediments to be falsely predicted as toxic; and the analytical uncertainty in the PCB analysis in the toxicity sample dataset confounds the extrapolation of the sediment chemistry data to more recent samples. Therefore, the primary issue is the reliance on this single line of evidence to characterize risk to the assessment endpoint; the health of the benthic community. In this case, the uncertainty in the model, which is primarily a function of the uncertainty in the data used to develop the model, does not lead to a sufficient level of confidence in the model to define the magnitude and extent of risks to the benthic community. As noted in Appendix E, we agree that there is some apparent association between Site COPC and sediment toxicity for a subset of COPC, including PCBs, PAHs, lead, and BEHP. However, the data, in conjunction with site conditions, do not provide a predictive dose-response model capable of defining the magnitude of risks to the benthic community as a function of sediment COPC concentrations for the four COPCs noted above. Per the discussions during the December 10, 2012 meeting, we als
27	8.1.2	Benthic Predictive Model Risk Characterization - The discussion states that since a dose-response relationship for the sediment-toxicity dataset was not observed, this suggests that other mechanisms are acting to cause the observed mortality in test species. This is too broad of a statement. The toxicity demonstrated in the historical toxicity tests could have been a result of unique site COPCs or a combination thereof, or as a result of a combination of site COPCs and non-site COPCs. Please revise the BERA text.	See response to Comment 26 above.
		<u>l</u>	

Comments and Responses Matrix for TCEQ, NOAA, and USEPA

Draft Baseline Ecological Risk Assessment

		Draft Baseline Ecological Risk Assessi	ment
28	8.1.3	Applicability of Benthic Toxicity Predictive Model - This section states that Leptocheirus is not considered a representative receptor for the species expected to be found at the site. We disagree with this statement. Amphipods are an important component of estuarine macrobenthic communities, even in tidal streams. In 110 tidal stream samples collected by TCEQ between 1988 and 2008, amphipods were present in 21% of those samples (TCEQ unpublished data). In the Patrick Bayou samples presented in the BERA, amphipods were present in 18% of the samples. In the Dobberstine (2007) study (presented in the BERA), amphipods were only present in 8% of the samples. The most likely reason for this difference in abundance is that less sediment volume was collected in the Dobberstine (2007) study compared to the benthic evaluations performed as part of the TMDL effort and the TCEQ's benthic evaluations. As a result, these less abundant organisms were simply missed.	This text will be revised. Text will included to state that <i>Leptocheirus</i> , relative to other species tested, is considered a sensitive indicator of sediment toxicity and, therefore, a conservative measurement endpoint for assessing the response of the benthic community to Site COPC. With regards to the frequency of amphipods in reference area samples, please see the response to TCEQ Comment #3.
29	8.2.1.1	Setting and General Conditions - This discussion should be revised so that it is clear that the areas of Patrick Bayou that are channelized or that have patchy benthic habitat are limited to the upper gunite-lined portion. This only affects four benthic sampling locations. The remainder of Patrick Bayou exhibits typical benthic habitat, which is not patchy or overly scoured. In fact, the benthic habitat in the lower portion of Patrick Bayou is less disturbed than that of other comparison tidal streams (e.g., Carpenter Bayou and Cedar Bayou are both channelized for much of their lower length and both are used for barge traffic). Additionally, Patrick Bayou is not subject to boat traffic (and its associated prop washing) because of the low bridge near the mouth of the bayou.	The text on patchy benthic habitat has been removed and the overall discussion on the benthic habitat has been revised as appropriate. The current approach no longer includes direct comparison to reference sites and, by extension, reference site conditions. The JDG maintains that understanding conditions between the Site and reference areas are important should such a direct comparison be included in any future evaluations (e.g., Feasibility Study).
30	8.2.2	Benthic Community Measurement Endpoints - This section of the BERA used two different benthic indices to evaluate the Patrick Bayou benthic community data compared with a group of reference sites sampled by Dobberstine (2007). Both of these benthic indices were developed using primarily open bay stations across the northern Gulf of Mexico. Neither of these indices have been used or tested in tidal streams. Similar to the Dobberstine et al. (2007) reference in Section 8.2.1.5, tidal streams are not expected to have the same macrobenthic community composition as open bay sites. It is not appropriate to use these open bay indexes to assess a tidal stream such as Patrick Bayou. In developing quantitative biocriteria, U.S. EPA guidance suggests using measures of diversity, richness, trophic structure, and the presence or absence of species that are considered tolerant or intolerant in comparison to a population of reference sites (U.S. EPA, 2000). In the absence of a tested index for tidal streams, we suggest evaluating the benthic community in Patrick Bayou and reference locations using five traditional metrics. The metrics we recommend are: species richness, diversity, percent dominance, percent tolerant individuals, and percent intolerant individuals. We evaluated the Patrick Bayou and comparison tidal stream benthic data using these five metrics in addition to a novel metric that evaluates the presence or absence of five major taxonomic groups that are generally present in tidal stream benthic communities. These six metrics performed well in evaluations of tidal stream reference and degraded benthic communities in the	The Gulf of Mexico (GOM) Biotic Index (BI) and Engle and Summers BI are peer-reviewed approaches that have been applied to estuarine site across the GOM. These BI's are currently the most appropriate standardized indices available for this Site. Per the concerns expressed by TCEQ during the December 10, 2012 meeting between USEPA, TCEQ, trustees, and the JDG, the GOM BI will not be used to evaluate the Site. Therefore, only the Engle and Summers BI will be used to evaluate the Site in the revised BERA. In addition, although not thoroughly peer-reviewed, the index proposed by TCEQ during this meeting will be evaluated as part of the uncertainty assessment for the benthic community line of evidence. In addition, per the discussions at the December 10, 2012 meeting, we will also reanalyze the Site-specific sediment chemistry data and the benthic community data to evaluate whether there are some potential areas which show a relatively stronger correspondence between chemistry and benthic community degradation that is not apparent in the benthic community analysis completed to date. This reanalysis will also be coordinated with a reanalysis of the Site-specific chemistry and toxicity data collected at the same time the benthic community data were collected (see response to Comment #26 above).
31	8.2.2.1	Benthic Indices (and Figures 8-8 and 8-11) - In the BERA evaluation of benthic community data (using these indices), benthic data from different seasons were combined. Macrobenthic communities show large seasonal fluctuations. For this reason, both of these indices were developed using summer data only. Summer data should not be combined with data from other seasons when evaluating macrobenthic communities. The statements that discuss the variability in the benthic scores over time and season should acknowledge that seasonal changes are expected. Further, the discussions should not attempt to use these differences as evidence that COPCs are not affecting the benthic community. Summer and non-summer scores are not expected to be the same.	We will update the assessment of the Site by separating the data into seasonal groups (Summer vs. Non-summer seasons) for the Engle and Summers BI based evaluations. Summer will include the months of June through September for purposes of this evaluation.
32	8.4.1	Exposure Assessment - The recommended total PCB 95% UCL was calculated to be 193 ng/L. Calculations are summarized in Appendix E-4, and the surface water data is presented in Appendix A-2. We calculated a slightly higher value (220 ng/L) based on the values in Table A-2 which totaled 21 values, rather than 25. Please verify the dataset and explain the discrepancy (if confirmed).	The output sheet for the total PCB congener surface water 95UCL is incorrect; the wrong sheet was included in the draft BERA. The correct 95UCL calculation sheet will be included in the revised BERA. Appendix A-2 includes results for 18 normal samples and three field duplicates (n=21). Treating field duplicates as normal samples does yield a 95UCL of 220 ng/L. When only the 18 normal samples are included, the approach specified in the Work Plan, a 95UCL of 193 ng/L is arrived at. This should be the appropriate EPC for the exposure assessment.

Comments and Responses Matrix for TCEQ, NOAA, and USEPA Draft Baseline Ecological Risk Assessment

		Draft baseline Ecological Risk Assessi	ment
33	8.4.2	Effects Assessment - The BERA proposed an alternate final chronic value (protective of benthos) of 540 ng/L based on the presentation in Fuchsman, et al. (2006). This paper relied on toxicity data for Aroclor 1254 to derive the final acute value (FAV) and acute and chronic voicity data for Aroclors 1242, 1248, and 1254 to calculate the acute-to-chronic ratio. Since the Patrick Bayou surface water data was based on total congeners rather than Aroclors, please discuss the uncertainty associated with this approach.	Although there is some uncertainty associated with the extrapolation between Aroclor and congener based PCB data, the uncertainty in this extrapolation is considered relatively low. Previous Aroclor analysis in Site sediments resulted in detected values for Aroclor 1248 and 1254, suggesting that the PCBs present in Site sediments are similar in composition to those used to derive the FAV and ACRs. Congener analysis in Site surface water supports this observation with congener composition profiles similar to what is found in commercial Aroclor 1248 mixtures. Thus, the congener compositions used in the studies to derive the alternate FCV are considered a reasonable match to what has been observed in Site media. The uncertainty associated with this approach and the information described above will be provided in the revised BERA.
34	8.5.2	Surface Water - Based on the alternate PCB surface water chronic value used, the BERA concludes that risk to benthic invertebrates from PCBs in surface water is determined to be negligible and no surface water COPCs are identified. We understand the basis for this statement. However, the PCB concentration (193 ng/L 95% UCL) exceeds the Texas Surface Water Quality Standard chronic criterion of 30 ng/L. This should be addressed later in the Superfund process.	Noted. The 95UCL does exceed the TSWQS saltwater chronic value which is equal to the USEPA saltwater CCC. The ecological relevance of the USEPA saltwater CCC is described in the BERA.
35	10	Conclusions and Risk Management Recommendations - We disagree with the conclusions regarding the evaluation of risks to the benthic invertebrate community and will revisit the risk management recommendations for this pathway after revision of the BERA and/or development of a consensus-based remedial target for the COPC risk drivers for this pathway. By consensus we mean a collective decision from the JDG, regulators, and Trustee agencies, where possible.	The final BERA, including Section 10, will incorporate and reflect all responses herein.
1	Section 8	BERA, General Comment, Section 8 - Benthic Invertebrate Risk Assessment: The benthic community risk assessment should be revised to include the Work Plan approved toxicity modeling approach, or at least include a toxicity based line of evidence in the evaluation. The BERA indicated that the PEC-Q metric met the acceptability criteria (particularly after modifications were made), and yet this approach was discarded in favor of the community indices – an explanation or justification would be helpful to understand this. The benthic assessment should not rely only on the community population data included in the draft BERA, particularly since that approach can be improved to provide a more accurate assessment (based on discussion with Linda Broach of TCEQ).	Please see response to TCEQ Comment #26.
2	Section 7	BERA, Specific Comment, Section 7 – Fish Risk Assessment: It is not clear why some of the fish tissue EPCs were presented as NA in table 7-2. Please provide more information regarding the fish EPCs, and in particular either the values depicted as NA in table 7-2 or justification for not including them. It would seem that "not applicable" does not apply to the PCB congener TEQ for Gulf Menhaden, at the least, since all 17 were detected in at least one sample and 11 were detected in all samples. Please provide the EPCs for all fish for the PCB congener TEQ.	
1	Page 5, Executive Summary, Conclusions and Risk Management Recommendations, Item 1, and Page 42, Risk Conclusions:	The risk assessment found that the risk to sediment-probing birds (e.g., sandpiper) and piscivorous birds (e.g., kingfisher) exceeded the threshold associated with potential adverse effects (with hazard quotients (HQs) as high as three) for individuals. The distinction of the risk to individuals vs. local populations is artificial since the toxicity tests used to develop the toxicity reference values (TRVs) are performed on groups of animals and the exposure assumptions are based upon populations of animals. Superfund remedial actions are expected to protect local populations (OSWER Directive 9285.7-28 P). The characterization of risk to sediment-probing birds and piscivorous birds would indicate that local populations would not be protected with HQs as high as three.	The BERA will be revised to clarify the conclusions associated with HQs > 1.0 based on a LOAEL TRV. Language in the risk characterization and conclusions stating that HQ > 1.0 based on the LOAEL TRV would not result in adverse population effects will be removed. The uncertainty discussion will address the significance of HQ > 1.0 based on a deterministic evaluation using a single TRV to assess population level effects and the range of HQs associated with variability in home range (i.e., AUFs).

Comments and Responses Matrix for TCEQ, NOAA, and USEPA

Draft Baseline Ecological Risk Assessment

2	Page 72,	In the second paragraph, the potential unacceptable ecological risks associated with the protection	Similar to the response to USEPA Comment #1, language will be removed / modified in Section
	Conclusions and	and maintenance of shorebird (sediment- probing birds) and piscivorous bird populations is	10 regarding the significance of HQs > 1.0 for wildlife receptors.
	Risk Management	acknowledged. Then the remaining part of the paragraph minimizes the ecological risk with a	
	Recommendations:	discussion of risk to individual receptors. Superfund ecological risk assessment are designed to	
		characterize the risk to a local population of receptors therefore risks found to exceed the threshold	
		associated with potential adverse effects would indicate that a local population of receptors would	
		not be protected.	

ATTACHMENT 1 ECOLOGICAL FISH AND INVERTEBRATE TISSUE SAMPLING DATA REPORT

TABLE OF CONTENTS

1	INT	RODUCTION AND PURPOSE	1
	1.1	Site Description	1
	1.2	Sampling Objectives	2
2	TISS	SUE COLLECTION AND PROCESSING	3
	2.1	Sample Identification Scheme	
	2.2	Sample Locations	
	2.3	Sample Collection Methods	5
	2.4	Invertebrate Depuration Methods	6
	2.5	Sample Processing Methods	7
	2.6	Sample Compositing Methods	7
	2.7	Field Deviations from the Quality Assurance Project Plan (QAPP) or SAP	8
3	LAB	SORATORY METHODS	9
	3.1	Methods for Chemical Analysis	9
	3.2	Laboratory Deviations from the QAPP or SAP	
4	ANA	ALYTICAL DATA VALIDATION AND DATA MANAGEMENT	11
	4.1	Overall Data Quality	
	4.2	Sample Transport and Holding Times.	
	4.3	Laboratory Quality Control	
5	RFS	ULTS	
•	5.1	General Summary	
	5.2	Blue Crab Size Class A	
	5.2.		
	5.2.		
	5.2.		
	5.2.	.4 SVOCs	14
	5.2.	.5 PCB Congeners	15
	5.2.	-	
	5.3	Blue Crab Size Class B	
	5.3.	.1 Conventional Parameters	15
	5.3.	.2 Mercury	15

5.3.3	SVOCs	16
5.3.4	PCBs	16
5.4 B	rown Shrimp	16
5.4.1	Conventional Parameters	16
5.4.2	Mercury	16
5.4.3	SVOCs	16
5.4.4	PCB Congeners	17
5.5 C)yster	17
5.5.1	Conventional Parameters	17
5.5.2	Metals	17
5.5.3	PAHs	17
5.5.4	SVOCs	18
5.5.5	PCB Congeners	18
5.5.6	Dixon/Furan Congeners	18
5.6 V	Vhite Shrimp Size Class A	18
5.6.1	Conventional Parameters	18
5.6.2	Metals	18
5.6.3	PAHs	19
5.6.4	SVOCs	19
5.6.5	PCB Congeners	19
5.6.6	Dixon/Furan Congeners	19
5.7 V	Vhite Shrimp Size Class B	19
5.7.1	Conventional Parameters	19
5.7.2	Mercury	20
5.7.3	SVOCs	20
5.7.4	PCB Congeners	20
5.8 G	fulf Killifish	
5.8.1	Conventional Parameters	20
5.8.2	Mercury	21
5.8.3	SVOCs	21
5.8.4	PCB Congeners	21
5.8.5	Dioxin/Furan Congeners	
59 G	fulf Menhaden	

5.9.1	Conventional Parameters	22
5.9.2	Mercury	22
5.9.3	SVOCs	22
5.9.4	PCB Congeners	22
5.9.5	Dioxin/Furan Congeners	22
5.10 Pi	nfish	22
5.10.1	Conventional Parameters	23
5.10.2	Mercury	23
5.10.3	SVOCs	23
5.10.4	PCB Congeners	23
5.10.5	Dioxin/Furan Congeners	23
5.11 Sa	nd Seatrout	24
5.11.1	Conventional Parameters	24
5.11.2	Mercury	24
5.11.3	SVOCs	24
5.11.4	PCB Congeners	24
5.11.5	Dioxin/Furan Congeners	25
5.12 St	riped Mullet	25
5.12.1	Conventional Parameters	25
5.12.2	Mercury	25
5.12.3	SVOCs	25
5.12.4	PCB Congeners	25
5.12.5	Dioxin/Furan Congeners	26
5.13 Su	mmary of Chemical Concentrations in Tissue	26
6 SUMM	ARY	27
7 REFER	ENCES	วด
, Krrry	LITULU	∠0

List of Tables

Table 1	Sample Ana	lysis Summary

Table 2 Sample List

Table 3 Blue Crab Size Class A Tissue Sample Summary

Table 4	Blue Crab Size Class B Tissue Sample Summary
Table 5	Brown Shrimp Tissue Sample Summary
Table 6	Oyster Tissue Sample Summary
Table 7	White Shrimp Size Class A Tissue Sample Summary
Table 8	White Shrimp Size Class B Tissue Sample Summary
Table 9	Gulf Killifish Tissue Sample Summary
Table 10	Gulf Menhaden Tissue Sample Summary
Table 11	Pinfish Tissue Sample Summary
Table 12	Sand Seatrout Tissue Sample Summary
Table 13	Striped Mullet Tissue Sample Summary
Table 14	Average Chemical Concentration for Each Sampled Species

List of Figures

Figure 1	Patrick Bayou Superfund Site Vicinity Map
Figure 2	Invertebrate Tissue Sample Locations
Figure 3	Fish Tissue Sample Locations

List of Appendices

Appendix A	Field Documentation on CD
Appendix B	Laboratory Data Packages on CD
Appendix C	Data Validation Report on CD
Appendix D	Analytical Results

LIST OF ACRONYMS AND ABBREVIATIONS

μg/kg microgram/kilogram

μm micrometer

Alpha Alpha Analytical

AOC Administrative Order on Consent
BERA Baseline Ecological Risk Assessment
BSAF biota-sediment accumulation factors

cm centimeter

COPC contaminant of potential concern

DGPS differential global positioning system

HpCDF 1,2,3,4,7,8,9-heptachlorodibenzofuran

HSC Houston Ship Channel

HxCDF 1,2,3,7,8,9-hexachlorodibenzofuran JDG Patrick Bayou Joint Defense Group

LCS/LSCD laboratory control sample/laboratory control sample duplicate

LDC Laboratory Data Consultants, Inc.

LELAP Louisiana Environmental Lab Accreditation Program

mg/kg milligram/kilogram

MS/MSD matrix spike/matrix spike duplicate

NELAC National Environmental Laboratory Accreditation Certified

ng/kg nanograms/kilogram

PAH polyaromatic hydrocarbons

PB Patrick Bayou

PCB polychlorinated biphenyl PQL practical quantitation limit

QA quality assurance

QAPP Quality Assurance Project Plan

RI/FS Remedial Investigation/Feasibility Study

SAP Final Fish and Invertebrate Tissue Sampling and Analysis Plan

SDG sample delivery group SGS SGS North America

SH State Highway

Site Patrick Bayou Superfund Site
SVOC semivolatile organic carbons
USACE U.S. Army Corps of Engineers

USEPA U.S. Environmental Protection Agency

1 INTRODUCTION AND PURPOSE

This data report summarizes the results of the tissue collection activities conducted during the implementation of the *Final Fish and Invertebrate Tissue Sampling and Analysis Plan* (SAP; Anchor QEA 2011a). This work was performed as part of the Remedial Investigation/Feasibility Study (RI/FS) being conducted by the Patrick Bayou Joint Defense Group (JDG) in response to an Administrative Order on Consent (AOC) and Settlement Agreement with the U.S. Environmental Protection Agency (USEPA), dated January 31, 2006.

1.1 Site Description

Patrick Bayou is a tidally influenced tributary of the Houston Ship Channel (HSC) in Harris County, Texas (Figure 1). The Patrick Bayou Superfund Site (Site) originates south of State Highway (SH) 225 in the City of Deer Park, Texas, and flows approximately 2.5 miles in a northerly direction, discharging into the south side of the HSC approximately 2.3 miles upstream of its confluence with the San Jacinto River. From the south side of SH 225, the bayou is contained in box culverts, which emerge into a gunite-lined channel approximately 0.3 miles downstream. The gunite-lined portion of the bayou extends approximately 0.45 miles downstream to a more natural channel bottom that makes up the remainder of the Site. A tributary of Patrick Bayou, referred to as the East Fork, joins the bayou approximately 1.4 miles downstream of SH 225.

Patrick Bayou has historically been utilized for residential, municipal, and industrial stormwater and wastewater discharge. Its urbanized/industrial nature is evident from many of the physical modifications to the channel, banks, and surrounding topography. The most obvious physical modifications are the box culvert and gunite channel in the upstream portion of the bayou. Downstream of these modifications, bulkheads and riprap revetments armor some stream banks. Stream bank heights in areas with bulkhead and riprap are generally steep. Areas without bank modifications, which include much of the middle section of the bayou, typically have low, sloping banks. Bank cover in areas without riprap or bulkheads is generally mowed grass with some low shrubs and bare earth. In many areas, impervious surfaces, such as parking lots and roads, are located adjacent to the bank of Patrick Bayou. A detailed description of the Site and its physical features is provided in the

Preliminary Site Characterization Report (Anchor 2006), and the Work Package 2
Hydrodynamic Field Data Collection and Contaminant Source Evaluation Data Report
(Anchor 2007a).

1.2 Sampling Objectives

The primary focus of this investigation was to support the Baseline Ecological Risk Assessment (BERA). Existing Site data and data collected during previous RI/FS tasks for the Site (Anchor 2007b, 2007c) were used to perform preliminary and refined assessments of ecological contaminants of potential concern (COPCs; Anchor 2007d, 2008a, and 2008b). These assessments identified several bioaccumulative COPCs for fish and aquatic-dependent wildlife receptors, such as birds and mammals. The site-specific exposure data provided by this study is intended to reduce uncertainty in the risk estimates performed as part of the BERA. In particular, site-specific exposure data are required to evaluate potential risks to fish and wildlife receptors in the BERA. This includes biota that represents both the fish species targeted as endpoints in the BERA and the typical dietary components of aquaticdependent wildlife receptors that may be present at the Site. These data will be used to refine the baseline exposure assessment for fish and wildlife receptors that consume aquaticbased prey items from the Site. In addition, assessing the benefits of alternative remediation plans will require an understanding of the relationship between contaminant concentrations in the sediments of Patrick Bayou and contaminant concentrations in the target species (fish bodies and wildlife diet). For organisms tied primarily to local sediments, the relationship between sediment and organism can be used to estimate how changes in Site sediment concentrations due to various remedial actions would affect changes in tissue concentrations (and subsequently exposure and risk) to ecological receptors. Finally, a robust preremediation dataset is required for monitoring the consequences of remedial decisions.

The specific objectives of this investigation were to:

- Collect sufficient data to support exposure assessments in the BERA
- Collect sufficient data to develop site-specific biota-sediment accumulation factors (BSAF) for modeling future conditions and remedial scenarios
- Establish baseline tissue concentrations of COPCs to provide a point of comparison for long-term (post-remediation) monitoring data

2 TISSUE COLLECTION AND PROCESSING

This section describes the methods used to collect and process fish and invertebrate tissue samples at the Site. Additional details regarding collection methods are presented in the SAP. A summary of all fish and invertebrate tissue samples and analyses is provided in Table 1. Copies of field notes, tissue sample collection forms, sample processing forms, and completed chain-of-custody forms are provided in Appendix A.

Four Prey Groups (two invertebrate and two fish) were established to reflect differences in body size, habitat preference, and distribution. Targeted fish and invertebrate species were assigned to a Prey Group based on these factors in the SAP. Predators were assumed to not prefer one species over another within a prey group. To further reflect variation in movement patterns and trophic levels, these numeric Prey Groups were subdivided into alphabetical (A and B) groups. Prey Groups are listed below and a complete description is provided in the SAP.

Group 1 Invertebrate. This group includes small, sedentary epibenthic invertebrates with localized distribution. The group is further subdivided into two groups (A and B), reflecting their feeding strategies, life cycles, and other attributes. Group 1A includes bivalves and Group 1B includes small crustaceans, such as juvenile blue crabs.

Group 2 Invertebrate. This group includes medium-sized invertebrates with a broader distribution and migratory patterns. This group is further subdivided into two groups reflecting trophic level. Group 2A includes lower trophic level organisms, such as shrimp, while Group 2B includes higher trophic level invertebrates (blue crabs).

Group 1 Fish. This group includes small bentho-pelagic fish (e.g., killifish and minnows) that inhabit shallow water and have localized distribution. Total length of an individual fish in Group 1 is less than 15 centimeters (cm).

Group 2 Fish. This group includes small pelagic fish that inhabit any water depth and have a broad foraging range relative to Group 1 fish. This group is further subdivided into two groups (A and B) based on their trophic level. Group 2A fish include lower trophic level

fishes, such as mullet and menhaden, while Group 2B includes higher trophic level fishes, such as spot and croaker. Total length of an individual fish in Group 2 is less than 15 cm.

2.1 Sample Identification Scheme

Each tissue sample was assigned a unique alphanumeric identifier using the format described below:

- Each location was identified by "PB" to depict the project location (Patrick Bayou), and a number, 01 through 04, identifying the reach (e.g., PB01 for the mouth of Patrick Bayou). A complete description of how the bayou was divided into reaches is provided in the SAP.
- Individual organisms to comprise each sample were identified by the same alphanumeric identifier used to identify the location followed by a three digit species identifier (e.g., "GKF" for Gulf killifish).
- Individual organisms that comprise one sample were frozen together in one package. Identifiers for each package were numbered using the following approach:

Where:

PB## = Patrick Bayou sample reach (## = 01, 02, 03, 04)

X = I or F (Invertebrate or Fish, respectively)

Y = Size class (A or B)

ZZZ = Species code

= Sequential sample number (001, 002, 003, ..., 999)

YYYYMMDD = Year, month, day sample was collected

In some instances, individual samples were composited together to meet minimum tissue requirements for laboratory analyses. In these cases, the composite sample was given a new name, following the nomenclature below:

Where:

PB## = Patrick Bayou sample reach (## = 01, 02, 03, 04)

X = I or F (Invertebrate or Fish, respectively)

Y = Size class (A or B)

ZZZ = Species code

= Sequential composite sample number (001, 002, 003, ..., 999)

COMP = Composite sample

YYYYMM = Year and month that samples within the composite were collected

2.2 Sample Locations

Locations of stationary sampling gear were recorded using a differential global positioning system (DGPS). For active sampling gear (i.e., trawls), the sampling track was recorded using DGPS. Prey Group 1A Fish and Group 1A/B Invertebrates sample locations were selected randomly from among the set of sediment sample locations previously sampled as part of the sediment and surface water COPC delineation effort (Anchor QEA 2010). Five randomly selected stations were sampled for Group 1A Fish and Group 1 Invertebrates in each of Reaches 1, 2, and 3. Reach 4 only had three sampling sites, due to its restricted size. Samples were collected from within a 15-meter radius of the designated coordinates.

Collection efforts for Group 2 Fish and Group 2A Invertebrates were performed on a reach-wide basis (i.e., sampling locations were selected opportunistically based on Site conditions such as substrate, water depth, etc.).

2.3 Sample Collection Methods

Organisms were collected using a combination of stationary and semi-stationary sampling gear, including: crab traps, minnow traps, gillnets, cast nets, and seine nets. The number and types of gear deployed was determined in the field based on targeted organisms, habitat type, and initial sampling efficiency observations. Gear was deployed from shore or boat, depending on water depth and substrate type.

Sampling was conducted over a period of two weeks from June 15, 2011 to June 28, 2011. At the beginning of the sampling event, baited minnow traps (three per station) were set at five randomly selected stations within Reaches 1, 2, and 3. Only three stations were set in Reach 4, due to the size of this reach and access restrictions within the gunite channel.

Baited minnow traps and crab traps were checked at regular intervals (i.e., daily) for three days. If sufficient fish were collected at a station to create at least two composite samples during that time, then the traps were moved to the next randomly selected station within a sample reach. If, after three days, insufficient numbers and volume of fish were collected at a station for two composite samples, then the traps were moved to the next randomly selected station within the sample reach. This cycling of traps was continued until the completion of the sampling event.

For active sampling and other passive gear aside from minnow or crab traps (e.g., gill nets), four days of sampling were performed in Reaches 1, 2, and 3, while two days of sampling were performed in Reach 4. The sampling crew completed the circuit of the four reaches in this manner.

To ensure that discrete locations were being sampled, a sampling area was specified for each Prey Group in order for sampling efforts targeting those Prey Groups to be considered the same "location". This was for the purpose of compositing samples, if necessary, to meet analytical tissue volume requirements. For Group 1A Fish and Group 1A/B Invertebrates, samples were collected within 30 meters of each other. For Group 2A/B Fish and Group 2A Invertebrates, samples were collected within 300 meters of each other. Organisms within a Prey Group collected outside of these limits were not combined across samples.

2.4 Invertebrate Depuration Methods

Consistent with USEPA/U.S. Army Corps of Engineers (USACE) guidance (1998), digestive tracts of benthic invertebrates and bivalves were emptied prior to homogenization and analysis. Organisms were allowed to void the material in their digestive tracts by being held for 24 hours in filtered (0.45 micrometer $[\mu m]$) Site surface water. No sediment was included in the containers and temperature conditions were maintained at field collection conditions. Only organisms collected from within the same reach of the bayou were placed together in a depuration container.

2.5 Sample Processing Methods

Collected organisms were kept alive in live wells on a boat supplied with Site water or wrapped in aluminum foil and placed in polyethylene bags (e.g., Ziplocs) on ice until they could be taken to the processing area. Invertebrates requiring depuration (oysters and blue crabs) were placed into depuration tanks to begin the required 24 hour depuration period. All other organisms were processed immediately. Organisms collected were identified to species.

The following steps, as recommended by USEPA (2000), were taken to process the organisms (with the exception of Group 1A Invertebrates) for shipping to the lab:

- 1. Whole organisms were euthanized by packing in ice and were then rinsed with distilled water to remove any debris.
- 2. Each organism was inspected for acceptability. Any specimens showing any severe skin, shell, or carapace laceration or fin deterioration, due to sampling gear were not included in the sample.
- 3. Each organism within a target species was measured to determine total body length to the nearest millimeter and weighed to the nearest 0.1 gram. If apparent, sex was determined (i.e., blue crabs). For small species such as killifish that require many individuals per sample, a subset of ten individuals from the sample were measured and weighed.
- 4. Organisms were composited according to the criteria outlined in the SAP (compositing methods are summarized in the following section), wrapped in aluminum foil, and placed in zip-sealed plastic bags with an identification label containing all relevant sample information. Bags were stored in a freezer at -4°C.

Bivalves were dissected prior to analysis. Shells were removed and only the soft tissue was retained for analysis. Shell length and soft tissue mass were measured.

2.6 Sample Compositing Methods

Composite samples are homogeneous mixtures of samples from two or more individual organisms of the same species collected at a particular location and analyzed as a single sample. Composite samples were limited to the minimum number of organisms needed to

obtain sufficient sample volume (190 grams) required to meet the laboratory analytical reporting limits.

Organisms used in a composite sample were required to meet the following criteria:

- Must all have been of the same species.
- Must have been collected from the same location. Depending on the Prey Group, samples were considered from the same location if they met the proximity requirements presented in Section 2.3.
- Must be within the same size class.
- Needed to include sufficient numbers of individuals to meet the sample tissue volume requirements for analysis of recommended target analytes.

These requirements were met in most cases. However, the number of samples for the Invertebrate Prey Groups was below the minimum targeted number of samples using these criteria. To achieve the minimum sampling objective identified in the SAP for Prey Group 1 Invertebrates, a modification to the proximity limits for compositing purposes was necessary. An increase in the proximity limit from 30 meters to 90 meters in areas of Site downstream of the gunite channel provided an additional seven samples. Samples from locations approximately 400 meters apart from within the gunite channel were composited together to create an additional sample. These modifications to the compositing scheme were outlined in a Memorandum to USEPA (Anchor QEA 2011b) and approved by USEPA.

2.7 Field Deviations from the Quality Assurance Project Plan (QAPP) or SAP

The only deviation from the SAP during the implementation of this investigation was the modifications made to the compositing scheme described in the previous section.

3 LABORATORY METHODS

This section briefly describes the analytical methods used to generate the chemistry data for the tissue samples in this report. A detailed description of the analytical methods can be found in the SAP (Anchor QEA 2011a). This section also summarizes any deviations by the laboratory from the SAP. Analyses included conventional parameters (lipid and percent moisture), metals, polyaromatic hydrocarbons (PAH), semivolatile organic carbons (SVOC), polychlorinated biphenyls (PCB), and dioxins and furans.

3.1 Methods for Chemical Analysis

Table 1 summarizes the number of tissue samples that were analyzed for each of the chemical methods listed below. This report includes data analyses performed by SGS North America (SGS) in Wilmington, North Carolina and Alpha Analytical (Alpha) in Mansfield, Massachusetts. SGS performed the PCB congener and dioxin/furan analyses and the remaining analyses were performed by Alpha. Alpha is National Environmental Laboratory Accreditation Certified (NELAC) accredited by the Louisiana Environmental Lab Accreditation Program (LELAP), Certificate No. 03090. SGS is NELAC accredited by the State of Florida Department of Health, Bureau of Laboratories, Certificate No. E87634. Fourteen data reports were received and validated from the laboratories. The laboratory data reports are provided in Appendix B. Eleven of the data reports were sent to Laboratory Data Consultants, Inc. (LDC) for data validation and three were validated by Anchor QEA staff. All data presented in this report reflect any findings reported in the validation reports.

3.2 Laboratory Deviations from the QAPP or SAP

This section discusses each of the laboratory's deviations from the SAP (Anchor QEA 2011a). SGS and Alpha adhered to the methods and procedures outlined in the SAP and as listed in SAP Table 4-2 with the following exceptions:

- The lipids analyses were done according to Alpha's internally developed method instead of the Bligh & Dyer method specified in the SAP. The methods are similar, however, so data are comparable, and this is not expected to affect data quality.
- Mercury analyses were performed using USEPA method 7474 instead of method
 7471A. The two methods are comparable and all sample results were above the

- detection limits.
- All of the Project Target Reporting Limits for non-detected SVOC results were not achieved, however; all non-detected practical quantitation limits (PQL) were well below the Risk-based Target Tissue PQL values, so data quality is not expected to be affected.
- Ten dioxin/furan non-detected detection limits were slightly above the Project Reporting Limit and nine of these results were above the Risk-Based Tissue Target PQL values, due to high concentrations of target and non-target analytes.
- Several PCB congener non-detected detection limits were above the Project Reporting Limits and the Risk-Based Tissue Target PQL values, due to high concentrations of target and non-target analytes.

4 ANALYTICAL DATA VALIDATION AND DATA MANAGEMENT

Third party data validation was performed on each dataset of chemical and physical analyses with the exceptions of two data reports from SGS and one report from Alpha which were validated by Anchor QEA staff. Data validation verified the accuracy and precision of chemical and physical determinations performed during this investigation. This section presents the data validation results conducted by LDC and Anchor QEA. The following sections summarize the overall data quality and do not necessarily address each individual sample result affected by data qualification. Detailed information regarding sample result qualifications is available in the data validation report in Appendix C.

4.1 Overall Data Quality

A review of the validation reports indicates that the overall data quality of the chemistry data generated was considered acceptable. Eighty-three tissue samples and two equipment blanks were collected and submitted to Alpha for SVOCs, mercury, lipids, and moisture analyses. Of these 85 samples, 21 were analyzed for lead and PAHs by Alpha. The data for these analyses were reported under seven sample delivery groups (SDGs): L1109178, L1109180, L1109182, L1109487, L1112532, L1112545, and L1120213. Eighty-three tissue samples were sent to SGS for PCB congener analyses and 76 samples were analyzed for dioxin/furan analyses. The data for these analyses were reported under seven SDGs: 31101717, 31101719, 31101798, 31101800, 31102315, 31103442, and 31103475. All of these data reports were validated under six validation reports. The four from LDC are: 26320-1, 26460, 25915, and 26422. Two validation reports were prepared by Anchor QEA.

Detailed data quality objectives and quality assurance (QA) procedures are provided in the SAP (Anchor QEA 2011a). Data packages were validated by LDC and Anchor QEA under USEPA National Functional Guidelines (USEPA 1999, 2004, 2005, 2008), as described in the SAP. The validation report is in Appendix C. Any data qualifiers applied to the data during the final validation procedures have been incorporated into the final database for this project. Data qualifiers assigned as a result of the data validation and their definitions are shown on the analytical results table (Appendix D). All data were considered useable as reported or as qualified with the exceptions of two PCB congener results. The PCB-169 result for sample PB03-F-A-GKF-W-002-20110615 was rejected, due to no recovery in the

matrix spike duplicate analyzed on this sample. The PCB-002 result was rejected for sample PB03-F-A-GKF-W-001-20110615, due to no recovery in the associated labeled compound. These rejections are not expected to impact overall data quality and completeness goals were met. Other data may have been qualified as estimated for a particular analysis based on method or technical criterion as stated in the functional guidelines (USEPA 1999, 2004, 2005, 2008). Data qualified with a "J" indicates that the associated numerical value is the approximate concentration of the analyte. Data qualified with a "UJ" indicates the approximate reporting limit below which the analyte was not detected. Data reported as "R" are rejected and unusable for any purpose. In some cases, reporting limits were raised to account for method blank contamination or matrix interference.

4.2 Sample Transport and Holding Times

Samples were received at the laboratories intact and appropriately preserved with two exceptions. Samples logged in under SDG L1109182 and all samples sent to SGS were received at a temperature above the -20° C requirement listed in the SAP. These discrepancies did not impact the data and no data were qualified because of them. All analyses were performed within recommended holding times.

4.3 Laboratory Quality Control

The validation reports indicate the majority of the data results did not require qualification. The laboratories followed the specified analytical methods or used comparable methods and all requested analyses were completed. Accuracy was acceptable as demonstrated by the surrogate, laboratory control sample/laboratory control sample duplicate (LCS/LCSD), matrix spike/matrix spike duplicate (MS/MSD), and internal standard recoveries with some exceptions. Precision was acceptable as demonstrated by the duplicate, LCS/LCSD, and MS/MSD relative percent difference values with some exceptions. Two results were rejected based on failure to meet the data quality objectives. One PCB congener result was rejected in two samples. Some reporting limits were elevated, due to method blank contamination. Most data are acceptable as reported or as qualified.

5 RESULTS

This section presents the results of the chemical analyses conducted on the fish and invertebrate tissue samples collected as part of this investigation. Only the final reporting limits and analytical results are presented in the tables associated with this section. The complete data table is presented in Appendix D.

For samples with laboratory duplicates, only the 'normal' sample value is presented and discussed. One-half of the detection limit was used for calculating total values for PCB congeners, PAHs, and dioxin/furans.

Table 2 presents a list of the analyzed samples, stations from which the samples were collected, species, and size class. Tables 3 through 13 present a summary of the analytical results for each invertebrate and fish species collected. For species where more than one size class was collected (i.e., blue crab and white shrimp), the size classes were summarized individually. Also included in these tables is the number of detected samples, range of detected concentrations, average detected concentrations, range of reporting limits for chemicals with reported non-detects, and average concentrations, including non-detected values (non-detected values are included in the calculations at values equal to one-half the detection limit). Data tables containing the results for each sample, including laboratory duplicates, are presented in Appendix D.

5.1 General Summary

A total of 83 samples were analyzed in fulfillment of this investigation. These samples were comprised of 33 invertebrate samples and 50 fish samples. Four different invertebrate species were collected and analyzed. These included blue crab (*Callinectes sapidus*), brown shrimp (*Penaeus aztecus*), oyster (*Crassostrea virginica*), and white shrimp (*Penaeus setiferus*) (Figure 2). Five different fish species were collected and analyzed. These included Gulf killifish (*Fundulus grandis*), Gulf menhaden (*Brevoortia patronus*), pinfish (*Lagodon rhomboids*), sand seatrout (*Cynoscion arenarius*), and striped mullet (*Mugil cephalus*) (Figure 3). The number of samples for individual species is provided in Table 1.

5.2 Blue Crab Size Class A

This section presents a summary of the analytical results from tissue samples comprised of blue crabs ranging in size from 2.5 cm to 7.5 cm. Eighteen samples were analyzed. Table 3 presents a summary of the results for the 18 crab tissue samples analyzed for conventional parameters, metals, PAHs, SVOCs, PCBs, and dioxin/furans.

5.2.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 0.555 percent to 3.73 percent. The average lipid content for all samples was 1.59 percent. Moisture ranged from 66.1 percent to 74.0 percent. The average moisture content for all samples was 70.3 percent.

5.2.2 Metals

Lead and mercury were included in the metal analyses. Both were detected in all 18 samples. Lead concentrations ranged from 0.092 milligrams/kilogram (mg/kg) to 0.518 mg/kg. The average lead concentration for all samples was 0.178 mg/kg. Mercury concentrations ranged from 0.009 mg/kg to 0.072 mg/kg. The average mercury concentration for all samples was 0.0475 mg/kg.

5.2.3 PAHs

Four individual PAHs were detected in at least one sample (acenaphthene, naphthalene, phenanthrene, and pyrene) and 13 were not detected in any of the samples. Five samples were reported with at least one detected PAH. All PAHs were reported as non-detect in 13 samples. Total PAH concentrations for samples with at least one detected PAH ranged from 60.9 micrograms/kilogram (μ g/kg) to 124 μ g/kg. The average total PAH concentration for all samples (including non-detects) was 25.9 μ g/kg.

5.2.4 SVOCs

Four chemicals were included in the SVOC analysis: 1,3-dichlorobenzene, 1,4-dichlorobenzene, hexachlorobenzene, and hexachlorobutadiene. Hexachlorobenzene was the only detected SVOC (detected in nine samples). Concentrations of detected

hexachlorobenzene ranged from 191 μ g/kg to 5,430 μ g/kg. The average concentration of hexachlorobenzene for all samples (including non-detects) was 612 μ g/kg.

5.2.5 PCB Congeners

One-hundred and ninety-two PCB congeners were detected in at least one sample. One-hundred and twenty-five were detected in all 18 samples. Concentrations of total PCB congeners ranged from 229 μ g/kg to 1,264 μ g/kg. The average concentration of total PCB congeners was 655 μ g/kg.

5.2.6 Dioxin/Furan Congeners

Sixteen dioxin/furan congeners were detected in at least one sample. 1,2,3,7,8,9-hexachlorodibenzofuran (HxCDF) was the only congener not detected in any samples. Five were detected in all 18 samples. Concentrations of total dioxin/furan congeners ranged from 17.3 nanograms/kilogram (ng/kg) to 235 ng/kg. The average concentration of total dioxin/furan congeners was 63.6 ng/kg.

5.3 Blue Crab Size Class B

This section presents a summary of the analytical results from tissue samples comprised of blue crabs ranging in size from 7.5 cm to 13 cm. Three samples were analyzed. Table 4 presents a summary of the results for the three crab tissue samples that were analyzed for conventional parameters, mercury, SVOCs, and PCBs.

5.3.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 0.575 percent to 0.943 percent. The average lipid content for all samples was 0.725 percent. Moisture ranged from 69.0 percent to 76.5 percent. The average moisture content for all samples was 73.1 percent.

5.3.2 Mercury

Mercury was detected in all three blue crab samples. Mercury concentrations ranged from 0.058 mg/kg to 0.083 mg/kg. The average mercury concentration was 0.067 mg/kg.

5.3.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. Neither chemical was detected in any of the samples.

5.3.4 PCBs

One-hundred and eighty-five PCB congeners were detected in at least one sample. One-hundred and sixty-one were detected in all three samples. Total PCB congener concentrations ranged from 722 μ g/kg to 936 μ g/kg. The average total PCB congener concentration for all samples was 804 μ g/kg.

5.4 Brown Shrimp

This section presents a summary of the analytical results from brown shrimp tissue samples made up of brown shrimp ranging in size from 7.5 cm to 13 cm. Eight samples were analyzed. Table 5 presents a summary of the results for the eight brown shrimp tissue samples that were analyzed for conventional parameters, mercury, SVOCs, and PCBs.

5.4.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 1.15 percent to 1.64 percent. The average lipid content for all samples was 1.37 percent. Moisture ranged from 74.0 percent to 76.2 percent. The average moisture content for all samples was 75.1 percent.

5.4.2 Mercury

Mercury was detected in all eight brown shrimp samples. Mercury concentrations ranged from 0.025 mg/kg to 0.065 mg/kg. The average mercury concentration was 0.041 mg/kg.

5.4.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in all samples. Concentration of hexachlorobenzene

ranged from 537 $\mu g/kg$ to 913 $\mu g/kg$. The average concentration of hexachlorobenzene for all samples was 710 $\mu g/kg$.

5.4.4 PCB Congeners

One-hundred and ninety-nine PCB congeners were detected in at least one sample. One-hundred and sixty-one were detected in all eight samples. Total PCB congener concentrations ranged from 997 μ g/kg to 2,192 μ g/kg. The average total PCB congener concentration for all samples was 1,716 μ g/kg.

5.5 Oyster

This section presents a summary of the analytical results from an oyster tissue sample made up of oysters ranging in size from 2.5 cm to 7.5 cm. One sample was analyzed. Table 6 presents a summary of the results for the oyster sample that was analyzed for conventional parameters, metals, PAHs, SVOCs, PCBs, and dioxin/furans.

5.5.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. The lipid content was reported at 1.1 percent. Moisture content was reported at 82.7 percent.

5.5.2 Metals

Lead and mercury were included in the metal analyses. Both were detected in the sample. The reported concentration of lead was 0.331 mg/kg. The reported concentration of mercury was 0.051 mg/kg.

5.5.3 PAHs

Eleven individual PAH chemicals were detected. The concentration of total PAHs was 552 $\mu g/kg$.

5.5.4 SVOCs

Four chemicals were included in the SVOC analysis: 1,3-dichlorobenzene, 1,4-dichlorobenzene, hexachlorobenzene, and hexachlorobutadiene. None of these chemicals were detected in the oyster sample.

5.5.5 PCB Congeners

One hundred and eighty eight PCB congeners were detected in the oyster sample. The concentration of total PCB congeners was $1,429 \mu g/kg$.

5.5.6 Dixon/Furan Congeners

Eight dioxin/furan congeners were detected in the oyster sample. The concentration of total dioxin/furan congeners was 41.7 ng/kg.

5.6 White Shrimp Size Class A

This section presents a summary of the analytical results of a tissue sample comprised of white shrimp ranging in size from 2.5 cm to 7.5 cm. One sample was analyzed. Table 7 presents a summary of the results for the white shrimp sample that was analyzed for conventional parameters, metals, PAHs, SVOCs, PCBs, and dioxin/furans.

5.6.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. The lipid content was reported at 0.914 percent. Moisture content was reported at 76.9 percent.

5.6.2 Metals

Lead and mercury were included in the metal analyses. Both were detected in the sample. The reported concentration of lead was 0.116 mg/kg. The reported concentration of mercury was 0.041 mg/kg.

5.6.3 PAHs

Three individual PAHs were detected: fluoranthene, phenanthrene, and pyrene. The concentration of total PAHs was $162 \mu g/kg$.

5.6.4 SVOCs

Four chemicals were included in the SVOC analysis: 1,3-dichlorobenzene, 1,4-dichlorobenzene, hexachlorobenzene, and hexachlorobutadiene. Hexachlorobenzene was the only chemical detected in the sample and had a concentration of $483 \mu g/kg$.

5.6.5 PCB Congeners

One-hundred and seventy-six PCB congeners were detected in the white shrimp sample. The concentration of total PCB congeners was 837 µg/kg.

5.6.6 Dixon/Furan Congeners

Eight dioxin/furan congeners were detected in the sample. The concentration of total dioxin/furan congeners was 69.4 ng/kg.

5.7 White Shrimp Size Class B

This section presents a summary of the analytical results from tissue samples comprised of white shrimp ranging in size from 7.5 cm to 13 cm. Two samples were analyzed. Table 8 presents a summary of the results for the two white shrimp tissue samples that were analyzed for conventional parameters, mercury, SVOCs, and PCBs.

5.7.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 1.13 percent to 1.30 percent. The average lipid content for all samples was 1.22 percent. Moisture ranged from 74.6 percent to 75.3 percent. The average moisture content for all samples was 75.0 percent.

5.7.2 Mercury

Mercury was detected in both white shrimp samples. Mercury concentrations ranged from 0.026 mg/kg to 0.042 mg/kg. The average mercury concentration for the two samples was 0.034 mg/kg.

5.7.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-Dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in both samples. Concentrations of hexachlorobenzene ranged from 380 μ g/kg to 682 μ g/kg. The average concentration of hexachlorobenzene for both samples was 531 μ g/kg.

5.7.4 PCB Congeners

One-hundred and eighty-three PCB congeners were detected in at least one sample. One-hundred and seventy-seven were detected in both samples. Total PCB congener concentrations ranged from $584 \, \mu g/kg$ to $919 \, \mu g/kg$. The average total PCB congener concentration for both samples was $733 \, \mu g/kg$.

5.8 Gulf Killifish

This section presents a summary of the analytical results from tissue samples comprised of Gulf killifish with total lengths less than 15 cm. Twenty five samples were analyzed. Table 9 presents a summary of the results for the Gulf killifish tissue samples that were analyzed for conventional parameters, mercury, SVOCs, PCBs, and dioxin/furans.

5.8.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 1.65 percent to 3.64 percent. The average lipid content for all samples was 2.72 percent. Moisture ranged from 71.9 percent to 75.9 percent. The average moisture content for all samples was 74.2 percent.

5.8.2 Mercury

Mercury was detected in all 25 Gulf killifish samples. Concentrations ranged from 0.025 mg/kg to 0.109 mg/kg. The average mercury concentration for all samples was 0.0524 mg/kg.

5.8.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in ten samples. Detected concentrations of hexachlorobenzene ranged from 200 μ g/kg to 2,360 μ g/kg. The average concentration of hexachlorobenzene for all samples (including non-detects) was 441 μ g/kg.

5.8.4 PCB Congeners

Two-hundred and four PCB congeners were detected in at least one sample. One-hundred and forty-seven congeners were detected in all samples. Total PCB congener concentrations ranged from 1,965 μ g/kg to 8,848 μ g/kg. The average total PCB congener concentration for all samples was 5,216 μ g/kg.

5.8.5 Dioxin/Furan Congeners

Fifteen dioxin/furan congeners were detected in at least one sample. 1,2,3,4,7,8,9-heptachlorodibenzofuran (HpCDF) and 1,2,3,7,8,9- HxCDF were not detected in any sample. Five were detected in all samples. Total dioxin/furan congener concentrations ranged from 19.1 ng/kg to 98.6 ng/kg. The average total dioxin/furan congener concentration for all samples was 36.2 ng/kg.

5.9 Gulf Menhaden

This section presents a summary of the analytical results from tissue samples comprised of Gulf menhaden with total lengths less than 15 cm. Ten samples were analyzed. Table 10 presents a summary of the results for the Gulf menhaden tissue samples that were analyzed for conventional parameters, mercury, SVOCs, PCBs, and dioxin/furans.

5.9.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 1.23 percent to 2.02 percent. The average lipid content for all samples was 1.67 percent. Moisture ranged from 77.5 percent to 80.0 percent. The average moisture content for all samples was 78.8 percent.

5.9.2 Mercury

Mercury was detected in all ten Gulf menhaden samples. Concentrations ranged from 0.068 mg/kg to 0.199 mg/kg. The average mercury concentration for all samples was 0.116 mg/kg.

5.9.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in six samples. Detected concentrations of hexachlorobenzene ranged from 398 μ g/kg to 671 μ g/kg. The average concentration of hexachlorobenzene for all samples (including non-detects) was 440 μ g/kg.

5.9.4 PCB Congeners

Two hundred and five PCB congeners were detected in at least one sample. One-hundred and sixty-one congeners were detected in all samples. Total PCB congener concentrations ranged from 1,259 μ g/kg to 4,005 μ g/kg. The average total PCB congener concentration for all samples was 2,642 μ g/kg.

5.9.5 Dioxin/Furan Congeners

All 17 dioxin/furan congeners were detected in at least one sample. Eleven were detected in all samples. Total dioxin/furan congener concentrations ranged from 154 ng/kg to 1,946 ng/kg. The average total dioxin/furan congener concentration for all samples was 763 ng/kg.

5.10 Pinfish

This section presents a summary of the analytical results from tissue samples comprised of pinfish with total lengths less than 15 cm. Four samples were analyzed. Table 11 presents a

summary of the results for the pinfish tissue samples that were analyzed for conventional parameters, mercury, SVOCs, PCBs, and dioxin/furans.

5.10.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 2.34 percent to 3.93 percent. The average lipid content for all samples was 2.86 percent. Moisture ranged from 73.2 percent to 74.9 percent. The average moisture content for all samples was 73.9 percent.

5.10.2 *Mercury*

Mercury was detected in all four pinfish samples. Concentrations ranged from 0.025 mg/kg to 0.083 mg/kg. The average mercury concentration for all samples was 0.0430 mg/kg.

5.10.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in three samples. Detected concentrations of hexachlorobenzene ranged from 397 μ g/kg to 2,290 μ g/kg. The average concentration of hexachlorobenzene for all samples (including non-detects) was 872 μ g/kg.

5.10.4 PCB Congeners

One-hundred and ninety-seven PCB congeners were detected in at least one sample. One-hundred and sixty-eight congeners were detected in all samples. Total PCB congener concentrations ranged from 2,114 μ g/kg to 4,334 μ g/kg. The average total PCB congener concentration for all samples was 3,157 μ g/kg.

5.10.5 Dioxin/Furan Congeners

Eleven dioxin/furan congeners were detected in at least one sample. Six were detected in all samples. Total dioxin/furan congener concentrations ranged from 14.6 ng/kg to 210 ng/kg. The average total dioxin/furan congener concentration for all samples was 74.5 ng/kg.

5.11 Sand Seatrout

This section presents a summary of the analytical results from tissue samples comprised of sand seatrout with total lengths less than 15 cm. Two samples were analyzed. Table 12 presents a summary of the results for the two sand seatrout tissue samples that were analyzed for conventional parameters, mercury, SVOCs, PCBs, and dioxin/furans.

5.11.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 0.931 percent to 1.18 percent. The average lipid content for all samples was 1.06 percent. Moisture ranged from 78.7 percent to 79.2 percent. The average moisture content for all samples was 79.0 percent.

5.11.2 *Mercury*

Mercury was detected in both sand seatrout samples. Concentrations ranged from 0.045 mg/kg to 0.063 mg/kg. The average mercury concentration for all samples was 0.0540 mg/kg.

5.11.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in both samples. Concentrations of hexachlorobenzene ranged from 240 μ g/kg to 325 μ g/kg. The average concentration of hexachlorobenzene for both samples was 283 μ g/kg.

5.11.4 PCB Congeners

One-hundred and ninety-six PCB congeners were detected in at least one sample. One-hundred and eighty-one congeners were detected in all samples. Total PCB congener concentrations ranged from 1,449 μ g/kg to 2,512 μ g/kg. The average total PCB congener concentration for all samples was 1,980 μ g/kg.

5.11.5 Dioxin/Furan Congeners

Twelve dioxin/furan congeners were detected in at least one sample. Nine were detected in all samples. Total dioxin/furan congener concentrations ranged from 11.6 ng/kg to 13.1 ng/kg. The average total dioxin/furan congener concentration for all samples was 12.3 ng/kg.

5.12 Striped Mullet

This section presents a summary of the analytical results from tissue samples comprised of striped mullet with total lengths less than 15 cm. Nine samples were analyzed. Table 13 presents a summary of the results for the striped mullet tissue samples that were analyzed for conventional parameters, mercury, SVOCs, PCBs, and dioxin/furans.

5.12.1 Conventional Parameters

Lipids and moisture content were included in the conventional parameters analyses. Lipids ranged from 1.24 percent to 2.36 percent. The average lipid content for all samples was 1.74 percent. Moisture ranged from 73.1 percent to 75.4 percent. The average moisture content for all samples was 74.0 percent.

5.12.2 Mercury

Mercury was detected in all nine striped mullet samples. Concentrations ranged from 0.024 mg/kg to 0.122 mg/kg. The average mercury concentration for all samples was 0.0516 mg/kg.

5.12.3 SVOCs

Two chemicals were included in the SVOC analyses: 1,3-dichlorobenzene and hexachlorobenzene. 1,3-dichlorobenzene was not detected in any of the samples. Hexachlorobenzene was detected in seven samples. Detected concentrations of hexachlorobenzene ranged from 389 μ g/kg to 1,830 μ g/kg. The average concentration of hexachlorobenzene for all samples (including non-detects) was 721 μ g/kg.

5.12.4 PCB Congeners

Two-hundred and two PCB congeners were detected in at least one sample. One-hundred and seventy-eight congeners were detected in all samples. Total PCB congener

concentrations ranged from 596 μ g/kg to 4,539 μ g/kg. The average total PCB congener concentration for all samples was 2,246 μ g/kg.

5.12.5 Dioxin/Furan Congeners

Fourteen dioxin/furan congeners were detected in at least one sample. Ten were detected in all samples. Total dioxin/furan congener concentrations ranged from 70.5 ng/kg to 931 ng/kg. The average total dioxin/furan congener concentration for all samples was 332 ng/kg.

5.13 Summary of Chemical Concentrations in Tissue

The chemical analyses conducted on fish and invertebrate tissue samples included metals, PAHs, SVOCs, PCBs, and dioxin/furans. Table 14 presents the average chemical concentration for each species. Lead concentrations were measured for size class A invertebrates, including blue crab, oyster, and white shrimp. The highest average lead concentration (0.331 mg/kg) was reported in the oyster sample. Mercury concentrations were measured in all tissue samples, both invertebrate and fish. The highest average mercury concentration for the invertebrate samples was reported in the blue crab (size class B) samples (0.0667 mg/kg) and in the Gulf menhaden samples for fish (0.116 mg/kg). Chemical analyses for PAHs were conducted on the size class A Invertebrates. The highest average total PAH concentration (552 μg/kg) was reported in the oyster sample. Of the four SVOCs analyzed, only hexachlorobenzene was detected. The highest average hexachlorobenzene concentration for the invertebrate samples was reported in the brown shrimp samples (710 μ g/kg) and in the pinfish samples for fish (872 μ g/kg). PCB and dioxin/furan concentrations were measured in all tissue samples. The highest average total PCB concentration for the invertebrates was reported in the brown shrimp samples (1,716 μg/kg) and in the Gulf killifish samples for fish (5,216 μg/kg). The highest average total dioxin/furan concentration for the invertebrates was reported in the white shrimp (size class A) samples (69.4 ng/kg) and in the Gulf menhaden for fish (763 ng/kg).

6 SUMMARY

During this investigation, 33 invertebrate tissue samples and 50 fish tissue samples were successfully collected and analyzed. Species sampled included blue crab, brown shrimp, oyster, white shrimp, Gulf killifish, Gulf menhaden, pinfish, sand seatrout, and striped mullet. Overall, these samples were collected consistent with the SAP and are of sufficient quality to meet the sampling objectives outlined in Section 1.2.

The data collected during this investigation were used to support exposure assessment in the BERA. Sufficient data was also collected for the development of site-specific BSAF for modeling future conditions and remedial scenarios. Baseline tissue concentrations of COPCs have also been established, which will provide a point of comparison for long-term (post-remediation) monitoring data.

7 REFERENCES

- Anchor Environmental, LLC (Anchor), 2006. *Preliminary Site Characterization Report.*Patrick Bayou Superfund Site. Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. Prepared by Anchor Environmental, L.L.C. April 2006.
- Anchor, 2007a. *Remedial Investigation Work Plan.* Patrick Bayou Superfund Site, Deer Park, Texas. Prepared for USEPA and Patrick Bayou Joint Defense Group. January 2007.
- Anchor, 2007b. *Technical Memorandum: Preliminary Contaminant of Potential Concern Selection.* Patrick Bayou Superfund Site, Deer Park, Texas. Prepared for U.S. Environmental Protection Agency and the Patrick Bayou Joint Defense Group. Prepared by Anchor Environmental, L.L.C. February 16, 2007.
- Anchor, 2007c. Work Package 2 Vertical Profiling Hydrodynamic Field Data Collection and Contaminant Source Evaluation. Patrick Bayou Superfund Site, Deer Park, Texas. Prepared for U.S. Environmental Protection Agency and the Patrick Bayou Joint Defense Group. Prepared by Anchor Environmental, L.L.C. April 2007.
- Anchor, 2007d. *Draft Supplemental Work Plan.* Additional Hydrodynamic Data Collection. Geotechnical Data Collection. Sediment Accretion Evaluations. Porewater Sampling and Estimation of Distribution Coefficients. Patrick Bayou Superfund Site, Deer Park, Texas. Prepared for U.S. Environmental Protection Agency and the Patrick Bayou Joint Defense Group. Prepared by Anchor Environmental, L.L.C. May 2007,
- Anchor, 2008a. *Draft Selection of Chemicals of Potential Concern for Ecological Risk Assessment.* Prepared for USEPA and the Patrick Bayou Joint Defense Group. Prepared by Anchor Environmental, L.L.C. September 2008.
- Anchor, 2008b. *Draft Selection of Chemicals of Potential Concern for Ecological Risk Assessment Addendum.* Prepared for USEPA and the Patrick Bayou Joint Defense Group. Prepared by Anchor Environmental, L.L.C. December 2008.
- Anchor QEA, 2010. *Patrick Bayou Sediment and Surface Water Contaminant of Potential Concern Delineation Data Report.* Prepared for the Patrick Bayou Joint Defense Group. Prepared by Anchor QEA, LLC. April 2010.

- Anchor QEA, 2011a. Final Fish and Invertebrate Tissue Sampling and Analysis Plan.

 Prepared for Patrick Bayou Joint Defense Group. Prepared by Anchor QEA, LLC.

 May 2011.
- Anchor QEA, 2011b. *Draft Memorandum: Request to Modify Composite Scheme for Patrick Bayou Tissue Samples.* Prepared for U.S. Environmental Protection Agency. Prepared by Anchor QEA, LLC. July 29, 2011.
- USEPA (U.S. Environmental Protection Agency), 1999. USEPA Contract Laboratory
 Program National Functional Guidelines for Organic Data Review.

 U.S. Environmental Protection Agency, Office of Emergency Response. EPA 540/R-99/008. October.
- USEPA, 2004. USEPA Contract Laboratory Program National Functional Guidelines for Inorganic Data Review. U.S. Environmental Protection Agency, Office of Superfund Remediation and Technology Innovation (OSRTI). EPA 540-R-04-004. October 2004.
- USEPA, 2005. USEPA Contract Laboratory Program National Functional Guidelines for Chlorinated Dibenzo-p-Dioxins (CDDs) and Chlorinated Dibenzofurans (CDFs) Data Review. U.S. Environmental Protection Agency, Office of Superfund Remediation and Technology Innovation (OSRTI). EPA 540-R-05-001. September 2005.
- USEPA, 2008. USEPA Contract Laboratory Program National Functional Guidelines for Superfund Organic Methods Data Review. U.S. Environmental Protection Agency, Office of Superfund Remediation and Technology Innovation. USEPA 540-R-08-01. June.
- USEPA/USACE, 1998. Evaluation of dredged material proposed for discharge in waters of the U.S. Testing Manual (Inland Testing Manual). EPA-823-B-98-004.

TABLES

Table 1
Sample Analysis Summary

			-				Ana	lysis					
Species	Number of Samples Collected	Lipid	Percent Moisture	Mercury	Lead	PAHs	1,3-Dichlorobenzene	1,4-Dichlororbenzene	2-Methylnapthalene	Hexachlorobenzene	Hexachlorobutadiene	Dioxins/Furans	PCB Congeners
Fish													
Gulf Killifish	25	Χ	Χ	Χ			Χ			Χ		Χ	Χ
Gulf Menhaden	10	Χ	Χ	Χ			Χ			Χ		Χ	Χ
Sand Seatrout	2	Х	Х	Х			Χ			Χ		Χ	Χ
Striped Mullet	9	Χ	Χ	Χ			Χ			Χ		Χ	Χ
Pinfish	4	Χ	Χ	Χ			Χ			Χ		Χ	Χ
Invertebrate (Size Class	s A)												
Blue Crab	18	Χ	Χ	Χ	Χ	Х	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Oyster	1	Χ	Χ	Χ	Χ	Х	Χ	Χ	Χ	Χ	Χ	Χ	Χ
White Shrimp	1	Χ	Χ	Χ	Χ	Х	Χ	Χ	Χ	Χ	Х	Χ	Χ
Invertebrate (Size Class	s B)												
Blue Crab	3	Х	Х	Х			Χ			Х			Χ
White Shrimp	2	Х	Х	Х			Χ			Χ			Χ
Brown Shrimp	8	Χ	Χ	Χ			Χ			Χ			Χ

Table 2
Sample List

Station ID	Reach	Sample ID	Species Group	Species	Size Class
PB015 A		·	FISH	Gulf Killifish	
PB015_A PB015.1	1	PB01-F-A-GKF-W-001-COMP-201106	FISH	Gulf Killifish	A A
PB015.1 PB016.2		PB01-F-A-GKF-W-002-COMP-201106	ļ		
	1	PB01-F-A-GKF-W-003-COMP-201106	FISH	Gulf Killifish	A
PB013.3	1	PB01-F-A-GKF-W-004-COMP-201106	FISH	Gulf Killifish	A
PB019_A	1	PB01-F-A-GKF-W-005-COMP-201106	FISH	Gulf Killifish	A
PB024.1	1	PB01-F-A-GM-W-001-20110615	FISH	Gulf Menhaden	A
PB024.1	1	PB01-F-A-GM-W-002-20110615	FISH	Gulf Menhaden	A
PB025_A	1	PB01-F-A-GM-W-004-20110615	FISH	Gulf Menhaden	A
PB010	1	PB01-F-A-GM-W-008-20110618	FISH	Gulf Menhaden	A
PB019.1	1	PB01-F-A-SAS-W-005-20110622	FISH	Sand Seatrout	A
PB026.1	1	PB01-F-A-STM-W-002-20110616	FISH	Striped Mullet	А
PB013.3	1	PB01-F-A-STM-W-003-20110616	FISH	Striped Mullet	А
PB003.1	1	PB01-F-A-STM-W-006-20110618	FISH	Striped Mullet	Α
PB002.1	1	PB01-I-A-BCR-W-001-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB005.1	1	PB01-I-A-BCR-W-002-COMP-201106	INVERTEBRATE	Blue Crab	А
PB010.3	1	PB01-I-A-BCR-W-003-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB015.2	1	PB01-I-A-BCR-W-004-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB024.4	1	PB01-I-A-BCR-W-005-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB015_A	1	PB01-I-A-BCR-W-006-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB015_A	1	PB01-I-A-BCR-W-007-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB019_A	1	PB01-I-A-BCR-W-008-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB015_A	1	PB01-I-A-BCR-W-009-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB013.1_A	1	PB01-I-A-BCR-W-010-COMP-201106	INVERTEBRATE	Blue Crab	Α
PB019.2	1	PB01-I-A-OYS-W-001-COMP-201106	INVERTEBRATE	Oyster	Α
PB010.1	1	PB01-I-A-WHS-W-001-COMP-201106	INVERTEBRATE	White Shrimp	Α
PB015_A	1	PB01-I-B-BRS-W-001-COMP-201106	INVERTEBRATE	Brown Shrimp	В
PB019.1	1	PB01-I-B-BRS-W-012-20110621	INVERTEBRATE	Brown Shrimp	В
PB019.1	1	PB01-I-B-BRS-W-014-20110621	INVERTEBRATE	Brown Shrimp	В
PB019.1	1	PB01-I-B-BRS-W-015-20110621	INVERTEBRATE	Brown Shrimp	В
PB019.1	1	PB01-I-B-BRS-W-019-20110621	INVERTEBRATE	Brown Shrimp	В
PB019.1	1	PB01-I-B-BRS-W-022-20110622	INVERTEBRATE	Brown Shrimp	В
PB019.1	1	PB01-I-B-BRS-W-023-20110622	INVERTEBRATE	Brown Shrimp	В
PB008.2	1	PB01-I-B-WHS-W-001-COMP-201106	INVERTEBRATE	White Shrimp	В
PB006.1	1	PB01-I-B-WHS-W-002-COMP-201106	INVERTEBRATE	White Shrimp	В
PB049_A	2	PB02-F-A-GKF-W-001-COMP-201106	FISH	Gulf Killifish	Α
PB056_A	2	PB02-F-A-GKF-W-010-20110618	FISH	Gulf Killifish	А
PB056_A	2	PB02-F-A-GKF-W-011-20110618	FISH	Gulf Killifish	А
PB053_B	2	PB02-F-A-GKF-W-013-20110618	FISH	Gulf Killifish	А
PB053.2	2	PB02-F-A-GKF-W-025-20110621	FISH	Gulf Killifish	А
PB053.2	2	PB02-F-A-GKF-W-029-20110622	FISH	Gulf Killifish	А
PB048.3	2	PB02-F-A-GM-W-001-20110616	FISH	Gulf Menhaden	Α
PB048.3	2	PB02-F-A-GM-W-002-20110616	FISH	Gulf Menhaden	A
PB047.1 A	2	PB02-F-A-PNF-W-002-20110615	FISH	Pinfish	A
PB032 A	2	PB02-F-A-PNF-W-009-20110617	FISH	Pinfish	A
PB052_7\	2	PB02-F-A-SAS-W-002-20110616	FISH	Sand Seatrout	A
PB048.3	2	PB02-F-A-STM-W-001-20110616	FISH	Striped Mullet	A
PB052.1	2	PB02-F-A-STM-W-003-20110616	FISH	Striped Mullet	A

Table 2
Sample List

Sample List												
Station ID	Reach	Sample ID	Species Group	Species	Size Class							
PB037.2	2	PB02-F-A-STM-W-006-20110616	FISH	Striped Mullet	А							
PB047.3	2	PB02-I-A-BCR-W-001-COMP-201106	INVERTEBRATE	Blue Crab	Α							
PB037.6	2	PB02-I-A-BCR-W-002-COMP-201106	INVERTEBRATE	Blue Crab	А							
PB049_A	2	PB02-I-A-BCR-W-003-COMP-201106	INVERTEBRATE	Blue Crab	А							
PB037_A	2	PB02-I-A-BCR-W-004-COMP-201106	INVERTEBRATE	Blue Crab	А							
PB037.4	2	PB02-I-B-BCR-W-001-COMP-201106	INVERTEBRATE	Blue Crab	В							
PB037.5	2	PB02-I-B-BRS-W-001-COMP-201106	INVERTEBRATE	Brown Shrimp	В							
PB057_B	3	PB03-F-A-GKF-W-001-20110615	FISH	Gulf Killifish	А							
PB069_A	3	PB03-F-A-GKF-W-002-20110615	FISH	Gulf Killifish	Α							
PB059.1_A	3	PB03-F-A-GKF-W-003-20110615	FISH	Gulf Killifish	А							
PB063.2_A	3	PB03-F-A-GKF-W-005-20110615	FISH	Gulf Killifish	А							
PB057_B	3	PB03-F-A-GKF-W-006-20110617	FISH	Gulf Killifish	А							
PB069_A	3	PB03-F-A-GKF-W-007-20110617	FISH	Gulf Killifish	Α							
PB063.1_A	3	PB03-F-A-GKF-W-009-20110617	FISH	Gulf Killifish	А							
PB063.1_A	3	PB03-F-A-GKF-W-010-20110617	FISH	Gulf Killifish	Α							
PB063.2_A	3	PB03-F-A-GKF-W-013-20110617	FISH	Gulf Killifish	А							
PB057.2	3	PB03-F-A-GM-W-001-20110617	FISH	Gulf Menhaden	А							
PB058.1	3	PB03-F-A-GM-W-003-20110618	FISH	Gulf Menhaden	А							
PB058.1	3	PB03-F-A-GM-W-004-20110618	FISH	Gulf Menhaden	А							
PB058.1	3	PB03-F-A-GM-W-005-20110618	FISH	Gulf Menhaden	А							
PB057_B	3	PB03-F-A-PNF-W-001-COMP-201106	FISH	Pinfish	А							
PB057.2	3	PB03-F-A-STM-W-001-20110617	FISH	Striped Mullet	А							
PB057.2	3	PB03-F-A-STM-W-001-COMP-201106	FISH	Striped Mullet	Α							
PB057.2	3	PB03-F-A-STM-W-002-20110617	FISH	Striped Mullet	А							
PB059.5	3	PB03-I-A-BCR-W-001-COMP-201106	INVERTEBRATE	Blue Crab	А							
PB069_A	3	PB03-I-A-BCR-W-002-COMP-201106	INVERTEBRATE	Blue Crab	А							
PB059.1_A	3	PB03-I-A-BCR-W-003-COMP-201106	INVERTEBRATE	Blue Crab	А							
PB059.3	3	PB03-I-B-BCR-W-002-20110627	INVERTEBRATE	Blue Crab	В							
PB059.4	3	PB03-I-B-BCR-W-003-20110628	INVERTEBRATE	Blue Crab	В							
PB101_D	4	PB04-F-A-GKF-W-001-20110627	FISH	Gulf Killifish	Α							
PB101_D	4	PB04-F-A-GKF-W-001-COMP-201106	FISH	Gulf Killifish	А							
PB084.1	4	PB04-F-A-GKF-W-005-20110627	FISH	Gulf Killifish	А							
PB101_D	4	PB04-F-A-GKF-W-009-20110627	FISH	Gulf Killifish	А							
PB084.1	4	PB04-F-A-GKF-W-013-20110627	FISH	Gulf Killifish	А							
PB102	4	PB04-F-A-PNF-W-001-COMP-201106	FISH	Pinfish	Α							
PB095.2	4	PB04-I-A-BCR-W-001-COMP-201106	INVERTEBRATE	Blue Crab	А							

Table 3
Blue Crab Size Class A Tissue Sample Summary

Blue Crab Size Class A Tissue Sample Summary													
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average				
Conventional Parameters (percent)		,						<u> </u>					
Lipids	18	18	0	0.555	3.73	1.59			1.59				
Moisture, percent	18	18	0	66.1	74.0	70.3			70.3				
Metals (mg/kg)													
Lead	18	18	0	0.092	0.518	0.178			0.178				
Mercury	18	18	0	0.009	0.072	0.0475			0.0475				
Polyaromatic Hydrocarbons (μg/kg)													
2-Methylnaphthalene	18	0	18				6.93	14.8	6.56				
Acenaphthene	18	2	16	5.72	11	8.36	6.93	14.8	5.98				
Acenaphthylene	18	0	18				6.93	14.8	5.65				
Anthracene	18	0	18				6.93	14.8	5.65				
Benzo(a)anthracene	18	0	18				6.93	14.8	5.65				
Benzo(a)pyrene	18	0	18				6.93	14.8	5.65				
Benzo(b)fluoranthene	18	0	18				6.93	14.8	5.65				
Benzo(g,h,i)perylene	18	0	18	1			6.93	14.8	5.65				
Benzo(k)fluoranthene	18	0	18	1			6.93	14.8	5.65				
Chrysene	18	0	18	1			6.93	14.8	5.65				
Dibenzo(a,h)anthracene	18	0	18				6.93	14.8	5.65				
Fluoranthene	18	0	18				6.93	14.8	5.65				
Fluorene	18	0	18				6.93	14.8	5.65				
Indeno(1,2,3-c,d)pyrene	18	0	18				6.93	14.8	5.65				
Naphthalene	18	4	14	4.19	8.23	5.91	6.93	14.8	6.13				
Phenanthrene	18	1	17	3.81	3.81	3.81	6.93	14.8	5.63				
Pyrene	18	1	17	6.08	6.08	6.08	6.93	14.8	5.75				
Total HPAH (U = 1/2)	18	1	17	39.6	39.6	39.6	6.93	14.8	7.62				
Total LPAH (U = 1/2)	18	5	13	25.5	53.3	33.4	6.93	14.8	13.7				
Total PAH (U = 1/2)	18	5	13	60.9	124	77.2	6.93	14.8	25.9				
Semivolatile Organic Carbons (μg/kg)	•												
1,3-Dichlorobenzene	18	0	18				341	745	213				
1,4-Dichlorobenzene	18	0	18				341	745	213				
Hexachlorobenzene	18	9	9	191	5,430	1,021	347	727	612				
Hexachlorobutadiene	26	0	26				341	745	213				
PCBs (μg/kg)	•				•								
PCB-001	18	2	16	0.00249	0.00384	0.00317	0.00131	0.00783	0.00379				
PCB-002	18	2	16	0.000956	0.001	0.000978	0.000847	0.00719	0.00149				
PCB-003	18	0	18				0.000816	0.00768	0.00164				
PCB-004	18	3	15	0.012	0.147	0.0690	0.00701	0.108	0.022				
PCB-005	18	0	18				0.00165	0.0181	0.00318				
PCB-006	18	5	13	0.00635	0.0786	0.0257	0.00229	0.0114	0.00948				
PCB-007	18	1	17	0.00665	0.00665	0.00665	0.00147	0.0166	0.00314				
PCB-008	18	17	1	0.0387	0.369	0.0854	0.0289	0.0289	0.0815				
PCB-009	18	1	17	0.00477	0.00477	0.00477	0.00151	0.0167	0.00312				
PCB-010	18	0	18				0.00233	0.0313	0.00394				
PCB-011	18	0	18				0.0112	0.0578	0.016				
PCB-012/013	18	15	3	0.00855	0.0567	0.0189	0.0112	0.0117	0.0167				

Table 3
Blue Crab Size Class A Tissue Sample Summary

Number of Number of State Sample Summary													
Chemical	Number of Samples	Number of Detected Samples	Non-Detected	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average				
PCB-014	18	0	18				0.00142	0.0158	0.00275				
PCB-015	18	18	0	0.393	3.99	1.72			1.72				
PCB-016	18	10	8	0.00507	0.0944	0.0232	0.00489	0.0106	0.0146				
PCB-017	18	18	0	0.0245	0.549	0.0875			0.0875				
PCB-018/030	18	18	0	0.0269	1.03	0.144			0.144				
PCB-019	18	4	14	0.00657	0.052	0.0265	0.00299	0.0126	0.00893				
PCB-020/028	18	18	0	18.6	185	79.1			79.1				
PCB-021/033	18	18	0	0.0218	1.03	0.186			0.186				
PCB-022	18	18	0	0.0559	1.12	0.343			0.343				
PCB-023	18	0	18				0.00101	0.00769	0.00164				
PCB-024	18	0	18				0.00107	0.00798	0.00169				
PCB-025	18	18	0	0.0125	0.209	0.0619			0.0619				
PCB-026/029	18	18	0	0.0175	0.603	0.130			0.130				
PCB-027	18	13	5	0.00352	0.0764	0.0133	0.00253	0.00721	0.0102				
PCB-031	18	18	0	0.286	5.33	1.48			1.48				
PCB-032	18	18	0	0.0272	0.819	0.148			0.148				
PCB-034	18	17	1	0.00432	0.0605	0.0163	0.00485	0.00485	0.0156				
PCB-035	18	12	6	0.00558	0.0436	0.0174	0.00402	0.012	0.0131				
PCB-036	18	1	17	0.0852	0.0852	0.0852	0.0032	0.0116	0.00795				
PCB-037	18	18	0	1.35	14.3	6.26			6.26				
PCB-038	18	7	11	0.0129	0.0399	0.0286	0.00336	0.0118	0.0133				
PCB-039	18	13	5	0.0212	0.108	0.0584	0.00429	0.0105	0.0431				
PCB-040/071	18	18	0	0.135	3.18	0.660			0.660				
PCB-041	18	8	10	0.00592	0.0507	0.0279	0.00408	0.0281	0.0151				
PCB-042	18	18	0	0.0527	1.41	0.264			0.264				
PCB-043	18	15	3	0.0205	0.416	0.0942	0.00506	0.0146	0.0794				
PCB-044/047/065	18	18	0	12.5	87.5	38.4			38.4				
PCB-045/051	18	17	1	0.0179	0.229	0.0555	0.013	0.013	0.0528				
PCB-046	18	5	13	0.00291	0.0487	0.0185	0.00256	0.018	0.00789				
PCB-048	18	18	0	0.0238	0.752	0.120			0.120				
PCB-049/069	18	18	0	0.0885	2.67	0.514			0.514				
PCB-050/053	18	18	0	0.00824	0.213	0.0399			0.0399				
PCB-052	18	18	0	0.113	3.32	0.614			0.614				
PCB-054	18	0	18				0.00135	0.0135	0.00283				
PCB-055	18	4	14	0.0757	0.238	0.158	0.00313	0.0265	0.04				
PCB-056	18	18	0	0.0282	1.96	0.442			0.442				
PCB-057	18	9	9	0.00623	0.0442	0.0164	0.00564	0.0238	0.0117				
PCB-058	18	7	11	0.0128	0.037	0.0248	0.00568	0.0243	0.0136				
PCB-059/062/075	18	18	0	1.16	6.26	3.08			3.08				
PCB-060	18	18	0	8.53	59.9	29.4			29.4				
PCB-061/070/074/076	18	18	0	19.3	249	74.3			74.3				
PCB-063	18	18	0	0.26	137	8.66			8.66				
PCB-064	18	18	0	0.149	4.18	1.06			1.06				
PCB-066	18	17	1	34.3	259	112	0.024	0.024	105				
PCB-067	18	17	1	0.0242	0.194	0.0771	0.0172	0.0172	0.0733				

Table 3
Blue Crab Size Class A Tissue Sample Summary

Number of Number of State Sample Summary													
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average				
PCB-068	18	18	0	0.0863	0.435	0.252			0.252				
PCB-072	18	17	1	0.0309	0.226	0.0802	0.0177	0.0177	0.0762				
PCB-073	18	7	11	0.0058	0.0497	0.0177	0.00269	0.0185	0.00799				
PCB-077	18	18	0	1.63	13.4	6.12			6.12				
PCB-078	18	0	18				0.00309	0.0252	0.00578				
PCB-079	18	9	9	0.0405	0.173	0.102	0.0051	0.0215	0.0534				
PCB-080	18	3	15	0.00442	0.0119	0.00848	0.0049	0.0239	0.00626				
PCB-081	18	18	0	0.0787	0.683	0.313			0.313				
PCB-082	18	11	7	0.00589	0.103	0.0296	0.00506	0.0212	0.0205				
PCB-083	18	0	18				0.00238	0.0277	0.00487				
PCB-084	18	18	0	0.0259	0.328	0.0842			0.0842				
PCB-085/116/117	18	18	0	4.79	34.9	15.3			15.3				
PCB-086/087/097/108/119/125	18	18	0	1.3	5.21	2.90			2.90				
PCB-088/091	18	18	0	0.0452	0.817	0.223			0.223				
PCB-089	18	2	16	0.00256	0.00835	0.00546	0.00356	0.0264	0.00505				
PCB-090/101/113	18	18	0	0.441	5.62	1.86			1.86				
PCB-092	18	18	0	0.0236	0.74	0.142			0.142				
PCB-093/100	18	18	0	0.297	1.05	0.541			0.541				
PCB-094	18	16	2	0.00561	0.0583	0.0212	0.00542	0.0198	0.0196				
PCB-095	18	18	0	0.0603	0.949	0.236			0.236				
PCB-096	18	1	17	0.00296	0.00296	0.00296	0.000738	0.011	0.00211				
PCB-098/102	18	17	1	0.014	0.224	0.0559	0.00542	0.00542	0.0529				
PCB-099	18	18	0	18.5	95.4	42.6			42.6				
PCB-103	18	8	10	0.00292	0.0609	0.0167	0.00424	0.0184	0.00992				
PCB-104	18	0	18				0.00093	0.00974	0.0019				
PCB-105	18	18	0	8.74	60.3	28.5			28.5				
PCB-106	18	0	18				0.00382	0.0278	0.00601				
PCB-107/124	18	17	1	0.0227	0.234	0.0888	0.0192	0.0192	0.0844				
PCB-109	18	18	0	1.08	6.54	3.46			3.46				
PCB-110/115	18	18	0	0.825	5.96	3.15			3.15				
PCB-111	18	17	1	0.0241	0.109	0.0452	0.0097	0.0097	0.043				
PCB-112	18	2	16	0.0264	0.0444	0.0354	0.00147	0.0189	0.00692				
PCB-114	18	18	0	0.62	4.09	2.05			2.05				
PCB-118	18	18	0	22.7	119	62.2			62.2				
PCB-120	18	18	0	0.0873	0.421	0.253			0.253				
PCB-121	18	10	8	0.0134	0.061	0.0234	0.00146	0.014	0.0146				
PCB-122	18	4	14	0.00788	0.0495	0.0224	0.0055	0.0308	0.0107				
PCB-123	18	18	0	0.476	3.45	1.54			1.54				
PCB-126	18	18	0	0.0688	0.419	0.209			0.209				
PCB-127	18	5	13	0.0141	0.0675	0.0413	0.00357	0.028	0.0155				
PCB-128/166	18	18	0	1.25	5.74	2.98			2.98				
PCB-129/138/163	18	18	0	8.94	38.2	22.7			22.7				
PCB-130	18	18	0	0.115	1.26	0.489			0.489				
PCB-131	18	11	7	0.0077	0.0351	0.0184	0.00717	0.0407	0.015				
PCB-132	18	14	4	0.0142	0.12	0.0327	0.0181	0.0378	0.0283				

Table 3
Blue Crab Size Class A Tissue Sample Summary

Number of													
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average				
PCB-133	18	18	0	0.183	0.908	0.407			0.407				
PCB-134	18	1	17	0.013	0.013	0.013	0.00317	0.0408	0.00752				
PCB-135/151	18	17	1	0.0392	0.754	0.116	0.0132	0.0132	0.11				
PCB-136	18	16	2	0.00562	0.0885	0.0188	0.00764	0.0113	0.0177				
PCB-137	18	18	0	0.511	3.03	1.53			1.53				
PCB-139/140	18	18	0	0.273	1.22	0.637			0.637				
PCB-141	18	10	8	0.00755	0.416	0.110	0.00335	0.033	0.0636				
PCB-142	18	2	16	0.0131	0.015	0.0141	0.00298	0.0373	0.00695				
PCB-143	18	1	17	0.00322	0.00322	0.00322	0.00351	0.0346	0.00602				
PCB-144	18	17	1	0.0131	0.134	0.0394	0.00972	0.00972	0.0375				
PCB-145	18	0	18				0.00052	0.0118	0.00182				
PCB-146	18	18	0	1.54	8.87	4.66			4.66				
PCB-147/149	18	18	0	0.153	4.01	0.966			0.966				
PCB-148	18	18	0	0.0169	0.301	0.0836			0.0836				
PCB-150	18	0	18				0.00048	0.0114	0.00175				
PCB-152	18	0	18				0.000469	0.0111	0.00171				
PCB-153/168	18	18	0	9.79	40.5	26.1			26.1				
PCB-154	18	18	0	0.211	1.68	0.721			0.721				
PCB-155	18	18	0	0.0132	0.397	0.0993			0.0993				
PCB-156/157	18	18	0	1.35	6.3	3.35			3.35				
PCB-158	18	18	0	1.1	5.06	2.93			2.93				
PCB-159	18	5	13	0.00727	0.0131	0.0101	0.00356	0.023	0.00642				
PCB-160	18	0	18				0.00195	0.0288	0.00475				
PCB-161	18	0	18				0.00183	0.024	0.00421				
PCB-162	18	18	0	0.0483	0.207	0.133			0.133				
PCB-164	18	10	8	0.0126	0.261	0.0853	0.0019	0.0238	0.0493				
PCB-165	18	15	3	0.00473	0.0463	0.0167	0.0138	0.0261	0.0155				
PCB-167	18	18	0	0.601	2.2	1.28			1.28				
PCB-169	18	4	14	0.00826	0.0295	0.0209	0.00174	0.0342	0.00865				
PCB-170	18	18	0	0.61	4.04	2.41			2.41				
PCB-171/173	18	18	0	0.287	1.63	1.02			1.02				
PCB-172	18	18	0	0.207	1	0.605			0.605				
PCB-174	18	16	2	0.0149	0.454	0.0631	0.0112	0.0209	0.057				
PCB-175	18	18	0	0.0517	0.37	0.171			0.171				
PCB-176	18	16	2	0.00535	0.112	0.0275	0.00158	0.00659	0.0247				
PCB-177	18	18	0	0.0947	1.96	0.681			0.681				
PCB-178	18	18	0	0.283	1.81	0.758			0.758				
PCB-179	18	15	3	0.00453	0.118	0.0153	0.0044	0.00726	0.0133				
PCB-180/193	18	18	0	2.7	13.1	8.69			8.69				
PCB-181	18	18	0	0.0237	0.12	0.0638			0.0638				
PCB-182	18	17	1	0.0247	0.165	0.0557	0.0085	0.0085	0.0528				
PCB-183/185	18	18	0	0.861	4.39	2.64			2.64				
PCB-184	18	18	0	0.0117	0.183	0.0562			0.0562				
PCB-186	18	0	18				0.000463	0.00806	0.00141				
PCB-187	18	18	0	1.34	9.74	5.25			5.25				

Table 3
Blue Crab Size Class A Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-188	18	18	0	0.011	0.167	0.0557			0.0557
PCB-189	18	18	0	0.0451	0.238	0.151			0.151
PCB-190	18	18	0	0.152	0.869	0.579			0.579
PCB-191	18	18	0	0.0511	0.249	0.158			0.158
PCB-192	18	2	16	0.00175	0.0144	0.00808	0.00126	0.0763	0.0055
PCB-194	18	18	0	0.282	1.72	1.06			1.06
PCB-195	18	18	0	0.0732	0.603	0.331			0.331
PCB-196	18	18	0	0.261	1.59	0.892			0.892
PCB-197/200	18	18	0	0.0376	0.665	0.235			0.235
PCB-198/199	18	18	0	0.505	3.32	1.68			1.68
PCB-201	18	18	0	0.0722	0.803	0.307			0.307
PCB-202	18	18	0	0.148	3.2	0.759			0.759
PCB-203	18	18	0	0.25	1.55	0.879			0.879
PCB-204	18	17	1	0.00767	0.157	0.0361	0.00738	0.00738	0.0343
PCB-205	18	17	1	0.0107	0.141	0.0424	0.0124	0.0124	0.0404
PCB-206	18	18	0	0.111	3.44	0.928			0.928
PCB-207	18	18	0	0.0958	3.24	0.888			0.888
PCB-208	18	18	0	0.119	4.92	1.12			1.12
PCB-209	18	18	0	1.64	175	27.8			27.8
Total PCB Congener (U = 1/2)	18	18	0	229	1,264	655			655
Dioxins and Furans (ng/kg)	•	•	•		•	•		•	
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	18	18	0	1.71	184	21.6			21.6
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	18	18	0	3.97	50.6	14.1			14.1
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	18	18	0	0.49	13.7	3.16			3.16
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	18	12	6	0.808	5.08	2.00	1.02	2.71	1.63
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	18	2	16	0.447	1.39	0.919	0.191	0.48	0.245
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	18	12	6	0.668	2.48	1.58	0.417	1.91	1.23
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	18	2	16	0.535	0.819	0.677	0.156	1.21	0.299
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	18	10	8	0.352	1.21	0.750	0.156	1.12	0.539
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	18	5	13	0.408	0.5	0.451	0.178	1.17	0.333
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	18	0	18				0.121	0.44	0.135
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	18	1	17	0.425	0.425	0.425	0.169	1.22	0.265
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	18	15	3	1.15	6.65	3.10	0.334	1.46	2.66
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	18	9	9	0.309	1.12	0.536	0.133	0.488	0.333
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	18	2	16	0.294	0.434	0.364	0.118	0.634	0.16
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	18	16	2	1.28	6.19	2.90	0.554	1.14	2.63
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	18	18	0	1.65	33	11.3			11.3
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	18	18	0	0.398	4.75	2.92			2.92
Total Dioxin/Furan (U = 1/2)	18	18	0	17.3	235	63.6			63.6

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram PCB - polychlorinated biphenyl

-- Results not applicable

Table 4
Blue Crab Size Class B Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)	<u> </u>		Samples		1				
Lipids	3	3	0	0.575	0.943	0.725			0.725
Moisture, percent	3	3	0	69.0	76.5	73.1			73.1
Metals (mg/kg)					I .			l .	
Mercury	3	3	0	0.058	0.083	0.067			0.067
Semivolatile Organic Carbons (μg/kg)					1			<u> </u>	
1,3-Dichlorobenzene	3	0	3				357	748	304
Hexachlorobenzene	3	0	3				357	748	304
PCBs (μg/kg)		<u> </u>		<u> </u>	1				
PCB-001	3	2	1	0.00439	0.0049	0.00465	0.00317	0.00317	0.00363
PCB-002	3	1	2	0.00252	0.00252	0.00252	0.000766	0.00292	0.00145
PCB-003	3	0	3				0.00276	0.00419	0.00166
PCB-004	3	2	1	0.0303	0.0398	0.0351	0.0392	0.0392	0.0299
PCB-005	3	0	3				0.00201	0.00987	0.00237
PCB-006	3	2	1	0.0156	0.0234	0.0195	0.00797	0.00797	0.0143
PCB-007	3	0	3				0.00171	0.00831	0.002
PCB-008	3	3	0	0.0654	0.119	0.100			0.100
PCB-009	3	0	3				0.00437	0.00895	0.00307
PCB-010	3	0	3				0.0013	0.0155	0.00306
PCB-011	3	0	3				0.0178	0.0292	0.0116
PCB-012/013	3	3	0	0.0159	0.0255	0.0207			0.0207
PCB-014	3	0	3				0.0017	0.00839	0.00201
PCB-015	3	3	0	1.58	1.92	1.8			1.8
PCB-016	3	3	0	0.0128	0.0255	0.0209			0.0209
PCB-017	3	3	0	0.0619	0.226	0.139			0.139
PCB-018/030	3	3	0	0.184	0.459	0.328			0.328
PCB-019	3	3	0	0.00903	0.0203	0.0165			0.0165
PCB-020/028	3	3	0	69.2	74.2	71.9			71.9
PCB-021/033	3	3	0	0.14	0.281	0.195			0.195
PCB-022	3	3	0	0.307	0.66	0.515			0.515
PCB-023	3	0	3				0.000778	0.00242	0.000705
PCB-024	3	0	3				0.000809	0.00231	0.000698
PCB-025	3	3	0	0.0748	0.0869	0.0796			0.0796
PCB-026/029	3	3	0	0.142	0.294	0.206			0.206
PCB-027	3	3	0	0.00793	0.0507	0.0264			0.0264
PCB-031	3	3	0	1.64	2.05	1.80			1.80
PCB-032	3	3	0	0.107	0.549	0.290			0.290
PCB-034	3	3	0	0.0107	0.0289	0.0169			0.0169
PCB-035	3	3	0	0.0108	0.0211	0.0164			0.0164
PCB-036	3	0	3				0.00205	0.0065	0.00182
PCB-037	3	3	0	5.99	7.65	6.62			6.62
PCB-038	3	2	1	0.0209	0.0252	0.0231	0.00677	0.00677	0.0165
PCB-039	3	2	1	0.0396	0.0585	0.0491	0.00677	0.00677	0.0338
PCB-040/071	3	3	0	0.486	3.62	1.77			1.77
PCB-041	3	1	2	0.0423	0.0423	0.0423	0.00378	0.00636	0.0175
PCB-042	3	3	0	0.316	0.681	0.508			0.508
PCB-043	3	3	0	0.0665	0.444	0.232			0.232
PCB-044/047/065	3	3	0	44.3	50.2	47.0			47.0

Table 4
Blue Crab Size Class B Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-045/051	3	3	0	0.04	0.165	0.109			0.109
PCB-046	3	3	0	0.00654	0.0215	0.0157			0.0157
PCB-048	3	3	0	0.139	0.356	0.268			0.268
PCB-049/069	3	3	0	0.478	1.04	0.819			0.819
PCB-050/053	3	3	0	0.0366	0.149	0.0972			0.0972
PCB-052	3	3	0	0.808	1.9	1.40			1.40
PCB-054	3	1	2	0.00187	0.00187	0.00187	0.00102	0.00396	0.00145
PCB-055	3	1	2	0.129	0.129	0.129	0.00448	0.005	0.0446
PCB-056	3	3	0	0.693	0.974	0.854			0.854
PCB-057	3	2	1	0.0153	0.0334	0.0244	0.0116	0.0116	0.0182
PCB-058	3	2	1	0.0169	0.0262	0.0216	0.00437	0.00437	0.0151
PCB-059/062/075	3	3	0	3.36	3.67	3.48			3.48
PCB-060	3	3	0	34.9	38.8	36.9			36.9
PCB-061/070/074/076	3	3	0	72.1	89.1	80.6			80.6
PCB-063	3	3	0	1.3	1.72	1.46			1.46
PCB-064	3	3	0	1.42	2.45	1.89			1.89
PCB-066	3	3	0	122	163	140			140
PCB-067	3	3	0	0.0948	0.127	0.111			0.111
PCB-068	3	3	0	0.192	0.305	0.241			0.241
PCB-072	3	3	0	0.0572	0.117	0.0795			0.0795
PCB-073	3	0	3			-	0.00231	0.00435	0.00164
PCB-077	3	3	0	7.25	8.4	7.94			7.94
PCB-078	3	0	3				0.0041	0.0124	0.00351
PCB-079	3	1	2	0.101	0.101	0.101	0.00403	0.0102	0.036
PCB-080	3	0	3				0.00393	0.011	0.00322
PCB-081	3	3	0	0.4	0.454	0.430			0.430
PCB-082	3	3	0	0.0373	0.0528	0.0431			0.0431
PCB-083	3	0	3				0.00458	0.00769	0.00282
PCB-084	3	3	0	0.106	0.463	0.283			0.283
PCB-085/116/117	3	3	0	23.9	25.6	24.6			24.6
PCB-086/087/097/108/119/125	3	3	0	3.96	4.31	4.08			4.08
PCB-088/091	3	3	0	0.179	0.735	0.444			0.444
PCB-089	3	2	1	0.0146	0.0162	0.0154	0.00663	0.00663	0.0114
PCB-090/101/113	3	3	0	2.06	3.13	2.63			2.63
PCB-092	3	3	0	0.135	0.519	0.323			0.323
PCB-093/100	3	3	0	0.716	0.797	0.748			0.748
PCB-094	3	3	0	0.0187	0.109	0.0553			0.0553
PCB-095	3	3	0	0.274	1.33	0.718			0.718
PCB-096	3	2	1	0.00374	0.00457	0.00416	0.0022	0.0022	0.00314
PCB-098/102	3	3	0	0.0566	0.248	0.147			0.147
PCB-099	3	3	0	59	66.1	61.8			61.8
PCB-103	3	3	0	0.00935	0.0145	0.0121			0.0121
PCB-104	3	1	2	0.00634	0.00634	0.00634	0.0015	0.00323	0.0029
PCB-105	3	3	0	36.9	51	42.37			42.37
PCB-106	3	0	3				0.00498	0.0138	0.00397
PCB-107/124	3	3	0	0.106	0.154	0.123			0.123

Table 4
Blue Crab Size Class B Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-109	3	3	0	4.17	5.2	4.63			4.63
PCB-110/115	3	3	0	4.87	5.75	5.19			5.19
PCB-111	3	3	0	0.0309	0.0565	0.0408			0.0408
PCB-112	3	0	3				0.00286	0.00458	0.00173
PCB-114	3	3	0	2.84	3.78	3.20			3.20
PCB-118	3	3	0	81.2	110	91.0			91.0
PCB-120	3	3	0	0.199	0.372	0.264			0.264
PCB-121	3	3	0	0.0141	0.0231	0.0171			0.0171
PCB-122	3	3	0	0.0191	0.101	0.053			0.053
PCB-123	3	3	0	1.97	2.82	2.29			2.29
PCB-126	3	3	0	0.26	0.366	0.302			0.302
PCB-127	3	2	1	0.0519	0.0536	0.0528	0.0135	0.0135	0.0374
PCB-128/166	3	3	0	4.11	5.82	4.72			4.72
PCB-129/138/163	3	3	0	26.1	44.7	32.8			32.8
PCB-130	3	3	0	0.397	0.66	0.518			0.518
PCB-131	3	2	1	0.0165	0.0237	0.0201	0.0112	0.0112	0.0153
PCB-132	3	3	0	0.037	0.0475	0.0408			0.0408
PCB-133	3	3	0	0.34	0.672	0.465			0.465
PCB-134	3	1	2	0.0159	0.0159	0.0159	0.0119	0.0121	0.0093
PCB-135/151	3	3	0	0.0779	0.204	0.138			0.138
PCB-136	3	3	0	0.021	0.0492	0.0349			0.0349
PCB-137	3	3	0	2.21	2.81	2.45			2.45
PCB-139/140	3	3	0	0.816	0.923	0.853			0.853
PCB-141	3	2	1	0.105	0.109	0.107	0.00983	0.00983	0.073
PCB-142	3	0	3				0.00608	0.0101	0.00436
PCB-143	3	1	2	0.0058	0.0058	0.00580	0.00911	0.00938	0.00502
PCB-144	3	3	0	0.0278	0.0702	0.0465			0.0465
PCB-145	3	0	3				0.00195	0.00366	0.00129
PCB-146	3	3	0	4.24	7.64	5.42			5.42
PCB-147/149	3	3	0	0.868	1.37	1.04			1.04
PCB-148	3	3	0	0.0553	0.09	0.0672			0.0672
PCB-150	3	0	3				0.00187	0.00354	0.00124
PCB-152	3	2	1	0.000974	0.00252	0.00175	0.00353	0.00353	0.00175
PCB-153/168	3	3	0	26.1	46.5	33.3			33.3
PCB-154	3	3	0	0.573	1.1	0.753			0.753
PCB-155	3	3	0	0.0343	0.0889	0.0526			0.0526
PCB-156/157	3	3	0	3.98	6.94	5.02			5.02
PCB-158	3	3	0	3.72	5.56	4.41			4.41
PCB-159	3	1	2	0.00921	0.00921	0.00921	0.00418	0.0142	0.00613
PCB-160	3	0	3				0.00472	0.00772	0.00327
PCB-161	3	0	3				0.00403	0.00696	0.00293
PCB-162	3	3	0	0.106	0.163	0.134			0.134
PCB-164	3	3	0	0.0461	0.101	0.0680			0.0680
PCB-165	3	3	0	0.0107	0.0201	0.0141			0.0141
PCB-167	3	3	0	1.41	2.51	1.81			1.81
PCB-169	3	0	3				0.00451	0.0122	0.00354

Table 4
Blue Crab Size Class B Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-170	3	3	0	2.29	5.92	3.52			3.52
PCB-171/173	3	3	0	0.949	2.08	1.34			1.34
PCB-172	3	3	0	0.482	1.43	0.806			0.806
PCB-174	3	3	0	0.0407	0.0587	0.0490			0.0490
PCB-175	3	3	0	0.142	0.311	0.200			0.200
PCB-176	3	2	1	0.0109	0.026	0.0185	0.00172	0.00172	0.0126
PCB-177	3	3	0	0.36	1.13	0.642			0.642
PCB-178	3	3	0	0.579	1.45	0.893			0.893
PCB-179	3	3	0	0.00727	0.0109	0.00898			0.00898
PCB-180/193	3	3	0	7.45	21	12.2			12.2
PCB-181	3	3	0	0.062	0.118	0.0826			0.0826
PCB-182	3	3	0	0.0398	0.0883	0.0567			0.0567
PCB-183/185	3	3	0	2.54	5.18	3.43			3.43
PCB-184	3	3	0	0.0284	0.0369	0.0317			0.0317
PCB-186	3	0	3				0.00182	0.00425	0.00142
PCB-187	3	3	0	3.67	12.6	6.75			6.75
PCB-188	3	3	0	0.0207	0.0495	0.0319			0.0319
PCB-189	3	3	0	0.133	0.321	0.196			0.196
PCB-190	3	3	0	0.481	1.42	0.812			0.812
PCB-191	3	3	0	0.139	0.4	0.230			0.230
PCB-192	3	0	3				0.00273	0.00485	0.00175
PCB-194	3	3	0	0.824	2.74	1.47			1.47
PCB-195	3	3	0	0.257	0.714	0.414			0.414
PCB-196	3	3	0	0.76	1.99	1.17			1.17
PCB-197/200	3	3	0	0.136	0.25	0.175			0.175
PCB-198/199	3	3	0	1.32	3.15	1.94			1.94
PCB-201	3	3	0	0.208	0.405	0.275			0.275
PCB-202	3	3	0	0.329	0.771	0.480			0.480
PCB-203	3	3	0	0.712	1.85	1.10			1.10
PCB-204	3	3	0	0.0146	0.0234	0.0178			0.0178
PCB-205	3	2	1	0.0176	0.0197	0.0187	0.0571	0.0571	0.022
PCB-206	3	3	0	0.405	0.988	0.609			0.609
PCB-207	3	3	0	0.408	0.827	0.556			0.556
PCB-208	3	3	0	0.458	1.01	0.648			0.648
PCB-209	3	3	0	8.14	22.2	13.3			13.3
Total PCB Congener (U = 1/2)	3	3	0	722	936	804			804

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram -- Results not applicable

PCB - polychlorinated biphenyl

Table 5
Brown Shrimp Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)		<u> </u>	Samples			l			
Lipids	8	8	0	1.15	1.64	1.37			1.37
Moisture, percent	8	8	0	74.0	76.2	75.1			75.1
Metals (mg/kg)	<u> </u>	<u> </u>	, i	70	70.2	75.2			, 3.1
Mercury	8	8	0	0.025	0.065	0.041			0.041
Semivolatile Organic Carbons (µg/kg)				0.020	0.000	1 0.0.1		l l	0.0.1
1,3-Dichlorobenzene	8	0	8				340	709	223
Hexachlorobenzene	8	8	0	537	913	710			710
PCBs (µg/kg)				557	3 2 3	, , , ,			, 10
PCB-001	8	8	0	0.0552	0.446	0.163			0.163
PCB-002	8	3	5	0.00536	0.0611	0.0349	0.00197	0.0472	0.0206
PCB-003	8	6	2	0.0317	0.216	0.0913	0.0265	0.0422	0.0727
PCB-004	8	8	0	0.239	0.698	0.461			0.461
PCB-005	8	1	7	0.0557	0.0557	0.0557	0.00676	0.0971	0.0245
PCB-006	8	8	0	0.262	0.64	0.484			0.484
PCB-007	8	7	1	0.0595	0.119	0.092	0.0797	0.0797	0.0854
PCB-008	8	8	0	1.81	4.33	3.06			3.06
PCB-009	8	8	0	0.082	0.204	0.147			0.147
PCB-010	8	3	5	0.0132	0.0333	0.027	0.0196	0.072	0.0213
PCB-011	8	8	0	0.0835	0.286	0.195			0.195
PCB-012/013	8	8	0	0.18	0.33	0.274			0.274
PCB-014	8	0	8				0.00568	0.0805	0.0157
PCB-015	8	8	0	1.46	2.49	1.96			1.96
PCB-016	8	8	0	0.573	2.03	1.33			1.33
PCB-017	8	8	0	2.63	5.5	4.35			4.35
PCB-018/030	8	8	0	4.11	12.3	8.67			8.67
PCB-019	8	8	0	0.113	0.317	0.223			0.223
PCB-020/028	8	8	0	64.4	121	91.3			91.3
PCB-021/033	8	8	0	7.95	15.7	12.3			12.3
PCB-022	8	8	0	9.79	18.9	15.3			15.3
PCB-022	8	2	6	0.00591	0.0152	0.0106	0.00281	0.0569	0.0135
PCB-024	8	8	0	0.0358	0.0821	0.0647			0.0133
PCB-025	8	8	0	1.32	3.06	2.37			2.37
PCB-026/029	8	8	0	5.84	11	8.75			8.75
PCB-027	8	8	0	0.263	0.662	0.514			0.514
PCB-031	8	8	0	52.1	107	81.7			81.7
PCB-032	8	8	0	2.19	5.47	4.03			4.03
PCB-032	8	8	0	0.107	0.298	0.215			0.215
PCB-035	8	8	0	0.484	0.912	0.745			0.745
PCB-036	8	0	8				0.0052	0.13	0.743
PCB-037	8	8	0	6.72	13	10.1			10.1
PCB-038	8	1	7	0.0781	0.0781	0.0781	0.00629	0.131	0.0351
PCB-039	8	2	6	0.504	0.609	0.0781	0.00629	0.0639	0.152
PCB-040/071	8	8	0	4.8	13.4	10.1		0.0039	10.1
PCB-040/071 PCB-041	8	8	0	0.797	1.87	1.27		+	1.27
F CD=041	٥ -	0	U	0./3/	1.0/	1.4/			1.27

Table 5
Brown Shrimp Tissue Sample Summary

			Brown Snrimp 118	sue Sample Sumn	nary				
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-043	8	2	6	0.381	1.03	0.706	0.0106	0.285	0.207
PCB-044/047/065	8	8	0	40	102	79.3			79.3
PCB-045/051	8	8	0	0.647	2.36	1.65			1.65
PCB-046	8	8	0	0.107	0.487	0.339			0.339
PCB-048	8	8	0	3.65	10.5	7.54			7.54
PCB-049/069	8	8	0	32.1	90.9	66.1			66.1
PCB-050/053	8	8	0	0.693	3.02	2.09			2.09
PCB-052	8	8	0	44.7	124	89.0			89.0
PCB-054	8	4	4	0.0124	0.0276	0.0184	0.0369	0.11	0.0246
PCB-055	8	5	3	0.391	1.82	1.29	0.0312	0.232	0.826
PCB-056	8	8	0	22.3	45.3	37.2			37.2
PCB-057	8	8	0	0.325	0.877	0.553			0.553
PCB-058	8	8	0	0.221	0.648	0.440			0.440
PCB-059/062/075	8	8	0	3.41	7.91	6.15			6.15
PCB-060	8	8	0	20.9	41	32.9			32.9
PCB-061/070/074/076	8	8	0	152	356	275			275
PCB-063	8	8	0	4.23	8.82	6.70			6.70
PCB-064	8	8	0	15.7	36.8	28.7			28.7
PCB-066	8	8	0	104	205	167			167
PCB-067	8	8	0	1.44	3.45	2.48			2.48
PCB-068	8	8	0	0.404	1.13	0.806			0.806
PCB-072	8	8	0	0.816	2.2	1.64			1.64
PCB-073	8	6	2	0.357	0.913	0.684	0.0044	0.0884	0.518
PCB-077	8	8	0	6.31	12	9.32			9.32
PCB-078	8	0	8				0.00884	0.218	0.0372
PCB-079	8	2	6	0.938	1.1	1.02	0.00771	0.183	0.277
PCB-080	8	0	8				0.00838	0.197	0.0337
PCB-081	8	8	0	0.321	0.57	0.429			0.429
PCB-082	8	8	0	1.12	2.87	2.19			2.19
PCB-083	8	3	5	1.14	1.87	1.47	0.0136	0.223	0.579
PCB-084	8	8	0	0.68	2.28	1.66			1.66
PCB-085/116/117	8	8	0	14.7	30.2	23.3			23.3
PCB-086/087/097/108/119/125	8	8	0	20.1	43.3	33.2			33.2
PCB-088/091	8	8	0	1.32	3.82	2.87			2.87
PCB-089	8	7	1	0.0629	0.203	0.132	0.195	0.195	0.128
PCB-090/101/113	8	8	0	43.9	105	77.8			77.8
PCB-092	8	8	0	5.5	12	9.23			9.23
PCB-093/100	8	8	0	0.401	1.19	0.823			0.823
PCB-094	8	6	2	0.0282	0.0872	0.0658	0.104	0.202	0.0685
PCB-095	8	8	0	4.29	12	9.18			9.18
PCB-096	8	7	1	0.0187	0.0911	0.0594	0.127	0.127	0.0599
PCB-098/102	8	8	0	0.376	1.22	0.882			0.882
PCB-099	8	8	0	33.9	69.3	55.8			55.8
PCB-103	8	8	0	0.29	0.908	0.622			0.622
PCB-104	8	0	8				0.00297	0.126	0.0191
PCB-105	8	8	0	27.2	45.8	37.2			37.2

Table 5
Brown Shrimp Tissue Sample Summary

			Brown Shrimp Tis	suc sample sum				1	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-106	8	0	8				0.0107	0.171	0.032
PCB-107/124	8	8	0	2.31	4.44	3.44			3.44
PCB-109	8	8	0	4.89	9.19	7.33			7.33
PCB-110/115	8	8	0	19.9	43.3	33.9			33.9
PCB-111	8	7	1	0.0397	0.111	0.0780	0.133	0.133	0.0766
PCB-112	8	1	7	0.415	0.415	0.415	0.00436	0.134	0.0691
PCB-114	8	8	0	1.95	3.3	2.62			2.62
PCB-118	8	8	0	59.2	113	84.9			84.9
PCB-120	8	8	0	0.213	0.565	0.424			0.424
PCB-121	8	2	6	0.0305	0.0318	0.0312	0.00421	0.135	0.0275
PCB-122	8	8	0	0.804	1.64	1.19			1.19
PCB-123	8	8	0	1.48	2.69	2.15			2.15
PCB-126	8	6	2	0.17	0.309	0.226	0.0512	0.2	0.185
PCB-127	8	2	6	0.101	0.106	0.104	0.0101	0.168	0.0536
PCB-128/166	8	8	0	3.09	9.82	6.32			6.32
PCB-129/138/163	8	8	0	23.6	66.4	44.5			44.5
PCB-130	8	8	0	1.48	4.7	3.17			3.17
PCB-131	8	7	1	0.0575	0.237	0.126	0.269	0.269	0.127
PCB-132	8	8	0	0.659	2.24	1.49			1.49
PCB-133	8	8	0	0.438	1.31	0.874			0.874
PCB-134	8	7	1	0.298	1.09	0.690	0.0431	0.0431	0.606
PCB-135/151	8	8	0	2.39	7.65	5.01			5.01
PCB-136	8	8	0	0.102	0.508	0.299			0.299
PCB-137	8	8	0	1.25	3.39	2.21			2.21
PCB-139/140	8	8	0	0.417	1.32	0.836			0.836
PCB-141	8	8	0	3.36	9.22	5.99			5.99
PCB-142	8	0	8				0.00614	0.268	0.0346
PCB-143	8	1	7	0.0157	0.0157	0.0157	0.00768	0.25	0.0339
PCB-144	8	8	0	0.581	1.59	1.08			1.08
PCB-145	8	1	7	0.00104	0.00104	0.00104	0.00124	0.114	0.0159
PCB-146	8	8	0	0.0368	11.2	6.49			6.49
PCB-147/149	8	8	0	6.09	20.5	13.0			13.0
PCB-148	8	8	0	0.0506	0.208	0.122			0.122
PCB-150	8	4	4	0.00827	0.0273	0.0183	0.02	0.11	0.0238
PCB-152	8	4	4	0.00434	0.0133	0.00827	0.0197	0.108	0.0185
PCB-153/168	8	8	0	18.7	59.9	37.55			37.55
PCB-154	8	8	0	0.457	1.65	1.01			1.01
PCB-155	8	8	0	0.0615	0.252	0.169			0.169
PCB-156/157	8	8	0	3.21	6.23	4.68			4.68
PCB-158	8	8	0	2.68	6.7	4.73			4.73
PCB-159	8	2	6	0.16	0.2	0.180	0.00422	0.158	0.603
PCB-160	8	0	8				0.00443	0.193	0.251
PCB-161	8	1	7	8.24	8.24	8.24	0.00443	0.186	1.05
PCB-161	8	7	1	0.135	0.326	0.236	0.0165	0.180	0.208
PCB-102 PCB-164	8	8	0	1.13	3.19	2.13		0.0103	2.13
PCB-104 PCB-165	8	4	4	0.0115	0.0358	0.0195	0.0304	0.206	0.0336
L CD-103		4	4	0.0112	0.0556	0.0132	0.0304	0.200	0.0550

Table 5
Brown Shrimp Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-167	8	8	0	1.14	2.16	1.73			1.73
PCB-169	8	0	8	-			0.00376	0.288	0.0411
PCB-170	8	8	0	2.95	7.29	5.68			5.68
PCB-171/173	8	8	0	0.862	2.11	1.59			1.59
PCB-172	8	8	0	0.605	1.63	1.21			1.21
PCB-174	8	8	0	1.33	3.03	2.27			2.27
PCB-175	8	8	0	0.159	0.429	0.315			0.315
PCB-176	8	8	0	0.0532	0.261	0.179			0.179
PCB-177	8	8	0	1.77	4.68	3.52			3.52
PCB-178	8	8	0	0.742	2.08	1.52			1.52
PCB-179	8	8	0	0.18	0.618	0.433			0.433
PCB-180/193	8	8	0	7.89	17.7	13.8			13.8
PCB-181	8	6	2	0.0555	0.148	0.104	0.0457	0.0874	0.0863
PCB-182	8	5	3	0.0323	0.114	0.0683	0.00526	0.0828	0.0502
PCB-183/185	8	8	0	2.32	5.66	4.33			4.33
PCB-184	8	7	1	0.0211	0.091	0.0656	0.0338	0.0338	0.0595
PCB-186	8	0	8				0.00119	0.0679	0.00964
PCB-187	8	8	0	5.64	13.9	11.2			11.2
PCB-188	8	7	1	0.0275	0.0899	0.0641	0.102	0.102	0.0625
PCB-189	8	8	0	0.143	0.35	0.256			0.256
PCB-190	8	8	0	0.754	1.81	1.36			1.36
PCB-191	8	8	0	0.15	0.387	0.293			0.293
PCB-192	8	2	6	0.00415	0.00785	0.00600	0.00597	0.0739	0.0135
PCB-194	8	8	0	1.06	2.85	2.13			2.13
PCB-195	8	8	0	0.351	0.88	0.689			0.689
PCB-196	8	8	0	0.564	1.28	1.04			1.04
PCB-197/200	8	8	0	0.15	0.418	0.324			0.324
PCB-198/199	8	8	0	1.19	2.94	2.31			2.31
PCB-201	8	8	0	0.166	0.402	0.313			0.313
PCB-202	8	8	0	0.494	1.37	0.991			0.991
PCB-203	8	8	0	0.704	1.62	1.30			1.30
PCB-204	8	6	2	0.014	0.0443	0.0322	0.0334	0.0627	0.0302
PCB-205	8	8	0	0.0543	0.14	0.107			0.107
PCB-206	8	8	0	0.623	1.59	1.22			1.22
PCB-207	8	8	0	0.343	1.09	0.812			0.812
PCB-208	8	8	0	0.483	1.39	1.05			1.05
PCB-209	8	8	0	14.2	44.8	29.5			29.5
Total PCB Congener (U = 1/2)	8	8	0	997	2192	1716			1716

mg/kg - milligrams per kilogram

μg/kg - micrograms per kilogram

-- Results not applicable

PCB - polychlorinated biphenyl

Table 6
Oyster Tissue Sample Summary

			Oyster Hissae	Sample Summary					
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)									
Lipids	1	1	0	1.1	1.1	1.1			1.1
Moisture, percent	1	1	0	82.7	82.7	82.7			82.7
Metals (mg/kg)									
Lead	1	1	0	0.331	0.331	0.331			0.331
Mercury	1	1	0	0.051	0.051	0.051			0.051
Polyaromatic Hydrocarbons (µg/kg)									
2-Methylnaphthalene	1	0	1				13.7	13.7	13.7
Acenaphthene	1	0	1				13.7	13.7	13.7
Acenaphthylene	1	1	0	16.1	16.1	16.1			16.1
Anthracene	1	1	0	14.5	14.5	14.5			14.5
Benzo(a)anthracene	1	1	0	38.5	38.5	38.5			38.5
Benzo(a)pyrene	1	1	0	13.8	13.8	13.8			13.8
Benzo(b)fluoranthene	1	1	0	37.5	37.5	37.5			37.5
Benzo(g,h,i)perylene	1	1	0	9.32	9.32	9.32			9.32
Benzo(k)fluoranthene	1	1	0	13.8	13.8	13.8			13.8
Chrysene	1	1	0	68.8	68.8	68.8			68.8
Dibenzo(a,h)anthracene	1	0	1				13.7	13.7	13.7
Fluoranthene	1	1	0	118	118	118			118
Fluorene	1	0	1				13.7	13.7	13.7
Indeno(1,2,3-c,d)pyrene	1	0	1				13.7	13.7	13.7
Naphthalene	1	0	1				13.7	13.7	13.7
Phenanthrene	1	1	0	18.7	18.7	18.7			18.7
Pyrene	1	1	0	162	162	162			162
Total HPAH (U = 1/2)	1	1	0	475	475	475			475
Total LPAH (U = 1/2)	1	1	0	76.7	76.7	76.7			76.7
Total PAH (U = 1/2)	1	1	0	552	552	552			552
Semivolatile Organic Carbons (μg/kg)			U	332	332	332			332
1,3-Dichlorobenzene	1	0	1		l	T I	704	704	704
1,4-Dichlorobenzene	1	0	1				704	704	704
Hexachlorobenzene	1	0	1				704	704	704
Hexachlorobutadiene	1	0	1	 			704	704	704
PCBs (µg/kg)	1	U	1				704	704	704
PCB-001	1	1 1	0	0.0392	0.0392	0.0392			0.0392
PCB-001 PCB-002			0	0.00943	0.00943	0.00943			0.00943
PCB-002 PCB-003	1 1	1	0	0.00943	0.00943	0.00943			0.00943
		1		0.0241					
PCB-004	1	1	0	0.0216	0.0316	1 0.0216			0.0216
PCB-005	1	1	0	0.0316		0.0316			0.0316
PCB-006	1	1	0	0.506	0.506	0.506			0.506
PCB-007	1	1	0	0.0845	0.0845	0.0845			0.0845
PCB-008	1	1	0	2.34	2.34	2.34			2.34
PCB-009	1	1	0	0.188	0.188	0.188			0.188
PCB-010	1	1	0	0.0235	0.0235	0.0235			0.0235
PCB-011	1	1	0	0.169	0.169	0.169			0.169
PCB-012/013	1	1	0	0.287	0.287	0.287			0.287
PCB-014	1	0	1				0.00608	0.00608	0.00608

Table 6
Oyster Tissue Sample Summary

			Oyster rissue	Sample Summary					
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-015	1	1	0	1.67	1.67	1.67			1.67
PCB-016	1	1	0	3.65	3.65	3.65			3.65
PCB-017	1	1	0	11.8	11.8	11.8			11.8
PCB-018/030	1	1	0	22.8	22.8	22.8			22.8
PCB-019	1	1	0	1.83	1.83	1.83			1.83
PCB-020/028	1	1	0	69.7	69.7	69.7			69.7
PCB-021/033	1	1	0	18.9	18.9	18.9			18.9
PCB-022	1	1	0	14.4	14.4	14.4			14.4
PCB-023	1	1	0	0.0252	0.0252	0.0252			0.0252
PCB-024	1	0	1				0.00376	0.00376	0.00376
PCB-025	1	1	0	2.49	2.49	2.49			2.49
PCB-026/029	1	1	0	8.89	8.89	8.89			8.89
PCB-027	1	1	0	1.04	1.04	1.04			1.04
PCB-031	1	1	0	62.2	62.2	62.2			62.2
PCB-032	1	1	0	8.7	8.7	8.7			8.7
PCB-034	1	1	0	0.402	0.402	0.402			0.402
PCB-035	1	1	0	0.864	0.864	0.864			0.864
PCB-036	1	0	1				0.0118	0.0118	0.0118
PCB-037	1	1	0	6.96	6.96	6.96			6.96
PCB-038	1	1	0	0.13	0.13	0.13			0.13
PCB-039	1	0	1				0.0124	0.0124	0.0124
PCB-040/071	1	1	0	29.8	29.8	29.8			29.8
PCB-041	1	1	0	6.41	6.41	6.41			6.41
PCB-042	1	1	0	16.4	16.4	16.4			16.4
PCB-043	1	1	0	2.47	2.47	2.47			2.47
PCB-044/047/065	1	1	0	75.6	75.6	75.6			75.6
PCB-045/051	1	1	0	9.42	9.42	9.42			9.42
PCB-046	1	1	0	1.65	1.65	1.65			1.65
PCB-048	1	1	0	16.3	16.3	16.3			16.3
PCB-049/069	1	1	0	60.8	60.8	60.8			60.8
PCB-050/053	1	1	0	8.35	8.35	8.35			8.35
PCB-052	1	1	0	97.9	97.9	97.9			97.9
PCB-054	1	1	0	0.193	0.193	0.193			0.193
PCB-055	1	1	0	0.637	0.637	0.637			0.637
PCB-056	1	1	0	37.3	37.3	37.3			37.3
PCB-057	1	1	0	0.335	0.335	0.335			0.335
PCB-058	1	1	0	0.333	0.333	0.333			0.333
PCB-059/062/075	1	1	0	6.5	6.5	6.5			6.5
PCB-060	1	1	0	19	19	19			19
PCB-060 PCB-061/070/074/076	1	1	0	177	177	177			177
PCB-063	1	1	0	4.05	4.05	4.05			4.05
PCB-063 PCB-064		+	0	37.7	37.7	37.7			37.7
	1	1							
PCB-066	1	1	0	113	113	113			113
PCB-067	1	1	0	2.23	2.23	2.23			2.23
PCB-068	1	1	0	0.553	0.553	0.553			0.553
PCB-072	1	1	0	1.09	1.09	1.09			1.09

Table 6
Oyster Tissue Sample Summary

•				Sample Summary		, ,			
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-073	1	0	1				0.0125	0.0125	0.0125
PCB-077	1	1	0	6.24	6.24	6.24			6.24
PCB-078	1	0	1				0.0132	0.0132	0.0132
PCB-079	1	0	1				0.0115	0.0115	0.0115
PCB-080	1	0	1				0.0125	0.0125	0.0125
PCB-081	1	1	0	0.288	0.288	0.288			0.288
PCB-082	1	1	0	5.71	5.71	5.71			5.71
PCB-083	1	1	0	2.99	2.99	2.99			2.99
PCB-084	1	1	0	9.59	9.59	9.59			9.59
PCB-085/116/117	1	1	0	12.8	12.8	12.8			12.8
PCB-086/087/097/108/119/125	1	1	0	38	38	38			38
PCB-088/091	1	1	0	9.49	9.49	9.49			9.49
PCB-089	1	1	0	0.995	0.995	0.995			0.995
PCB-090/101/113	1	1	0	56.3	56.3	56.3			56.3
PCB-092	1	1	0	9.13	9.13	9.13			9.13
PCB-093/100	1	1	0	0.885	0.885	0.885			0.885
PCB-094	1	1	0	0.383	0.383	0.383			0.383
PCB-095	1	1	0	29.5	29.5	29.5			29.5
PCB-096	1	1	0	0.479	0.479	0.479			0.479
PCB-098/102	1	1	0	2.64	2.64	2.64			2.64
PCB-099	1	1	0	35.9	35.9	35.9			35.9
PCB-103	1	1	0	0.677	0.677	0.677			0.677
PCB-104	1	1	0	0.0369	0.0369	0.0369			0.0369
PCB-105	1	1	0	20.3	20.3	20.3			20.3
PCB-106	1	0	1				0.011	0.011	0.011
PCB-107/124	1	1	0	1.85	1.85	1.85			1.85
PCB-109	1	1	0	4.79	4.79	4.79			4.79
PCB-110/115	1	1	0	47.9	47.9	47.9			47.9
PCB-111	1	1	0	0.0456	0.0456	0.0456			0.0456
PCB-112	1	0	1				0.00794	0.00794	0.00794
PCB-114	1	1	0	1.29	1.29	1.29			1.29
PCB-118	1	1	0	56.1	56.1	56.1			56.1
PCB-120	1	1	0	0.253	0.253	0.253			0.253
PCB-121	1	0	1				0.00766	0.00766	0.00766
PCB-122	1	1	0	0.758	0.758	0.758			0.758
PCB-123	1	1	0	1.22	1.22	1.22			1.22
PCB-126	1	1	0	0.175	0.175	0.175			0.175
PCB-127	1	0	1				0.0104	0.0104	0.0104
PCB-128/166	1	1	0	2.4	2.4	2.4			2.4
PCB-129/138/163	1	1	0	21	21	21			21
PCB-130	1	1	0	1.31	1.31	1.31			1.31
PCB-131	1	1	0	0.173	0.173	0.173			0.173
PCB-132	1	1	0	5.8	5.8	5.8			5.8
PCB-133	1	1	0	0.452	0.452	0.452			0.452
PCB-134	1	1	0	0.973	0.973	0.973			0.973
PCB-135/151	1	1	0	6.39	6.39	6.39			6.39

Table 6
Oyster Tissue Sample Summary

			•	Sample Summary					
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-136	1	1	0	1.96	1.96	1.96			1.96
PCB-137	1	1	0	0.138	0.138	0.138			0.138
PCB-139/140	1	1	0	0.334	0.334	0.334			0.334
PCB-141	1	1	0	0.235	0.235	0.235			0.235
PCB-142	1	0	1				0.00848	0.00848	0.00848
PCB-143	1	0	1				0.00773	0.00773	0.00773
PCB-144	1	1	0	0.502	0.502	0.502			0.502
PCB-145	1	1	0	0.0094	0.0094	0.0094			0.0094
PCB-146	1	1	0	4.16	4.16	4.16			4.16
PCB-147/149	1	1	0	16.3	16.3	16.3			16.3
PCB-148	1	1	0	0.0619	0.0619	0.0619			0.0619
PCB-150	1	1	0	0.0657	0.0657	0.0657			0.0657
PCB-152	1	1	0	0.0243	0.0243	0.0243			0.0243
PCB-153/168	1	1	0	22.3	22.3	22.3			22.3
PCB-154	1	1	0	0.599	0.599	0.599			0.599
PCB-155	1	1	0	0.126	0.126	0.126			0.126
PCB-156/157	1	1	0	1.65	1.65	1.65			1.65
PCB-158	1	1	0	0.253	0.253	0.253			0.253
PCB-159	1	0	1				0.00579	0.00579	0.00579
PCB-160	1	0	1				0.00613	0.00613	0.00613
PCB-161	1	0	1				0.00581	0.00581	0.00581
PCB-162	1	1	0	0.205	0.205	0.205			0.205
PCB-164	1	1	0	1.15	1.15	1.15			1.15
PCB-165	1	1	0	0.0165	0.0165	0.0165			0.0165
PCB-167	1	1	0	0.762	0.762	0.762			0.762
PCB-169	1	0	1				0.00606	0.00606	0.00606
PCB-170	1	1	0	0.565	0.565	0.565			0.565
PCB-171/173	1	1	0	0.54	0.54	0.54			0.54
PCB-172	1	1	0	0.101	0.101	0.101			0.101
PCB-174	1	1	0	0.208	0.208	0.208			0.208
PCB-175	1	1	0	0.0788	0.0788	0.0788			0.0788
PCB-175	1	1	0	0.316	0.316	0.316			0.316
PCB-170	1	1	0	2.09	2.09	2.09			2.09
PCB-177	1	1	0	0.804	0.804	0.804			0.804
PCB-178	1	1	0	1.29	1.29	1.29			1.29
PCB-179 PCB-180/193	1	1	0	0.628	0.628	0.628			0.628
PCB-180/193 PCB-181	1	1	0	0.028	0.00768	0.00768			0.00768
						1			
PCB-182	1	0	0	1 16	1 16	1 16	0.00262	0.00262	0.00262
PCB-183/185	1	1		1.16	1.16	1.16			1.16
PCB-184	1	1	0	0.0238	0.0238	0.0238		0.0010	0.0238
PCB-186	1	0	1				0.0019	0.0019	0.0019
PCB-187	1	1	0	6.63	6.63	6.63			6.63
PCB-188	1	1	0	0.0416	0.0416	0.0416			0.0416
PCB-189	1	1	0	0.053	0.053	0.053			0.053
PCB-190	1	1	0	0.087	0.087	0.087			0.087
PCB-191	1	1	0	0.00888	0.00888	0.00888			0.00888

Table 6
Oyster Tissue Sample Summary

Oyster rissue sample summary									
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-192	1	0	1				0.00297	0.00297	0.00297
PCB-194	1	1	0	0.0706	0.0706	0.0706			0.0706
PCB-195	1	1	0	0.0269	0.0269	0.0269			0.0269
PCB-196	1	1	0	0.0155	0.0155	0.0155			0.0155
PCB-197/200	1	1	0	0.0418	0.0418	0.0418			0.0418
PCB-198/199	1	1	0	0.167	0.167	0.167			0.167
PCB-201	1	1	0	0.11	0.11	0.11			0.11
PCB-202	1	1	0	0.917	0.917	0.917			0.917
PCB-203	1	1	0	0.0992	0.0992	0.0992			0.0992
PCB-204	1	1	0	0.00312	0.00312	0.00312			0.00312
PCB-205	1	1	0	0.00742	0.00742	0.00742			0.00742
PCB-206	1	1	0	0.0342	0.0342	0.0342			0.0342
PCB-207	1	1	0	0.0327	0.0327	0.0327			0.0327
PCB-208	1	1	0	0.0459	0.0459	0.0459			0.0459
PCB-209	1	1	0	0.911	0.911	0.911			0.911
Total PCB Congener (U = 1/2)	1	1	0	1,429	1,429	1,429			1,429
Dioxins and Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	1	1	0	10.9	10.9	10.9			10.9
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	1	1	0	6.89	6.89	6.89			6.89
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	1	1	0	0.794	0.794	0.794			0.794
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	1	1	0	0.988	0.988	0.988			0.988
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	1	0	1				0.454	0.454	0.454
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.257	0.257	0.257
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	1	0	1		-		0.328	0.328	0.328
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.251	0.251	0.251
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	1	0	1		-		0.359	0.359	0.359
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.425	0.425	0.425
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	1	0	1				0.349	0.349	0.349
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	1	1	0	1.05	1.05	1.05			1.05
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	1	0	1				0.325	0.325	0.325
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.361	0.361	0.361
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	1	1	0	2.62	2.62	2.62			2.62
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	1	1	0	14.5	14.5	14.5			14.5
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1	1	0	2.36	2.36	2.36			2.36
Total Dioxin/Furan (U = 1/2)	1	1	0	41.7	41.7	41.7			41.7

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram -- Results not applicable PCB - polychlorinated biphenyl

Table 7
White Shrimp Size Class A Tissue Sample Summary

		Wille	e Shrimp Size Clas	3 A 1133ac 3ampic	Janimary				
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)	_								
Lipids	1	1	0	0.914	0.914	0.914			0.914
Moisture, percent	1	1	0	76.9	76.9	76.9			76.9
Metals (mg/kg)									
Lead	1	1	0	0.116	0.116	0.116			0.116
Mercury	1	1	0	0.041	0.041	0.041			0.041
Polyaromatic Hydrocarbons (μg/kg)									
2-Methylnaphthalene	1	0	1				14.2	14.2	14.2
Acenaphthene	1	0	1				14.2	14.2	14.2
Acenaphthylene	1	0	1				14.2	14.2	14.2
Anthracene	1	0	1				14.2	14.2	14.2
Benzo(a)anthracene	1	0	1				14.2	14.2	14.2
Benzo(a)pyrene	1	0	1				14.2	14.2	14.2
Benzo(b)fluoranthene	1	0	1				14.2	14.2	14.2
Benzo(g,h,i)perylene	1	0	1	-			14.2	14.2	14.2
Benzo(k)fluoranthene	1	0	1	-			14.2	14.2	14.2
Chrysene	1	0	1				14.2	14.2	14.2
Dibenzo(a,h)anthracene	1	0	1				14.2	14.2	14.2
Fluoranthene	1	1	0	14.8	14.8	14.8			14.8
Fluorene	1	0	1				14.2	14.2	14.2
Indeno(1,2,3-c,d)pyrene	1	0	1				14.2	14.2	14.2
Naphthalene	1	0	1				14.2	14.2	14.2
Phenanthrene	1	1	0	18.2	18.2	18.2			18.2
Pyrene	1	1	0	29.9	29.9	29.9			29.9
Total HPAH (U = 1/2)	1	1	0	102	102	102			102
Total LPAH (U = 1/2)	1	1	0	60.8	60.8	60.8			60.8
Total PAH (U = 1/2)	1	1	0	162	162	162			162
Semivolatile Organic Carbons (μg/kg)									
1,3-Dichlorobenzene	1	0	1				714	714	714
1,4-Dichlorobenzene	1	0	1				714	714	714
Hexachlorobenzene	1	1	0	483	483	483			483
Hexachlorobutadiene	1	0	1				714	714	714
PCBs (µg/kg)	•	•				<u>'</u>			
PCB-001	1	1	0	0.0948	0.0948	0.0948			0.0948
PCB-002	1	0	1				0.00401	0.00401	0.00401
PCB-003	1	1	0	0.0493	0.0493	0.0493			0.0493
PCB-004	1	1	0	0.148	0.148	0.148			0.148
PCB-005	1	0	1				0.0244	0.0244	0.0244
PCB-006	1	1	0	0.18	0.18	0.180			0.180
PCB-007	1	0	1				0.0204	0.0204	0.0204
PCB-008	1	1	0	1.19	1.19	1.19			1.19
PCB-009	1	1	0	0.0657	0.0657	0.0657			0.0657
PCB-010	1	0	1				0.0178	0.0178	0.0178
PCB-011	1	1	0	0.0735	0.0735	0.0735			0.0735
PCB-012/013	1	1	0	0.119	0.119	0.119			0.119
PCB-014	1	0	1				0.0202	0.0202	0.0202
	1 *	. · ·	±		l .		5.0202	5.0202	5.0202

Table 7
White Shrimp Size Class A Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
	·	·	Samples				non Detection	rton Detection	
PCB-015	1	1	0	0.899	0.899	0.899			0.899
PCB-016	1	1	0	0.45	0.45	0.45			0.45
PCB-017	1	1	0	1.68	1.68	1.68			1.68
PCB-018/030	1	1	0	3.61	3.61	3.61			3.61
PCB-019	1	1	0	0.0617	0.0617	0.0617			0.0617
PCB-020/028	1	1	0	40.4	40.4	40.4			40.4
PCB-021/033	1	1	0	6.72	6.72	6.72			6.72
PCB-022	1	1	0	7.36	7.36	7.36			7.36
PCB-023	1	0	1				0.00546	0.00546	0.00546
PCB-024	1	1	0	0.0193	0.0193	0.0193			0.0193
PCB-025	1	1	0	1.15	1.15	1.15			1.15
PCB-026/029	1	1	0	4.36	4.36	4.36			4.36
PCB-027	1	1	0	0.233	0.233	0.233			0.233
PCB-031	1	1	0	37.3	37.3	37.3			37.3
PCB-032	1	1	0	2.19	2.19	2.19			2.19
PCB-034	1	1	0	0.133	0.133	0.133			0.133
PCB-035	1	1	0	0.402	0.402	0.402			0.402
PCB-036	1	1	0	0.0276	0.0276	0.0276			0.0276
PCB-037	1	1	0	4.6	4.6	4.6			4.6
PCB-038	1	0	1				0.0161	0.0161	0.0161
PCB-039	1	0	1				0.0140	0.0140	0.0140
PCB-040/071	1	1	0	5.35	5.35	5.35			5.35
PCB-041	1	1	0	0.597	0.597	0.597			0.597
PCB-042	1	1	0	4.43	4.43	4.43			4.43
PCB-043	1	1	0	0.474	0.474	0.474			0.474
PCB-044/047/065	1	1	0	39.5	39.5	39.5			39.5
PCB-045/051	1	1	0	0.576	0.576	0.576			0.576
PCB-046	1	1	0	0.112	0.112	0.112			0.112
PCB-048	1	1	0	2.18	2.18	2.18			2.18
PCB-049/069	1	1	0	33.1	33.1	33.1			33.1
PCB-050/053	1	1	0	1.07	1.07	1.07			1.07
PCB-052	1	1	0	57.6	57.6	57.6			57.6
PCB-054	1	0	1				0.0100	0.0100	0.0100
PCB-055	1	1	0	0.420	0.420	0.420			0.420
PCB-056	1	1	0	19.1	19.1	19.1			19.1
PCB-057	1	1	0	0.208	0.208	0.208			0.208
PCB-058	1	1	0	0.206	0.206	0.206			0.206
PCB-059/062/075	1	1	0	3.16	3.16	3.16			3.16
PCB-060	1	1	0	13.9	13.9	13.9			13.9
PCB-061/070/074/076	1	1	0	124	124	124			124
PCB-063	1	1	0	2.7	2.7	2.7			2.7
PCB-064	1	1	0	18.5	18.5	18.5			18.5
PCB-066	1	1	0	72.3	72.3	72.3			72.3
PCB-067	1	1	0	1.06	1.06	1.06			1.06
PCB-068	1	1	0	0.364	0.364	0.364			0.364
PCB-072	1	1	0	0.753	0.753	0.753			0.753

Table 7
White Shrimp Size Class A Tissue Sample Summary

		VVIIICE	e Shrimp Size Clas	s A Tissue Sample	Julilliary	Т Т		Г Т	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-073	1	0	1				0.0170	0.0170	0.0170
PCB-077	1	1	0	4.76	4.76	4.76			4.76
PCB-078	1	0	1				0.0515	0.0515	0.0515
PCB-079	1	0	1				0.0430	0.0430	0.0430
PCB-080	1	0	1				0.0462	0.0462	0.0462
PCB-081	1	1	0	0.217	0.217	0.217			0.217
PCB-082	1	1	0	0.905	0.905	0.905			0.905
PCB-083	1	1	0	1.11	1.11	1.11			1.11
PCB-084	1	1	0	0.778	0.778	0.778			0.778
PCB-085/116/117	1	1	0	10.8	10.8	10.8			10.8
PCB-086/087/097/108/119/125	1	1	0	15.9	15.9	15.9			15.9
PCB-088/091	1	1	0	1.82	1.82	1.82			1.82
PCB-089	1	1	0	0.0692	0.0692	0.0692			0.0692
PCB-090/101/113	1	1	0	37.8	37.8	37.8			37.8
PCB-092	1	1	0	4.78	4.78	4.78			4.78
PCB-093/100	1	1	0	0.341	0.341	0.341			0.341
PCB-094	1	1	0	0.0348	0.0348	0.0348			0.0348
PCB-095	1	1	0	4.85	4.85	4.85			4.85
PCB-096	1	1	0	0.0170	0.0170	0.0170			0.0170
PCB-098/102	1	1	0	0.388	0.388	0.388			0.388
PCB-099	1	1	0	25.3	25.3	25.3			25.3
PCB-103	1	1	0	0.249	0.249	0.249			0.249
PCB-104	1	0	1				0.00993	0.00993	0.00993
PCB-105	1	1	0	20.0	20.0	20.0			20.0
PCB-106	1	0	1				0.0219	0.0219	0.0219
PCB-107/124	1	1	0	1.62	1.62	1.62			1.62
PCB-109	1	1	0	2.99	2.99	2.99			2.99
PCB-110/115	1	1	0	24.3	24.3	24.3			24.3
PCB-111	1	1	0	0.0391	0.0391	0.0391			0.0391
PCB-112	1	0	1				0.0133	0.0133	0.0133
PCB-114	1	1	0	1.33	1.33	1.33			1.33
PCB-118	1	1	0	46.2	46.2	46.2			46.2
PCB-120	1	1	0	0.218	0.218	0.218			0.218
PCB-121	1	0	1				0.0131	0.0131	0.0131
PCB-122	1	1	0	0.573	0.573	0.573			0.573
PCB-123	1	1	0	1.12	1.12	1.12			1.12
PCB-126	1	1	0	0.147	0.147	0.147			0.147
PCB-127	1	0	1				0.0219	0.0219	0.0219
PCB-128/166	1	1	0	2.74	2.74	2.74			2.74
PCB-129/138/163	1	1	0	19.2	19.2	19.2			19.2
PCB-130	1	1	0	1.32	1.32	1.32			1.32
PCB-131	1	0	1				0.0212	0.0212	0.0212
PCB-132	1	1	0	0.825	0.825	0.825			0.825
						0.382			0.382
PCR-133	1	l !			II 3× /				
PCB-133 PCB-134	1	1 1	0	0.382 0.383	0.382 0.383	0.383			0.383

Table 7
White Shrimp Size Class A Tissue Sample Summary

White Shrimp Size Class A Tissue Sample Summary Number of Summary											
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average		
PCB-136	1	1	0	0.114	0.114	0.114			0.114		
PCB-137	1	1	0	0.959	0.959	0.959			0.959		
PCB-139/140	1	1	0	0.344	0.344	0.344			0.344		
PCB-141	1	1	0	2.97	2.97	2.97			2.97		
PCB-142	1	0	1				0.0213	0.0213	0.0213		
PCB-143	1	0	1				0.0189	0.0189	0.0189		
PCB-144	1	1	0	0.409	0.409	0.409			0.409		
PCB-145	1	0	1				0.00542	0.00542	0.00542		
PCB-146	1	1	0	3.77	3.77	3.77	==		3.77		
PCB-147/149	1	1	0	6.47	6.47	6.47			6.47		
PCB-148	1	1	0	0.0442	0.0442	0.0442			0.0442		
PCB-150	1	0	1				0.00546	0.00546	0.00546		
PCB-152	1	0	1				0.00519	0.00519	0.00519		
PCB-153/168	1	1	0	14.9	14.9	14.9			14.9		
PCB-154	1	1	0	0.435	0.435	0.435			0.435		
PCB-155	1	1	0	0.0963	0.0963	0.0963			0.0963		
PCB-156/157	1	1	0	2.57	2.57	2.57			2.57		
PCB-158	1	1	0	2.05	2.05	2.05			2.05		
PCB-159	1	0	1				0.0119	0.0119	0.0119		
PCB-160	1	0	1				0.0157	0.0157	0.0157		
PCB-161	1	0	1				0.0151	0.0151	0.0151		
PCB-162	1	1	0	0.172	0.172	0.172			0.172		
PCB-164	1	1	0	1.02	1.02	1.02			1.02		
PCB-165	1	0	1				0.0161	0.0161	0.0161		
PCB-167	1	1	0	0.892	0.892	0.892			0.892		
PCB-169	1	1	0	0.0231	0.0231	0.0231			0.0231		
PCB-170	1	1	0	2.74	2.74	2.74			2.74		
PCB-171/173	1	1	0	0.854	0.854	0.854			0.854		
PCB-172	1	1	0	0.598	0.598	0.598			0.598		
PCB-174	1	1	0	1.5	1.5	1.5			1.5		
PCB-175	1	1	0	0.16	0.16	0.16			0.16		
PCB-176	1	1	0	0.0709	0.0709	0.0709			0.0709		
PCB-177	1	1	0	1.61	1.61	1.61			1.61		
PCB-178	1	1	0	0.779	0.779	0.779			0.779		
PCB-179	1	1	0	0.289	0.289	0.289			0.289		
PCB-180/193	1	1	0	7.19	7.19	7.19			7.19		
PCB-181	1	1	0	0.0526	0.0526	0.0526			0.0526		
PCB-182	1	1	0	0.0428	0.0428	0.0428			0.0428		
PCB-183/185	1	1	0	1.90	1.90	1.90			1.90		
PCB-184	1	1	0	0.0358	0.0358	0.0358			0.0358		
PCB-186	1	0	1				0.00447	0.00447	0.00447		
PCB-187	1	1	0	6.13	6.13	6.13			6.13		
PCB-187	1	1	0	0.0329	0.0329	0.0329		 	0.0329		
		+									
PCB-189	1	1	0	0.138	0.138	0.138			0.138 0.670		
PCB-190	1	1	0	0.670	0.670	0.670					
PCB-191	1	1	0	0.156	0.156	0.156			0.156		

Table 7
White Shrimp Size Class A Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-192	1	0	1				0.00727	0.00727	0.00727
PCB-194	1	1	0	0.919	0.919	0.919			0.919
PCB-195	1	0	1				0.718	0.718	0.718
PCB-196	1	1	0	0.538	0.538	0.538			0.538
PCB-197/200	1	1	0	0.177	0.177	0.177			0.177
PCB-198/199	1	1	0	1.11	1.11	1.11			1.11
PCB-201	1	1	0	0.137	0.137	0.137			0.137
PCB-202	1	1	0	0.545	0.545	0.545			0.545
PCB-203	1	1	0	0.598	0.598	0.598			0.598
PCB-204	1	0	1				0.0115	0.0115	0.0115
PCB-205	1	0	1				0.517	0.517	0.517
PCB-206	1	1	0	0.654	0.654	0.654			0.654
PCB-207	1	1	0	0.440	0.440	0.440			0.440
PCB-208	1	1	0	0.600	0.600	0.600			0.600
PCB-209	1	1	0	17.3	17.3	17.3			17.3
Total PCB Congener (U = 1/2)	1	1	0	837	837	837			837
Dioxins and Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	1	1	0	25.2	25.2	25.2			25.2
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	1	1	0	10.9	10.9	10.9			10.9
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	1	1	0	3.08	3.08	3.08			3.08
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	1	0	1				1.00	1.00	1.00
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	1	0	1				0.705	0.705	0.705
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	1	1	0	1.66	1.66	1.66			1.66
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	1	0	1				0.526	0.526	0.526
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.493	0.493	0.493
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	1	0	1				0.567	0.567	0.567
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.608	0.608	0.608
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	1	0	1				0.556	0.556	0.556
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	1	1	0	4.65	4.65	4.65			4.65
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	1	0	1				0.480	0.480	0.480
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	1	0	1				0.622	0.622	0.622
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	1	1	0	4.07	4.07	4.07			4.07
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	1	1	0	13.7	13.7	13.7			13.7
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1	1	0	3.36	3.36	3.36			3.36
Total Dioxin/Furan (U = 1/2)	1	1	0	69.4	69.4	69.4			69.4

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram -- Results not applicable PCB - polychlorinated biphenyl

Table 8
White Shrimp Size Class B Tissue Sample Summary

White Shrimp Size Class B Tissue Sample Summary											
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average		
Conventional Parameters (percent)											
Lipids	2	2	0	1.13	1.3	1.22			1.22		
Moisture, percent	2	2	0	74.6	75.3	75.0			75.0		
Metals (mg/kg)											
Mercury	2	2	0	0.026	0.042	0.034			0.034		
Semivolatile Organic Carbons (μg/kg)											
1,3-Dichlorobenzene	2	0	2				691	758	362		
Hexachlorobenzene	2	2	0	380	682	531			531		
PCBs (μg/kg)											
PCB-001	2	2	0	0.0270	0.113	0.0700			0.0700		
PCB-002	2	0	2				0.00229	0.00581	0.00203		
PCB-003	2	2	0	0.0169	0.0616	0.0393			0.0393		
PCB-004	2	1	1	0.22	0.22	0.220	0.0354	0.0354	0.119		
PCB-005	2	0	2				0.0126	0.022	0.00865		
PCB-006	2	2	0	0.144	0.246	0.195			0.195		
PCB-007	2	2	0	0.0315	0.0583	0.0449			0.0449		
PCB-008	2	2	0	0.941	1.69	1.32			1.32		
PCB-009	2	2	0	0.0498	0.0894	0.0696			0.0696		
PCB-010	2	0	2				0.0145	0.0186	0.00828		
PCB-011	2	2	0	0.0596	0.0751	0.0674			0.0674		
PCB-012/013	2	2	0	0.0918	0.135	0.113			0.113		
PCB-014	2	0	2				0.0104	0.0182	0.00715		
PCB-015	2	2	0	0.604	1.23	0.917			0.917		
PCB-016	2	2	0	0.241	0.61	0.426			0.426		
PCB-017	2	2	0	0.971	2.28	1.63			1.63		
PCB-018/030	2	2	0	1.93	4.38	3.16			3.16		
PCB-019	2	2	0	0.0343	0.0898	0.0621			0.0621		
PCB-020/028	2	2	0	25.7	49.3	37.5			37.5		
PCB-021/033	2	2	0	4.59	8.02	6.31			6.31		
PCB-022	2	2	0	3.91	9.13	6.52			6.52		
PCB-023	2	0	2				0.00245	0.00629	0.00219		
PCB-024	2	2	0	0.0101	0.0224	0.0163			0.0163		
PCB-025	2	2	0	0.738	1.4	1.07			1.07		
PCB-026/029	2	2	0	2.74	4.67	3.71			3.71		
PCB-027	2	2	0	0.131	0.258	0.195			0.195		
PCB-031	2	2	0	25.6	50.4	38			38		
PCB-032	2	2	0	1.23	2.69	1.96			1.96		
PCB-034	2	2	0	0.0843	0.158	0.121			0.121		
PCB-035	2	2	0	0.244	0.458	0.351			0.351		
PCB-036	2	0	2				0.00977	0.0128	0.00564		
PCB-037	2	2	0	2.59	5.17	3.88			3.88		
PCB-038	2	0	2				0.0108	0.0142	0.00625		
PCB-039	2	0	2				0.00946	0.0124	0.00547		
PCB-040/071	2	2	0	2.72	6.03	4.38			4.38		
PCB-041	2	2	0	0.189	0.542	0.366			0.366		
PCB-042	2	2	0	2.15	5.06	3.61			3.61		
1 00 074			U	2.13	5.00	3.01	· -	_	3.01		

Table 8
White Shrimp Size Class B Tissue Sample Summary

White Shrimp Size Class B Tissue Sample Summary											
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average		
PCB-043	2	2	0	0.196	0.53	0.363			0.363		
PCB-044/047/065	2	2	0	23.3	41.8	32.6			32.6		
PCB-045/051	2	2	0	0.234	0.516	0.375			0.375		
PCB-046	2	2	0	0.0428	0.103	0.0729			0.0729		
PCB-048	2	2	0	0.955	2.71	1.83			1.83		
PCB-049/069	2	2	0	19.8	34.8	27.3			27.3		
PCB-050/053	2	2	0	0.357	0.85	0.604			0.604		
PCB-052	2	2	0	29.8	56.2	43.0			43.0		
PCB-054	2	0	2				0.00627	0.0125	0.00469		
PCB-055	2	1	1	0.32	0.32	0.320	0.0242	0.0242	0.166		
PCB-056	2	2	0	9.61	20.2	14.9			14.9		
PCB-057	2	2	0	0.131	0.248	0.190			0.190		
PCB-058	2	2	0	0.138	0.232	0.185			0.185		
PCB-059/062/075	2	2	0	1.89	3.28	2.59			2.59		
PCB-060	2	2	0	7.9	14.8	11.4			11.4		
PCB-061/070/074/076	2	2	0	81.8	147	114			114		
PCB-063	2	2	0	1.7	3.11	2.41			2.41		
PCB-064	2	2	0	10	19.7	14.9			14.9		
PCB-066	2	2	0	46.7	80.1	63.4			63.4		
PCB-067	2	2	0	0.681	1.2	0.941			0.941		
PCB-068	2	2	0	0.275	0.422	0.349			0.349		
PCB-072	2	2	0	0.525	0.885	0.705			0.705		
PCB-073	2	0	2				0.00557	0.0114	0.00424		
PCB-077	2	2	0	3.1	5.45	4.28			4.28		
PCB-078	2	0	2				0.0163	0.0241	0.0101		
PCB-079	2	0	2				0.0136	0.0201	0.00843		
PCB-080	2	0	2				0.0146	0.0216	0.00905		
PCB-081	2	2	0	0.135	0.233	0.184			0.184		
PCB-082	2	2	0	0.448	0.935	0.692			0.692		
PCB-083	2	2	0	0.515	1.11	0.813			0.813		
PCB-084	2	2	0	0.344	0.727	0.536			0.536		
PCB-085/116/117	2	2	0	7.85	11.8	9.83			9.83		
PCB-086/087/097/108/119/125	2	2	0	10.6	18.1	14.4			14.4		
PCB-088/091	2	2	0	1.12	1.95	1.54			1.54		
PCB-089	2	2	0	0.0225	0.0601	0.0413			0.0413		
PCB-090/101/113	2	2	0	29.8	38.9	34.4			34.4		
PCB-092	2	2	0	2.36	4.55	3.46			3.46		
PCB-093/100	2	2	0	0.272	0.313	0.293			0.293		
PCB-094	2	2	0	0.015	0.0346	0.0248			0.0248		
PCB-095	2	2	0	2.69	4.81	3.75			3.75		
PCB-096	2	2	0	0.00719	0.0134	0.0103			0.0103		
PCB-098/102	2	2	0	0.203	0.437	0.320			0.320		
PCB-099	2	2	0	21	27.3	24.2			24.2		
PCB-103	2	2	0	0.156	0.248	0.202			0.202		
PCB-103 PCB-104	2	0	2	0.150		0.202	0.00419	0.00731	0.00288		
P(B-11)4											

Table 8
White Shrimp Size Class B Tissue Sample Summary

White Shrimp Size Class B Tissue Sample Summary											
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average		
PCB-106	2	0	2				0.0153	0.0411	0.0141		
PCB-107/124	2	2	0	1.26	1.72	1.49			1.49		
PCB-109	2	2	0	2.4	3.46	2.93			2.93		
PCB-110/115	2	2	0	17.6	27.6	22.6			22.6		
PCB-111	2	2	0	0.0284	0.0424	0.0354			0.0354		
PCB-112	2	0	2				0.00559	0.0077	0.00332		
PCB-114	2	2	0	0.978	1.47	1.22			1.22		
PCB-118	2	2	0	32.7	48.8	40.8			40.8		
PCB-120	2	2	0	0.173	0.22	0.197			0.197		
PCB-121	2	0	2				0.0055	0.00757	0.00327		
PCB-122	2	2	0	0.418	0.629	0.524			0.524		
PCB-123	2	2	0	0.758	1.1	0.929			0.929		
PCB-126	2	2	0	0.0696	0.133	0.101			0.101		
PCB-127	2	0	2				0.0153	0.0411	0.0141		
PCB-128/166	2	2	0	1.8	2.7	2.25			2.25		
PCB-129/138/163	2	2	0	14	19.8	16.9			16.9		
PCB-130	2	2	0	1.01	1.45	1.23			1.23		
PCB-131	2	1	1	0.0241	0.0241	0.0241	0.0692	0.0692	0.0294		
PCB-132	2	2	0	0.414	0.885	0.650			0.650		
PCB-133	2	2	0	0.3	0.375	0.338			0.338		
PCB-134	2	2	0	0.159	0.276	0.218			0.218		
PCB-135/151	2	2	0	1.29	2.14	1.72			1.72		
PCB-136	2	2	0	0.0552	0.103	0.0791			0.0791		
PCB-137	2	2	0	0.73	1.03	0.880			0.880		
PCB-139/140	2	2	0	0.229	0.304	0.267			0.267		
PCB-141	2	2	0	2.25	2.98	2.62			2.62		
PCB-142	2	0	2				0.00867	0.0695	0.0195		
PCB-143	2	1	1	0.0120	0.0120	0.0120	0.0619	0.0619	0.0215		
PCB-144	2	2	0	0.228	0.405	0.317			0.317		
PCB-145	2	0	2				0.0025	0.00297	0.00137		
PCB-146	2	2	0	3.01	3.57	3.29			3.29		
PCB-147/149	2	2	0	4.27	6.19	5.23			5.23		
PCB-148	2	2	0	0.0299	0.0429	0.0364			0.0364		
PCB-150	2	2	0	0.00588	0.00814	0.00701			0.00701		
PCB-152	2	0	2				0.00239	0.00284	0.00131		
PCB-153/168	2	2	0	12.4	15.1	13.8			13.8		
PCB-154	2	2	0	0.36	0.419	0.390			0.390		
PCB-155	2	2	0	0.0602	0.0933	0.0768			0.0768		
PCB-156/157	2	2	0	1.84	2.62	2.23			2.23		
PCB-158	2	2	0	1.52	2.1	1.81			1.81		
PCB-159	2	0	2				0.00452	0.0105	0.00376		
PCB-159	2	0	2				0.00432	0.0514	0.0145		
PCB-161	2	0	2				0.00618	0.0495	0.0139		
PCB-161 PCB-162	2	2	0	0.052	0.163	0.108			0.0139		
PCB-162 PCB-164	2	2	0	0.052	1.03	0.108			0.108		
PCB-104 PCB-165	2		2								
rcp-105		0	2				0.00656	0.0526	0.0148		

Table 8
White Shrimp Size Class B Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-167	2	2	0	0.676	0.908	0.792			0.792
PCB-169	2	2	0	0.0151	0.0177	0.0164			0.0164
PCB-170	2	2	0	1.93	2.45	2.19			2.19
PCB-171/173	2	2	0	0.615	0.743	0.679			0.679
PCB-172	2	2	0	0.51	0.582	0.546			0.546
PCB-174	2	2	0	0.984	1.39	1.19			1.19
PCB-175	2	2	0	0.11	0.131	0.121			0.121
PCB-176	2	2	0	0.0358	0.0621	0.0490			0.0490
PCB-177	2	2	0	1.13	1.45	1.29			1.29
PCB-178	2	2	0	0.581	0.673	0.627			0.627
PCB-179	2	2	0	0.178	0.214	0.196			0.196
PCB-180/193	2	2	0	5.67	6.7	6.19			6.19
PCB-181	2	2	0	0.0401	0.0427	0.0414			0.0414
PCB-182	2	2	0	0.0239	0.0288	0.0264			0.0264
PCB-183/185	2	2	0	1.62	1.88	1.75			1.75
PCB-184	2	2	0	0.0321	0.0326	0.0324			0.0324
PCB-186	2	0	2				0.00227	0.00396	0.00156
PCB-187	2	2	0	3.89	6.13	5.01			5.01
PCB-188	2	2	0	0.0222	0.0308	0.0265			0.0265
PCB-189	2	2	0	0.093	0.13	0.112			0.112
PCB-190	2	2	0	0.512	0.635	0.574			0.574
PCB-191	2	2	0	0.116	0.136	0.126			0.126
PCB-192	2	1	1	0.00600	0.00600	0.00600	0.00443	0.00443	0.00411
PCB-194	2	2	0	0.707	1.02	0.864			0.864
PCB-195	2	2	0	0.227	0.352	0.290			0.290
PCB-196	2	2	0	0.362	0.53	0.446			0.446
PCB-197/200	2	2	0	0.13	0.173	0.152			0.152
PCB-198/199	2	2	0	0.994	1.23	1.11			1.11
PCB-201	2	2	0	0.0963	0.134	0.115			0.115
PCB-202	2	2	0	0.526	0.614	0.570			0.570
PCB-203	2	2	0	0.55	0.704	0.627			0.627
PCB-204	2	2	0	0.0115	0.0182	0.0149			0.0149
PCB-205	2	2	0	0.0388	0.0554	0.0471			0.0471
PCB-206	2	2	0	0.473	0.623	0.548			0.548
PCB-207	2	2	0	0.276	0.406	0.341			0.341
PCB-208	2	2	0	0.46	0.607	0.534			0.534
PCB-209	2	2	0	13	16.9	15.0			15.0
Total PCB Congener (U = 1/2)	2	2	0	548	919	733			733

mg/kg - milligrams per kilogram

μg/kg - micrograms per kilogram

-- Results not applicable

PCB - polychlorinated biphenyl

Table 9
Gulf Killifish Tissue Sample Summary

		1	ulf Killifish Tissue		T	1		 	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)									
Lipids	25	25	0	1.65	3.64	2.72			2.72
Moisture, percent	25	25	0	71.9	75.9	74.2			74.2
Metals (mg/kg)									
Mercury	25	25	0	0.025	0.109	0.0524			0.0524
Semivolatile Organic Carbons (μg/kg)									
1,3-Dichlorobenzene	25	0	25				351	763	283
Hexachlorobenzene	25	10	15	200	2,360.00	664	351	763	441
PCBs (μg/kg)									
PCB-001	25	25	0	0.104	0.462	0.276			0.276
PCB-002	25	16	9	0.00632	0.0292	0.0134	0.00448	0.0152	0.0101
PCB-003	25	25	0	0.0217	0.113	0.0549			0.0549
PCB-004	25	25	0	1.17	7.72	4.12			4.12
PCB-005	25	20	5	0.0332	0.226	0.114	0.0343	0.0843	0.0962
PCB-006	25	25	0	0.512	3.33	1.83			1.83
PCB-007	25	25	0	0.0619	0.483	0.237			0.237
PCB-008	25	25	0	3.08	19.1	11.2			11.2
PCB-009	25	25	0	0.162	1.2	0.619			0.619
PCB-010	25	24	1	0.0309	0.217	0.121	0.0356	0.0356	0.117
PCB-011	25	20	5	0.0808	1.18	0.220	0.0367	0.171	0.185
PCB-012/013	25	25	0	0.0471	0.379	0.160			0.160
PCB-014	25	0	25				0.00387	0.0704	0.00878
PCB-015	25	25	0	0.538	2.31	1.45			1.45
PCB-016	25	25	0	3.38	21.2	12.7			12.7
PCB-017	25	25	0	10.3	52	32.7			32.7
PCB-018/030	25	25	0	17.8	112	71.5			71.5
PCB-019	25	25	0	1.02	4.6	3.10			3.10
PCB-020/028	25	25	0	71.9	441	231			231
PCB-021/033	25	25	0	6.78	45.3	25.7			25.7
PCB-022	25	25	0	11.9	79.7	41.0			41.0
PCB-023	25	25	0	0.0229	0.168	0.0915			0.0915
PCB-023	25	24	1	0.137	0.694	0.0913	0.00411	0.00411	0.427
PCB-025	25	25	0	2.06	12.1	7.07			7.07
PCB-025 PCB-026/029	25	25	0	7.41	44.9	25.0			25.0
PCB-026/029 PCB-027	25	25	0	1.05	6.1	3.73			3.73
PCB-027 PCB-031	25	25	0	66	366	196			196
PCB-031 PCB-032	25	25	0	10.6	54.7	31.5			31.5
PCB-032 PCB-034	25	25	0	0.205	1.09	0.634			0.634
		7		ł — — — — — — — — — — — — — — — — — — —	 	+	0.0106	1.06	
PCB-035	25		18	0.0124	0.0472	0.0252	0.0106	1.06	0.0481
PCB-036	25	0	25		7.64	 	0.00331	1.07	0.042
PCB-037	25	25	0	2.12	7.64	5.16		1.00	5.16
PCB-038	25	20	5	0.0799	0.497	0.275	0.00719	1.08	0.252
PCB-039	25	8	17	0.493	2.45	1.04	0.00314	1.05	0.371
PCB-040/071	25	24	1	32.9	165	99.2	0.127	0.127	95.2
PCB-041	25	24	1	3.84	24.7	13.4	0.166	0.166	12.9
PCB-042	25	24	1	16.1	89.1	48.9	0.14	0.14	47.0

Table 9
Gulf Killifish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Sample Summary Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-043	25	17	8	2.8	21.1	10.9	0.0178	0.463	7.44
PCB-044/047/065	25	24	1	103	457	271	0.124	0.124	260
PCB-045/051	25	25	0	7.57	44.2	26.5			26.5
PCB-046	25	25	0	0.855	5.23	2.94			2.94
PCB-048	25	24	1	11.1	82.8	42.1	0.137	0.137	40.4
PCB-049/069	25	24	1	109	440	260	0.118	0.118	250
PCB-050/053	25	25	0	8.75	46.8	27.1			27.1
PCB-052	25	24	1	182	648	417	0.14	0.14	400
PCB-054	25	25	0	0.0717	0.32	0.217			0.217
PCB-055	25	19	6	0.495	4.69	1.93	0.0161	0.149	1.47
PCB-056	25	25	0	21	126	63.8			63.8
PCB-057	25	24	1	0.383	2.6	1.26	0.868	0.868	1.23
PCB-058	25	24	1	0.277	5.35	0.917	0.932	0.932	0.899
PCB-059/062/075	25	24	1	8.46	40.3	23.1	0.106	0.106	22.2
PCB-060	25	25	0	31.5	174	93.9			93.9
PCB-061/070/074/076	25	25	0	227	1190	636			636
PCB-063	25	25	0	6.32	38.8	19.4			19.4
PCB-064	25	24	1	55.9	262	150	0.0991	0.0991	144
PCB-066	25	25	0	175	805	438			438
PCB-067	25	25	0	1.17	7.23	3.69			3.69
PCB-068	25	25	0	0.612	3.02	1.59			1.59
PCB-072	25	25	0	1.31	6.4	3.40			3.40
PCB-073	25	8	17	0.562	14.8	9.13	0.00564	0.111	2.93
PCB-077	25	25	0	4.4	19.1	12.9			12.9
PCB-078	25	0	25				0.0148	0.91	0.0404
PCB-079	25	9	16	0.541	2.82	1.92	0.0128	0.766	0.718
PCB-080	25	0	25				0.0141	0.798	0.0363
PCB-081	25	24	1	0.257	1.08	0.700	1.08	1.08	0.694
PCB-082	25	25	0	2.04	14.1	7.97			7.97
PCB-083	25	21	4	1.98	10.1	6.31	0.0189	0.0604	5.31
PCB-084	25	25	0	4.03	19.8	12.6			12.6
PCB-085/116/117	25	25	0	22.6	106	65.0			65.0
PCB-086/087/097/108/119/125	25	24	1	38.2	192	119	0.0185	0.0185	115
PCB-088/091	25	25	0	10.4	45.5	29.1			29.1
PCB-089	25	25	0	0.568	3.39	2.04			2.04
PCB-090/101/113	25	25	0	78.7	339	212			212
PCB-092	25	25	0	14.2	64.1	38.8			38.8
PCB-093/100	25	25	0	1.06	4.63	2.68			2.68
PCB-094	25	25	0	0.31	1.6	0.961			0.961
PCB-095	25	25	0	33.1	131	90.3			90.3
PCB-096	25	25	0	0.213	1.16	0.671			0.671
PCB-098/102	25	25	0	2.13	10.9	6.75			6.75
PCB-099	25	25	0	63.2	229	151			151
PCB-103	25	25	0	0.825	2.84	1.85			1.85
PCB-104	25	23	2	0.0148	0.0794	0.0395	0.0203	0.0238	0.0372
PCB-105	25	25	0	34.1	143	98.6			98.6

Table 9
Gulf Killifish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-106	25	2	23	0.0621	0.101	0.0816	0.00611	0.0973	0.0224
PCB-107/124	25	25	0	1.98	12.2	6.88			6.88
PCB-109	25	25	0	6.62	31.6	19.0			19.0
PCB-110/115	25	25	0	54.8	257	162			162
PCB-111	25	25	0	0.0509	0.193	0.120			0.120
PCB-112	25	6	19	0.617	1.42	0.904	0.00415	0.0788	0.224
PCB-114	25	25	0	2.75	12.2	7.84			7.84
PCB-118	25	25	0	78.5	282	204			204
PCB-120	25	25	0	0.276	1.24	0.710			0.710
PCB-121	25	15	10	0.0225	0.0635	0.0410	0.00408	0.08	0.0294
PCB-122	25	25	0	0.294	1.59	0.910			0.910
PCB-123	25	25	0	1.82	8.84	5.59			5.59
PCB-126	25	24	1	0.2	1.22	0.675	0.0473	0.0473	0.649
PCB-127	25	18	7	0.0852	0.397	0.267	0.00617	0.0504	0.196
PCB-128/166	25	25	0	5.19	23.1	15.3			15.3
PCB-129/138/163	25	25	0	42.8	162	106			106
PCB-130	25	25	0	2.57	10.7	6.82			6.82
PCB-131	25	24	1	0.0815	0.51	0.281	0.0243	0.0243	0.270
PCB-132	25	25	0	4.96	24	14.2			14.2
PCB-133	25	25	0	0.851	3.06	1.84			1.84
PCB-134	25	24	1	0.629	2.87	1.85	0.0199	0.0199	1.78
PCB-135/151	25	25	0	8.96	33.8	20.6			20.6
PCB-136	25	25	0	1.27	5.12	3.23			3.23
PCB-137	25	25	0	2.63	11.5	7.31			7.31
PCB-139/140	25	25	0	0.93	4.2	2.41			2.41
PCB-141	25	25	0	5.29	24.3	15.3			15.3
PCB-142	25	3	22	0.0135	0.28	0.103	0.00762	0.066	0.0238
PCB-143	25	17	8	0.0122	0.277	0.175	0.0146	0.0613	0.124
PCB-144	25	25	0	1.52	6.22	3.76			3.76
PCB-145	25	24	1	0.00924	0.0366	0.0230	0.0124	0.0124	0.0223
PCB-146	25	24	1	0.0604	23.2	13.4	0.0327	0.0327	12.8
PCB-147/149	25	25	0	18.4	77.3	44.4			44.4
PCB-148	25	25	0	0.083	0.309	0.177			0.177
PCB-150	25	25	0	0.0565	0.193	0.115			0.115
PCB-152	25	25	0	0.0303	0.0991	0.0591			0.0591
PCB-152 PCB-153/168	25	25	0	43.4	161	96.9			96.9
PCB-154	25	25	0	0.799	3.01	1.68			1.68
PCB-155	25	25	0	0.0527	0.224	0.135			0.135
PCB-156/157	25	25	0	4.76	18.8	12.7			12.7
PCB-158	25	25	0	4.38	17.4	11.3		 	11.3
PCB-158	25	9	16	0.16	0.561	0.386	0.00571	0.0492	0.145
PCB-159		0	25			+	0.00571	0.0492	0.145
	25					10.1		+	
PCB-161	25	5	20	14.2	23.8	18.1	0.00495	0.0469	3.62
PCB-162	25	25	0	0.224	1.07	0.599			0.599
PCB-164 PCB-165	25 25	25 23	0 2	1.9 0.0222	7.53 0.101	4.84 0.0551	0.0306	0.0506	4.84 1.31

Table 9
Gulf Killifish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Sample Summary Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-167	25	25	0	1.96	7.39	4.93			4.93
PCB-169	25	8	17	0.024	0.222	0.0856	0.00852	0.0853	0.0345
PCB-170	25	25	0	6.02	22	15.3			15.3
PCB-171/173	25	25	0	2.02	7.17	4.60			4.60
PCB-172	25	25	0	1.19	4.38	2.84			2.84
PCB-174	25	25	0	3.1	13.6	8.08			8.08
PCB-175	25	25	0	0.369	1.37	0.840			0.840
PCB-176	25	25	0	0.463	2.02	1.09			1.09
PCB-177	25	25	0	3.05	10.9	6.75			6.75
PCB-178	25	25	0	1.71	5.71	3.63			3.63
PCB-179	25	25	0	1.51	5.61	3.26			3.26
PCB-180/193	25	25	0	16	58	38.6			38.6
PCB-181	25	25	0	0.102	0.429	0.255			0.255
PCB-182	25	24	1	0.0501	0.266	0.151	0.00431	0.00431	3.63
PCB-183/185	25	25	0	5.21	20.9	12.3			12.3
PCB-184	25	25	0	0.0646	0.203	0.122			0.122
PCB-186	25	8	17	0.0031	0.00649	0.00465	0.0017	0.0123	0.0034
PCB-187	25	25	0	8.82	33.1	22.0			22.0
PCB-188	25	25	0	0.0349	0.146	0.0889			0.0889
PCB-189	25	25	0	0.212	0.78	0.500			0.500
PCB-190	25	25	0	1.01	3.72	2.50			2.50
PCB-191	25	25	0	0.297	1.14	0.755			0.755
PCB-192	25	7	18	0.00536	0.0155	0.00846	0.00256	0.0246	0.00629
PCB-194	25	25	0	2	8.7	5.78			5.78
PCB-195	25	25	0	0.943	3.89	2.51			2.51
PCB-196	25	25	0	1.55	7.13	4.30			4.30
PCB-197/200	25	25	0	0.503	2.25	1.16			1.16
PCB-198/199	25	25	0	3.19	15.1	8.67			8.67
PCB-201	25	25	0	0.604	2.65	1.42			1.42
PCB-202	25	25	0	1.03	3.86	2.50			2.50
PCB-203	25	25	0	1.48	7.68	4.36			4.36
PCB-204	25	25	0	0.025	0.18	0.0789			0.0789
PCB-205	25	24	1	0.0964	0.345	0.225	0.32	0.32	0.222
PCB-206	25	25	0	1.26	4.51	2.99			2.99
PCB-207	25	25	0	1.43	6.45	3.55			3.55
PCB-208	25	25	0	1.63	7.08	4.29			4.29
PCB-209	25	25	0	33.4	172	83.0			83.0
Total PCB Congener (U = 1/2)	25	25	0	1,965	8,848	5,216			5,216
Dioxins and Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	25	18	7	0.467	24.7	4.38	0.204	0.49	3.20
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	25	25	0	1.93	48.2	10.0			10.0
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	25	18	7	0.291	3.59	1.06	0.107	0.42	0.789
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	25	21	4	0.401	5.55	1.01	0.252	4.71	1.05
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	25	0	25				0.142	1.14	0.174
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	25	15	10	0.147	0.789	0.339	0.095	0.534	0.246
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	25	1	24	0.348	0.348	0.348	0.0871	3.05	0.256

Table 9
Gulf Killifish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	25	13	12	0.12	0.529	0.277	0.0981	0.529	0.195
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	25	8	17	0.204	0.569	0.314	0.152	3.06	0.331
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	25	0	25		-		0.11	0.76	0.132
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	25	1	24	0.185	0.185	0.185	0.0913	3.16	0.262
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	25	25	0	0.559	2.88	1.17			1.17
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	25	19	6	0.425	2.37	0.976	0.116	1.22	0.810
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	25	1	24	0.118	0.118	0.118	0.112	0.666	0.122
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	25	25	0	1.31	4.55	2.43			2.43
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	25	25	0	6.08	15.6	9.96			9.96
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	25	25	0	2.85	8.53	5.00			5.00
Total Dioxin/Furan (U = 1/2)	25	25	0	19.1	98.6	36.2			36.2

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram -- Results not applicable

Table 10
Gulf Menhaden Tissue Sample Summary

Guit Mennagen Tissue Sample Summary									
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)									
Lipids	10	10	0	1.23	2.02	1.67			1.67
Moisture, percent	10	10	0	77.5	80.0	78.8			78.8
Metals (mg/kg)									
Mercury	10	10	0	0.068	0.199	0.116			0.116
Semivolatile Organic Carbons (µg/kg)	•	•							
1,3-Dichlorobenzene	10	0	10				341	765	267
Hexachlorobenzene	10	6	4	398	671	491	699	765	440
PCBs (μg/kg)	•	•							
PCB-001	10	10	0	0.0902	0.283	0.169			0.169
PCB-002	10	7	3	0.0091	0.0134	0.0116	0.00556	0.00814	0.00919
PCB-003	10	10	0	0.0301	0.0582	0.0419			0.0419
PCB-004	10	10	0	0.907	3.33	2.11			2.11
PCB-005	10	7	3	0.0357	0.111	0.0709	0.0165	0.0287	0.0532
PCB-006	10	10	0	0.508	1.62	1.16			1.16
PCB-007	10	10	0	0.0865	0.256	0.175			0.175
PCB-008	10	10	0	2.1	9.42	5.47			5.47
PCB-009	10	10	0	0.18	0.562	0.414			0.414
PCB-010	10	10	0	0.0265	0.114	0.0689			0.0689
PCB-011	10	2	8	0.0495	0.0703	0.0599	0.0232	0.149	0.0407
PCB-012/013	10	9	1	0.0763	0.245	0.145	0.0234	0.0234	0.132
PCB-014	10	0	10				0.00423	0.025	0.00648
PCB-015	10	10	0	0.215	1.56	0.690			0.690
PCB-016	10	10	0	2.59	20.8	9.64			9.64
PCB-017	10	10	0	7.35	36.5	20.3			20.3
PCB-018/030	10	10	0	12.7	92.7	46.9			46.9
PCB-019	10	10	0	0.923	4.62	2.50			2.50
PCB-020/028	10	10	0	47.6	149	110			110
PCB-021/033	10	10	0	13.6	51	32.6			32.6
PCB-022	10	10	0	9.63	41	25.7			25.7
PCB-023	10	10	0	0.0201	0.0722	0.0504			0.0504
PCB-024	10	10	0	0.0978	0.584	0.281			0.281
PCB-025	10	10	0	1.75	6.55	4.40			4.40
PCB-026/029	10	10	0	5.56	19.7	13.6			13.6
PCB-027	10	10	0	0.737	4.44	2.34			2.34
PCB-031	10	10	0	44.6	149	107			107
PCB-032	10	10	0	5.66	29.8	17.0			17.0
PCB-034	10	10	0	0.305	0.567	0.486			0.486
PCB-035	10	7	3	0.0388	0.125	0.0821	0.0184	0.0551	0.0624
PCB-036	10	0	10				0.00814	0.0533	0.0108
PCB-037	10	10	0	1.66	8.36	4.40			4.40
PCB-038	10	10	0	0.0677	0.189	0.126			0.126
PCB-039	10	3	7	0.544	1.19	0.968	0.0152	0.0505	0.299

Table 10
Gulf Menhaden Tissue Sample Summary

				i Tissue Sample St	, '				
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-040/071	10	8	2	23.3	104	61.0	0.01	0.0136	48.8
PCB-041	10	8	2	3.62	23.5	11.9	0.0118	0.017	9.48
PCB-042	10	8	2	14.8	63.6	37.7	0.0106	0.0147	30.2
PCB-043	10	3	7	1.99	9.76	7.15	0.0117	0.0523	2.16
PCB-044/047/065	10	8	2	58	233	142	0.0095	0.0129	113
PCB-045/051	10	10	0	6.04	41.8	19.5			19.5
PCB-046	10	10	0	1.24	11.3	4.68			4.68
PCB-048	10	8	2	13.3	55.5	34.5	0.0104	0.0141	27.6
PCB-049/069	10	8	2	51.8	175	117	0.00903	0.0122	93.3
PCB-050/053	10	10	0	5.69	40.2	18.7			18.7
PCB-052	10	8	2	78.9	312	190	0.0102	0.0143	152
PCB-054	10	10	0	0.0813	0.311	0.188			0.188
PCB-055	10	8	2	0.487	2.3	1.64	0.03	0.031	1.32
PCB-056	10	10	0	24	81.2	60.6			60.6
PCB-057	10	10	0	0.274	0.942	0.672			0.672
PCB-058	10	8	2	0.253	0.562	0.407	0.0145	0.0167	0.327
PCB-059/062/075	10	8	2	4.54	17.4	10.7	0.00812	0.0112	8.54
PCB-060	10	10	0	15.1	53.4	39.5			39.5
PCB-061/070/074/076	10	10	0	137	410	315			315
PCB-063	10	10	0	3.21	11	7.80			7.80
PCB-064	10	8	2	29.2	120	73.4	0.00757	0.0106	58.7
PCB-066	10	10	0	84.6	245	183			183
PCB-067	10	10	0	1.34	4.76	3.48			3.48
PCB-068	10	9	1	0.505	0.867	0.717	0.0213	0.0213	0.647
PCB-072	10	10	0	1.03	1.9	1.60			1.60
PCB-073	10	5	5	2.01	4.75	3.40	0.00826	0.0137	1.70
PCB-077	10	10	0	3.39	12.4	7.50			7.50
PCB-078	10	0	10				0.0134	0.0524	0.0133
PCB-079	10	3	7	0.628	1.15	0.959	0.0202	0.0448	0.297
PCB-080	10	0	10				0.0121	0.0477	0.0122
PCB-081	10	10	0	0.147	1.22	0.428			0.428
PCB-082	10	10	0	4.07	21.8	12.0			12.0
PCB-083	10	7	3	1.39	9.52	5.01	0.00989	0.0428	3.51
PCB-084	10	10	0	5.86	42.8	19.2			19.2
PCB-085/116/117	10	10	0	10.7	41.4	26.6			26.6
PCB-086/087/097/108/119/125	10	10	0	26.2	109	67.4			67.4
PCB-088/091	10	10	0	6.71	33.1	18.0			18.0
PCB-089	10	10	0	0.571	4.12	1.94			1.94
PCB-090/101/113	10	10	0	41.1	142	93.3			93.3
PCB-092	10	10	0	6.9	25.5	16.2			16.2
PCB-093/100	10	10	0	0.55	2.46	1.40			1.40
PCB-094	10	10	0	0.253	1.59	0.794			0.794
PCB-095	10	10	0	19	109	55.0			55.0

Table 10
Gulf Menhaden Tissue Sample Summary

				i i issue sample si	, ,				
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-096	10	10	0	0.236	2.29	0.912			0.912
PCB-098/102	10	10	0	1.5	8.1	4.21			4.21
PCB-099	10	10	0	28.7	87.8	60.4			60.4
PCB-103	10	10	0	0.504	1.62	1.01			1.01
PCB-104	10	9	1	0.0147	0.0511	0.0303	0.0102	0.0102	0.0278
PCB-105	10	10	0	16.9	57.5	38.7			38.7
PCB-106	10	1	9	0.0326	0.0326	0.0326	0.0155	0.0441	0.014
PCB-107/124	10	10	0	1.56	4.73	3.67			3.67
PCB-109	10	10	0	3.42	9.68	7.55			7.55
PCB-110/115	10	10	0	36.6	139	87.3			87.3
PCB-111	10	10	0	0.0405	0.0754	0.0615			0.0615
PCB-112	10	2	8	0.0936	0.473	0.283	0.00591	0.0289	0.0629
PCB-114	10	10	0	1.15	4.12	2.73			2.73
PCB-118	10	10	0	37.8	112	80.1			80.1
PCB-120	10	10	0	0.26	0.466	0.359			0.359
PCB-121	10	9	1	0.0164	0.0397	0.0263	0.0282	0.0282	0.0251
PCB-122	10	10	0	0.569	1.75	1.30			1.30
PCB-123	10	10	0	0.866	2.71	1.98			1.98
PCB-126	10	10	0	0.104	0.392	0.261			0.261
PCB-127	10	9	1	0.0518	0.136	0.103	0.0426	0.0426	0.0952
PCB-128/166	10	10	0	3.59	9.39	7.15			7.15
PCB-129/138/163	10	10	0	25.7	64	49.1			49.1
PCB-130	10	10	0	1.87	4.3	3.42			3.42
PCB-131	10	10	0	0.238	0.981	0.589			0.589
PCB-132	10	10	0	5.01	19.4	11.9			11.9
PCB-133	10	10	0	0.55	1.11	0.902			0.902
PCB-134	10	10	0	0.8	3.68	2.04			2.04
PCB-135/151	10	10	0	6.18	18.1	11.9			11.9
PCB-136	10	10	0	1.64	7.15	3.79			3.79
PCB-137	10	10	0	1.29	3.74	2.82			2.82
PCB-139/140	10	10	0	0.514	1.5	1.09			1.09
PCB-141	10	10	0	3.7	10.6	7.78			7.78
PCB-142	10	8	2	0.0113	0.0349	0.0267	0.0185	0.0329	0.0240
PCB-143	10	2	8	0.0458	0.109	0.0774	0.00914	0.0322	0.0231
PCB-144	10	10	0	0.919	2.82	1.87			1.87
PCB-145	10	10	0	0.00599	0.0281	0.0131			0.0131
PCB-146	10	9	1	0.0239	10.2	3.85	0.027	0.027	3.46
PCB-147/149	10	10	0	15.2	46.3	30.8			30.8
PCB-148	10	10	0	0.0797	0.134	0.109			0.109
PCB-150	10	10	0	0.0763	0.177	0.120			0.120
PCB-152	10	10	0	0.0162	0.0856	0.0441			0.0441
PCB-153/168	10	10	0	25.4	56.5	44.0			44.0
PCB-154	10	10	0	0.674	1.2	0.967			0.967

Table 10
Gulf Menhaden Tissue Sample Summary

				i rissue sample su					
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-155	10	10	0	0.0745	0.13	0.103			0.103
PCB-156/157	10	10	0	2.8	7.11	5.37			5.37
PCB-158	10	10	0	2.44	6.38	4.90			4.90
PCB-159	10	3	7	0.206	0.323	0.276	0.00722	0.0235	0.0878
PCB-160	10	0	10				0.00663	0.0229	0.00674
PCB-161	10	6	4	5.02	8.68	6.84	0.00607	0.0172	4.11
PCB-162	10	10	0	0.106	0.314	0.224			0.224
PCB-164	10	10	0	1.66	3.77	2.99			2.99
PCB-165	10	10	0	0.018	0.0633	0.0421			0.0421
PCB-167	10	10	0	1.09	2.64	1.99			1.99
PCB-169	10	2	8	0.00912	0.0519	0.0305	0.00581	0.016	0.0111
PCB-170	10	10	0	4.93	11.4	8.08			8.08
PCB-171/173	10	10	0	1.61	3.64	2.66			2.66
PCB-172	10	10	0	1.02	2.34	1.73			1.73
PCB-174	10	10	0	4.03	9.01	7.08			7.08
PCB-175	10	10	0	0.318	0.596	0.465			0.465
PCB-176	10	10	0	0.623	1.26	1.01			1.01
PCB-177	10	10	0	2.82	6.32	4.64			4.64
PCB-178	10	10	0	1.43	2.69	2.09			2.09
PCB-179	10	10	0	1.78	3.97	3.01			3.01
PCB-180/193	10	10	0	12.5	28.8	20.6			20.6
PCB-181	10	10	0	0.0768	0.18	0.137			0.137
PCB-182	10	10	0	0.0489	0.117	0.0849			0.0849
PCB-183/185	10	10	0	4.17	9.13	6.81			6.81
PCB-184	10	10	0	0.061	0.096	0.0806			0.0806
PCB-186	10	3	7	0.00257	0.00353	0.00301	0.00255	0.0062	0.00229
PCB-187	10	10	0	8.39	18.3	12.5			12.5
PCB-188	10	10	0	0.0456	0.0788	0.0595			0.0595
PCB-189	10	10	0	0.215	0.423	0.327			0.327
PCB-190	10	10	0	1.03	2.33	1.64			1.64
PCB-191	10	10	0	0.253	0.538	0.390			0.390
PCB-192	10	7	3	0.0069	0.0147	0.0106	0.0082	0.0124	0.00888
PCB-194	10	10	0	2.74	6.01	4.28			4.28
PCB-195	10	10	0	0.999	2.07	1.51			1.51
PCB-196	10	10	0	1.55	2.94	2.27			2.27
PCB-197/200	10	10	0	0.531	1.1	0.875			0.875
PCB-198/199	10	10	0	3.55	6.69	5.17			5.17
PCB-201	10	10	0	0.568	0.922	0.765			0.765
PCB-202	10	10	0	1.31	2.05	1.61			1.61
PCB-203	10	10	0	1.97	3.73	2.89			2.89
PCB-204	10	10	0	0.0397	0.0668	0.0509			0.0509
PCB-205	10	10	0	0.21	0.357	0.274			0.274
PCB-206	10	10	0	2.79	4.68	3.67			3.67

Table 10
Gulf Menhaden Tissue Sample Summary

duit Methiaden Tissue Sample Summary										
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average	
PCB-207	10	10	0	1.92	3.27	2.68			2.68	
PCB-208	10	10	0	2.64	4.54	3.67			3.67	
PCB-209	10	10	0	71.8	152	110			110	
Total PCB Congener (U = 1/2)	10	10	0	1,259	4,005	2,642			2,642	
Dioxins and Furans (ng/kg)										
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	10	10	0	57.5	1500	512			512	
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	10	10	0	57.7	293	165			165	
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	10	10	0	4.46	54.8	21.8			21.8	
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	10	10	0	6.42	30.6	17.9			17.9	
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	10	10	0	0.697	13.4	4.32			4.32	
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	10	10	0	1.58	6.41	3.40			3.40	
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	10	2	8	0.343	0.362	0.353	0.189	2.83	0.419	
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	10	10	0	0.712	2.94	1.70			1.70	
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	10	8	2	0.547	1.59	1.05	2.35	2.76	1.10	
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	10	3	7	1.01	1.39	1.15	0.3	0.679	0.507	
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	10	3	7	0.733	0.981	0.863	0.198	2.89	0.602	
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	10	10	0	3.11	5.86	4.56			4.56	
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	10	8	2	0.673	1.08	0.855	0.618	0.687	0.749	
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	10	9	1	0.552	2.2	1.25	0.671	0.671	1.16	
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	10	10	0	3.1	6.39	4.91			4.91	
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	10	10	0	9.79	17	13.8			13.8	
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	10	10	0	5.86	11	8.66			8.66	
Total Dioxin/Furan (U = 1/2)	10	10	0	154	1,946	763			763	

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram -- Results not applicable

Table 11
Pinfish Tissue Sample Summary

	T	1	misii rissue samp	· · · · · ·	1	1		1	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)		.							
Lipids	4	4	0	2.34	3.93	2.86			2.86
Moisture, percent	4	4	0	73.2	74.9	73.9			73.9
Metals (mg/kg)									
Mercury	4	4	0	0.025	0.083	0.0430			0.0430
Semivolatile Organic Carbons (µg/kg)									
1,3-Dichlorobenzene	4	0	4				671	765	359
Hexachlorobenzene	4	3	1	397	2,290	1,045	708	708	872
PCBs (μg/kg)									
PCB-001	4	4	0	0.106	0.441	0.204			0.204
PCB-002	4	0	4				0.0013	0.00938	0.00264
PCB-003	4	3	1	0.011	0.0195	0.0164	0.00824	0.00824	0.0133
PCB-004	4	4	0	1.12	5.21	2.19			2.19
PCB-005	4	2	2	0.0204	0.086	0.0532	0.0265	0.0358	0.0344
PCB-006	4	4	0	0.345	1.5	0.661			0.661
PCB-007	4	3	1	0.044	0.164	0.0862	0.0311	0.0311	0.0685
PCB-008	4	4	0	1.36	5.27	2.38			2.38
PCB-009	4	4	0	0.237	0.808	0.423			0.423
PCB-010	4	4	0	0.0317	0.116	0.0560			0.0560
PCB-011	4	0	4				0.0243	0.111	0.0246
PCB-012/013	4	3	1	0.0238	0.0551	0.0392	0.0318	0.0318	0.0334
PCB-014	4	0	4				0.0033	0.0308	0.00746
PCB-015	4	4	0	0.113	0.496	0.224			0.224
PCB-016	4	4	0	1.64	8.81	3.59			3.59
PCB-017	4	4	0	4.27	18.5	7.92			7.92
PCB-018/030	4	4	0	16.4	71.1	32.8			32.8
PCB-019	4	4	0	0.64	3.11	1.36			1.36
PCB-020/028	4	4	0	92.6	259	148			148
PCB-021/033	4	4	0	2.33	9.76	4.49			4.49
PCB-022	4	4	0	7.32	22.5	11.8			11.8
PCB-023	4	4	0	0.0245	0.0884	0.0411			0.0411
PCB-024	4	4	0	0.161	0.687	0.326			0.326
PCB-025	4	4	0	1.12	3.93	1.91			1.91
PCB-026/029	4	4	0	9.41	29.7	15.7			15.7
PCB-027	4	4	0	0.388	1.77	0.752			0.752
PCB-031	4	4	0	81.2	244	137			137
PCB-032	4	4	0	1.78	8.83	3.70			3.70
PCB-034	4	4	0	0.144	0.511	0.254			0.254
PCB-035	4	1	3	0.0103	0.0103	0.0103	0.0281	0.173	0.0340
PCB-036	4	0	4				0.00415	0.166	0.0305
PCB-037	4	4	0	0.903	1.94	1.34			1.34
PCB-038	4	3	1	0.133	0.205	0.160	0.169	0.169	0.141
PCB-039	4	1	3	0.148	0.148	0.148	0.0248	0.159	0.0656

Table 11
Pinfish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-040/071	4	4	0	7.43	29.2	14.4			14.4
PCB-041	4	3	1	0.786	0.852	0.812	6.55	6.55	1.43
PCB-042	4	4	0	13.8	50.1	26.7			26.7
PCB-043	4	2	2	1.8	1.98	1.89	0.0829	6.73	1.80
PCB-044/047/065	4	4	0	122	320	200			200
PCB-045/051	4	4	0	5.97	25	11.9			11.9
PCB-046	4	4	0	0.41	1.77	0.791			0.791
PCB-048	4	4	0	4.08	14.6	7.29			7.29
PCB-049/069	4	4	0	105	279	177			177
PCB-050/053	4	4	0	2.16	8.62	3.85			3.85
PCB-052	4	4	0	190	395	284			284
PCB-054	4	4	0	0.104	0.348	0.180			0.180
PCB-055	4	2	2	0.399	0.738	0.569	0.0919	0.532	0.362
PCB-056	4	4	0	11.3	41.9	21.9			21.9
PCB-057	4	4	0	0.259	0.895	0.556			0.556
PCB-058	4	3	1	0.256	0.464	0.330	0.512	0.512	0.312
PCB-059/062/075	4	4	0	9.86	22.2	15.0			15.0
PCB-060	4	4	0	37.8	82.9	61.0			61.0
PCB-061/070/074/076	4	4	0	282	629	450			450
PCB-063	4	4	0	8.18	17.8	12.4			12.4
PCB-064	4	4	0	58.2	174	107			107
PCB-066	4	4	0	190	428	311			311
PCB-067	4	4	0	0.829	2.24	1.42			1.42
PCB-068	4	4	0	0.623	1.39	1.07			1.07
PCB-072	4	4	0	1.28	2.9	2.17			2.17
PCB-073	4	1	3	1.73	1.73	1.73	0.024	3.95	0.934
PCB-077	4	4	0	0.602	1.65	0.972			0.972
PCB-078	4	0	4				0.0184	0.492	0.0797
PCB-079	4	1	3	1.00	1.00	1.00	0.0156	0.428	0.315
PCB-080	4	0	4				0.0167	0.465	0.0747
PCB-081	4	3	1	0.265	0.494	0.387	0.479	0.479	0.350
PCB-082	4	4	0	3.63	9.39	6.48			6.48
PCB-083	4	4	0	1.69	3.13	2.25			2.25
PCB-084	4	4	0	5.58	14.9	9.57			9.57
PCB-085/116/117	4	4	0	24.5	59.1	38.2			38.2
PCB-086/087/097/108/119/125	4	4	0	46.9	111	73.3			73.3
PCB-088/091	4	4	0	11.3	24.1	17.9			17.9
PCB-089	4	4	0	0.148	0.538	0.282			0.282
PCB-090/101/113	4	4	0	83.5	191	129			129
PCB-092	4	4	0	9.31	21.7	15.3			15.3
PCB-093/100	4	4	0	0.888	2	1.46			1.46
PCB-094	4	3	1	0.0326	0.102	0.0637	0.0899	0.0899	0.0590
PCB-095	4	4	0	24.6	54.6	40.9			40.9

Table 11
Pinfish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-096	4	4	0	0.196	0.694	0.377			0.377
PCB-098/102	4	4	0	0.288	0.885	0.524			0.524
PCB-099	4	4	0	59	123	87.0			87.0
PCB-103	4	4	0	0.577	1.45	1.03			1.03
PCB-104	4	1	3	0.0467	0.0467	0.0467	0.00159	0.0206	0.0165
PCB-105	4	4	0	35.8	86.4	58.6			58.6
PCB-106	4	0	4				0.00453	0.0868	0.0211
PCB-107/124	4	4	0	3.01	7.08	4.50			4.50
PCB-109	4	4	0	7.03	16.2	10.2			10.2
PCB-110/115	4	4	0	54.5	129	89.7			89.7
PCB-111	4	3	1	0.0401	0.0694	0.0522	0.0581	0.0581	0.0464
PCB-112	4	0	4				0.00304	0.0649	0.0139
PCB-114	4	4	0	2.66	6.68	4.50			4.50
PCB-118	4	4	0	87.1	188	128			128
PCB-120	4	4	0	0.245	0.637	0.411			0.411
PCB-121	4	1	3	0.0304	0.0304	0.0304	0.00302	0.061	0.0176
PCB-122	4	4	0	0.525	0.939	0.731			0.731
PCB-123	4	4	0	2.16	4.91	3.32			3.32
PCB-126	4	4	0	0.186	0.407	0.292			0.292
PCB-127	4	2	2	0.0913	0.191	0.141	0.00424	0.0454	0.0768
PCB-128/166	4	4	0	5.53	14.3	8.17			8.17
PCB-129/138/163	4	4	0	41.6	93.7	57.0			57.0
PCB-130	4	4	0	1.93	4.82	2.85			2.85
PCB-131	4	4	0	0.263	0.605	0.399			0.399
PCB-132	4	4	0	5.68	11.6	7.85			7.85
PCB-133	4	3	1	0.399	0.92	0.64	0.00418	0.00418	0.481
PCB-134	4	4	0	0.224	1.02	0.580			0.580
PCB-135/151	4	4	0	4.27	10.5	6.62			6.62
PCB-136	4	4	0	1.26	3.08	2.07			2.07
PCB-137	4	4	0	2.24	5.78	3.46			3.46
PCB-139/140	4	4	0	0.838	1.82	1.18			1.18
PCB-141	4	4	0	5.41	12.5	7.57			7.57
PCB-142	4	0	4				0.00456	0.0518	0.0120
PCB-143	4	0	4				0.00423	0.0481	0.0111
PCB-144	4	4	0	1.29	2.78	1.76			1.76
PCB-145	4	2	2	0.00267	0.00613	0.00440	0.00924	0.0132	0.00500
PCB-146	4	4	0	0.0406	7.81	5.51			5.51
PCB-147/149	4	4	0	14.4	33.7	22.1			22.1
PCB-148	4	4	0	0.0525	0.136	0.0948			0.0948
PCB-150	4	4	0	0.0456	0.113	0.0797			0.0797
PCB-152	4	3	1	0.00525	0.018	0.0121	0.0127	0.0127	0.0107
PCB-153/168	4	4	0	35.8	82.7	51.8			51.8
PCB-154	4	4	0	0.474	1.13	0.760			0.760

Table 11
Pinfish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-155	4	4	0	0.0379	0.199	0.0985			0.0985
PCB-156/157	4	4	0	4.66	11.1	6.92			6.92
PCB-158	4	4	0	4.17	10	5.94			5.94
PCB-159	4	1	3	0.181	0.181	0.181	0.00352	0.0329	0.0503
PCB-160	4	0	4				0.00298	0.0356	0.00832
PCB-161	4	1	3	12.1	12.1	12.1	0.0028	0.0157	3.03
PCB-162	4	4	0	0.165	0.363	0.243			0.243
PCB-164	4	4	0	1.69	4.07	2.44			2.44
PCB-165	4	3	1	0.0138	0.0423	0.0307	0.0116	0.0116	0.0245
PCB-167	4	4	0	1.8	4.16	2.56			2.56
PCB-169	4	2	2	0.0412	0.0465	0.0439	0.0033	0.00384	0.0228
PCB-170	4	4	0	5.91	15.5	8.76			8.76
PCB-171/173	4	4	0	1.82	4.24	2.54			2.54
PCB-172	4	4	0	0.927	2.64	1.44			1.44
PCB-174	4	4	0	2.26	6.02	3.51			3.51
PCB-175	4	4	0	0.287	0.735	0.439			0.439
PCB-176	4	4	0	0.389	1.06	0.631			0.631
PCB-177	4	4	0	1.44	4.34	2.49			2.49
PCB-178	4	4	0	0.727	1.98	1.17			1.17
PCB-179	4	4	0	0.958	2.43	1.50			1.50
PCB-180/193	4	4	0	15	38.1	21.6			21.6
PCB-181	4	4	0	0.0941	0.249	0.144			0.144
PCB-182	4	3	1	0.0486	0.0902	0.0629	0.00805	0.00805	0.0482
PCB-183/185	4	4	0	4.33	11.3	6.52			6.52
PCB-184	4	4	0	0.0215	0.0902	0.0524			0.0524
PCB-186	4	0	4				0.000818	0.00960	0.00237
PCB-187	4	4	0	7.64	22.2	12.5			12.5
PCB-188	4	4	0	0.0311	0.0979	0.0619			0.0619
PCB-189	4	4	0	0.202	0.464	0.282			0.282
PCB-190	4	4	0	1.18	3.35	1.85			1.85
PCB-191	4	4	0	0.266	0.721	0.415			0.415
PCB-192	4	3	1	0.00531	0.0164	0.00905	0.019	0.019	0.00916
PCB-194	4	4	0	1.79	7.58	3.69			3.69
PCB-195	4	4	0	0.897	2.61	1.44			1.44
PCB-196	4	4	0	1.38	3.93	2.21			2.21
PCB-197/200	4	4	0	0.354	0.931	0.600			0.600
PCB-198/199	4	4	0	2.15	6.46	3.58			3.58
PCB-201	4	4	0	0.418	1.02	0.645			0.645
PCB-202	4	4	0	0.477	1.71	1.06			1.06
PCB-203	4	4	0	1.71	5.14	2.78			2.78
PCB-204	4	4	0	0.0183	0.0717	0.0401			0.0401
PCB-205	4	4	0	0.149	0.389	0.241			0.241
PCB-206	4	4	0	1.24	4.34	2.65			2.65

Table 11
Pinfish Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-207	4	4	0	0.776	2.75	1.80			1.80
PCB-208	4	4	0	0.728	3.16	1.94			1.94
PCB-209	4	4	0	14.3	92.3	48.9			48.9
Total PCB Congener (U = 1/2)	4	4	0	2,114	4,334	3,157			3,157
Dioxins and Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	4	4	0	1.56	169	45.3			45.3
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	4	4	0	3.46	15	8.60			8.60
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	4	3	1	0.255	5.41	2.16	0.140	0.140	1.64
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	4	2	2	1.23	1.24	1.24	0.253	0.357	0.694
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	4	0	4				0.207	0.573	0.176
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	4	3	1	0.405	0.62	0.506	0.110	0.110	0.393
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	4	0	4				0.135	0.235	0.0930
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	4	2	2	0.251	0.276	0.264	0.102	0.153	0.164
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	4	0	4				0.141	0.242	0.0990
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	4	0	4				0.156	0.419	0.130
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	4	0	4				0.140	0.242	0.0980
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	4	4	0	0.678	2.38	1.50			1.50
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	4	3	1	0.262	0.652	0.394	0.152	0.152	0.314
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	4	0	4				0.145	0.364	0.115
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	4	4	0	0.916	3.19	1.84			1.84
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	4	4	0	4.27	11.4	9.27			9.27
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	4	4	0	2.29	7.96	4.09			4.09
Total Dioxin/Furan (U = 1/2)	4	4	0	14.6	210	74.5			74.5

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram -- Results not applicable

Table 12
Sand Seatrout Tissue Sample Summary

Sand Seatrout Tissue Sample Summary										
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average	
Conventional Parameters (percent)										
Lipids	2	2	0	0.931	1.18	1.06			1.06	
Moisture, percent	2	2	0	78.7	79.2	79.0			79.0	
Metals (mg/kg)										
Mercury	2	2	0	0.045	0.063	0.0540			0.0540	
Semivolatile Organic Carbons (μg/kg)										
1,3-Dichlorobenzene	2	0	2				353	370	181	
Hexachlorobenzene	2	2	0	240	325	283			283	
PCBs (μg/kg)										
PCB-001	2	2	0	0.0597	0.176	0.118			0.118	
PCB-002	2	1	1	0.0191	0.0191	0.0191	0.00834	0.00834	0.0116	
PCB-003	2	1	1	0.072	0.072	0.0720	0.00642	0.00642	0.0376	
PCB-004	2	2	0	0.76	0.767	0.764			0.764	
PCB-005	2	0	2				0.00389	0.0197	0.00590	
PCB-006	2	2	0	0.403	0.57	0.487			0.487	
PCB-007	2	2	0	0.0659	0.0914	0.0787			0.0787	
PCB-008	2	2	0	2.38	3.07	2.73			2.73	
PCB-009	2	2	0	0.143	0.217	0.180			0.180	
PCB-010	2	2	0	0.0277	0.0293	0.0285			0.0285	
PCB-011	2	0	2				0.0288	0.0922	0.0303	
PCB-012/013	2	1	1	0.0282	0.0282	0.0282	0.0164	0.0164	0.0182	
PCB-014	2	0	2				0.00339	0.016	0.00485	
PCB-015	2	2	0	0.088	0.191	0.140			0.140	
PCB-016	2	2	0	3.09	4.2	3.65			3.65	
PCB-017	2	2	0	7.68	12.4	10.0			10.0	
PCB-018/030	2	2	0	15.7	28	21.9			21.9	
PCB-019	2	2	0	0.708	1.08	0.894			0.894	
PCB-020/028	2	2	0	54.5	101	77.8			77.8	
PCB-021/033	2	2	0	11.8	21.9	16.9			16.9	
PCB-022	2	2	0	9.77	17.8	13.8			13.8	
PCB-023	2	2	0	0.0231	0.0509	0.0370			0.0370	
PCB-024	2	2	0	0.132	0.208	0.170			0.170	
PCB-025	2	2	0	1.36	2.62	1.99			1.99	
PCB-026/029	2	2	0	5.94	11	8.47			8.47	
PCB-027	2	2	0	0.785	1.35	1.07			1.07	
PCB-031	2	2	0	42	81.6	61.8			61.8	
PCB-032	2	2	0	6.4	11.3	8.85			8.85	
PCB-032	2	2	0	0.250	0.389	0.320			0.320	
PCB-035	2	0	2				0.00715	0.0314	0.00964	
PCB-035	2	0	2				0.00713	0.0314	0.00916	
PCB-030	2	2	0	0.704	1.16	0.932			0.932	
PCB-037 PCB-038	2	2	0	0.0626	0.168	0.932			0.932	
	+									
PCB-039	2	0	2				0.00656	0.0281	0.00867	

Table 12
Sand Seatrout Tissue Sample Summary

		7	nd Seatrout Hissue	. Sample Samina	y	 		 	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-040/071	2	2	0	23.3	43.9	33.6			33.6
PCB-041	2	2	0	4.58	6.74	5.66			5.66
PCB-042	2	2	0	12.4	19.4	15.9			15.9
PCB-043	2	1	1	4.01	4.01	4.01	0.0212	0.0212	2.01
PCB-044/047/065	2	2	0	69.7	115	92.4			92.4
PCB-045/051	2	2	0	5.67	7.58	6.63			6.63
PCB-046	2	2	0	1.01	1.48	1.25			1.25
PCB-048	2	2	0	15.8	28.2	22.0			22.0
PCB-049/069	2	2	0	53	86	69.5			69.5
PCB-050/053	2	2	0	3.3	5	4.15			4.15
PCB-052	2	2	0	85.3	148	117			117
PCB-054	2	2	0	0.0851	0.165	0.125			0.125
PCB-055	2	1	1	1.43	1.43	1.43	0.0171	0.0171	0.719
PCB-056	2	2	0	15.9	31.6	23.8			23.8
PCB-057	2	2	0	0.302	0.567	0.435			0.435
PCB-058	2	2	0	0.225	0.357	0.291			0.291
PCB-059/062/075	2	2	0	5.84	9.59	7.72			7.72
PCB-060	2	2	0	25	48.7	36.9			36.9
PCB-061/070/074/076	2	2	0	168	307	238			238
PCB-063	2	2	0	4.87	8.4	6.64			6.64
PCB-064	2	2	0	35.6	61.4	48.5			48.5
PCB-066	2	2	0	121	217	169			169
PCB-067	2	2	0	1.12	2.06	1.59			1.59
PCB-068	2	2	0	0.524	0.655	0.590			0.590
PCB-072	2	2	0	1.07	1.53	1.30			1.30
PCB-073	2	1	1	1.33	1.33	1.33	0.0634	0.0634	0.681
PCB-077	2	2	0	1.37	2.94	2.16			2.16
PCB-078	2	0	2				0.0162	0.0567	0.0182
PCB-079	2	1	1	0.969	0.969	0.969	0.0139	0.0139	0.488
PCB-080	2	0	2				0.0146	0.0521	0.0167
PCB-081	2	2	0	0.0462	0.138	0.0921			0.0921
PCB-082	2	2	0	4.06	8.13	6.10			6.10
PCB-083	2	1	1	3.68	3.68	3.68	0.012	0.012	1.84
PCB-084	2	2	0	4.81	8.55	6.68			6.68
PCB-085/116/117	2	2	0	18.1	31.5	24.8			24.8
PCB-086/087/097/108/119/125	2	2	0	29.2	51.9	40.6			40.6
PCB-088/091	2	2	0	5.13	7.97	6.55			6.55
PCB-089	2	2	0	0.409	0.836	0.623			0.623
PCB-090/101/113	2	2	0	46.7	75.2	61.0			61.0
PCB-092	2	2	0	7.7	12.9	10.3			10.3
PCB-093/100	2	2	0	0.810	1.13	0.970			0.970
PCB-094	2	2	0	0.078	0.177	0.128			0.128
PCB-095	2	2	0	12	19.4	15.7			15.7

Table 12
Sand Seatrout Tissue Sample Summary

			Number of	•	<u> </u>				
Chemical	Number of Samples	Number of Detected Samples	Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-096	2	2	0	0.327	0.512	0.420			0.420
PCB-098/102	2	2	0	0.926	1.5	1.21			1.21
PCB-099	2	2	0	44.8	66.2	55.5			55.5
PCB-103	2	2	0	0.591	0.790	0.691			0.691
PCB-104	2	2	0	0.0131	0.0254	0.0193			0.0193
PCB-105	2	2	0	25.2	48.7	37.0			37.0
PCB-106	2	0	2				0.0114	0.0506	0.0155
PCB-107/124	2	2	0	1.71	3.42	2.57			2.57
PCB-109	2	2	0	5.03	8.85	6.94			6.94
PCB-110/115	2	2	0	44.1	82.6	63.4			63.4
PCB-111	2	2	0	0.0507	0.0631	0.0569			0.0569
PCB-112	2	1	1	0.360	0.360	0.360	0.00718	0.00718	0.182
PCB-114	2	2	0	1.84	3.81	2.83			2.83
PCB-118	2	2	0	55.6	107	81.3			81.3
PCB-120	2	2	0	0.317	0.409	0.363			0.363
PCB-121	2	1	1	0.0174	0.0174	0.0174	0.0223	0.0223	0.0143
PCB-122	2	2	0	0.137	0.568	0.353			0.353
PCB-123	2	2	0	1.04	2.28	1.66			1.66
PCB-126	2	1	1	0.202	0.202	0.202	0.0139	0.0139	0.104
PCB-127	2	2	0	0.0748	0.136	0.105			0.105
PCB-128/166	2	2	0	5.12	9.2	7.16			7.16
PCB-129/138/163	2	2	0	36.4	59.5	48.0			48.0
PCB-130	2	2	0	1.4	2.49	1.95			1.95
PCB-131	2	2	0	0.222	0.401	0.312			0.312
PCB-132	2	2	0	3.96	7.33	5.65			5.65
PCB-133	2	2	0	0.628	0.916	0.772			0.772
PCB-134	2	2	0	0.57	1.14	0.855			0.855
PCB-135/151	2	2	0	4.3	6.59	5.45			5.45
PCB-136	2	2	0	1.08	1.69	1.39			1.39
PCB-137	2	2	0	2.23	3.77	3.00			3.00
PCB-139/140	2	2	0	0.858	1.26	1.06			1.06
PCB-141	2	2	0	4.22	7.46	5.84			5.84
PCB-142	2	1	1	0.0154	0.0154	0.0154	0.0315	0.0315	0.0156
PCB-143	2	1	1	0.0582	0.0582	0.0582	0.0109	0.0109	0.0318
PCB-144	2	2	0	1.04	1.53	1.29			1.29
PCB-145	2	2	0	0.00528	0.0119	0.00859			0.00859
PCB-146	2	2	0	0.0256	0.0429	0.0343			0.0343
PCB-147/149	2	2	0	7.78	11.6	9.69			9.69
PCB-148	2	2	0	0.073	0.0871	0.0801			0.0801
PCB-150	2	2	0	0.0602	0.0675	0.0639			0.0639
PCB-152	2	2	0	0.0186	0.0298	0.0242			0.0242
PCB-153/168	2	2	0	38.5	55.9	47.2			47.2
PCB-154	2	2	0	0.877	0.996	0.937			0.937

Table 12
Sand Seatrout Tissue Sample Summary

		7	nd Seatrout Hissue	- Jampie Jaminai	y	 		1	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-155	2	2	0	0.110	0.125	0.118			0.118
PCB-156/157	2	2	0	3.95	7.26	5.61			5.61
PCB-158	2	2	0	3.76	6.35	5.06			5.06
PCB-159	2	0	2				0.00852	0.0195	0.00701
PCB-160	2	0	2				0.00841	0.0216	0.00750
PCB-161	2	2	0	3.54	5.02	4.28			4.28
PCB-162	2	2	0	0.165	0.344	0.255			0.255
PCB-164	2	2	0	1.36	2.47	1.92			1.92
PCB-165	2	2	0	0.0176	0.0286	0.0231			0.0231
PCB-167	2	2	0	1.39	2.45	1.92			1.92
PCB-169	2	0	2				0.00704	0.0135	0.00514
PCB-170	2	2	0	6.92	12	9.46			9.46
PCB-171/173	2	2	0	2.2	3.46	2.83			2.83
PCB-172	2	2	0	1.32	2.22	1.77			1.77
PCB-174	2	2	0	2.11	3.9	3.01			3.01
PCB-175	2	2	0	0.341	0.500	0.421			0.421
PCB-176	2	2	0	0.416	0.622	0.519			0.519
PCB-177	2	2	0	1.47	2.59	2.03			2.03
PCB-178	2	2	0	1.46	2.09	1.78			1.78
PCB-179	2	2	0	1.03	1.74	1.39			1.39
PCB-180/193	2	2	0	19	31.1	25.1			25.1
PCB-181	2	2	0	0.124	0.194	0.159			0.159
PCB-182	2	2	0	0.0829	0.109	0.0960			0.0960
PCB-183/185	2	2	0	6.27	8.88	7.58			7.58
PCB-184	2	2	0	0.0948	0.0955	0.0952			0.0952
PCB-186	2	0	2				0.00127	0.00395	0.00131
PCB-187	2	2	0	5.52	9.98	7.75			7.75
PCB-188	2	2	0	0.0585	0.0758	0.0672			0.0672
PCB-189	2	2	0	0.248	0.428	0.338			0.338
PCB-190	2	2	0	1.43	2.45	1.94			1.94
PCB-191	2	2	0	0.359	0.566	0.463			0.463
PCB-192	2	2	0	0.0107	0.0172	0.0140			0.0140
PCB-194	2	2	0	4.33	7.61	5.97			5.97
PCB-195	2	2	0	1.42	2.43	1.93			1.93
PCB-196	2	2	0	2.3	3.48	2.89			2.89
PCB-197/200	2	2	0	0.736	1	0.868			0.868
PCB-198/199	2	2	0	3.78	6.33	5.06			5.06
PCB-201	2	2	0	0.744	0.997	0.871			0.871
PCB-202	2	2	0	1.43	2.18	1.81			1.81
PCB-203	2	2	0	3.03	4.61	3.82			3.82
PCB-204	2	2	0	0.0636	0.0748	0.0692			0.0692
PCB-205	2	1	1	0.38	0.38	0.380	0.207	0.207	0.242
PCB-206	2	2	0	4.25	6.59	5.42			5.42

Table 12
Sand Seatrout Tissue Sample Summary

		Jul	na scatioat 11ssat	Sample Summar	7				
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-207	2	2	0	3.67	5.4	4.54			4.54
PCB-208	2	2	0	4.5	6.72	5.61			5.61
PCB-209	2	2	0	112	175	144			144
Total PCB Congener (U = 1/2)	2	2	0	1,449	2,512	1,980			1,980
Dioxins and Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	2	2	0	3.2	5.39	4.30			4.30
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	2	2	0	2.81	3.08	2.95			2.95
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	2	2	0	0.441	0.469	0.455			0.455
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	2	2	0	0.549	0.662	0.606			0.606
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	2	0	2				0.240	0.294	0.134
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	2	2	0	0.221	0.403	0.312			0.312
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	2	0	2				0.121	0.136	0.0643
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	2	2	0	0.155	0.243	0.199			0.199
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	2	1	1	0.285	0.285	0.285	0.145	0.145	0.179
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	2	0	2				0.165	0.181	0.0865
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	2	0	2				0.129	0.143	0.0680
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	2	2	0	0.871	1.08	0.976			0.976
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	2	0	2				0.108	0.162	0.0675
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	2	1	1	0.181	0.181	0.181	0.147	0.147	0.127
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	2	2	0	0.216	0.356	0.286			0.286
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	2	1	1	0.877	0.877	0.877	1.05	1.05	0.701
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	2	2	0	0.800	15.5	0.825			0.825
Total Dioxin/Furan (U = 1/2)	2	2	0	11.6	13.1	12.3			12.3

mg/kg - milligrams per kilogram
μg/kg - micrograms per kilogram
ng/kg - nanograms per kilogram
-- Results not applicable

Table 13
Striped Mullet Tissue Sample Summary

		Julipe	a Mullet Hissue Sa	ampic Sammary	1			1	
Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
Conventional Parameters (percent)									
Lipids	9	9	0	1.24	2.36	1.74			1.74
Moisture, percent	9	9	0	73.1	75.4	74.0			74.0
Metals (mg/kg)									
Mercury	9	9	0	0.024	0.122	0.0516			0.0516
Semivolatile Organic Carbons (μg/kg)									
1,3-Dichlorobenzene	9	0	9				342	769	304
Hexachlorobenzene	9	7	2	389	1,830	823	727	742	721
PCBs (μg/kg)	•	•	•		•	•		•	
PCB-001	9	9	0	0.0785	0.237	0.127			0.127
PCB-002	9	3	6	0.00426	0.0133	0.00999	0.00543	0.0164	0.0062
PCB-003	9	8	1	0.0275	0.0625	0.0392	0.0142	0.0142	0.0356
PCB-004	9	9	0	0.91	2.87	1.49			1.49
PCB-005	9	5	4	0.046	0.127	0.0735	0.017	0.0414	0.0461
PCB-006	9	9	0	0.342	1.64	0.818			0.818
PCB-007	9	9	0	0.0901	0.303	0.149			0.149
PCB-008	9	9	0	2.42	8.4	4.33			4.33
PCB-009	9	9	0	0.147	0.757	0.360			0.360
PCB-010	9	9	0	0.0246	0.11	0.0530			0.0530
PCB-010	9	0	9				0.0333	0.176	0.0403
PCB-011 PCB-012/013	9	8	1	0.0654	0.248	0.114	0.0335	0.0335	0.103
PCB-012/013 PCB-014	9	+	9					+	0.00802
		0					0.00252	0.0327	
PCB-015	9	9	0	0.438	2.01	0.904			0.904
PCB-016	9	9	0	1.13	12	5.63			5.63
PCB-017	9	9	0	3.63	27.4	13.3			13.3
PCB-018/030	9	9	0	6.22	74.3	34.1			34.1
PCB-019	9	9	0	0.478	3.53	1.71			1.71
PCB-020/028	9	9	0	24.4	218	107			107
PCB-021/033	9	9	0	6.49	40	18.4			18.4
PCB-022	9	9	0	4.26	42.4	19.5			19.5
PCB-023	9	9	0	0.012	0.091	0.0458			0.0458
PCB-024	9	9	0	0.0566	0.493	0.234			0.234
PCB-025	9	9	0	1.00	7.49	3.68			3.68
PCB-026/029	9	9	0	3.00	26.6	13.0			13.0
PCB-027	9	9	0	0.378	3.18	1.53			1.53
PCB-031	9	9	0	22.1	212	99.4			99.4
PCB-032	9	9	0	2.71	24.2	11.6			11.6
PCB-034	9	9	0	0.165	0.69	0.388			0.388
PCB-035	9	7	2	0.044	0.131	0.0630	0.0356	0.0403	0.0532
PCB-036	9	0	9				0.00614	0.0464	0.0133
PCB-037	9	9	0	1.80	10.8	4.53			4.53
PCB-038	9	9	0	0.0463	0.294	0.133			0.133
PCB-039	9	1	8	0.592	0.592	0.592	0.00581	0.044	0.0781

Table 13
Striped Mullet Tissue Sample Summary

Chemical PCD 040/074	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-040/071	9	9	0	9.95	77.5	37.0			37.0
PCB-041	9	9	0	1.09	9.99	4.42			4.42
PCB-042	9	9	0	6.67	45.1	22.3			22.3
PCB-043	9	4	5	0.732	9.41	4.96	0.0235	0.0805	2.22
PCB-044/047/065	9	9	0	29.4	252	126			126
PCB-045/051	9	9	0	2.84	26.6	13.2			13.2
PCB-046	9	9	0	0.632	5.35	2.50			2.50
PCB-048	9	9	0	5.45	45.8	22.1			22.1
PCB-049/069	9	9	0	25.5	208	107			107
PCB-050/053	9	9	0	2.58	23.5	11.4			11.4
PCB-052	9	9	0	38.2	330	167			167
PCB-054	9	9	0	0.0547	0.345	0.176			0.176
PCB-055	9	9	0	0.167	3.28	1.39			1.39
PCB-056	9	9	0	10.5	116	51.7			51.7
PCB-057	9	9	0	0.121	1.53	0.710			0.710
PCB-058	9	9	0	0.138	0.807	0.395			0.395
PCB-059/062/075	9	9	0	2.41	20.3	10.2			10.2
PCB-060	9	9	0	5.83	82.3	36.5			36.5
PCB-061/070/074/076	9	9	0	64.5	623	296			296
PCB-063	9	9	0	1.27	16	7.54			7.54
PCB-064	9	9	0	13	121	59.3			59.3
PCB-066	9	9	0	41.2	372	179			179
PCB-067	9	9	0	0.68	7.52	3.50			3.50
PCB-068	9	9	0	0.31	1.19	0.704			0.704
PCB-072	9	9	0	0.6	2.63	1.49			1.49
PCB-073	9	5	4	1.62	4.57	2.58	0.00782	0.0611	1.44
PCB-077	9	9	0	1.81	13.1	5.90			5.90
PCB-078	9	0	9				0.00907	0.0783	0.0217
PCB-079	9	3	6	0.379	1.82	1.13	0.0127	0.0575	0.389
PCB-080	9	0	9				0.00819	0.0719	0.0198
PCB-081	9	9	0	0.0789	0.815	0.387			0.387
PCB-082	9	9	0	1.67	11.1	5.06			5.06
PCB-083	9	7	2	0.76	7.1	3.50	0.0139	0.0396	2.73
PCB-084	9	9	0	2.89	25.2	11.8			11.8
PCB-085/116/117	9	9	0	4.49	47.8	22.4			22.4
PCB-086/087/097/108/119/125	9	9	0	12.3	103	49.0			49.0
PCB-088/091	9	9	0	3.31	23.1	11.2			11.2
PCB-089	9	9	0	0.222	2.19	0.952			0.952
PCB-090/101/113	9	9	0	20.9	158	78.9			78.9
PCB-092	9	9	0	3.23	26.9	13.5			13.5
PCB-093/100	9	9	0	0.322	2.3	1.22			1.22
PCB-094	9	9	0	0.105	0.904	0.400			0.400
PCB-095	9	9	0	9.52	71.1	34.9			34.9

Table 13
Striped Mullet Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-096	9	9	0	0.122	1.25	0.641			0.641
PCB-098/102	9	9	0	0.847	6.05	2.91			2.91
PCB-099	9	9	0	14.1	95.6	50.5			50.5
PCB-103	9	9	0	0.281	1.44	0.835			0.835
PCB-104	9	9	0	0.0131	0.0528	0.0349			0.0349
PCB-105	9	9	0	7.36	74.6	34.3			34.3
PCB-106	9	3	6	0.0136	0.269	0.138	0.00702	0.0561	0.0552
PCB-107/124	9	9	0	0.725	7.25	3.32			3.32
PCB-109	9	9	0	1.52	14.1	6.75			6.75
PCB-110/115	9	9	0	18.2	143	68.1			68.1
PCB-111	9	9	0	0.023	0.093	0.0521			0.0521
PCB-112	9	3	6	0.0345	0.732	0.391	0.0027	0.0284	0.136
PCB-114	9	9	0	0.487	5.78	2.56			2.56
PCB-118	9	9	0	17.6	151	72.0			72.0
PCB-120	9	9	0	0.148	0.45	0.303			0.303
PCB-121	9	5	4	0.0111	0.0365	0.0222	0.00268	0.0387	0.0178
PCB-122	9	9	0	0.261	2.69	1.13			1.13
PCB-123	9	9	0	0.44	4.15	1.98			1.98
PCB-126	9	8	1	0.0418	0.423	0.193	0.00919	0.00919	0.172
PCB-127	9	7	2	0.0273	0.152	0.0803	0.00656	0.0545	0.0659
PCB-128/166	9	9	0	1.82	11.3	5.77			5.77
PCB-129/138/163	9	9	0	13.3	72.1	39.2			39.2
PCB-130	9	9	0	0.949	4.86	2.62			2.62
PCB-131	9	9	0	0.11	0.708	0.380			0.380
PCB-132	9	9	0	2.74	11.3	6.02			6.02
PCB-133	9	9	0	0.291	1.09	0.687			0.687
PCB-134	9	9	0	0.433	2.58	1.303			1.303
PCB-135/151	9	9	0	3.07	12.9	7.97			7.97
PCB-136	9	9	0	0.958	4.13	2.52			2.52
PCB-137	9	9	0	0.66	4.64	2.32			2.32
PCB-139/140	9	9	0	0.285	1.40	0.785			0.785
PCB-141	9	9	0	1.92	11.3	6.05			6.05
PCB-142	9	1	8	0.0127	0.0127	0.0127	0.00584	0.0389	0.0116
PCB-143	9	3	6	0.0273	0.173	0.106	0.00326	0.0324	0.0412
PCB-144	9	9	0	0.482	2.37	1.38			1.38
PCB-145	9	9	0	0.00394	0.0328	0.0156			0.0156
PCB-146	9	8	1	0.0187	10.1	2.89	0.0177	0.0177	2.57
PCB-147/149	9	9	0	8.6	30.3	17.6			17.6
PCB-148	9	9	0	0.0414	0.121	0.0756			0.0756
PCB-150	9	9	0	0.0486	0.118	0.0807			0.0807
PCB-152	9	9	0	0.0115	0.078	0.0413			0.0413
PCB-153/168	9	9	0	13.7	60.1	35.1			35.1
PCB-154	9	9	0	0.363	0.953	0.687			0.687

Table 13
Striped Mullet Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-155	9	9	0	0.0305	0.141	0.0683			0.0683
PCB-156/157	9	9	0	1.3	8.83	4.47			4.47
PCB-158	9	9	0	1.29	7.49	3.98			3.98
PCB-159	9	0	9				0.00376	0.0286	0.00833
PCB-160	9	0	9				0.0023	0.0267	0.00724
PCB-161	9	5	4	3.64	7.46	5.53	0.00215	0.026	3.07
PCB-162	9	9	0	0.0513	0.302	0.1727			0.1727
PCB-164	9	9	0	0.845	3.50	1.89			1.89
PCB-165	9	9	0	0.011	0.0631	0.0367			0.0367
PCB-167	9	9	0	0.513	3.18	1.64			1.64
PCB-169	9	1	8	0.0161	0.0161	0.0161	0.00722	0.021	0.00791
PCB-170	9	9	0	2.1	11.9	6.52			6.52
PCB-171/173	9	9	0	0.689	3.16	1.87			1.87
PCB-172	9	9	0	0.418	2.24	1.27			1.27
PCB-174	9	9	0	1.62	5.45	3.21			3.21
PCB-175	9	9	0	0.135	0.505	0.331			0.331
PCB-176	9	9	0	0.284	1.05	0.678			0.678
PCB-177	9	9	0	1.27	4.98	2.99			2.99
PCB-178	9	9	0	0.615	2.15	1.42			1.42
PCB-179	9	9	0	0.84	3.25	2.10			2.10
PCB-180/193	9	9	0	5.22	28.5	16.0			16.0
PCB-181	9	8	1	0.0348	0.194	0.112	0.0312	0.0312	0.101
PCB-182	9	9	0	0.0182	0.100	0.0608			0.0608
PCB-183/185	9	9	0	1.86	8.14	5.02			5.02
PCB-184	9	9	0	0.0202	0.0876	0.0440			0.0440
PCB-186	9	2	7	0.00281	0.00571	0.00426	0.00169	0.00716	0.00253
PCB-187	9	9	0	3.67	16.0	9.24			9.24
PCB-188	9	9	0	0.019	0.0691	0.0421			0.0421
PCB-189	9	9	0	0.0783	0.396	0.226			0.226
PCB-190	9	9	0	0.43	2.34	1.30			1.30
PCB-191	9	9	0	0.107	0.542	0.304			0.304
PCB-192	9	1	8	0.00282	0.00282	0.00282	0.00425	0.025	0.00567
PCB-194	9	9	0	0.871	5.00	2.71			2.71
PCB-195	9	9	0	0.37	1.72	0.980			0.980
PCB-196	9	9	0	0.567	2.48	1.48			1.48
PCB-197/200	9	9	0	0.256	0.731	0.490			0.490
PCB-198/199	9	9	0	1.28	5.24	3.05			3.05
PCB-201	9	9	0	0.23	0.671	0.448			0.448
PCB-202	9	9	0	0.501	1.59	1.00			1.00
PCB-203	9	9	0	0.721	3.14	1.83			1.83
PCB-204	9	9	0	0.011	0.0518	0.0271			0.0271
PCB-205	9	9	0	0.0777	0.250	0.162			0.162
PCB-206	9	9	0	0.736	2.63	1.55			1.55

Table 13
Striped Mullet Tissue Sample Summary

Chemical	Number of Samples	Number of Detected Samples	Number of Non-Detected Samples	Minimum Detection	Maximum Detection	Average Detection	Minimum Non-Detection	Maximum Non-Detection	Average
PCB-207	9	9	0	0.445	1.96	0.966			0.966
PCB-208	9	9	0	0.586	2.53	1.28			1.28
PCB-209	9	9	0	12.3	92.1	35.6			35.6
Total PCB Congener (U = 1/2)	9	9	0	596	4,539	2,246			2,246
Dioxins and Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	9	9	0	15.2	747	234			234
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	9	9	0	21.7	75.8	47.2			47.2
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	9	9	0	1.31	22.6	7.91			7.91
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	9	9	0	2.38	7.91	5.02			5.02
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	9	4	5	0.721	3.28	2.44	0.211	0.702	1.23
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	9	9	0	0.767	2.17	1.34			1.34
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	9	0	9				0.136	0.513	0.143
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	9	9	0	0.34	0.99	0.638			0.638
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	9	4	5	0.289	0.462	0.364	0.236	0.591	0.271
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	9	0	9				0.145	0.681	0.202
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	9	0	9				0.146	0.561	0.152
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	9	9	0	1.16	3.48	2.22			2.22
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	9	6	3	0.382	0.501	0.448	0.163	0.322	0.336
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	9	4	5	0.223	0.739	0.522	0.273	0.617	0.358
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	9	9	0	1.37	4.15	2.73			2.73
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	9	9	0	8.12	59.2	22.5			22.5
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	9	9	0	1.43	15.5	6.10			6.10
Total Dioxin/Furan (U = 1/2)	9	9	0	70.5	931	332			332

mg/kg - milligrams per kilogram μg/kg - micrograms per kilogram ng/kg - nanograms per kilogram -- Results not applicable

Table 14
Average Chemical Concentration for Each Sampled Species

	Average Concentration									
Tissue Sample Type Lead (mg/kg) Mercury (mg/kg) PAHs (μg/kg)		PAHs	1,3- Dichlorobenzene (μg/kg)	1,4- Dichlorobenzene (μg/kg)	HCB (μg/kg)	HCBD (μg/kg)	Total PCB Congeners (µg/kg)	Total Dioxin/Furan Congeners (ng/kg)		
Blue Crab Size Class A	0.178	0.0475	25.9	All ND	All ND	612	All ND	655	63.6	
Blue Crab Size Class B		0.0667		All ND		All ND		804		
Brown Shrimp		0.041		All ND		710		1,716		
Oyster	0.331	0.051	552	All ND	All ND	All ND	All ND	1,429	41.7	
White Shrimp Size Class A	0.116	0.041	162	All ND	All ND	483	All ND	837	69.4	
White Shrimp Size Class B		0.034		All ND		531		733		
Gulf Killifish		0.0524		All ND		441		5,216	36.2	
Gulf Menhaden		0.116		All ND		440		2,642	763	
Pinfish		0.043		All ND		872		3,157	74.5	
Sand Seatrout		0.054	-	All ND	-	283		1,980	12.3	
Striped Mullet		0.0516		All ND		721		2,246	332	

mg/kg - milligrams per kilogram

μg/kg - micrograms per kilogram

ng/kg - nanograms per kilogram

-- Results not applicable


All ND = All samples were non-detect

HCB = hexachlorobenzene

HCBD - hexachlorobutadiene

PAH - polyaromatic hydrocarbons


FIGURES













Notes:
Multiple samples may be associated with some sample locations.

Station numbers from Patrick Bayou PSCR indicate length along channel in hundreds of feet. Aerial orthoimagery: December 2007.





Invertebrate Tissue Sample Locations Ecological Fish and Invertebrate Tissue Sampling Data Report Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas

Figure 2





Multiple samples may be associated with some sample locations.

Station numbers from Patrick Bayou PSCR indicate length along channel in hundreds of feet. Aerial orthoimagery: December 2007.





Fish Tissue Sample Locations Ecological Fish and Invertebrate Tissue Sampling Data Report Patrick Bayou Superfund Site Remedial Investigation, Deer Park, Texas

Figure 3

APPENDIX A (AVAILABLE ON CD)

APPENDIX B (AVAILABLE ON CD)

APPENDIX C (AVAILABLE ON CD)

APPENDIX D TISSUE ANALYTICAL RESULTS

Appendix D Tissue Analytical Results

Sample Date 6/84/2011 6/2	Tissue Analytical Results										
Second S		Location ID	Field QC	Field QC	PB002.1	PB003.1	PB005.1	PB006.1	PB008.2	PB010	PB010.1
Part		Sample ID	EB-110621-A01	EB-110624-A01	PB01-I-A-BCR-W-001-COMP-201106	PB01-F-A-STM-W-006-20110618	PB01-I-A-BCR-W-002-COMP-201106	PB01-I-B-WHS-W-002-COMP-201106	PB01-I-B-WHS-W-001-COMP-201106	PB01-F-A-GM-W-008-20110618	PB01-I-A-WHS-W-001-COMP-201106
		Sample Date	6/24/2011	6/24/2011	6/20/2011	6/18/2011	6/20/2011	6/20/2011	6/19/2011	6/18/2011	6/20/2011
Part		Sample Type	EB	EB	N	N	N	N	N	N	N
Page Page		х			3202691	3202445.49	3202312	3202026.63	3202032	3201710.87	3201704.8
1968 1969		Υ			13836499	13836263.53	13836201	13836356.8	13836072	13836029.16	13836071.32
1968 1969		Scientific_Name			Callinectes sapidus	Mugil cephalus	Callinectes sapidus	Penaeus setiferus	Penaeus setiferus	Brevoortia patronus	Penaeus setiferus
Process		_			· '	• .	· ·	>7.5 to =<13	>7.5 to =<13	·	>2 to =<7.5
Perfect Pe					· ·	= -	·			- ·	
1960 1970	Chemical			Result	·			•	•	· · · · · · · · · · · · · · · · · · ·	•
Montaining Month	Conventional Parameters (pct)	ı									
March	Lipids	LIPID	4.24		3.02	1.24	3.73	1.13	1.3	1.23	0.914
Second 1969 1967	Moisture, percent	MOIST	71.2		69.1	73.1	70.5	75.3	74.6	77.5	76.9
Mary Mary	Metals (mg/kg)	<u>l</u>									
Proposed p	Lead	7439-92-1	0.474		0.121		0.111				0.116
## Accordance \$1.50	Mercury		0.042			0.085		0.042	0.026	0.078	
Accomplement March	Polyaromatic Hydrocarbons (μg/kg)										
Accomplement March	2-Methylnaphthalene	91-57-6	3.44 U		6.93 U		7.23 U				14.2 U
Martinary Mart			3.44 U		6.93 U		7.23 U				14.2 U
Manuscapinstratemen	Acenaphthylene	208-96-8	3.44 U		6.93 U		7.23 U				14.2 U
Marie Mari	Anthracene	120-12-7	3.44 U		6.93 U		7.23 U				14.2 U
Mary State 1969 1964 1964 1969 1	Benzo(a)anthracene	56-55-3	3.44 U		6.93 U		7.23 U				14.2 U
Money Mone	Benzo(a)pyrene	50-32-8	3.44 U		6.93 U		7.23 U				14.2 U
Personal Processes	Benzo(b)fluoranthene	205-99-2	3.44 U		6.93 U	-	7.23 U				14.2 U
Depart	Benzo(g,h,i)perylene	191-24-2	3.44 U		6.93 U		7.23 U				14.2 U
Description 19.00	Benzo(k)fluoranthene	207-08-9	3.44 U		6.93 U		7.23 U				14.2 U
Househole 2004-00 3.64 0 0 - 5.90 - 7.23 - - - - - - 5.90 - 1.42 - 1.4	Chrysene	218-01-9	3.44 U	-	6.93 U		7.23 U				14.2 U
Pubmer P	Dibenzo(a,h)anthracene	53-70-3	3.44 U	-	6.93 U		7.23 U				14.2 U
Indees 123-124 123-1	Fluoranthene	206-44-0	3.44 U		6.93 U		7.23 U				14.8
Naphthisine	Fluorene	86-73-7	3.44 U		6.93 U		7.23 U				14.2 U
Phenethorhere	Indeno(1,2,3-c,d)pyrene	193-39-5	3.44 U		6.93 U		7.23 U				14.2 U
Pyrene	Naphthalene	91-20-3	3.44 U		6.93 U		7.23 U				14.2 U
Trail Plant (10 15 17) Plant (17 15 11) Plant (Phenanthrene	85-01-8	3.44 U		6.93 U		7.23 U				18.2
Total PAM (16) (19 17)	Pyrene	129-00-0	3.44 U		6.93 U		7.23 U				29.9
Total PAM (16) (19 17)	Total HPAH (9 of 16) (U = 1/2)	tPAH 17 HM N	3.44 U		6.93 U		7.23 U				101.5
Total PAN [16] (U = 1/2)			3.44 U			-					
Semi-sublik Organic Carbons (Lig/Ng)			3.44 U								162.3
1,3-behiprobenzene 54.73+ 721 0 721 0 721 0 721 0 738 0 388 0 344 0 372 0 691 758 0 357 0 714 0 14-behiprobenzene 16.46-7 721 0 721 0 721 0 721 0 721 0 738 0 389 372 0 662 380 380 443 483 348											
1.4-Fichlorobenzene 16-46-7 721 U U U U U U U U U		541-73-1	721 U		388 U	344 U	372 U	691 U	758 U	357 U	714 U
Hexachlorobenzene 118.74 72 1											
Hexachlorobutadene											
PCB-001 PCB-002 PCB-002 PCB-002 PCB-002 PCB-002 PCB-003 PCB-004 PCB-004 PCB-005 PCB-006 PCB-00											
PCB-001 PCB-002 5.03 U 1.86 U 110 6.33 U 113 27 90.9 J 94.8 J PCB-002 PCB-002 6.05 U 1.7 U 5.43 U 1.33 U 5.81 U 2.29 U 8.14 U 4.01 U PCB-003 PCB-004 PCB-003 4.68 U 1.8 U 34.1 1.4 U 616 J 16.9 J 1.0 J 9.7 J 1.48 J PCB-004 PCB-004 4.68 U 1.8 U 34.1 1.4 U 616 J 16.9 J 30.1 J 4.93 PCB-004 PCB-005 10.5 U 4.17 U 17 UJ 2.9 U 22 U 13.6 U 907 J 148 J PCB-005 PCB-006 PCB-006 16.1 J 13.3 J 34.2 J 4.86 U 246 144 508 J 180 PCB-007 PCB-007 9.79 U 3.82 U 95.3 J 2.66 U 58.3 J 31.5 J 86.5 J <t< td=""><td></td><td>0. 00 0</td><td>, 0</td><td></td><td>333 0</td><td></td><td>5.2 0</td><td></td><td></td><td></td><td>72.0</td></t<>		0. 00 0	, 0		333 0		5.2 0				72.0
PCB-002 PCB-003 - 6.05 U 1.7 U 5.43 U 1.33 U 5.81 U 2.29 U 8.14 U 4.01 U PCB-003 PCB-003 - 4.68 U U 1.8 U 34.1 1.42 U 61.6 J 1.69 J 30.1 J 49.3 PCB-004 PCB-005 - 22.3 J 22.9 U 910 J 11.9 U 220 U 35.4 U 907 J 148 J PCB-005 PCB-005 - 10.5 U 4.17 U 17 UJ 2.9 U 22 U 12.6 U 26.3 UJ 24.4 U PCB-006 PCB-006 - 16.1 J 13.8 J 342 J 4.86 U 246 144 508 J 180 PCB-007 PCB-007 - 9.79 U 3.82 U 95.3 J 2.66 U 58.3 J 31.5 J 86.5 J 20.4 U PCB-008 PCB-007 - 42.6 J 2.89 U 3190 J 42 1690 U 941 U 2100 J 1190 U PCB-010 PCB-009 -		PCR-001		5.03 111	1 86 U	110	6 33 11	113	27	90 9 1	94 8 1
PCB-003 PCB-003 4.68 U 1.8 U 34.1 1.42 U 61.6 J 16.9 J 30.1 J 49.3 PCB-004 PCB-004 22.3 J 22.9 U 910 J 11.9 U 220 35.4 U 907 J 148 J PCB-005 PCB-005 10.5 U 4.17 U 17 UJ 2.9 U 22 U 12.6 U 26.3 UJ 24.4 U PCB-006 PCB-006 16.1 J 13.8 J 342 J 4.86 U 26 U 12.6 U 26.3 UJ 24.4 U PCB-007 PCB-007 PCB-007 10.1 J 13.8 J 342 J 4.86 U 26 U 58.3 J 31.5 J 86.5 J 20.4 U PCB-007 PCB-008 42.6 J 28.9 U 95.9 J 42 1690 941 2100 J 1190 J PCB-008 PCB-008 42.6 J 147 J 2.68 U 89.4 J 49.8 180 J 65.7 J PCB-010 PCB-010											
PCB-004 PCB-004 - 22.3 J 22.9 910 J 11.9 U 220 35.4 U 907 J 148 J PCB-005 PCB-005 - 10.5 U 4.17 U 17 UJ 2.9 U 22 U 12.6 U 26.3 UJ 24.4 U PCB-006 PCB-006 - 16.1 J 13.8 J 342 J 4.86 U 246 144 508 J 180 PCB-007 PCB-007 - 9.79 U 3.82 U 95.3 J 2.66 U 58.3 J 31.5 J 86.5 J 20.4 U PCB-007 PCB-008 - 4.26 J 2.89 U 3190 J 42 1690 941 2100 J 1190 J PCB-008 PCB-009 - 10.2 U 3.86 U 147 J 2.68 U 89.4 J 49.8 180 J 65.7 J PCB-010 PCB-010 - 6.3 U 3.83 U 24.6 J 2.74 U 18.6 U 14.5 U 26.5 J 17.8 U											
PCB-005 PCB-005 - 10.5 U 4.17 U 17 UJ 2.9 U 22 U 12.6 U 26.3 UJ 24.4 U PCB-006 PCB-006 - 16.1 J 13.8 J 342 J 4.86 U 246 144 508 J 180 PCB-007 PCB-007 - 9.79 U 3.82 U 95.3 J 2.66 U 58.3 J 31.5 J 86.5 J 20.4 U PCB-008 PCB-008 - 42.6 J 28.9 U 3190 J 42 1690 941 2100 J 1190 PCB-009 PCB-009 - 10.2 U 3.86 U 147 J 2.68 U 89.4 J 49.8 U 180 J 65.7 J PCB-010 PCB-010 - 63 U 3.83 U 24.6 J 2.74 U 18.6 U 14.5 U 26.5 J 17.8 U PCB-011 PCB-011 - 51.2 U 55.5 UJ 30.4 U 75.1 59.6 J 23.2 UJ 73.5 PCB-012/013 PCB-012/013 - 9.48 U<											
PCB-006 PCB-006 16.1 J 13.8 J 342 J 4.86 U 246 144 508 J 180 PCB-007 PCB-007 9.79 U 3.82 U 95.3 J 2.66 U 58.3 J 31.5 J 86.5 J 20.4 U PCB-008 PCB-008 42.6 J 28.9 U 3190 J 42 1690 941 2100 J 1190 J PCB-009 PCB-009 10.2 U 3.86 U 147 J 2.68 U 89.4 J 49.8 180 J 65.7 J PCB-010 PCB-010 6.3 U 3.83 U 24.6 J 2.74 U 18.6 U 14.5 U 26.5 J 17.8 U PCB-011 PCB-012/013 PCB-014 51.9 U 22.7 U 55.5 UJ 30.4 U 75.1 59.6 J 23.2 UJ 73.5 U PCB-014 PCB-014 9.48 U 10.6 J 65.4 J 8.55 J 135 91.8 23.4 UJ 119 PCB-014 PCB-014											
PCB-007 9.79 U 3.82 U 95.3 J 2.66 U 58.3 J 31.5 J 86.5 J 20.4 U PCB-008 PCB-008 42.6 J 28.9 U 3190 J 42 1690 941 2100 J 1190 PCB-009 PCB-009 10.2 U 3.86 U 147 J 2.68 U 89.4 J 49.8 180 J 65.7 J PCB-010 PCB-010 6.3 U 3.83 U 24.6 J 2.74 U 18.6 U 14.5 U 26.5 J 17.8 U PCB-011 51.9 U 22.7 U 55.5 UJ 30.4 U 75.1 59.6 J 23.2 UJ 73.5 PCB-012/013 PCB-012/013 9.48 U 10.6 J 65.4 J 8.55 J 135 91.8 23.4 UJ 119 PCB-014 9.45 U 3.66 U 14.7 UJ 2.54 U 18.2 U 10.4 U 22.7 UJ 20.2 U PCB-015 7.73 U 393 844 429 <											
PCB-008 PCB-008 42.6 J 28.9 3190 J 42 1690 941 2100 J 1190 PCB-009 PCB-009 10.2 U 3.86 U 147 J 2.68 U 89.4 J 49.8 180 J 65.7 J PCB-010 PCB-010 6.3 U 3.83 U 24.6 J 2.74 U 18.6 U 14.5 U 26.5 J 17.8 U PCB-011 PCB-011 51.9 22.7 U 55.5 UJ 30.4 U 75.1 59.6 J 23.2 UJ 73.5 PCB-012/013 PCB-012/013 9.48 U 10.6 J 65.4 J 8.55 J 135 91.8 23.4 UJ 119 PCB-014 PCB-014 9.45 U 3.66 U 14.7 UJ 2.54 U 18.2 U 10.4 U 22.7 UJ 20.2 U PCB-015 7.73 U 393 844 429 1230 604 215 899											
PCB-009 PCB-009 PCB-010 PCB-010 PCB-010 PCB-010 PCB-010 PCB-010 PCB-010 PCB-010 PCB-011 PCB-011 PCB-011 PCB-012 PCB-012 <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>											
PCB-010 PCB-010 6.3 U 3.83 U 24.6 J 2.74 U 18.6 U 14.5 U 26.5 J 17.8 U PCB-011 PCB-012 51.9 U 22.7 U 55.5 UJ 30.4 U 75.1 59.6 J 23.2 UJ 73.5 PCB-012/013 PCB-012/013 9.48 U 10.6 J 65.4 J 8.55 J 135 91.8 23.4 UJ 119 PCB-014 PCB-014 9.45 U 3.66 U 14.7 UJ 2.54 U 18.2 U 10.4 U 22.7 UJ 20.2 U PCB-015 PCB-015 7.73 U 393 844 429 1230 604 215 899											
PCB-011 PCB-011 - 51.9 22.7 U 55.5 UJ 30.4 U 75.1 59.6 J 23.2 UJ 73.5 PCB-012/013 PCB-012/013 - 9.48 U 10.6 J 65.4 J 8.55 J 135 91.8 23.4 UJ 119 PCB-014 PCB-014 - 9.45 U 3.66 U 14.7 UJ 2.54 U 18.2 U 10.4 U 22.7 UJ 20.2 U PCB-015 PCB-015 7.73 U 393 844 429 1230 604 215 899											
PCB-012/013 PCB-012/013 - 9.48 U 10.6 J 65.4 J 8.55 J 135 91.8 23.4 UJ 119 PCB-014 PCB-014 - 9.45 U 3.66 U 14.7 UJ 2.54 U 18.2 U 10.4 U 22.7 UJ 20.2 U PCB-015 PCB-015 7.73 U 393 844 429 1230 604 215 899											
PCB-014 PCB-014 - 9.45 U 3.66 U 14.7 UJ 2.54 U 18.2 U 10.4 U 22.7 UJ 20.2 U PCB-015 PCB-015 7.73 U 393 844 429 1230 604 215 899											
PCB-015 PCB-015 7.73 U 393 844 429 1230 604 215 899	·										
PCB-016 PCB-016 40.2 4.89 U 1130 6.65 U 610 241 2590 450											
	PCB-016	PCB-016		40.2	4.89 U	1130	6.65 U	610	241	2590	450

Appendix D Tissue Analytical Results

				1133UE AII	alytical Results				
	Location ID	PB010.3	PB013.1_A	PB013.3	PB013.3	PB015.1	PB015.2	PB015_A	PB015_A
	Sample ID	PB01-I-A-BCR-W-003-COMP-201106		PB01-F-A-GKF-W-004-COMP-201106	PB01-F-A-STM-W-003-20110616	PB01-F-A-GKF-W-002-COMP-201106	PB01-I-A-BCR-W-004-COMP-201106	PB01-F-A-GKF-W-001-COMP-20110	_
	Sample Date	6/19/2011	6/22/2011	6/21/2011	6/15/2011	6/15/2011	6/16/2011	6/25/2011	6/22/2011
	Sample Type	N	N	N	N	N	N	N N	N
	х	3201743	3201306.47	3201433.88	3201433.88	3201175	3201180	3201168.73	3201168.73
	V	13836152	13836191.23	13836075.12	13836075.12	13836251	13836183	13836254.66	13836254.66
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Fundulus grandis	Mugil cephalus	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Callinectes sapidus
	Size_Class	>2 to =<7.5	>2 to =<7.5	=<15	=<15	=<15	>2 to =<7.5	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Sedentary	Sedentary	Migratory	Sedentary	Sedentary	Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Conventional Parameters (pct)	5.6.145						1.000.0		1
Lipids	LIPID	1.73	1.15	2.94	1.94	3.57	2.84	3.64	0.983
Moisture, percent	MOIST	70.3	71.3	73.6	73.4	73.5	69.6	74.2	69.6
Metals (mg/kg)			-		-				
Lead	7439-92-1	0.345	0.13				0.114		0.518
Mercury	7439-97-6	0.037	0.046	0.109	0.04	0.074	0.046	0.107	0.053
Polyaromatic Hydrocarbons (μg/kg)	+					1			
2-Methylnaphthalene	91-57-6	6.97 U	14.1 U				7.17 U		14.6 U
Acenaphthene	83-32-9	6.97 U	11 J				7.17 U		14.6 U
Acenaphthylene	208-96-8	6.97 U	14.1 U				7.17 U		14.6 U
Anthracene	120-12-7	6.97 U	14.1 U				7.17 U		14.6 U
Benzo(a)anthracene	56-55-3	6.97 U	14.1 U				7.17 U		14.6 U
Benzo(a)pyrene	50-32-8	6.97 U	14.1 U				7.17 U		14.6 U
Benzo(b)fluoranthene	205-99-2	6.97 U	14.1 U				7.17 U		14.6 U
Benzo(g,h,i)perylene	191-24-2	6.97 U	14.1 U				7.17 U		14.6 U
Benzo(k)fluoranthene	207-08-9	6.97 U	14.1 U				7.17 U		14.6 U
Chrysene	218-01-9	6.97 U	14.1 U				7.17 U		14.6 U
Dibenzo(a,h)anthracene	53-70-3	6.97 U	14.1 U				7.17 U		14.6 U
Fluoranthene	206-44-0	6.97 U	14.1 U				7.17 U		14.6 U
Fluorene	86-73-7	6.97 U	14.1 U				7.17 U		14.6 U
Indeno(1,2,3-c,d)pyrene	193-39-5	6.97 U	14.1 U				7.17 U		14.6 U
Naphthalene	91-20-3	5.17 J	14.1 U				7.17 U		14.6 U
Phenanthrene	85-01-8	6.97 U	14.1 U				7.17 U		14.6 U
Pyrene	129-00-0	6.97 U	14.1 U				7.17 U		14.6 U
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N	6.97 U	14.1 U				7.17 U		14.6 U
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N	26.08 J	53.3 J				7.17 U		14.6 U
Total PAH (16) (U = 1/2)	tPAH_17_N	60.93 J	123.8 J				7.17 U		14.6 U
Semi-volatile Organic Carbons (μg/kg)	11.71127_11	00.50 1	12510 0				7.127 0		1
1,3-Dichlorobenzene	541-73-1	382 U	377 U	707 U	372 U	686 U	341 U	752 U	745 U
1,4-Dichlorobenzene	106-46-7	382 U	377 U				341 U		745 U
Hexachlorobenzene	118-74-1	268 J	345 J	759	1110	822	465	920	542 J
Hexachlorobutadiene	87-68-3	382 U	377 U				341 U		745 U
PCB Congeners (ng/kg)	07-00-3	302 0	377 0	-		-	341 0		743 0
PCB-001	PCB-001	1.47 U	2.49 J	199	99.5	235	5.82 U	201	5.93 U
PCB-002	PCB-002	1.38 U	0.847 U	8.1 J 40.9	12.4 J	12.1 J	1.6 U	6.32 J	5.47 U
PCB-003	PCB-003	1.5 U	0.816 U		38.4	50.3	4.67 U	28	5.66 U
PCB-004	PCB-004	13.1 U	16.3 U	2390	1180	2630	9.15 U	3020	33.6 U
PCB-005	PCB-005	3.74 U	3.21 U	64.1 J	46 J	80.6 J	4.38 U	87 J	7.15 U
PCB-006	PCB-006	7.72 U	2.74 U	1210	680	1210	3.82 U	1420	17.1 J
PCB-007	PCB-007	3.43 U	2.92 U	152	108 J	160	4.02 U	178	6.02 U
PCB-008	PCB-008	119	61.3	6950	3280	6890	54 J	7660	90.5
PCB-009	PCB-009	3.46 U	3.1 U	402	271	384	4.06 U	523	6.48 U
PCB-010	PCB-010	3.24 U	5.41 U	60.6	39.4	65.7	4.44 U	81.5 J	13.4 U
PCB-011	PCB-011	54.2 U	23.9 U	88.4	64.3 U	98.1 J	42.4 U	118	28.1 U
PCB-012/013	PCB-012/013	22.8	14.2 J	89.6	82.2 J	86.3	15.6 J	84.5 J	13.6
PCB-014	PCB-014	3.28 U	2.76 U	4.72 U	10.2 U	6.82 U	3.85 U	6.19 U	6.08 U
PCB-015	PCB-015	1110	1420	1300	606	1200	1550	1170	1210
PCB-016	PCB-016	10.4 U	7.09	8970	3960	7610	7.56 U	11000	24.5

				1133uc Alle	alytical Results				
	Location ID	PB015_A	PB015_A	PB015_A	PB016.2	PB019.1	PB019.1	PB019.1	PB019.1
			PB01-I-A-BCR-W-009-COMP-201106	PB01-I-B-BRS-W-001-COMP-201106	PB01-F-A-GKF-W-003-COMP-201106	PB01-F-A-SAS-W-005-20110622	PB01-I-B-BRS-W-012-20110621	PB01-I-B-BRS-W-014-20110621	PB01-I-B-BRS-W-015-20110621
	Sample Date	6/22/2011	6/22/2011	6/20/2011	6/15/2011	6/22/2011	6/21/2011	6/21/2011	6/21/2011
	Sample Type	N	N N	N	N	N	N	N	N
	х	3201168.73	3201168.73	3201168.73	3201190	3201152.87	3201152.87	3201152.87	3201152.87
	v	13836254.66	13836254.66	13836254.66	13836042	13835753.58	13835753.58	13835753.58	13835753.58
	Colontific Name								
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Penaeus aztecus	Fundulus grandis	Cynoscion arenarius	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus
	Size_Class		>2 to =<7.5	>7.5 to =<13	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13
	Tissue_Category	Sedentary	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Conventional Parameters (pct)	Lunin	1.00				0.004	1.51		
Lipids	LIPID	1.08	1.64	1.19	2.31	0.931	1.64	1.54	1.17
Moisture, percent	MOIST	72.1	66.9	74.6	73.9	79.2	74.4	75.1	76.2
Metals (mg/kg)	I=	2.107	0.455						
Lead	7439-92-1	0.107	0.157						
Mercury	7439-97-6	0.058	0.065	0.045	0.077	0.063	0.036	0.034	0.029
Polyaromatic Hydrocarbons (μg/kg)	la:								
2-Methylnaphthalene	91-57-6	14 U	14.6 U						
Acenaphthene	83-32-9	14 U	14.6 U						
Acenaphthylene	208-96-8	14 U	14.6 U		-	-			
Anthracene	120-12-7	14 U	14.6 U		-	-			
Benzo(a)anthracene	56-55-3	14 U	14.6 U						
Benzo(a)pyrene	50-32-8	14 U	14.6 U						
Benzo(b)fluoranthene	205-99-2	14 U	14.6 U						
Benzo(g,h,i)perylene	191-24-2	14 U	14.6 U						
Benzo(k)fluoranthene	207-08-9	14 U	14.6 U						
Chrysene	218-01-9	14 U	14.6 U						
Dibenzo(a,h)anthracene	53-70-3	14 U	14.6 U						
Fluoranthene	206-44-0	14 U	14.6 U						
Fluorene	86-73-7	14 U	14.6 U						
Indeno(1,2,3-c,d)pyrene	193-39-5	14 U	14.6 U						
Naphthalene	91-20-3	14 U	14.6 U	-	-	-			
Phenanthrene	85-01-8	14 U	14.6 U						
Pyrene	129-00-0	14 U	14.6 U						
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N	14 U	14.6 U						
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N	14 U	14.6 U						
Total PAH (16) (U = 1/2)	tPAH_17_N	14 U	14.6 U						
Semi-volatile Organic Carbons (μg/kg)	•								
1,3-Dichlorobenzene	541-73-1	348 U	734 U	709 U	714 U	353 U	341 U	344 U	340 U
1,4-Dichlorobenzene	106-46-7	348 U	734 U						
Hexachlorobenzene	118-74-1	386	638 J	913	470 J	325 J	710	618	591
Hexachlorobutadiene	87-68-3	348 U	734 U		-				
PCB Congeners (ng/kg)	<u> </u>	-	-						
PCB-001	PCB-001	4.06 U	3.82 U	55.2	104	176 J	128	109 J	92
PCB-002	PCB-002	3.55 U	3.73 U	5.36 J	6.6	19.1	47.2 U	18.4 U	25.6 U
PCB-002	PCB-002	3.53 U	4.08 U	31.7	29.7	72 J	47.2 U	62.6	52.3 J
PCB-003	PCB-003	16.1 U	108 U	239	1170	760 J	469	505	414
PCB-004	PCB-004	3.12 U	108 U	20.4 U	33.2	3.89 U	97.1 U	35.1 U	59.4 U
				262			640		
PCB-006	PCB-006	6.35 J	10.2 U		512	403		495	432
PCB-007	PCB-007	2.63 U	10.6 U	59.5 J	61.9	65.9	79.7 U	107 J	100
PCB-008	PCB-008	38.7	48.7 J	1810	3080	2380	4330	3340	2600
PCB-009	PCB-009	2.83 U	11.4 U	91.3 J	162	143	204	159	140
PCB-010	PCB-010	4.71 U	31.3 U	19.6 U	30.9	27.7	72 U	21.7 U	31.1 U
PCB-011	PCB-011	27.4 U	11.2 U	96.5	36.7 U	28.8 U	286 J	193 J	223
PCB-012/013	PCB-012/013	15.2	11.2 U	207	47.1 J	28.2	330	256	274
PCB-014	PCB-014	2.66 U	10.7 U	17 U	5.58 U	3.39 U	80.5 U	29.1 U	49.2 U
PCB-015	PCB-015	1410	2500	1560	538	191	2490	2140	1780
PCB-016	PCB-016	5.07	6.17	628	3380	3090	1450	1400	1230

				1.000.07	alytical Results				
	Location ID	PB019.1	PB019.1	PB019.1	PB019.2	PB019_A	PB019_A	PB024.1	PB024.1
	Sample ID	PB01-I-B-BRS-W-019-20110621	PB01-I-B-BRS-W-022-20110622	PB01-I-B-BRS-W-023-20110622	PB01-I-A-OYS-W-001-COMP-201106	PB01-F-A-GKF-W-005-COMP-201106	PB01-I-A-BCR-W-008-COMP-201106	PB01-F-A-GM-W-001-20110615	PB01-F-A-GM-W-002-20110615
	Sample Date	6/21/2011	6/22/2011	6/22/2011	6/28/2011	6/21/2011	6/23/2011	6/15/2011	6/15/2011
	Sample Type	N	N	N	N	N	N	N	N
	x	3201152.87	3201152.87	3201152.87	3201194.8	3201314.96	3201314.96	3201090.96	3201090.96
	v	13835753.58	13835753.58	13835753.58	13835735.1	13835750.59	13835750.59	13835160.37	13835160.37
	Scientific_Name	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus	Crassostrea virginica	Fundulus grandis	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus
	Size_Class	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Conventional Parameters (pct)	CAS IVAIIIDEI	nesure	Result	Result	Result	Result	nesan	nesure	resure
Lipids	LIPID	1.57	1.26	1.46	1.1	2.32	0.712	1.71	1.67
Moisture, percent	MOIST	74	76.1	74.9	82.7	74.9	70.8	78.3	77.6
Metals (mg/kg)	INOIST	,	70.1	74.3	02.7	74.3	76.5	70.3	77.5
Lead	7439-92-1				0.331		0.103		
Mercury	7439-97-6	0.025	0.049	0.048	0.051	0.091	0.071	0.186	0.199
Polyaromatic Hydrocarbons (μg/kg)	7433 37 0	0.023	0.043	0.040	0.031	0.031	0.071	0.100	0.193
2-Methylnaphthalene	91-57-6				13.7 U		14.5 U		
	83-32-9				13.7 U		14.5 U		
Acenaphthylone	208-96-8				13.7 U		14.5 U		
Acenaphthylene									
Anthracene	120-12-7				14.5		14.5 U		
Benzo(a)anthracene	56-55-3				38.5		14.5 U		
Benzo(a)pyrene	50-32-8				13.8	_	14.5 U		
Benzo(b)fluoranthene	205-99-2				37.5		14.5 U		
Benzo(g,h,i)perylene	191-24-2				9.32 J	-	14.5 U		
Benzo(k)fluoranthene	207-08-9				13.8		14.5 U		
Chrysene	218-01-9				68.8		14.5 U		
Dibenzo(a,h)anthracene	53-70-3				13.7 U		14.5 U		
Fluoranthene	206-44-0				118		14.5 U		
Fluorene	86-73-7				13.7 U		14.5 U		
Indeno(1,2,3-c,d)pyrene	193-39-5				13.7 U		14.5 U		
Naphthalene	91-20-3				13.7 U		14.5 U		
Phenanthrene	85-01-8				18.7		14.5 U		
Pyrene	129-00-0				162		14.5 U		
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N				475.42 J		14.5 U		
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N				76.7		14.5 U		
Total PAH (16) (U = 1/2)	tPAH_17_N				552.12 J		14.5 U		
Semi-volatile Organic Carbons (μg/kg)									
1,3-Dichlorobenzene	541-73-1	363 U	383 U	388 U	704 U	725 U	371 U	362 U	341 U
1,4-Dichlorobenzene	106-46-7				704 U		371 U		
Hexachlorobenzene	118-74-1	679	841	794	704 U	2360	927	671	552
Hexachlorobutadiene	87-68-3				704 U		371 U		
PCB Congeners (ng/kg)	1				10.0				
PCB-001	PCB-001	88.2 J	318	446	39.2	240	1.31 U	118	90.2
PCB-002	PCB-001	27 U	38.2 J	61.1 J	9.43	16.6 J	3.99 U	12.5	7.43 U
							3.99 U 4.02 U		
PCB-003	PCB-003	26.5 U	150	216	24.1	58.8		44	43.8
PCB-004	PCB-004	392	683	698	1000	2770	30.1 U	1070	1030
PCB-005	PCB-005	52.7 U	6.76 U	55.7 J	31.6 J	75.3	7.16 U	36.7	35.7
PCB-006	PCB-006	529 J	580	638	506	1070	5.78 U	646	627
PCB-007	PCB-007	94.5 J	103	119 J	84.5	148 J	6.03 U	97.4	95.5 J
PCB-008	PCB-008	3380	3280	3800	2340	6130	95.4	3400	3310
PCB-009	PCB-009	137 J	175	190	188	461	6.5 U	244	234
PCB-010	PCB-010	36.2 U	33.3	33.2 J	23.5	35.6 U	10.5 U	34.4 J	31.7
PCB-011	PCB-011	249	208	217	169	103	29.2 U	89.3 U	81.8 U
PCB-012/013	PCB-012/013	324	303	316 J	287	92.2 J	18.8 J	103	76.3 J
PCB-014	PCB-014	43.7 U	5.68 U	17.8 U	6.08 U	16.8 U	6.09 U	10.4 U	20.1 U
PCB-015	PCB-015	2260	1990	1990	1670	1080	1620	367 J	320
PCB-016	PCB-016	1330	2030	1970	3650	6400	11.9	4570	4480

				1133UC AII	alytical Results				
	Location ID	PB024.4	PB025_A	PB026.1	PB032_A	PB037.2	PB037.4	PB037.5	PB037.6
	Sample ID	PB01-I-A-BCR-W-005-COMP-201106	PB01-F-A-GM-W-004-20110615	PB01-F-A-STM-W-002-20110616	PB02-F-A-PNF-W-009-20110617	PB02-F-A-STM-W-006-20110616	PB02-I-B-BCR-W-001-COMP-201106	PB02-I-B-BRS-W-001-COMP-201106	PB02-I-A-BCR-W-002-COMP-20110
	Sample Date	6/20/2011	6/15/2011	6/15/2011	6/17/2011	6/16/2011	6/17/2011	6/17/2011	6/16/2011
	Sample Type		N	N	N	N	N	N	N
	х х	3201319	3201214.42	3201153.65	3201407.03	3201346.97	3201504	3201380	3201402
	Y	13835188	13835154.42	13835079.21	13834542.25	13834038.02	13833530	13834151	13833775
	Scientific_Name		Brevoortia patronus	Mugil cephalus	Lagodon rhomboides	Mugil cephalus	Callinectes sapidus	Penaeus aztecus	Callinectes sapidus
	Size_Class	· ·	=<15	=<15	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Migratory	Migratory	Migratory	Migratory	Migratory	Sedentary
	Tissue_Type	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Conventional Parameters (pct)	0.0.10	1111111			1.000.0				
Lipids	LIPID	1.76	1.78	2.36	2.69	2.06	0.943	1.15	1.68
Moisture, percent	MOIST	69.1	77.5	74.3	74.2	73.6	69	75.1	70.6
Metals (mg/kg)			-	-				-	
Lead	7439-92-1	0.212							0.292
Mercury	7439-97-6	0.065	0.142	0.036	0.083	0.028	0.083	0.065	0.05
Polyaromatic Hydrocarbons (µg/kg)	1								
2-Methylnaphthalene	91-57-6	7.45 U							7.77 U
Acenaphthene	83-32-9	5.72 J							7.77 U
Acenaphthylene	208-96-8	7.45 U							7.77 U
Anthracene	120-12-7	7.45 U							7.77 U
Benzo(a)anthracene	56-55-3	7.45 U							7.77 U
Benzo(a)pyrene	50-32-8	7.45 U							7.77 U
Benzo(b)fluoranthene	205-99-2	7.45 U							7.77 U
Benzo(g,h,i)perylene	191-24-2	7.45 U							7.77 U
Benzo(k)fluoranthene	207-08-9	7.45 U							7.77 U
Chrysene	218-01-9	7.45 U							7.77 U
Dibenzo(a,h)anthracene	53-70-3	7.45 U							7.77 U
Fluoranthene	206-44-0	7.45 U							7.77 U
Fluorene	86-73-7	7.45 U							7.77 U
Indeno(1,2,3-c,d)pyrene	193-39-5	7.45 U							7.77 U
Naphthalene	91-20-3	6.03 J							8.23
Phenanthrene	85-01-8	3.81 J							7.77 U
Pyrene	129-00-0	6.08 J							7.77 U
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N	39.61 J							7.77 U
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N	30.46 J							31.54
Total PAH (16) (U = 1/2)	tPAH_17_N	70.07 J							70.39
Semi-volatile Organic Carbons (µg/kg)	0.7.112711	70.07							7 0.03
1,3-Dichlorobenzene	541-73-1	371 U	348 U	342 U	765 U	694 U	357 U	699 U	354 U
1,4-Dichlorobenzene	106-46-7	371 U							354 U
Hexachlorobenzene	118-74-1	5430	416	1830	2290	997	357 U	537 J	354 U
Hexachlorobutadiene	87-68-3	371 U							354 U
PCB Congeners (ng/kg)	07-00-3	3/1 0		-		-			334 0
PCB-001	PCB-001	7.83 U	151	134 J	142 J	190	3.17 U	71.3	2.25 U
PCB-002	PCB-002	7.19 U	13.4	13.3 J 36.4	9.38 U	6.75 U	2.92 U	1.97 U	1.96 U
PCB-003	PCB-003	7.68 U	34.9		8.24 U	53.8 J	3.02 U	34.9	1.98 U
PCB-004	PCB-004	147	1730	1480	1190 J	2090	39.2 U	290	23.9 U
PCB-005	PCB-005	18.1 U	57.7 J	67.7	35.8 UJ	77.1 J	9.87 U	9.08 U	4.43 U
PCB-006	PCB-006	78.6 J	1040	829	358 J	1170	7.97 U	295	9.76 U
PCB-007	PCB-007	16.6 U	161	136	31.1 UJ	209	8.31 U	60.1	4.07 U
PCB-008	PCB-008	369	4970	4060	1480 J	6020	119	1940	103
PCB-009	PCB-009	16.7 U	392	353	275 J	495	8.95 U	82	4.1 U
PCB-010	PCB-010	12.2 U	51.1	54.7	36.5 J	78	15.5 U	13.2	3.91 U
PCB-011	PCB-011	57.8 U	58 U	88.3 U	31.5 UJ	107 U	29.2 U	83.5	47.3 U
PCB-012/013	PCB-012/013	56.7 J	109	89.1	31.8 UJ	126	25.5 J	180	17.1
PCB-014	PCB-014	15.8 U	9.34 U	12 U	30.8 UJ	16.4 U	8.39 U	7.59 U	3.89 U
PCB-015	PCB-015	1880	557	438 J	116	1120	1580	1460	1450
PCB-016	PCB-016	94.4	7810	5990	1640	7440	12.8 J	573	8.73 U

				TISSUE And	alytical Results				
	Location ID	PB037_A	PB047.1_A	PB047.3	PB048.3	PB048.3	PB048.3	PB049_A	PB049_A
	Sample ID	PB02-I-A-BCR-W-004-COMP-201106	PB02-F-A-PNF-W-002-20110615	PB02-I-A-BCR-W-001-COMP-201106	PB02-F-A-GM-W-001-20110616	PB02-F-A-GM-W-002-20110616	PB02-F-A-STM-W-001-20110616	PB02-F-A-GKF-W-001-COMP-201106	PB02-I-A-BCR-W-003-COMP-20110
	Sample Date	6/19/2011	6/15/2011	6/19/2011	6/16/2011	6/16/2011	6/16/2011	6/15/2011	6/22/2011
	Sample Type	N	N	N	N	N	N	N	N
	. ,,	3201325.97	3201461.79	3201645	3201519.15	3201519.15	3201519.15	3201468.39	3201468.39
	γ	13833885.45	13833007.68	13832925	13832869.89	13832869.89	13832869.89	13832722.8	13832722.8
	Scientific_Name		Lagodon rhomboides	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus	Mugil cephalus	Fundulus grandis	Callinectes sapidus
	Size_Class	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15	=<15	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Conventional Parameters (pct)									
Lipids	LIPID	1.14	2.34	1.7	2	1.73	1.78	3.14	1.63
Moisture, percent	MOIST	72.6	73.2	69.1	79.4	79.9	73.7	75.4	66.1
Metals (mg/kg)									
Lead	7439-92-1	0.138		0.099			-		0.263
Mercury	7439-97-6	0.033	0.034	0.046	0.071	0.078	0.041	0.068	0.072
Polyaromatic Hydrocarbons (μg/kg)	+								
2-Methylnaphthalene	91-57-6	14.2 U		7.09 U					14.8 U
Acenaphthene	83-32-9	14.2 U		7.09 U					14.8 U
Acenaphthylene	208-96-8	14.2 U		7.09 U	-				14.8 U
Anthracene	120-12-7	14.2 U		7.09 U					14.8 U
Benzo(a)anthracene	56-55-3	14.2 U		7.09 U					14.8 U
Benzo(a)pyrene	50-32-8	14.2 U		7.09 U					14.8 U
Benzo(b)fluoranthene	205-99-2	14.2 U		7.09 U					14.8 U
Benzo(g,h,i)perylene	191-24-2	14.2 U		7.09 U					14.8 U
Benzo(k)fluoranthene	207-08-9	14.2 U		7.09 U					14.8 U
Chrysene	218-01-9	14.2 U		7.09 U					14.8 U
Dibenzo(a,h)anthracene	53-70-3	14.2 U		7.09 U					14.8 U
Fluoranthene	206-44-0	14.2 U		7.09 U					14.8 U
Fluorene	86-73-7	14.2 U		7.09 U					14.8 U
Indeno(1,2,3-c,d)pyrene	193-39-5	14.2 U		7.09 U					14.8 U
Naphthalene	91-20-3	14.2 U		4.19 J					14.8 U
Phenanthrene	85-01-8	14.2 U		7.09 U					14.8 U
Pyrene	129-00-0	14.2 U		7.09 U					14.8 U
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N	14.2 U		7.09 U					14.8 U
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N	14.2 U		25.46 J					14.8 U
Total PAH (16) (U = 1/2)	tPAH_17_N	14.2 U		60.91 J					14.8 U
Semi-volatile Organic Carbons (μg/kg)	U.7.127_1	1.12 0		00.513					1
1,3-Dichlorobenzene	541-73-1	366 U	671 U	368 U	385 U	697 U	769 U	741 U	727 U
1,4-Dichlorobenzene	106-46-7	366 U		368 U					727 U
Hexachlorobenzene	118-74-1	366 U	 447 J	191 J	398	464 J	460 J	741 U	727 U
Hexachlorobutadiene	87-68-3	366 U	447 J 	368 U		404 J	400 J	741 0	727 U
PCB Congeners (ng/kg)	07-00-3	300 0		300 0	=		-		121 0
PCB-001	PCB-001	1.32 U	127	6.24 U	210	180	89.9	289	2.08 U
PCB-001 PCB-002	PCB-001	1.32 U 1.09 U	8.79 U	5.6 U	12.1 J	5.56 U	6.11 U	13.6 J	0.956 J
				5.84 U					0.956 J 1.86 U
PCB-003	PCB-003	1.75 U	19.5 J		41	35.8 J	29.6 J	58.1	
PCB-004	PCB-004	16.1 U	1230	21.3 U	2550	2260	1350	2910	7.01 U
PCB-005	PCB-005	2.68 U	26.5 U	13 U	76.2	75.9	18.6 U	95.2	1.65 U
PCB-006	PCB-006	2.29 U	441	11.4 U	1350	1260	855	1190	5.49 U
PCB-007	PCB-007	2.44 U	50.5	12 U	211	184	141	162	1.47 U
PCB-008	PCB-008	64.6	1360	70.2	6630	5920	4350	8020	47.6
PCB-009	PCB-009	2.58 U	371	12.1 U	509	469	396	499	1.51 U
PCB-010	PCB-010	5.46 U	39.7	12.2 U	81.5	78.1	57.2	81.5 J	3.08 U
PCB-011	PCB-011	19.6 U	111 U	35 U	149 U	72 U	110 U	98.9	31.3 U
PCB-012/013	PCB-012/013	13.9	38.6 J	11.7 U	133	118	109 J	110	16.5
PCB-014	PCB-014	2.31 U	21.5 U	11.4 U	9.56 U	8.96 U	15.1 U	6.29 U	1.42 U
PCB-015	PCB-015	3990	113 J	1780	440	363	1150	1230	2210
PCB-016	PCB-016	12.6 J	1900	10.6 U	9620	8370	6410	8590	5.76 U

				Tissue And	alytical Results				
	Location ID	PB052.1	PB052.1	PB053.2	PB053.2	PB053_B	PB056_A	PB056_A	PB057.2
	Sample ID	PB02-F-A-SAS-W-002-20110616	PB02-F-A-STM-W-003-20110616	PB02-F-A-GKF-W-025-20110621	PB02-F-A-GKF-W-029-20110622	PB02-F-A-GKF-W-013-20110618	PB02-F-A-GKF-W-010-20110618	PB02-F-A-GKF-W-011-20110618	PB03-F-A-GM-W-001-20110617
	Sample Date	6/16/2011	6/16/2011	6/21/2011	6/22/2011	6/18/2011	6/18/2011	6/18/2011	6/17/2011
	Sample Type	N	N	N N	N	N	N	N	N
	,	3201515.24	3201515.24	3201266.16	3201266.16	3201390.48	3201420.22	3201420.22	3201603.32
	γ	13832564.31	13832564.31	13832433.69	13832433.69	13832440.54	13832183.54	13832183.54	13832054.52
	Scientific_Name	Cynoscion arenarius	Mugil cephalus	Fundulus grandis	Brevoortia patronus				
	Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
	Tissue_Category	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Migratory
	Tissue_Type	Whole Body							
Chemical	CAS Number	Result							
Conventional Parameters (pct)									
Lipids	LIPID	1.18	1.76	3.33	2.14	2.75	2.36	2.68	2.02
Moisture, percent	MOIST	78.7	75.4	73	75.2	74.1	74.2	73.1	80
Metals (mg/kg)	l								
Lead	7439-92-1								
Mercury	7439-97-6	0.045	0.024	0.045	0.075	0.058	0.049	0.051	0.068
Polyaromatic Hydrocarbons (μg/kg)	'								
2-Methylnaphthalene	91-57-6								
Acenaphthene	83-32-9								
Acenaphthylene	208-96-8								
Anthracene	120-12-7								
Benzo(a)anthracene	56-55-3								
Benzo(a)pyrene	50-32-8								
Benzo(b)fluoranthene	205-99-2								
Benzo(g,h,i)perylene	191-24-2								
Benzo(k)fluoranthene	207-08-9								
Chrysene	218-01-9								
Dibenzo(a,h)anthracene	53-70-3								
Fluoranthene	206-44-0								
Fluorene	86-73-7								
Indeno(1,2,3-c,d)pyrene	193-39-5								
Naphthalene	91-20-3								
Phenanthrene	85-01-8								
Pyrene	129-00-0								
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N								
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N								
Total PAH (16) (U = 1/2)	tPAH_17_N								
Semi-volatile Organic Carbons (μg/kg)	'								
1,3-Dichlorobenzene	541-73-1	370 U	721 U	365 U	372 U	763 U	351 U	390 U	765 U
1,4-Dichlorobenzene	106-46-7								
Hexachlorobenzene	118-74-1	240 J	416 J	243 J	224 J	763 U	369	200 J	765 U
Hexachlorobutadiene	87-68-3								
PCB Congeners (ng/kg)	'								
PCB-001	PCB-001	59.7	237	313	291	308	312 J	296	147
PCB-002	PCB-002	8.34 U	9.78 U	13.2	12.2 J	5.88 UJ	11.4 U	12.2 U	11.9 J
PCB-003	PCB-003	6.42 U	62.5	51.2	55.6 J	66.9 J	62.2	61.5 J	31
PCB-004	PCB-004	767	2870	5070 J	4610	5280 J	4270 J	4450 J	2250 J
PCB-005	PCB-005	19.7 U	127 J	133 J	127	121 J	40.3 UJ	34.3 UJ	16.5 U
PCB-006	PCB-006	570	1640	2500 J	2130	2110 J	1480 J	1990 J	1360
PCB-007	PCB-007	91.4	303	316 J	267	290 J	214 J	269 J	214
PCB-008	PCB-008	3070	8400	19100 J	15500	13900 J	11400 J	13000 J	4520
PCB-009	PCB-009	217	757	852 J	714	756 J	503 J	698 J	506
PCB-009	PCB-010	29.3	110	138 J	121	118 J	97.9 J	128 J	75.1
PCB-010 PCB-011	PCB-010 PCB-011	92.2 U	176 U	110 J	115	80.8 J	1170 J	1180 J	75.1 55.7 U
PCB-011 PCB-012/013	PCB-012/013	16.4 U	248	170 J	180	191 J	170 J	160 J	129
PCB-012/013 PCB-014	PCB-012/013 PCB-014	16.4 U	248 25.2 U	9.23 UJ	7.58 U	191 J 11.5 UJ	34.8 UJ	29.6 UJ	129 14.4 U
PCB-014 PCB-015	PCB-014 PCB-015	88	25.2 U	1370	7.58 U 1240	11.5 UJ 1960 J	34.8 UJ 1640	1640	598
PCB-016	PCB-016	4200	12000	21200 J	16400	14300 J	9620	9740	8350

				1133UC Allu	lytical Results				
	Location ID Sample ID	PB057.2 PB03-F-A-STM-W-001-20110617	PB057.2 PB03-F-A-STM-W-001-COMP-201106	PB057.2 PB03-F-A-STM-W-002-20110617	PB057_B PB03-F-A-GKF-W-001-20110615	PB057_B PB03-F-A-GKF-W-006-20110617	PB057_B PB03-F-A-PNF-W-001-COMP-201106	PB058.1 PB03-F-A-GM-W-003-20110618	PB058.1 PB03-F-A-GM-W-004-20110618
	Sample Date	6/17/2011	6/17/2011	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/18/2011	6/18/2011
	Sample Type	0/17/2011 N	0/1//2011 N	0/17/2011 N	0/13/2011 N	0/1//2011 N	0/13/2011 N	0/18/2011 N	6/16/2011 N
	Sample Type	3201603.32	3201603.32	3201603.32	3201644.06	3201644.06	3201644.06	3201682.3	3201682.3
	Ŷ	13832054.52	13832054.52	13832054.52	13832071.63	13832071.63	13832071.63	13832011.93	13832011.93
	Coloratica Name		1						
	Scientific_Name	Mugil cephalus	Mugil cephalus	Mugil cephalus	Fundulus grandis	Fundulus grandis	Lagodon rhomboides	Brevoortia patronus	Brevoortia patronus
	Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Migratory	Migratory	Migratory
al	Tissue_Type	Whole Body Result	Whole Body	Whole Body Result	Whole Body Result	Whole Body	Whole Body	Whole Body Result	Whole Body
Chemical Conventional Parameters (pct)	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
.,	LIPID	1.4	1.78	1.34	2.68	2.52	2.49	1.68	1.26
Lipids Moisture persent		74.5				74.7	74.9		78.9
Moisture, percent	MOIST	74.5	73.5	74.6	74	74.7	74.9	79.2	78.9
Metals (mg/kg)	7439-92-1								
Lead	7439-92-1	0.047	0.122	0.041	0.044	0.044	0.03	0.075	0.116
Mercury	7439-97-6	0.047	0.122	0.041	0.044	0.044	0.03	0.075	0.116
Polyaromatic Hydrocarbons (µg/kg)	01 57 6		 						
2-Methylnaphthalene	91-57-6								
Acenaphthulana	83-32-9 208-96-8								
Acenaphthylene Anthracene	208-96-8 120-12-7								
Benzo(a)anthracene	56-55-3								
Benzo(a)pyrene	50-32-8 205-99-2								
Benzo(b)fluoranthene	191-24-2		-						
Benzo(g,h,i)perylene Benzo(k)fluoranthene	207-08-9								
	218-01-9								
Chrysene Dibenzo(a,h)anthracene	53-70-3								
Fluoranthene	206-44-0								
Fluorene	86-73-7								
Indeno(1,2,3-c,d)pyrene	193-39-5								
Naphthalene	91-20-3								
Phenanthrene	85-01-8								
Pyrene	129-00-0								
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N								
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N								
Total PAH (16) (U = 1/2)	tPAH_17_N			-					
Semi-volatile Organic Carbons (μg/kg)		-		_		-			-
1,3-Dichlorobenzene	541-73-1	727 U	753 U	742 U	360 U	716 U	708 U	699 U	718 UJ
1,4-Dichlorobenzene	106-46-7					-			-
Hexachlorobenzene	118-74-1	727 U	556 J	742 U	271 J	716 U	708 U	699 U	718 UJ
Hexachlorobutadiene	87-68-3								
PCB Congeners (ng/kg)									
PCB-001	PCB-001	78.5 J	125	80	271	313 J	106	175	283
PCB-002	PCB-002	16.4 U	4.26 J	7.23 U	R	14.4 J	1.3 U	11.8	9.1 J
PCB-003	PCB-003	14.2 U	27.5	31.3 J	44.5 J	61.7	11 J	43.3	57
PCB-004	PCB-004	992 J	1260	1240 J	4580	4980	1120	2610 J	3330
PCB-005	PCB-005	41.4 UJ	49.8	17.9 U	147	185	20.4	28.7 U	103
PCB-006	PCB-006	495 J	634	717	1950	2310	345	1600	1620
PCB-007	PCB-007	90.1 J	112	145	222 J	309	44 J	256	221
PCB-008	PCB-008	2420 J	3390	3860	12800	13600	1410	5250	9140
PCB-009	PCB-009	241 J	263	314	635	808	237	562	520
PCB-010	PCB-010	31.7 J	35.2	46.5	162 J	151	31.7 J	84.6	112
PCB-011	PCB-011	33.3 UJ	39.8 U	50.5 U	99.7	171 U	24.3 U	70.3 J	49.5
PCB-012/013	PCB-012/013	33.5 UJ	75.9	114 J	140 J	173	23.8 J	245	194
PCB-014	PCB-014	32.7 UJ	2.52 U	15.6 U	28 U	11 U	3.3 U	25 U	4.91 U
	PCB-015	610	805	554	977	1260	172	944	1540
PCB-015	PCD-013	010		334	3//		1/2	344	

				Tissue Alla	llytical Results				
	Location ID	PB058.1	PB059.1_A	PB059.1_A	PB059.3	PB059.4	PB059.4	PB059.5	PB063.1_A
	Sample ID	PB03-F-A-GM-W-005-20110618	PB03-F-A-GKF-W-003-20110615	PB03-I-A-BCR-W-003-COMP-201106	PB03-I-B-BCR-W-002-20110627	PB03-I-B-BCR-W-003-20110628	PB03-I-B-BCR-W-003-20110628DUP	PB03-I-A-BCR-W-001-COMP-201106	PB03-F-A-GKF-W-009-20110617
	Sample Date	6/18/2011	6/15/2011	6/19/2011	6/26/2011	6/27/2011	6/27/2011	6/27/2011	6/17/2011
	Sample Type	N	N	N	N	N	N	N	N
	x	3201682.3	3201501.32	3201501.32	3201714.16	3201714.16	3201714.16	3201714	3201509.99
	Υ	13832011.93	13831828.71	13831828.71	13831964.63	13831964.63	13831964.63	13831965	13831505.57
	Scientific_Name	Brevoortia patronus	Fundulus grandis	Callinectes sapidus	Callinectes sapidus	Callinectes sapidus		Callinectes sapidus	Fundulus grandis
	Size_Class	=<15	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13		>2 to =<7.5	=<15
	Tissue_Category	Migratory	Sedentary	Sedentary	Migratory	Migratory		Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body		Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Conventional Parameters (pct)									
Lipids	LIPID	1.63	2.26	1.04	0.575	0.658		0.555	2.81
Moisture, percent	MOIST	79.2	74.3	71.3	73.7	76.5		72.8	74.4
Metals (mg/kg)	L								
Lead	7439-92-1			0.092	,			0.113	
Mercury	7439-97-6	0.149	0.025	0.032	0.059	0.058		0.045	0.032
Polyaromatic Hydrocarbons (µg/kg)	•								
2-Methylnaphthalene	91-57-6			14.5 U			-	14.8 U	
Acenaphthene	83-32-9			14.5 U			-	14.8 U	
Acenaphthylene	208-96-8			14.5 U			-	14.8 U	
Anthracene	120-12-7			14.5 U			-	14.8 U	
Benzo(a)anthracene	56-55-3			14.5 U	,			14.8 U	-
Benzo(a)pyrene	50-32-8			14.5 U				14.8 U	
Benzo(b)fluoranthene	205-99-2			14.5 U				14.8 U	
Benzo(g,h,i)perylene	191-24-2			14.5 U				14.8 U	
Benzo(k)fluoranthene	207-08-9			14.5 U				14.8 U	
Chrysene	218-01-9			14.5 U				14.8 U	
Dibenzo(a,h)anthracene	53-70-3			14.5 U				14.8 U	
Fluoranthene	206-44-0			14.5 U				14.8 U	
Fluorene	86-73-7			14.5 U				14.8 U	
Indeno(1,2,3-c,d)pyrene	193-39-5			14.5 U				14.8 U	
Naphthalene	91-20-3			14.5 U				14.8 U	
Phenanthrene	85-01-8			14.5 U				14.8 U	
Pyrene	129-00-0			14.5 U				14.8 U	
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N			14.5 U				14.8 U	
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N			14.5 U				14.8 U	
Total PAH (16) (U = 1/2)	tPAH_17_N			14.5 U				14.8 U	
Semi-volatile Organic Carbons (μg/kg)									
1,3-Dichlorobenzene	541-73-1	727 UJ	377 U	363 U	718 UJ	748 UJ		347 U	351 U
1,4-Dichlorobenzene	106-46-7			363 U				347 U	
Hexachlorobenzene	118-74-1	727 UJ	377 U	363 U	718 UJ	748 UJ		347 U	351 U
Hexachlorobutadiene	87-68-3			363 U			-	347 U	
PCB Congeners (ng/kg)	!								
PCB-001	PCB-001	244	351	2.15 U	4.9 J	4.39 J	3.04 U	2.74 U	241
PCB-002	PCB-002	10.5 J	18.7	1 J	2.52 J	0.766 U	2.58 U	2.75 U	15.2 U
PCB-003	PCB-003	58.2	74.3	2.06 U	4.19 U	2.76 U	2.54 U	2.7 U	61.3 J
PCB-004	PCB-004	3320	5900	12 J	30.3	39.8	38	23.8 UJ	4480 J
PCB-005	PCB-005	111	84.3 U	3.98 U	2.31 U	2.01 U	4.23 U	6.53 UJ	45.2 U
PCB-006	PCB-006	1610	2660	3.52 U	15.6 J	23.4	22.9	5.58 UJ	2080
PCB-007	PCB-007	223	350 J	3.65 U	1.96 U	1.71 U	3.88 U	5.94 UJ	290
PCB-008	PCB-008	9420	18100	76.1	65.4	117	109	61.1 J	11800
PCB-009	PCB-009	522	872	3.72 U	4.37 U	5.11 U	5 U	6.3 UJ	787
PCB-010	PCB-010	114	187	2.33 U	1.55 U	1.3 U	2.3 U	7.88 UJ	157
PCB-010	PCB-010	45.9 U	71.4 U	2.33 U	22.8 U	1.3 U	22.5 U	21.4 UJ	113 J
PCB-011 PCB-012/013	PCB-011 PCB-012/013	198	71.4 U	19.1 J	20.6	17.8 0	16.6 J	16.1 J	207
PCB-012/013 PCB-014	PCB-012/013 PCB-014	4.23 U	70.4 U	3.33 U	1.95 U	15.9 1.7 U	3.73 U	5.62 UJ	37.5 U
PCB-014 PCB-015	PCB-014 PCB-015	1560	1500	3.33 U 1680	1.95 U	1.7 0	3.73 U 1690	1590	1850
PCB-016	PCB-016	20800	17400	9.6 J	24.4	25.5	25.4 J	17	12200

Lipids LiPiD 2.76 1.76 1.65 1.65 3.4 2.58 1.18 2.77 2.75 Moisture, percent MOIST 75.3 74.3 74.6 71.9 71.9 71.9 71.9 71.9 71.9 71.9 71.9					Hissue Alle	alytical Results				
Second Second		Location ID	PB063.1_A	PB063.2_A	PB063.2_A	PB069_A	PB069_A	PB069_A	PB084.1	PB084.1
Separate Parameter Param		Sample ID	PB03-F-A-GKF-W-010-20110617	PB03-F-A-GKF-W-005-20110615	PB03-F-A-GKF-W-013-20110617	PB03-F-A-GKF-W-002-20110615	PB03-F-A-GKF-W-007-20110617	PB03-I-A-BCR-W-002-COMP-201106	PB04-F-A-GKF-W-005-20110627	PB04-F-A-GKF-W-013-20110627
March Marc		Sample Date	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/17/2011	6/19/2011	6/27/2011	6/27/2011
March Marc		Sample Type	N	N	N	N	N	N	N	N
Profession Pr		х		3201561.28	3201561.28	3201132.36	3201132.36	3201132.36	3200727.85	3200727.85
Profession Pr		Υ	13831505.57	13831519.55	13831519.55	13830993.22	13830993.22	13830993.22	13829784.47	13829784.47
Second Second		Scientific Name								
Fig. Solicity So		_	•	Ĭ	_	1	1	·	-	_
Time										
Profession Pro			-	· · · · · · · · · · · · · · · · · · ·	•	· ·	· ·	1	•	· ·
Section 1900	Chemical		•	, , , , , , , , , , , , , , , , , , ,	,	·	· '	· ' +	•	•
March Supplied March March March March March March Supplied Conventional Parameters (pct)										
Memory	Lipids	LIPID	2.76	1.76	1.65	3.4	2.58	1.18	2.77	2.75
March Marc	Moisture, percent	MOIST	75.3	74.3	74.6	71.9	74.3	70	75.7	75.9
Money Mone	Metals (mg/kg)	l								
New New	Lead	7439-92-1	-					0.117		
Manufactories \$2.50 \$			0.04	0.044	0.036	0.027 J	0.028	0.025	0.046	0.042
Manual	Polyaromatic Hydrocarbons (µg/kg)									
Manual		91-57-6						14.3 U		
Montherwork 1988										
ABT ABT	·									
Person-Different Spi-296	Anthracene		-					14.3 U		
Bases Bases Bases Fine	Benzo(a)anthracene	56-55-3						14.3 U		
Process 19-52	Benzo(a)pyrene	50-32-8	-					14.3 U		
Second Progress 1949 1940 194		205-99-2						14.3 U		
Company Com	Benzo(g,h,i)perylene	191-24-2	-		-			14.3 U		
Demonsharemen 19-94 19-9	Benzo(k)fluoranthene	207-08-9						14.3 U		
Househole 10,440	Chrysene	218-01-9						14.3 U		
Process Proc	Dibenzo(a,h)anthracene	53-70-3						14.3 U		
Biother 13-29-5	Fluoranthene	206-44-0						14.3 U		
Nach Principle 12-0-3	Fluorene	86-73-7	-					14.3 U		
Note Note	Indeno(1,2,3-c,d)pyrene	193-39-5						14.3 U		
Promishing Sol	91-20-3						14.3 U			
Pyrene										
Total PAR (19 16) PAR 17 19 PAR 17 1	Pyrene									
Total Park (15) (11 - 17)	· ·									
Total PAM [18] U = 1/2)										
Semi-selfa Organic Carbons (ig/Ng)	, ,, ,,									
1.3-Dehinonbenzene \$41-73-1 384 U 361 U 735 U 364 U 360 U 370 U 722 UJ 748 UJ 1.4-Dehinonbenzene 118-74-1 384 U 361 U 735 U 364 U 366 U 370 U 722 UJ 748 UJ Hexachiorobutadiene 18-74-1 384 U 361 U 735 U 364 U 366 U 370 U 722 UJ 748 UJ Hexachiorobutadiene 87-68-3 -		10.7.11.227211						1.00		
1,4 Olchlorobenzene 106.46-7 — — — — — 370 U — <th< td=""><td></td><td>541-73-1</td><td>384 U</td><td>361 U</td><td>735 U</td><td>364 U</td><td>366 U</td><td>370 U</td><td>722 111</td><td>748 UI</td></th<>		541-73-1	384 U	361 U	735 U	364 U	366 U	370 U	722 111	748 UI
Hexachlorobutadene 18-741 384 U 361 U 735 U 364 U 366 U 370 U 722 U 748 U Hexachlorobutadene 87-63										
Hexachlorobutadiene 87-68-3 - </td <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>										
PCB Congeners (ng/kg) CRO-001 PCB-001 275 223 261 462 373 3.84 J 230 345 PCB-002 PCB-002 8.64 U 10.1 U 4.48 U 23.7 J 29.2 J 1.74 U 8.23 J 3.9 J PCB-003 PCB-003 59 37.9 J 50.7 113 94.1 1.63 U 3.14 J 55.7 PCB-004 PCB-004 5130 J 4040 4060 J 7720 J 6540 J 48.1 J 2820 4490 PCB-005 PCB-006 64.7 U 119 J 116 226 207 3.4 U 67.8 125 PCB-007 PCB-008 9.800 1820 1560 3330 2710 12.8 1320 193 PCB-007 PCB-008 15800 11600 1000 18100 13700 69.8 6390 10500 PCB-008 PCB-009 965 615 495 120 96.3 4.77 J 3.47 J 5.54										
PCB-001 PCB-001 275 223 261 462 373 3.84 J 230 345 PCB-002 PCB-002 8.64 U 10.1 U 4.48 U 23.7 J 29.2 J 1.74 U 8.23 J 9 J PCB-003 PCB-003 59 37.9 J 50.7 113 94.1 1.63 U 31.4 J 55.7 PCB-004 PCB-004 5130 J 4040 4060 J 7720 J 6540 J 48.1 J 2280 M 4490 PCB-005 PCB-006 64.7 U 119 J 116 226 207 3.4 U 67.8 125 PCB-006 PCB-006 2620 1820 1560 3330 2710 12.8 1320 1930 PCB-007 PCB-007 368 227 202 483 404 665 J 134 218 218 PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-		0, 00 3						5,00		
PCB-002 RCB-002 8.64 U 10.1 U 4.48 U 23.7 J 29.2 J 1.74 U 8.23 J 9 J PCB-003 PCB-003 PCB-003 PCB-003 PCB-004 PCB-005 PCB-004 PCB-004 PCB-004 PCB-005 PCB-005 PCB-005 PCB-005 PCB-005 PCB-005 PCB-006 PCB-007 PCB-006 PCB-006 PCB-007 PCB-007 PCB-007 PCB-007 PCB-007 PCB-007 PCB-007 PCB-007 PCB-007 PCB-008 PCB-009		PCR_∩∩1	275	222	261	462	272	3 84 1	230	2/15
PCB-003 PCB-003 59 37.9 J 50.7 113 94.1 1.63 U 31.4 J 55.7 PCB-004 PCB-004 5130 J 4040 4060 J 7720 J 6540 J 48.1 J 2820 4490 PCB-005 PCB-005 64.7 U 119 J 116 226 207 3.4 U 67.8 125 PCB-006 PCB-006 2620 1820 1560 3330 2710 12.8 1320 1930 PCB-007 PCB-007 368 227 202 483 404 6.65 J 134 218 PCB-008 PCB-008 11580 11000 18100 13700 69.8 6390 10500 PCB-019 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-012/013 PCB-011 54.8 U <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></td<>										
PCB-004 PCB-004 5130 J 4040 4060 J 7720 J 6540 J 48.1 J 2820 4490 PCB-005 PCB-005 64.7 U 119 J 116 226 207 3.4 U 67.8 125 PCB-006 PCB-006 2620 1820 1560 3330 2710 12.8 1320 1930 PCB-007 7CB-007 368 227 202 483 404 6.65 J 134 218 PCB-008 PCB-008 15800 11600 10000 18100 13700 69.8 6390 10500 PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-012 54.8 U 132 U 90.7 144 J 114 31.2 U 96.9 167 PCB-013 PCB-014 54										
PCB-005 PCB-005 64.7 U 119 J 116 226 207 3.4 U 67.8 125 PCB-006 PCB-006 2620 1820 1560 3330 2710 12.8 1320 1930 PCB-007 PCB-007 368 227 202 483 404 6.65 J 134 218 PCB-008 PCB-008 15800 11600 10000 18100 13700 69.8 699 699 10500 PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-012/013 PCB-013 132 U 90.7 144 J 114 31.2 U 112 146 PCB-014 PCB-017/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014										
PCB-006 PCB-006 2620 1820 1560 3330 2710 12.8 1320 1930 PCB-007 PCB-007 368 227 202 483 404 6.65 J 134 218 PCB-008 PCB-008 1580 11600 10000 18100 13700 69.8 6390 10500 PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-012 PCB-013 132 U 90.7 144 J 114 31.2 U 112 146 PCB-012/013 PCB-012/013 PCB-012/013 PCB-012/013 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015										
PCB-007 368 227 202 483 404 6.65 J 134 218 PCB-008 PCB-008 15800 11600 10000 18100 13700 69.8 6390 10500 PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-011 54.8 U 132 U 90.7 144 J 114 31.2 U 112 146 PCB-012/013 PCB-012/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 1560 1470 1340 2210 2310 3580 1150 2210										
PCB-008 PCB-008 15800 11600 10000 18100 13700 69.8 69.9 6390 10500 PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-011 54.8 U 132 U 90.7 144 J 114 31.2 U 112 146 PCB-012/013 PCB-012/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015 1500 1470 1340 2210 2310 3580 1150 2210										
PCB-009 PCB-009 965 615 495 1200 963 4.77 J 347 545 PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-011 54.8 U 132 U 90.7 144 J 114 31.2 U 112 146 PCB-012/013 PCB-012/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015 1560 1470 1340 2210 2310 3580 1150 2210										
PCB-010 PCB-010 177 130 J 93.8 217 191 3.62 U 87.2 136 PCB-011 PCB-012 54.8 U 132 U 90.7 144 J 114 31.2 U 112 146 PCB-012/013 PCB-012/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015 1560 1470 1340 2210 2310 3580 1150 2210										
PCB-011 PCB-012 54.8 U 132 U 90.7 144 J 114 J 31.2 U 112 M 146 M PCB-012/013 PCB-012/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015 1560 1470 1340 2210 2310 3580 1150 2210										
PCB-012/013 PCB-012/013 242 186 J 153 379 310 24.3 96.9 167 PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015 1560 1470 1340 2210 2310 3580 1150 2210										
PCB-014 PCB-014 54 U 23 U 9.83 U 22.9 U 17 U 2.93 U 3.87 U 5.13 U PCB-015 PCB-015 1560 1470 1340 2210 2310 3580 1150 2210										
PCB-015 PCB-015 1560 1470 1340 2210 2310 3580 1150 2210	PCB-012/013			186 J			310			
	PCB-014	PCB-014				22.9 U	17 U		3.87 U	
PCB-016 PCB-016 14700 13600 11900 17600 13400 43.3 11800 17900	PCB-015	PCB-015	1560	1470	1340	2210	2310		1150	
	PCB-016	PCB-016	14700	13600	11900	17600	13400	43.3	11800	17900

		Tissue Analytical Results									
	Location ID	PB095.2	PB101_D	PB101_D	PB101_D	PB102					
	Sample ID	PB04-I-A-BCR-W-001-COMP-201106	PB04-F-A-GKF-W-001-20110627	PB04-F-A-GKF-W-001-COMP-201106	PB04-F-A-GKF-W-009-20110627	PB04-F-A-PNF-W-001-COMP-201106					
	Sample Date	6/29/2011	6/27/2011	6/27/2011	6/27/2011	6/27/2011					
	Sample Type		N	N	N	N					
	x , , , ,	3200882	3201317.36	3201317.36	3201317.36	3201302.69					
	v.	13829528	13828141.83	13828141.83	13828141.83	13828160.48					
	Scientific_Name		Fundulus grandis	Fundulus grandis	Fundulus grandis	Lagodon rhomboides					
	Size_Class	· ·	=<15	=<15	=<15	=<15					
	Tissue_Category		Sedentary	Sedentary	Sedentary	Migratory					
			-	Whole Body	•						
Chemical	Tissue_Type	Whole Body Result	Whole Body Result	Result	Whole Body Result	Whole Body Result					
Conventional Parameters (pct)	CAS Number	Result	nesuit	nesuit	Result	Result					
" /	LIPID	1.02	3.36	3.37	2.13	3.93					
Lipids	MOIST	74	5.30 74	73.8	73.7	73.2					
Moisture, percent	IVIOIST	74	74	/3.6	73.7	/3.2					
Metals (mg/kg)	7439-92-1	0.173									
Lead	7439-92-1										
Mercury Polyaromatic Hydrocarbons (µg/kg)	/433-3/-0	0.009 J	0.029	0.04	0.028	0.025					
, , , , , , , , , , , , , , , , , , , ,	91-57-6	7 74 11									
2-Methylnaphthalene		7.71 U									
Acenaphthylana	83-32-9	7.71 U									
Acenaphthylene	208-96-8	7.71 U			-	-					
Anthracene	120-12-7	7.71 U				-					
Benzo(a)anthracene	56-55-3	7.71 U									
Benzo(a)pyrene	50-32-8	7.71 U				-					
Benzo(b)fluoranthene	205-99-2	7.71 U									
Benzo(g,h,i)perylene	191-24-2	7.71 U			-						
Benzo(k)fluoranthene	207-08-9	7.71 U									
Chrysene	218-01-9	7.71 U			-						
Dibenzo(a,h)anthracene	53-70-3	7.71 U			-						
Fluoranthene	206-44-0	7.71 U									
Fluorene	86-73-7	7.71 U									
Indeno(1,2,3-c,d)pyrene	193-39-5	7.71 U									
Naphthalene	91-20-3	7.71 U									
Phenanthrene	85-01-8	7.71 U	-		-	-					
Pyrene	129-00-0	7.71 U									
Total HPAH (9 of 16) (U = 1/2)	tPAH_17_HM_N	7.71 U									
Total LPAH (7 of 16) (U = 1/2)	tPAH_17_LM_N	7.71 U									
Total PAH (16) (U = 1/2)	tPAH_17_N	7.71 U									
Semi-volatile Organic Carbons (μg/kg)	•										
1,3-Dichlorobenzene	541-73-1	360 U	705 UJ	719 U	699 UJ	727 U					
1,4-Dichlorobenzene	106-46-7	360 U									
Hexachlorobenzene	118-74-1	360 U	705 UJ	719 U	699 UJ	397 J					
Hexachlorobutadiene	87-68-3	360 U									
PCB Congeners (ng/kg)		1									
PCB-001	PCB-001	6.69 U	205	417 J	145	441					
PCB-002	PCB-002	5.8 U	4.49 U	15 J	8.06 J	1.66 U					
PCB-003	PCB-002	5.81 U	37.3 J	67.7 J	21.7 J	18.6					
PCB-004	PCB-003	25 U	2960	4820	1840	5210					
						86 J					
PCB-005	PCB-005	12.2 U	75.9	138	48.3						
PCB-006	PCB-006	10.8 U	1470	2230	938	1500					
PCB-007	PCB-007	11.3 U	160	257	88.8	164					
PCB-008	PCB-008	41.1 J	7560	12200	5170	5270					
PCB-009	PCB-009	11.3 U	396	653	245	808					
PCB-010	PCB-010	11.7 U	88.7	139	57.8	116					
PCB-011	PCB-011	30 U	159	154	105	29.8 U					
PCB-012/013	PCB-012/013	11.4 U	103	164 J	69.3	55.1 J					
PCB-014	PCB-014	10.8 U	4.37 U	8.56 U	4.56 U	4.09 U					
PCB-015	PCB-015	1100	1300	1790	866	496					
PCB-016	PCB-016	7.55 U	12800	15600	9740	8810					

Appendix D
Tissue Analytical Results

					Tissue	Analytical Results				
	Location ID	Field QC	Field QC	PB002.1	PB003.1	PB005.1	PB006.1	PB008.2	PB010	PB010.1
	Sample ID	1				PB01-I-A-BCR-W-002-COMP-201106			PB01-F-A-GM-W-008-20110618	PB01-I-A-WHS-W-001-COMP-20110
	Sample Date	l	6/24/2011	6/20/2011	6/18/2011	6/20/2011	6/20/2011	6/19/2011	6/18/2011	6/20/2011
	Sample Type	l	EB	N	N N	N N	N	N	N	N
	х			3202691	3202445.49	3202312	3202026.63	3202032	3201710.87	3201704.8
	Y			13836499	13836263.53	13836201	13836356.8	13836072	13836029.16	13836071.32
	Scientific_Name			Callinectes sapidus	Mugil cephalus	Callinectes sapidus	Penaeus setiferus	Penaeus setiferus	Brevoortia patronus	Penaeus setiferus
	Size_Class	l		>2 to =<7.5	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13	=<15	>2 to =<7.5
	Tissue_Category			Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type			Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017		237	24.5	3630	32.9	2280	971	7350	1680
PCB-018/030	PCB-018/030		302	59.2	6220	27.3	4380	1930	12700	3610
PCB-019	PCB-019		24.3 J	9.64 U	478	3.47 U	89.8 J	34.3	923	61.7
PCB-020/028	PCB-020/028		1870	18600	24400	31300	49300	25700	47600	40400
PCB-021/033	PCB-021/033		203	22.3 J	6490	40.7	8020	4590	13600	6720
PCB-022	PCB-022		243	55.9 J	4260	132	9130	3910	9630	7360
PCB-023	PCB-022		7.99 U	1.63 U	12 J	1.7 U	6.29 U	2.45 U	20.1	5.46 U
PCB-024	PCB-023		8.05 U	1.69 U	56.6	1.76 U	22.4 J	10.1	97.8	19.3
PCB-024 PCB-025	PCB-024 PCB-025		190	28.6	1000	26.6	1400	738	1750	19.3
PCB-025 PCB-026/029	PCB-025		391	56.9	3000	35.7	4670	2740	5560	4360
PCB-026/029 PCB-027	PCB-026/029 PCB-027		39.9	56.9 5.27 J	378	35.7 2.65 U	258	131	737	233
PCB-027	PCB-027		1350	3.27 J	22100	469	50400	25600	44600	37300
PCB-032	PCB-031		183	27.2	2710	57.5	2690	1230	5660	2190
PCB-034	PCB-032		28 J	4.32 J	165	11.7	158	84.3	305	133
PCB-035	PCB-034 PCB-035		14.5 U	4.32 J 4.02 U	44	5.58	458	244	38.8 J	402
PCB-036	PCB-035		14.5 U	4.02 U 3.8 U	6.14 U	3.77 U	12.8 U	9.77 U	8.14 U	27.6
PCB-037	PCB-036		43.8 J	1350	1800	2020	5170	2590	1660	4600
PCB-037	PCB-037			12.9	46.3	2020	14.2 U	10.8 U	67.7	
	PCB-039					34.5		9.46 U	544	16.1 U
PCB-039			1	21.2	5.81 U		12.4 U			14 U
PCB-040/071	PCB-040/071		1410	150	9950 J	394	6030 J	2720	23300 J	5350
PCB-041	PCB-041		75.9	4.08 U	1090 J	4.81 U	542 J	189	3620 J	597
PCB-042	PCB-042		1040	62.4	6670 J	84.1	5060 J	2150	14800 J	4430
PCB-043	PCB-043		35.5 U	21.4	732 J	21.9	530 J	196	1990 J	474
PCB-044/047/065	PCB-044/047/065		4030	12500	29400 J	22100	41800 J	23300	58000 J	39500
PCB-045/051	PCB-045/051		234	19.6	2840 J	22	516 J	234	6040 J	576
PCB-046	PCB-046		39.4	4.46 U	632 J	3.87 U	103 J	42.8	1240 J	112
PCB-048	PCB-048		400	23.8 J	5450 J	36.2	2710 J	955	13300 J	2180
PCB-049/069	PCB-049/069		4330	109	25500 J	146	34800 J	19800	51800 J	33100
PCB-050/053	PCB-050/053		251	20.5	2580 J	8.7 J	850 J	357	5690 J	1070
PCB-052	PCB-052		5350	277	38200 J	113	56200 J	29800	78900 J	57600
PCB-054	PCB-054		15.1 U	3.27 U	54.7 J	2.33 U	12.5 U	6.27 U	81.3 J	10 U
PCB-055	PCB-055		43.9 J	5.79 U	219 J	5.45 U	320 J	24.2 U	487 J	420
PCB-056	PCB-056		840	64.6 5.64 U	10500 J	156 6.23	20200 J	9610	24000 J	19100
PCB-057	PCB-057		57.8	5.64 U 5.68 U	121 J	23.5	248 J	131	274 J	208
PCB-058	PCB-058		35.7 J		138 J	23.5	232 J	138	253 J	
PCB-059/062/075	PCB-059/062/075		345	1160	2410 J		3280 J	1890	4540 J	3160
PCB-060	PCB-060		558	8530	5830 J	12200	14800 J	7900	15100 J	13900
PCB-061/070/074/076	PCB-CE12		6960	19300	64500 J	31700	147000 J	81800	137000 J	124000
PCB-063	PCB-063		342	260	1270 J	469	3110 J	1700	3210 J	2700
PCB-064	PCB-064		1320	252	13000 J	310	19700 J	10000	29200 J	18500
PCB-066	PCB-066		4880	34300	41200 J	56400	80100 J	46700	84600 J	72300
PCB-067	PCB-067		123	24.2	680 J	34.4	1200 J	681	1340 J	1060
PCB-068	PCB-068		165	121	310 J	210	422 J	275	505 J	364
PCB-072	PCB-072		238	30.9	600 J	57.7	885	525	1030 J	753
PCB-073	PCB-073		85.5	2.69 U	7.82 UJ	5.8	11.4 U	5.57 U	13.1 UJ	17 U
PCB-077	PCB-077		265 J	1630	1810	2490	5450	3100	3390	4760
PCB-078	PCB-078		37.7 U	5.51 U	9.07 UJ	5.18 U	16.3 U	24.1 U	13.4 UJ	51.5 U
PCB-079	PCB-079		32.2 U	40.5	379 J	64.6	13.6 U	20.1 U	628 J	43 U

				lissue Ana	alytical Results				
	Location ID	PB010.3	PB013.1_A	PB013.3	PB013.3	PB015.1	PB015.2	PB015_A	PB015_A
	Sample ID	PB01-I-A-BCR-W-003-COMP-201106	_	PB01-F-A-GKF-W-004-COMP-201106	PB01-F-A-STM-W-003-20110616	PB01-F-A-GKF-W-002-COMP-201106	PB01-I-A-BCR-W-004-COMP-201106	_	PB01-I-A-BCR-W-006-COMP-20110
	Sample Date		6/22/2011	6/21/2011	6/15/2011	6/15/2011	6/16/2011	6/25/2011	6/22/2011
	Sample Type		N	N	N	N	N	N	N
	х х	3201743	3201306.47	3201433.88	3201433.88	3201175	3201180	3201168.73	3201168.73
	ν	13836152	13836191.23	13836075.12	13836075.12	13836251	13836183	13836254.66	13836254.66
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Fundulus grandis	Mugil cephalus	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Callinectes sapidus
	Size_Class	>2 to =<7.5	>2 to =<7.5	=<15	=<15	=<15	>2 to =<7.5	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Sedentary	Sedentary	Migratory	Sedentary	Sedentary	Sedentary	Sedentary
	Tissue_Type	-	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017	88.3	52.4	24700	9660	21000	41	34200	96
PCB-018/030	PCB-018/030	104	44	41400	23800	35300	26.9 J	57200	152
PCB-019	PCB-019	6.44 U	2.99 U	2370	1310	2600	4.59 U	3010	9.11 U
PCB-020/028	PCB-020/028	56200	73800	177000	76900	151000	83900	241000	64400
PCB-021/033	PCB-020/028	241	138	18000	12600	19200	21.8	21100	206
PCB-021/033	PCB-021/033	527	279	30500	13200	24600	222	35200	221
PCB-022 PCB-023	PCB-022	1.69 U	1.01 U	59.4	26.3 J	49.2	2.51 U	80.9	5.14 U
PCB-023 PCB-024	PCB-023	1.69 U 1.76 U	1.01 U 1.07 U	263	26.3 J 170	197	2.51 U 2.6 U	80.9 499	4.9 U
PCB-024 PCB-025	PCB-024 PCB-025	1.76 U	38.6	4810	2740	4110	53.3	6260	4.9 U 39.4
PCB-026/029	PCB-025 PCB-026/029	103	79.6	4810 17500	9780	13600	48.6	24700	103
•									
PCB-027	PCB-027	7.27 2500	7.13 948	2350 154000	1160	1980	2.53 U 1250	3650 188000	13.3 J
PCB-031	PCB-031				71900	122000			798
PCB-032	PCB-032	157	160	23100	8380	19800	56.2	29900	131
PCB-034	PCB-034	14.8	16.3	527	351	468	7.07	746	24.2
PCB-035	PCB-035	19.5	10.2	25.6	49.3 J	187 U	13.1	284 U	10.9 U
PCB-036	PCB-036	5.02 U	3.2 U	10.3 U	21.2 U	189 U	5.74 U	288 U	9.58 U
PCB-037	PCB-037	4000	5790	4820	2850	3830	5720	4640	4810
PCB-038	PCB-038	5.35 U	3.36 U	173	103	189 U	35.8	288 U	9.98 U
PCB-039	PCB-039	4.85 U	52.3	530	20.3 U	183 U	62.9	278 U	57
PCB-040/071	PCB-040/071	683	565	81200	25600	55800	259	79200	590
PCB-041	PCB-041	12.5 J	5.92	8460	2280	7390	5.96 U	9970	38.2
PCB-042	PCB-042	588	102	35000	14800	30400	180	38700	111
PCB-043	PCB-043	58.9	41.3	10700	40.8 U	8540	31	9480	14.6 U
PCB-044/047/065	PCB-044/047/065	38400	26500	103000	94600	181000	50000	266000	31300
PCB-045/051	PCB-045/051	46.7 J	26.5	19000	8790	18100	30.8 J	22600	39.1 J
PCB-046	PCB-046	6.63 J	2.91	2130	1670	2040	7.08 U	2640	12.4 U
PCB-048	PCB-048	119	50.4	21900	13500	24100	35.5	35200	88.4
PCB-049/069	PCB-049/069	1220	163	195000	81100	178000	243	267000	319
PCB-050/053	PCB-050/053	29.6 J	14.1	18600	8120	16000	12 J	25600	31.4
PCB-052	PCB-052	833	290	291000	128000	228000	115	413000	489
PCB-054	PCB-054	2.52 U	1.35 U	158	131	144	4.43 U	197	8.96 U
PCB-055	PCB-055	75.7	3.13 U	34.2 U	1070	851	10.9 U	1260	19.5 U
PCB-056	PCB-056	961	284	46700	33700	40200	252	46000	284
PCB-057	PCB-057	10.7	6.95	856	463	630 J	10.6 U	662 J	18.7 U
PCB-058	PCB-058	29.4	20.2	645	302	498	10.7 U	699	19.1 U
PCB-059/062/075	PCB-059/062/075	3170	2200	17300	7720	13100	4270	20300	2680
PCB-060	PCB-060	26400	22200	60100	23200	54800	32900	76200	21500
PCB-061/070/074/076	PCB-CE12	64400	50500	428000	217000	407000	78400	551000	53700
PCB-063	PCB-063	1610	759	12700	5050	11600	1490	16800	564
PCB-064	PCB-064	2440	494	109000	43700	102000	971	131000	472
PCB-066	PCB-066	104000	84300	311000	132000	299000	131000	410000	105000
PCB-067	PCB-067	111	42.4	1960	2330	2130	87.6	2610	45.1
PCB-068	PCB-068	333	158	1390	671	1120	376	1640	220
PCB-072	PCB-072	102	50.1	2960	1440	2470	62.6	3460	67.6
PCB-073	PCB-073	6.11	7.14	25.3 U	1620	18 U	3.93 U	562	49.7
PCB-077	PCB-077	4640	4480	9690	3880	8640	6110	11500	5070
PCB-078	PCB-078	8.46 U	3.09 U	32.4 U	34.5 U	19.3 U	10.4 U	48.4 U	19.9 U
PCB-079	PCB-079	7.22 U	80.2	27.2 U	29.6 U	16.3 U	8.85 U	40.9 U	16.4 U

Appendix D
Tissue Analytical Results

				rissue Ana	alytical Results				
	Location ID	PB015_A	PB015_A	PB015_A	PB016.2	PB019.1	PB019.1	PB019.1	PB019.1
		-	_	_	PB01-F-A-GKF-W-003-COMP-201106		PB01-I-B-BRS-W-012-20110621	PB01-I-B-BRS-W-014-20110621	PB01-I-B-BRS-W-015-20110621
	Sample Date	6/22/2011	6/22/2011	6/20/2011	6/15/2011	6/22/2011	6/21/2011	6/21/2011	6/21/2011
	Sample Type	N	N	N N	N	N	N	N	N
	x	3201168.73	3201168.73	3201168.73	3201190	3201152.87	3201152.87	3201152.87	3201152.87
	y	13836254.66	13836254.66	13836254.66	13836042	13835753.58	13835753.58	13835753.58	13835753.58
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Penaeus aztecus	Fundulus grandis	Cynoscion arenarius	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus
	Size_Class	>2 to =<7.5	>2 to =<7.5	>7.5 to =<13	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13
	Tissue_Category	Sedentary	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017	32.9	27.2	2790	10300	7680	4980	4620	3960
PCB-018/030	PCB-018/030	56.9	36.2	4200	17800	15700	9880	9070	7980
PCB-019	PCB-019	6.98 U	11.1 U	113 J	1020	708	253 J	226	189 J
PCB-020/028	PCB-020/028	62400	89800	76000	71900	54500	96200	92300	76200
PCB-021/033	PCB-021/033	122	165	8030	6780	11800	14300	12700	10400
PCB-022	PCB-022	201	221	11500	11900	9770	16300	15400	13400
PCB-023	PCB-023	2.43 U	3.57 U	2.81 U	22.9	23.1	56.9 U	21.1 U	37.5 U
PCB-024	PCB-024	2.32 U	3.4 U	47.5	137	132	74.8 J	60.6 J	67 J
PCB-025	PCB-025	34.7	34.1	1610	2060	1360	2850	2540	2150
PCB-026/029	PCB-026/029	62.4	71.6	6780	7410	5940	9510	9240	7530
PCB-027	PCB-027	4.36 J	3.7 U	274	1050	785	608	544	505
PCB-031	PCB-031	662	702	56200	66000	42000	89800	85400	73900
PCB-032	PCB-032	54.3	56.1	2190	10600	6400	4330	4110	3530
PCB-034	PCB-034	9.1	9.36	149	205	250	246	242	205
PCB-035	PCB-035	14.6	19.3 J	614	14.5 J	7.15 U	722	843	722
PCB-036	PCB-036	6.1 U	10.1 U	9.13 U	3.31 U	6.92 U	130 U	65.9 U	66.6 U
PCB-037	PCB-037	5670	9160	7600	2120	704	10400	11600	9540
PCB-038	PCB-038	6.35 U	10.6 U	11 U	80.2	62.6	131 U	66.5 U	67.2 U
PCB-039	PCB-039	59.2	10.5 U	9.6 U	3.14 U	6.56 U	504	63.2 U	63.9 U
PCB-040/071	PCB-040/071	249 J	262 J	5750	32900	23300 J	13400	10400	8730
PCB-041	PCB-041	6.83 UJ	6.5 UJ	862	3840	4580 J	1540	1170	1110 J
PCB-042	PCB-042	52.7 J	55.4 J	6030	16100	12400 J	13800	11200	9060
PCB-043	PCB-043	5.06 UJ	20.5 J	10.6 U	2800	21.2 UJ	285 U	71.4 U	57.9 U
PCB-044/047/065	PCB-044/047/065	21300 J	28500 J	49800	106000	69700 J	102000	82400	68000
PCB-045/051	PCB-045/051	18.3 J	17.9 J	806	7570	5670 J	2080	1640	1460
PCB-046	PCB-046	2.56 UJ	2.85 UJ	142	855	1010 J	478 J	327	327 J
PCB-048	PCB-048	43 J	33.9 J	4010	11100	15800 J	10500	7640	6420
PCB-049/069	PCB-049/069	123 J	97.7 J	38800	109000	53000 J	90900	64200	56400
PCB-050/053	PCB-050/053	10.8 J	9.12 J	946	8750	3300 J	3020	2000	2050
PCB-052	PCB-052	170 J	166 J	54300	182000	85300 J	124000	81900	80400
PCB-054	PCB-054	3.99 U	5.64 U	13.7	71.7	85.1	110 U	36.9 U	48.6 U
PCB-055	PCB-055	6.97 UJ	157 J	31.2 U	495	17.1 UJ	232 U	1020	85.6 U
PCB-056	PCB-056	201 J	288 J	26000	21000	15900 J	44700	39400	33100
PCB-057	PCB-057	6.68 UJ	9.49 UJ	363	383	302 J	877	559	466
PCB-058	PCB-058	6.81 UJ	12.8 J	284	277	225 J	648	501	343
PCB-059/062/075	PCB-059/062/075	1880 J	2480 J	4070	8460	5840 J	7910	5980	5470
PCB-060	PCB-060	22200 J	29200 J	22800	31500	25000 J	39800	33100	27900
PCB-061/070/074/076	PCB-CE12	48600 J	66200 J	172000	227000	168000 J	356000	289000	261000
PCB-063	PCB-063	798 Ј	993 J	4490	6320	4870 J	8820	6760	5760
PCB-064	PCB-064	356 J	353 J	18300	55900	35600 J	33800	28900	24800
PCB-066	PCB-066	88900 J	123000 J	117000	175000	121000 J	205000	174000	157000
PCB-067	PCB-067	40.7 J	50.7 J	1630	1170	1120 J	3450	2540	2150
PCB-068	PCB-068	186 J	239 J	546	753	524 J	1130	803	630
PCB-072	PCB-072	50.2 J	58 J	1090	1420	1070 J	2200	1700	1440
PCB-073	PCB-073	26.7 J	4.45 UJ	357	5.64 U	1330 J	913 J	571	549
PCB-077	PCB-077	5130 J	7540 J	7510	4400	1370 J	10400	10200	8830
PCB-078	PCB-078	7.11 UJ	10.1 UJ	31.2 U	15.3 U	16.2 UJ	218 U	55.2 U	80.4 U
PCB-079	PCB-079	5.85 UJ	8.32 UJ	27.2 U	13.2 U	13.9 UJ	183 U	46.2 U	67.3 U

				Tissue An	alytical Results				
	Location ID	PB019.1	PB019.1	PB019.1	PB019.2	PB019_A	PB019_A	PB024.1	PB024.1
	Sample ID		PB01-I-B-BRS-W-022-20110622	PB01-I-B-BRS-W-023-20110622		PB01-F-A-GKF-W-005-COMP-201106		PB01-F-A-GM-W-001-20110615	PB01-F-A-GM-W-002-20110615
	Sample Date		6/22/2011	6/22/2011	6/28/2011	6/21/2011	6/23/2011	6/15/2011	6/15/2011
	Sample Type		N	N	N	N	0,25,2011 N	N	N
	y x	3201152.87	3201152.87	3201152.87	3201194.8	3201314.96	3201314.96	3201090.96	3201090.96
	^ v	13835753.58	13835753.58	13835753.58	13835735.1	13835750.59	13835750.59	13835160.37	13835160.37
	Scientific_Name		Penaeus aztecus	Penaeus aztecus	Crassostrea virginica	Fundulus grandis	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus
	Size_Class	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Migratory	Migratory
	Tissue_Type		Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017	4970	5380	5500	11800	20100	94.2	11600	11600
PCB-018/030	PCB-018/030	10700	11100	12300	22800	27300	72.1	24700	25800
PCB-019	PCB-019	226	313	317	1830	2070	6.76 U	1180	1140
PCB-020/028	PCB-020/028	121000	99400	105000	69700	160000	75800	72300	67300
PCB-021/033	PCB-020/028	15700	14700	14800	18900	13600	237	20300	18500
PCB-022	PCB-021/033	18700	18900	18400	14400	26500	553	15000	13600
PCB-022 PCB-023	PCB-022	41.3 U	15.2	18400 14.1 U	25.2 J	58.6	2.76 U	31.5	28.5 J
PCB-023	PCB-023	70.6 J	82.1 J	79.4 J	25.2 J 3.76 U	4.11 U	2.76 U 2.63 U	145	28.5 J 155
PCB-025	PCB-024 PCB-025	3060	2660	2770	2490	4.11 0	92.5	2810	2670
	PCB-025 PCB-026/029	11000	9880	10200	2490 8890	14700	92.5 175	8680	8170
PCB-026/029		626	627	662			8.12	1290	
PCB-027	PCB-027	107000	92600		1040 62200	2110	8.12 2130		1260 64600
PCB-031	PCB-031			96200		124000		68800	
PCB-032	PCB-032	4740	5470	5450	8700	18900	211	9790	9550
PCB-034	PCB-034	298 J	222	250	402	411	30.2	389	373
PCB-035	PCB-035	912	838	828	864	10.6 U	17.8 J	80.5	66.1
PCB-036	PCB-036	73.4 U	14.5 U	49.9 U	11.8 U	10.4 U	5.54 U	17 U	39.5 U
PCB-037	PCB-037	13000	11300	10500	6960	4170	6040	2470	2330
PCB-038	PCB-038	74.1 U	78.1	49.2 U	130	204	5.77 U	94 J	94.3 J
PCB-039	PCB-039	609	13.4 U	46 U	12.4 U	873	5.77 U	16.3 U	37.9 U
PCB-040/071	PCB-040/071	12100	13000	12600	29800	64800	832	35000	33000
PCB-041	PCB-041	1200	1870	1650	6410	8420	9.28 U	5540	4690
PCB-042	PCB-042	13500	14100	13600	16400	30900	249	21800	20700
PCB-043	PCB-043	1030	19 U	44.2 U	2470	6940	51	25.6 U	52.3 U
PCB-044/047/065	PCB-044/047/065	99600	96800	95600	75600	181000	29100	84600	82000
PCB-045/051	PCB-045/051	1990	2360	2240	9420	13800	34.3 J	9660	8950
PCB-046	PCB-046	399 J	487	444	1650	1210	4.3 U	2020	1960
PCB-048	PCB-048	9100	9580	9400	16300	26200	92	19300	18600
PCB-049/069	PCB-049/069	84500	80300	81400	60800	188000	579	73500	72000
PCB-050/053	PCB-050/053	2460	2830	2710	8350	14300	18.6	9490	9060
PCB-052	PCB-052	110000	109000	108000	97900	312000	533	118000	116000
PCB-054	PCB-054	51.2 U	19.8	27.6 J	193	114	5.36 U	95.6	100
PCB-055 PCB-056	PCB-055 PCB-056	1720 43300	1480 45300	1820 J 43700	637 37300	16.1 U 43700	161 630	31 U 37100	1310 34300
PCB-056 PCB-057		43300 610	45300 612	610	37300	849	16.6	393	34300
PCB-057 PCB-058	PCB-057 PCB-058	525	505	491	333	545	37	274	390
		7720	7430	7190	6500	14600	2580	6380	6120
PCB-059/062/075 PCB-060	PCB-059/062/075 PCB-060	39300	41000	38700	19000	64100	2580	23200	21400
PCB-061/070/074/076	PCB-060 PCB-CE12	39300	324000	317000	177000	448000	60800	23200	196000
PCB-063	PCB-CE12 PCB-063	7930	7930	7660	4050	12400	1250	4990	4500
PCB-064	PCB-063 PCB-064	7930 35200	7930 36800	36000			1150		42600
PCB-066	PCB-064 PCB-066	200000			37700	105000 330000		45300	114000
		200000	191000	191000 2860	113000		108000	123000	
PCB-067	PCB-067		2780		2230	2460	86.1	2170	2020 559
PCB-068	PCB-068	1040	944	950	553	1310	316	593	
PCB-072	PCB-072	2110	1910	1890	1090	2290	127	1240	1260
PCB-073	PCB-073	88.4 U	801	910	12.5 U	17.2 U	8.65 J	2010	2090
PCB-077	PCB-077	12000	9820	9460	6240	9280	5700	4560	4080
PCB-078	PCB-078	116 U	21.2 U	64.9 U	13.2 U	15 U	11 U	29.6 U	52.4 U
PCB-079	PCB-079	1100	17.9 U	938	11.5 U	12.9 U	9.05 U	25.4 U	44.8 U

Tissue Analytical Results												
	Location ID	PB024.4	PB025_A	PB026.1	PB032_A	PB037.2	PB037.4	PB037.5	PB037.6			
		PB01-I-A-BCR-W-005-COMP-201106	_	PB01-F-A-STM-W-002-20110616	PB02-F-A-PNF-W-009-20110617	PB02-F-A-STM-W-006-20110616			PB02-I-A-BCR-W-002-COMP-201106			
	Sample Date		6/15/2011	6/15/2011	6/17/2011	6/16/2011	6/17/2011	6/17/2011	6/16/2011			
	Sample Type		N	N	N	N	N N	N	N			
	X	3201319	3201214.42	3201153.65	3201407.03	3201346.97	3201504	3201380	3201402			
	Y	13835188	13835154.42	13835079.21	13834542.25	13834038.02	13833530	13834151	13833775			
	Scientific_Name		Brevoortia patronus	Mugil cephalus	Lagodon rhomboides	Mugil cephalus	Callinectes sapidus	Penaeus aztecus	Callinectes sapidus			
	Size_Class		=<15	=<15	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5			
	Tissue_Category		Migratory	Migratory	Migratory	Migratory	Migratory	Migratory	Sedentary			
	Tissue_Type	1	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body			
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result			
PCB-017	PCB-017	549	18200	12100	4500	19100	61.9	2630	65			
PCB-018/030	PCB-018/030	1030	40200	33700	16400	47400	184	4110	133			
PCB-019	PCB-019	52 J	2140	1760	640	2210	9.03	146	8.87 U			
PCB-020/028	PCB-020/028	72000	113000	112000	92600	140000	74200	64400	64600			
PCB-021/033	PCB-021/033	1030	31000	15000	2980	26100	140	7950	142			
PCB-022	PCB-022	1120	24700	19600	7320	25600	660	9790	517			
PCB-023	PCB-023	7.69 U	52 J	43.9	25.4 J	64.3	2.42 U	5.91	1.58 U			
PCB-024	PCB-024	7.98 U	231	276	205	300 J	2.31 U	35.8 J	1.64 U			
PCB-025	PCB-025	209	4490	3900	1160	5070	74.8	1320	69.4			
PCB-026/029	PCB-026/029	603	13800	14100	9410	17500	142 J	5840	134			
PCB-027	PCB-027	76.4 J	2110	1490	388	2210	7.93	263	9.57			
PCB-031	PCB-031	5330	110000	103000	81200	133000	1640	52100	1740			
PCB-032	PCB-032	819	15700	10700	1780	16500	107	2420	113			
PCB-034	PCB-034	60.5	543	392	175	592	11.1	107	8.86 J			
PCB-035	PCB-035	43.6 J	51.1	49.9	28.1 U	35.6 U	21.1	484	13.2 J			
PCB-036	PCB-036	85.2	19.5 U	24.9 U	26.3 U	33.6 U	6.5 U	5.2 U	5.15 U			
PCB-037	PCB-037	5970	3420	3190	1070	6170	5990	6720	5670			
PCB-038	PCB-038	38.3 J	137	132	133	188 J	6.77 U	6.29 U	26.7 J			
PCB-039	PCB-039	108 J	18.7 U	23.9 U	24.8 U	31.9 U	6.77 U	5.47 U	47.4 J			
PCB-040/071	PCB-040/071	3180	58200	31000	7430	54000	486 J	4800	580			
PCB-041	PCB-041	50.7 J	9960	3180	852	6450	6.36 UJ	797	4.86 U			
PCB-042	PCB-042	1410	36100	18700	13800	32400	316 J	4850	362			
PCB-043	PCB-043	286	37.1 U	32 U	1800	48.4 U	66.5 J	381	81			
PCB-044/047/065	PCB-044/047/065	38700	137000	125000	122000	181000	44300 J	40000	35000			
PCB-045/051	PCB-045/051	229	15900	12800	5970	19100	40 J	647	49 J			
PCB-046	PCB-046	48.7 J	3450	2350	410	3510	6.54 J	107	4.72 U			
PCB-048	PCB-048	752	31600	16800	5140	32300	139 J	3650	91.9			
PCB-049/069	PCB-049/069	2670	117000	104000	105000	155000	478 J	32100	661			
PCB-050/053	PCB-050/053	213	16100	11300	2270	17600	36.6 J	693	30.5			
PCB-052	PCB-052	3320	189000	172000	190000	243000	808 J	44700	813			
PCB-054	PCB-054	13.5 U	163	183	104	261	3.96 U	12.4	2.73 U			
PCB-055	PCB-055	26.5 U	30 U	1340	738	2320	129 J	391 J	7.77 U			
PCB-056	PCB-056	1960	63000	47800	11300	69800	974 J	22300	631			
PCB-057	PCB-057	44.2 J	709	695	410	996	11.6 UJ	325	12.2 J			
PCB-058	PCB-058	33.4 J	429	379	271	564 J	26.2 J	221	17.3			
PCB-059/062/075	PCB-059/062/075	3400	10400	10100	9860	14500	3400 J	3410	2880			
PCB-060	PCB-060	26500	40400	36200	37800	47400	38800 J	20900	31900			
PCB-061/070/074/076	PCB-CE12	68000	333000	300000	282000	402000	89100 J	152000	64300			
PCB-063	PCB-063	1740	8260	7550	8180	10200	1720 J	4230	1340			
PCB-064	PCB-064	4180	73700	58400	58200	82900	1420 J	15700	1590			
PCB-066	PCB-066	107000	191000	180000	190000	243000	163000 J	104000	103000			
PCB-067	PCB-067	194	3560	3390	981	4750	127 J	1440	78			
PCB-068	PCB-068	435	804	804	991	1060	305 J	404	240			
PCB-072	PCB-072	226 17 U	1820 3450	1640	2000	2200	64.3 J	816 4.4 U	82.1 3.2 U			
PCB-073 PCB-077	PCB-073 PCB-077	5270	7200	2340 5170	36.4 U 784	4570 7590	4.35 UJ 8400 J	6310	3.2 U 5690			
PCB-077	PCB-077 PCB-078	25.2 U	7200 28.6 U	39.8 U	784 40.8 U	7590 61.7 U	12.4 UJ	8.84 U	7.39 U			
				39.8 U 34.1 U			12.4 UJ 10.2 UJ		104			
PCB-079	PCB-079	21.5 U	24.5 U	34.1 U	1000	53.4 U	10.2 UJ	7.71 U	104			

				lissue Alia	alytical Results				
	Location ID	PB037_A	PB047.1_A	PB047.3	PB048.3	PB048.3	PB048.3	PB049_A	PB049_A
	Sample ID	PB02-I-A-BCR-W-004-COMP-201106	PB02-F-A-PNF-W-002-20110615	PB02-I-A-BCR-W-001-COMP-201106	PB02-F-A-GM-W-001-20110616	PB02-F-A-GM-W-002-20110616	PB02-F-A-STM-W-001-20110616	PB02-F-A-GKF-W-001-COMP-201106	PB02-I-A-BCR-W-003-COMP-201106
	Sample Date	6/19/2011	6/15/2011	6/19/2011	6/16/2011	6/16/2011	6/16/2011	6/15/2011	6/22/2011
	Sample Type		N	N N	N	N	N	N	N N
		3201325.97	3201461.79	3201645	3201519.15	3201519.15	3201519.15	3201468.39	3201468.39
	Υ	13833885.45	13833007.68	13832925	13832869.89	13832869.89	13832869.89	13832722.8	13832722.8
	Scientific_Name	Callinectes sapidus	Lagodon rhomboides	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus	Mugil cephalus	Fundulus grandis	Callinectes sapidus
	Size_Class	· ·	=<15	>2 to =<7.5	=<15	=<15	=<15	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Sedentary	Sedentary
	Tissue_Type	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017	59.7	4390	53.3 J	20400	20300	16900	25700	36.4
PCB-018/030	PCB-018/030	79.2	25400	51.8	46400	46100	41100	49700	34.5
PCB-019	PCB-019	7.05 U	806	12.6 U	2700	2540	1810	2250	11.4 U
PCB-020/028	PCB-020/028	185000	132000	82700	124000	122000	120000	191000	78600
PCB-021/033	PCB-021/033	129	2330	97.3	35600	34900	23600	17600	209
PCB-022	PCB-022	506	9850	274	28100	27200	22800	30500	270
PCB-023	PCB-023	5.37 U	25.9 J	7.14 U	57	50.8	52.6	70.1	3.89 U
PCB-024	PCB-024	5.88 U	249	7.42 U	267	295	237	319	4.12 U
PCB-025	PCB-025	64	1430	44.3 J	4940	4620	4110	5020	42.8
PCB-026/029	PCB-026/029	86.6	13500	59.8	15400	14400	14900	19800	56.7
PCB-027	PCB-027	7.13	434	7.21 U	2410	2270	1860	2480	3.52 J
PCB-031	PCB-031	2250	120000	864	121000	119000	114000	159000	859
PCB-032	PCB-032	133	2050	105	18000	17200	14400	24700	79.1
PCB-034	PCB-034	11.8 J	186	15.2 J	543	513	463	496	10.8
PCB-035	PCB-035	12 U	50.6 U	14.7 J	18.4 U	24.6 U	40.3 U	1060 U	9.7 U
PCB-036	PCB-036	11.6 U	47.7 U	11.5 U	17.5 U	23.2 U	38 U	1070 U	8.67 U
PCB-037	PCB-037	14300	1430	6410	4270	3850	6320	4210	6060
PCB-038	PCB-038	11.8 U	205	39.9	160	189	144	1080 U	9.11 U
PCB-039	PCB-039	82.9	45.2 U	61.3	16.8 U	22 U	36 U	1050 U	8.17 U
PCB-040/071	PCB-040/071	416	12400	870	13.6 U	10 U	54900	109000	301
PCB-041	PCB-041	23.8	786	28.1 U	17 U	11.8 U	6600	11300	11.4 U
PCB-042	PCB-042	341	27800	169	14.7 U	10.6 U	33200	40600	113
PCB-043	PCB-043	102	82.9 U	70.7	15.1 U	11.7 U	5970	15300	33.4
PCB-044/047/065	PCB-044/047/065	87500	226000	52100	12.9 U	9.5 U	163000	272000	26500
PCB-045/051	PCB-045/051	61.7	9100	63.1	17500	16900	17400	23600	18.9
PCB-046	PCB-046	14.2 U	521	18 U	3910	3670	3220	2160	5.01 U
PCB-048	PCB-048	65.3	5320	60.2 J	14.1 U	10.4 U	34100	42500	27.8 J
PCB-049/069	PCB-049/069	381	201000	281	12.2 U	9.03 U	136000	283000	102
PCB-050/053	PCB-050/053	39.6	2330	24.5 J	16500	16400	15100	23900	8.24
PCB-052	PCB-052	342	341000	219	14.3 U	10.2 U	216000	437000	118
PCB-054	PCB-054	8.36 U	151 J	10.3 U	198	197	184	126	5.98 U
PCB-055	PCB-055	25.2 U	91.9 U	22 U	1770	2080	1970	1590	238
PCB-056	PCB-056	392	21900	247	69600	65900	68200	49400	340
PCB-057	PCB-057	23.8 U	661	21.5 U	799	732	885	868 U	9.2 U
PCB-058	PCB-058	24.3 U	464 J	21.6 U	562	474	434	932 U	9.51 U
PCB-059/062/075	PCB-059/062/075	6120	17600	4220	11.2 U	8.12 U	13000	22100	2180
PCB-060	PCB-060	59900	82900	43700	46700	43300	45900	86000	24700
PCB-061/070/074/076	PCB-CE12	249000	560000	92700	370000	345000	364000	563000	53700
PCB-063	PCB-063	137000	17800	1640	9430	8620	9330	18000	692
PCB-064	PCB-064	1320	120000	869	10.6 U	7.57 U	78100	154000	412
PCB-066	PCB-066	24 U	396000	154000	216000	204000	219000	425000	94400
PCB-067	PCB-067	95.6	1640	89.8 J	4050	3720	4460	2590	39.2 J
PCB-068	PCB-068	353	1390	288	838	796	762	1910	179
PCB-072	PCB-072	57	2900	78.9	1880	1820	1720	4020	38.2
PCB-073	PCB-073	9.2 U	1730	18.5 U	11.2 U	8.26 U	57.3 U	24.5 U	5.88 U
PCB-077	PCB-077	11700	850	7880	7960	7440	7230	11600	4900
PCB-078	PCB-078	23.3 U	86.1 U	20.9 U	29 U	37 U	63.9 U	910 U	9.28 U
PCB-079	PCB-079	173	74.5 U	126	24.9 U	32 U	1200 J	766 U	7.59 U

					Tissue An	alytical Results				
		Location ID	PB052.1	PB052.1	PB053.2	PB053.2	PB053 B	PB056 A	PB056 A	PB057.2
Section Sect							_	_	_	
Part		•								
		•								
Page		запіріс турс		· ·						
Part		^ v								
Page		Scientific Name								
Part			·		_	1	1	_	•	•
Column Tools (Name) Stock (Name) Work body (Name body (N		_								
March Marc					· ·	· ·	· ·	•	•	
March Marc	Chemical			· · · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · · ·	·	· · · · · · · · · · · · · · · · · · ·	•	•	•
Manufact Manufact										
Process Process 1900										
Production										
Feed										
Fig. 12										
Profess										
March Marc										
No. 10, 10, 10, 10, 10, 10, 10, 10, 10, 10,										
Procession Pro										
Per										
Manual M										
Page Page										
Post										
PAPEN										
MA-10										
PART										
Fig. 10 Fig.										
PAC 1989 1980 1981 1981 1982 1										
Part Part										
P.C. P.C.										
Page Page										
PGB-95/95 PGB										
PCR-048										
PCB-049/099 PCB-050/053 S000 22800 2440000 32900 32900 32900 23800 23800 23800 23800 339000 359000 499000 35900 499000 359000 499000 359000 499000 4	PCB-046	PCB-046	1480	5350	3960 J	3070	2960 J	1780 J	2780	3790
PCR-950/053 PCR-950/053 PCR-950/053 PCR-950 PC										
PCB-052 PCB-052 148000 330000 648000 537000 479000 359000 497000 194000 PCB-054 PCB-054 PCB-054 165 345 283 249 272 264 165 195 200 PCB-055 PCB-055 PCB-055 1430 3280 51.5 UI 33.7 U 2230 2150 1190 2110 PCB-056 PCB-056 31600 116000 126000 92500 86600 68800 70400 71100 PCB-057 PCB-057 567 1330 2600 1900 1790 1790 1380 1500 1540 843 PCB-058 PCB-058 PCB-058 337 807 1490 10600 11600 12500 25100 12500 12500 12500 11000 PCB-058 PCB-058 837 807 1490 1060 11600 12500 25100 25100 2500 12500 12500 11200 PCB-058 PCB-058 837 807 1490 1060 11600 12500 25100 25000 12500 12500 11200 PCB-060 PCB-060 48700 82300 119000 32800 25100 25100 25100 25000 11200 26500 11200 PCB-061/07074/076 PCB-0612 307000 623000 1190000 38800 37000 825000 825000 16000 38800 27000 27000 27000 27000 27000 381000 PCB-064 PCB-064 PCB-064 61400 112000 262000 263000 385000 575000 497000 501000 230000 PCB-066 PCB-066 217000 37200 885000 588000 575000 497000 43300 4240 PCB-067 PCB-068 PCB-068 21700 37200 885000 5190 2340 2190 3700 4330 4420 PCB-073 PCB-073 63.4 U 61.1 U 14800 15900 26.4 U 7410 9730 4710 PCB-073 PCB-075 PCB-077 2940 15100 18500 15500 10500 12500 12500 12500 PCB-078 PCB-078 PCB-077 2940 15100 18500 16500 16500 17000 12500 1	PCB-049/069	•								
PCB 054 PCB 054 165 345 283 283 249 272 246 159 195 200 PCB 055 PCB 055 1430 3280 51.5 U 33.7 U 2280 2150 1190 1190 2110 PCB 056 PCB 055 31600 116000 126000 92500 86600 68300 70400 70400 71100 PCB 057 PCB 057 567 1530 2600 1900 1790 1380 1580 1540 843 PCB 058 PCB 058 337 807 1490 1060 1160 1160 5350 717 480 PCB 059/062/075 PCB 059/062/075 9590 20300 40300 32800 23800 13800 11000 10000 10000 PCB 060 PCB 060 48700 82300 17400 18000 13800 10000 13800 10000 107000 48500 PCB 063/070/074/076 PCB -12 307000 623000 119000 138000 82500 19000 13000 10000 381000 PCB 064 PCB 064 61400 121000 262000 138000 17700 13800 133000 131000 161000 76400 PCB 066 PCB 064 61400 12100 22000 805000 17900 133000 131000 161000 76400 PCB 067 PCB 068 62570 2060 7520 7230 63000 13000 13000 13000 13000 13000 13000 13000 PCB 067 PCB 067 2060 7520 7230 5170 5280 3710 4370 4330 4240 PCB 067 PCB 067 2560 1530 2630 6400 3000 2340 2190 1740 3710 4390 2100 2130 PCB 067 PCB 067 2560 1530 2630 6400 13000 12000 12000 17400 11000 17400 11000 11000 11000 PCB 073 PCB 073 63.4 U 61.1 U 14800 16000 12000 12000 12000 11	PCB-050/053	•								
PCB-055 PCB-055 1430 3280 51.5 UJ 33.7 U 2380 J 2150 J 1190 J 2110 PCB-056 PCB-056 PCB-056 31600 116000 1126000 J 92500 86600 J 68800 J 70400 71100 PCB-057	PCB-052								497000	
PCB-056 PCB-056 RCB-057 RCB-										
PCB-057 S67 1530 2600 J 1900 1790 J 1380 J 1540 843 PCB-058 PCB-058 357 807 1490 J 1000 1160 J 5350 J 717 480 PCB-059/062/075 PCB-059/062/075 9590 20300 40300 J 32800 29100 J 21000 J 25000 11700 PCB-066 PCB-066 48700 82300 114000 J 14800 135000 J 110000 J 25000 J 107000 48500 PCB-061/070/074/076 PCB-062 48700 82300 119000 J 928000 825000 J 693000 J 73000 48500 PCB-063 PCB-064 61400 121000 J 38800 J 30700 825000 J 69300 J 73000 381000 PCB-064 PCB-066 51400 121000 2820000 J 227100 J 22100 J 22400 10300 PCB-067 PCB-066 217000 372000 855000 J 683000 J 576000 J 497000 J 4330 J <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>										
PCB-058 PCB-058 357 807 1490 J 1060 1160 J 5350 J 717 480 PCB-059/062/075 PCB-059/062/075 959 0 20300 40300 J 32800 29100 J 21000 J 26500 11200 PCB-060 PCB-060 PCB-060 48700 82300 174000 J 148000 135000 J 11000 J 107000 48500 PCB-061/07/07/07/076 PCB-061 37000 623000 119000 J 928000 825000 J 69300 J 73000 73000 831000 PCB-063 PCB-063 8400 16000 38800 J 30700 27100 J 22100 J 22400 10300 PCB-064 PCB-063 8400 11000 262000 J 220000 179000 J 13300 J 161000 76400 PCB-066 PCB-066 217000 37200 805000 J 63800 57600 J 497000 J 497000 J 43300 J 4440 PCB-076 PCB-068 655 1190 3020 J		PCB-056								
PCB-059/062/075 PCB-059/062/075 PCB-059/062/075 PCB-050 RCB-050	PCB-057	PCB-057								
PCB-060 PCB-060 48700 82300 174000 J 148000 135000 J 110000 J 107000 48500 PCB-061/070/074/076 PCB-0E12 307000 623000 1190000 J 928000 825000 J 693000 J 730000 381000 PCB-063 PCB-063 8400 16000 38800 J 30700 27100 J 22100 J 22400 10300 PCB-064 PCB-066 61400 121000 262000 J 220000 179000 J 497000 J 501000 230000 PCB-066 PCB-066 217000 372000 805000 J 638000 576000 J 497000 J 501000 230000 PCB-067 PCB-066 217000 37200 805000 J 5170 5280 J 3710 J 4330 4240 PCB-067 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 21.3 U PCB-072 PCB-073 63.4 U 61.1 U 14800 J 1900 26.4 U 7400 J <td>PCB-058</td> <td>PCB-058</td> <td>357</td> <td>807</td> <td>1490 J</td> <td></td> <td>1160 J</td> <td>5350 J</td> <td>717</td> <td>480</td>	PCB-058	PCB-058	357	807	1490 J		1160 J	5350 J	717	480
PCB-061/070/074/076 PCB-CE12 307000 623000 119000 J 928000 825000 J 693000 J 730000 381000 PCB-063 PCB-063 8400 16000 38800 J 30700 27100 J 22100 J 22400 10300 PCB-064 PCB-064 61400 121000 262000 J 220000 179000 J 133000 J 161000 76400 PCB-066 PCB-066 217000 372000 805000 J 638000 576000 J 497000 J 501000 230000 PCB-067 PCB-067 2060 7520 7230 J 5170 5280 J 3710 J 4330 4240 PCB-068 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 213 U PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 1990 26.4 UJ 7410 J 9730 </td <td>PCB-059/062/075</td> <td>PCB-059/062/075</td> <td>9590</td> <td>20300</td> <td>40300 J</td> <td>32800</td> <td>29100 J</td> <td>21000 J</td> <td>26500</td> <td>11200</td>	PCB-059/062/075	PCB-059/062/075	9590	20300	40300 J	32800	29100 J	21000 J	26500	11200
PCB-063 PCB-063 8400 16000 38800 J 30700 27100 J 22100 J 22400 10300 PCB-064 PCB-064 61400 121000 262000 J 220000 179000 J 133000 J 161000 76400 PCB-066 PCB-066 217000 372000 805000 J 638000 576000 J 497000 J 501000 230000 PCB-067 PCB-067 2060 7520 7230 J 5170 5280 J 3710 J 4330 4240 PCB-068 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 21.3 U PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U	PCB-060	PCB-060	48700	82300	174000 J	148000	135000 J	110000 J	107000	48500
PCB-064 61400 12100 262000 J 220000 179000 J 133000 J 161000 76400 PCB-066 PCB-066 217000 372000 805000 J 638000 576000 J 497000 J 501000 230000 PCB-067 PCB-067 2060 7520 7230 J 5170 5280 J 3710 J 4330 4240 PCB-068 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 21.3 U PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-061/070/074/076	PCB-CE12	307000	623000	1190000 J	928000	825000 J	693000 J	730000	381000
PCB-066 217000 372000 805000 J 638000 576000 J 497000 J 501000 230000 PCB-067 PCB-067 2060 7520 7230 J 5170 5280 J 3710 J 4330 4240 PCB-068 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 21.3 U PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-077 2940 13100 18500 J 16000 17400 J 15100 13900 7360 PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-063	PCB-063	8400	16000	38800 J	30700	27100 J	22100 J	22400	10300
PCB-067 PCB-067 2060 7520 7230 J 5170 5280 J 3710 J 4330 4240 PCB-068 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 21.3 U PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-078 13100 18500 J 16000 17400 J 15100 13900 7360 PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-064	PCB-064	61400	121000	262000 J	220000	179000 J	133000 J	161000	76400
PCB-068 PCB-068 655 1190 3020 J 2340 2190 J 1740 J 2010 21.3 U PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-077 2940 13100 18500 J 16000 17400 J 15100 13900 7360 PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-066	PCB-066	217000	372000	805000 J	638000	576000 J	497000 J	501000	230000
PCB-072 PCB-072 1530 2630 6400 J 5190 4720 J 3740 J 4000 1830 PCB-073 PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-077 2940 13100 18500 J 16000 17400 J 15100 13900 7360 PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-067	PCB-067	2060	7520	7230 J	5170	5280 J	3710 J	4330	4240
PCB-073 63.4 U 61.1 U 14800 J 10900 26.4 UJ 7410 J 9730 4710 PCB-077 PCB-077 2940 13100 18500 J 16000 17400 J 15100 13900 7360 PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-068	PCB-068	655	1190	3020 J	2340	2190 J	1740 J	2010	21.3 U
PCB-077 PCB-077 2940 13100 18500 J 16000 17400 J 15100 13900 7360 PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-072	PCB-072	1530	2630	6400 J	5190	4720 J	3740 J	4000	1830
PCB-078 PCB-078 56.7 U 78.3 U 47.2 UJ 30.9 U 28.6 UJ 37.5 UJ 69.7 U 23.5 U	PCB-073	PCB-073	63.4 U	61.1 U	14800 J	10900	26.4 UJ	7410 J	9730	4710
	PCB-077	PCB-077	2940	13100	18500 J	16000	17400 J	15100	13900	7360
PCB-079 PCB-079 969 1820 40.3 UJ 26.4 U 2590 J 2220 J 1870 20.2 U	PCB-078	PCB-078	56.7 U	78.3 U	47.2 UJ	30.9 U	28.6 UJ	37.5 UJ	69.7 U	23.5 U
	PCB-079	PCB-079	969	1820	40.3 UJ	26.4 U	2590 J	2220 J	1870	20.2 U

				Tissue Ana	lytical Results				
	Location ID	PB057.2	PB057.2	PB057.2	PB057_B	PB057_B	PB057_B	PB058.1	PB058.1
	Sample ID			PB03-F-A-STM-W-002-20110617	PB03-F-A-GKF-W-001-20110615	PB03-F-A-GKF-W-006-20110617	PB03-F-A-PNF-W-001-COMP-201106		PB03-F-A-GM-W-004-20110618
	Sample Date		6/17/2011	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/18/2011	6/18/2011
	Sample Type		N	N	N	N	N	N	N
	y X	3201603.32	3201603.32	3201603.32	3201644.06	3201644.06	3201644.06	3201682.3	3201682.3
	^ V	13832054.52	13832054.52	13832054.52	13832071.63	13832071.63	13832071.63	13832011.93	13832011.93
	Scientific_Name	Mugil cephalus	Mugil cephalus	Mugil cephalus	Fundulus grandis	Fundulus grandis	Lagodon rhomboides	Brevoortia patronus	Brevoortia patronus
	Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017	6460	11900	12300 J	34800	47500	4270	22500	34400
PCB-018/030	PCB-018/030	17400	29700	33700 J	84600	112000	18300	49600	88100
PCB-019	PCB-019	846	1810	1600	3000	4110	882	3070	4620
PCB-020/028	PCB-020/028	77000	91100	99700 J	261000	312000	107000	138000	143000
PCB-021/033	PCB-020/028	10400	15700	15800 J	24300	40800	2890	39700	48200
PCB-022	PCB-021/033	13700	16600	17700 J	44600	56500	7540	31900	39000
PCB-022 PCB-023	PCB-022	33.8 J	36.8	51.5 J	44600	123	7540 24.5	72.2	66.2
PCB-023 PCB-024	PCB-023	33.8 J 147	178	251 J	499	675	161	288	468
PCB-025	PCB-025	2470	3050	3320 J	7070	9550	1120	5260	6240
PCB-026/029	PCB-026/029	8680	10800	11900 J	26400	34400	10300	16500	18800
PCB-027	PCB-027	791	1320	1410 J	4310	5410	416	2490	4160
PCB-031	PCB-031	67600	82400	88500 J	220000	275000	103000	132000	143000
PCB-032	PCB-032	6020	10200	11000 J	35800	43700	2130	19100	28500
PCB-034	PCB-034	224	277	338 J	647	863	144	567	539
PCB-035	PCB-035	54.8	47	65.1 J	43.4 U	35.4 U	10.3	125	107
PCB-036	PCB-036	21.4 U	6.58 U	40.3 UJ	41.7 U	33.5 U	4.15 U	16.1 U	10.7 U
PCB-037	PCB-037	2870	4070	2730 J	4330	6760	903	5510	8280
PCB-038	PCB-038	106	82.8	102 J	273	397	143	146	119
PCB-039	PCB-039	21.1 U	592	38.2 UJ	40.5 U	31.3 U	148	15.2 U	1190
PCB-040/071	PCB-040/071	21400 J	27900	30600 J	127 U	134000	8590	69100 J	103000
PCB-041	PCB-041	2620 J	3630	3980 J	166 U	22000	798	15100 J	23500
PCB-042	PCB-042	12400 J	17700	19400 J	140 U	70100	14900	43900 J	63600
PCB-043	PCB-043	23.5 UJ	3740	80.5 UJ	146 U	17.8 U	1980	35.6 UJ	9760
PCB-044/047/065	PCB-044/047/065	78400 J	99900	111000 J	124 U	385000	132000	164000 J	233000
PCB-045/051	PCB-045/051	7430 J	11900	11900 J	30600	37200	7490	20200 J	40700
PCB-046	PCB-046	1310 J	2310	2160 J	3130	4230	463	4530 J	10900
PCB-048	PCB-048	13000 J	17800	20100 J	137 U	62500	4080	43900 J	55500
PCB-049/069	PCB-049/069	67600 J	89400	96900 J	118 U	359000	122000	140000 J	175000
PCB-050/053	PCB-050/053	5920 J	9170	9680 J	32300	39400	2160	18500 J	38500
PCB-052	PCB-052	97100 J	140000	143000 J	140 U	612000	209000	209000 J	312000
PCB-054	PCB-054	128 J	144	149	212	271	115	237 J	311
PCB-055	PCB-055	167 J	613	1520 J	149 U	3790	399 J	2300 J	1720
PCB-056	PCB-056	34400 J	40300	44400 J	67900	87900	12300	80200 J	81200
PCB-057	PCB-057	529 J	504	669 J	1350	1690	259 J	942 J	817
PCB-058	PCB-058	270 J	294	371 J	713	878	256	467 J	16.7 U
PCB-059/062/075	PCB-059/062/075	6760 J	7570	9050 J	106 U	32400	10300	12600 J	17400
PCB-060	PCB-060	26800 J	27800	32900 J	113000	127000	42500	53400 J	52100
PCB-061/070/074/076	PCB-CE12	201000 J	224000	268000 J	730000	859000	328000	410000 J	391000
PCB-063	PCB-063	5720 J	5420	7300 J	22200	25900	9080	11000 J	8930
PCB-064	PCB-064	37600 J	47400	51800 J	99.1 U	203000	77600	82800 J	120000
PCB-066	PCB-066	121000 J	137000	166000 J	508000	582000	230000	245000 J	216000
PCB-067	PCB-067	2590 J	2550	3270 J	3520	5190	829	4760 J	4550
PCB-068	PCB-068	508 J	418	612 J	1720	1880	623	867 J	750
PCB-072	PCB-072	1020 J	920	1280 J	3760	4190	1280	1900 J	1630
PCB-073	PCB-073	1700 J	13.3 U	2680 J	111 U	12800	24 U	4750 J	9.72 U
PCB-077	PCB-077	4380	5600	4350 J	13500	15600	602	8430	12200
PCB-078	PCB-078	21.7 UJ	15 U	66.8 UJ	140 U	57 U	18.4 U	23.9 UJ	15.5 U
PCB-079	PCB-079	18.4 UJ	12.7 U	57.5 UJ	122 U	48.7 U	15.6 U	20.5 UJ	1150

Location ID Sample ID Sample ID Sample ID Sample Date Sample Type X Y Scientific_Name Size_Class Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-018/030 PCB-018/030 PCB-019 PCB-019 PCB-019 PCB-020/028 PCB-020/028 PCB-021/033 PCB-022 PCB-022 PCB-022 PCB-022 PCB-023 PCB-022 PCB-023 PCB-024 PCB-024 PCB-024 PCB-024 PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-027 PCB-027 PCB-031 PCB-031 PCB-031 PCB-031 PCB-031 PCB-032 PCB-034 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-036 PCB-036 PCB-036 PCB-037 PCB-038 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-041 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-044 PCB-044 PCB-045/0551 PCB-045/0551 PCB-045/0551 PCB-046 PCB-048 PCB-049/069 PB03-F-A-GM-W-005-20110618 6/18/2011 N 3201682.3 13832011.93 Brevoortia patronus =<15 Migratory	PB059.1_A PB03-F-A-GKF-W-003-20110615 6/15/2011	PB059.1_A PB03-I-A-BCR-W-003-COMP-201106 6/19/2011	PB059.3 PB03-I-B-BCR-W-002-20110627 6/26/2011 N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307 1.03 U	PB059.4 PB03-I-B-BCR-W-003-20110628 6/27/2011 N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200 281	PB059.4 PB03-I-B-BCR-W-003-20110628DUP 6/27/2011 N 3201714.16 13831964.63 Result 216 439 18.3 67600	PB059.5 PB03-I-A-BCR-W-001-COMP-201106 6/27/2011	PB063.1_A PB03-F-A-GKF-W-009-20110617 6/17/2011 N 3201509.99 13831505.57 Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900 3040	
Sample ID Sample Date Sample Type X Y Scientific_Name Size_Class Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-020/028 PCB-021/033 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-023 PCB-023 PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-039 PCB-041 PCB-039 PCB-041 PCB-042 PCB-042 PCB-044 PCB-045 PCB-045 PCB-045 PCB-045 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046	PB03-F-A-GM-W-005-20110618 6/18/2011 N 3201682.3 13832011.93 Brevoortia patronus =<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	PB03-F-A-GKF-W-003-20110615 6/15/2011 N 3201501.32 13831828.71 Fundulus grandis =<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	PB03-I-A-BCR-W-003-COMP-201106 6/19/2011 N 3201501.32 13831828.71 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	PB03-I-B-BCR-W-002-20110627 6/26/2011 N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	PB03-I-B-BCR-W-003-20110628 6/27/2011 N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200	PB03-I-B-BCR-W-003-20110628DUP 6/27/2011 N 3201714.16 13831964.63 Result 216 439 18.3	PB03-I-A-BCR-W-001-COMP-201106 6/27/2011 N 3201714 13831965 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 62.1 87.7	PB03-F-A-GKF-W-009-20110617 6/17/2011 N 3201509.99 13831505.57 Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900
Sample Date Sample Type X Y Scientific_Name Size_Class Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-019 PCB-019 PCB-020/028 PCB-021/033 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-031 PCB-032 PCB-034 PCB-035 PCB-034 PCB-035 PCB-036 PCB-036 PCB-037 PCB-036 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-041 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044 PCB-044 PCB-045 PCB-046	6/18/2011 N 3201682.3 13832011.93 Brevoortia patronus =<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	6/15/2011 N 3201501.32 13831828.71 Fundulus grandis =<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	6/19/2011 N 3201501.32 13831828.71 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	6/26/2011 N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	6/27/2011 N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200	6/27/2011 N 3201714.16 13831964.63 Result 216 439 18.3	6/27/2011 N 3201714 13831965 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 62.1	6/17/2011 N 3201509.99 13831505.57 Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900
Sample Type	N 3201682.3 13832011.93 Brevoortia patronus =<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	N 3201501.32 13831828.71 Fundulus grandis =<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	N 3201501.32 13831828.71 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	N 3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200	N 3201714.16 13831964.63 Result 216 439 18.3	N 3201714 13831965 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 62.1	N 3201509.99 13831505.57 Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900
Scientific_Name Size_Class	3201682.3 13832011.93 Brevoortia patronus =<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	3201501.32 13831828.71 Fundulus grandis =<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	3201501.32 13831828.71 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	3201714.16 13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200	3201714.16 13831964.63 Result 216 439 18.3	3201714 13831965 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 62.1 87.7	3201509.99 13831505.57 Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900
Size_Class Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-034 PCB-032 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-036 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-041 PCB-041 PCB-042 PCB-043 PCB-043 PCB-044 PCB-044 PCB-045 PCB-045 PCB-045 PCB-046	13832011.93 Brevoortia patronus =<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	13831828.71 Fundulus grandis =<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	13831828.71 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	13831964.63 Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2	Result 216 439 18.3	13831965 Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 62.1 87.7	13831505.57 Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900
Size_Class Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-034 PCB-032 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-036 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-041 PCB-041 PCB-042 PCB-043 PCB-043 PCB-044 PCB-044 PCB-045 PCB-045 PCB-045 PCB-046	Brevoortia patronus =<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	Fundulus grandis =<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	Callinectes sapidus >7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200	Result 216 439 18.3	Callinectes sapidus >2 to =<7.5 Sedentary Whole Body Result 62.1 87.7	Fundulus grandis =<15 Sedentary Whole Body Result 30500 72900
Size_Class Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-034 PCB-032 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-036 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-041 PCB-041 PCB-042 PCB-043 PCB-043 PCB-044 PCB-044 PCB-045 PCB-045 PCB-045 PCB-046	=<15 Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	=<15 Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	>2 to =<7.5 Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	>7.5 to =<13 Migratory Whole Body Result 128 341 20.3 72200 164 307	>7.5 to =<13 Migratory Whole Body Result 226 459 20.2 69200	216 439 18.3	>2 to =<7.5 Sedentary Whole Body Result 62.1 87.7	=<15 Sedentary Whole Body Result 30500 72900
Tissue_Category Tissue_Type Chemical CAS Number PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-020/028 PCB-021/033 PCB-022 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-032 PCB-034 PCB-035 PCB-036 PCB-036 PCB-037 PCB-036 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-041 PCB-039 PCB-041 PCB-041 PCB-041 PCB-041 PCB-041 PCB-041 PCB-041 PCB-041 PCB-042 PCB-043 PCB-044 PCB-043 PCB-044 PCB-045 PCB-045 PCB-046	Migratory Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	Sedentary Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	Sedentary Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	Migratory Whole Body Result 128 341 20.3 72200 164 307	Migratory Whole Body Result 226 459 20.2 69200	216 439 18.3	Sedentary Whole Body Result 62.1 87.7	Sedentary Whole Body Result 30500 72900
Chemical CAS Number PCB-017 PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-019 PCB-020/028 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-031 PCB-031 PCB-032 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-034 PCB-036 PCB-035 PCB-037 PCB-036 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-042 PCB-043 PCB-042 PCB-044 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	Whole Body Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	Whole Body Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	Whole Body Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	Whole Body Result 128 341 20.3 72200 164 307	Whole Body Result 226 459 20.2 69200	216 439 18.3	Whole Body Result 62.1 87.7	Whole Body Result 30500 72900
Chemical CAS Number PCB-017 PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-019 PCB-020/028 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-022 PCB-024 PCB-023 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-034 PCB-036 PCB-035 PCB-037 PCB-038 PCB-038 PCB-037 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-042 PCB-042 PCB-042 PCB-043 PCB-042 PCB-044 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	Result 36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	Result 44900 106000 2980 312000 33700 56000 143 582 8790 32000	Result 59.5 79.1 6.57 J 88600 128 320 1.26 U	Result 128 341 20.3 72200 164 307	Result 226 459 20.2 69200	216 439 18.3	Result 62.1 87.7	Result 30500 72900
PCB-017 PCB-018/030 PCB-018/030 PCB-019 PCB-019 PCB-020/028 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-035 PCB-037 PCB-036 PCB-038 PCB-037 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	36500 92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	44900 106000 2980 312000 33700 56000 143 582 8790 32000	59.5 79.1 6.57 J 88600 128 320 1.26 U	128 341 20.3 72200 164 307	226 459 20.2 69200	216 439 18.3	62.1 87.7	30500 72900
PCB-018/030 PCB-019 PCB-019 PCB-019 PCB-020/028 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-035 PCB-037 PCB-036 PCB-038 PCB-039 PCB-039 PCB-039 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-042 PCB-044 PCB-043 PCB-0440 PCB-044 PCB-044 PCB-045/051 PCB-045/051 PCB-045/051 PCB-046 PCB-048	92700 4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	106000 2980 312000 33700 56000 143 582 8790 32000	79.1 6.57 J 88600 128 320 1.26 U	341 20.3 72200 164 307	459 20.2 69200	439 18.3	87.7	72900
PCB-019 PCB-019 PCB-020/028 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-031 PCB-031 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-035 PCB-035 PCB-035 PCB-036 PCB-035 PCB-037 PCB-036 PCB-038 PCB-039 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044 PCB-044 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	4620 149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	2980 312000 33700 56000 143 582 8790 32000	6.57 J 88600 128 320 1.26 U	20.3 72200 164 307	20.2 69200	18.3		
PCB-020/028 PCB-020/028 PCB-021/033 PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-031 PCB-031 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-036 PCB-038 PCB-039 PCB-039 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	149000 51000 41000 66.3 584 J 6550 19700 4440 149000 29800	312000 33700 56000 143 582 8790 32000	88600 128 320 1.26 U	72200 164 307	69200		0.74	
PCB-021/033 PCB-022 PCB-022 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-027 PCB-031 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-035 PCB-036 PCB-037 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-044/047/065 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-046 PCB-048	51000 41000 66.3 584 J 6550 19700 4440 149000	33700 56000 143 582 8790 32000	128 320 1.26 U	164 307		6/600	93400	197000
PCB-022 PCB-023 PCB-023 PCB-023 PCB-024 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-044/047/065 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	41000 66.3 584 J 6550 19700 4440 149000 29800	56000 143 582 8790 32000	320 1.26 U	307	281	370		
PCB-023 PCB-024 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-041 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	66.3 584 J 6550 19700 4440 149000 29800	143 582 8790 32000	1.26 U		F70	278	122 J	27600
PCB-024 PCB-025 PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-042 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	584 J 6550 19700 4440 149000 29800	582 8790 32000		1113 11	578	553	308	34800
PCB-025 PCB-025 PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-044/047/065 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	6550 19700 4440 149000 29800	8790 32000	1.34 U		0.778 U	1.86 U	2.23 U	103
PCB-026/029 PCB-026/029 PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	19700 4440 149000 29800	32000	CC 1	1.07 U	0.809 U	1.97 U	2.37 U	413
PCB-027 PCB-027 PCB-031 PCB-031 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	4440 149000 29800		66.1	77.2	86.9	84.3	31.1	6790
PCB-031 PCB-032 PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	149000 29800	4020	146	183	294	291	61.1	23200
PCB-032 PCB-032 PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	29800	4830	4.53 J	20.6	50.7	46.3	5.76	3800
PCB-034 PCB-034 PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048		276000	1930	1710	2050	1960	1050	168000
PCB-035 PCB-035 PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048		44900	128	213	549	512	115	26900
PCB-036 PCB-036 PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-043 PCB-043 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048		772	14.2	10.7	28.9	28.6	8.58	627
PCB-037 PCB-037 PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	106	49.6 U	14.6 J	17.2	10.8 J	10.9 J	7.82 U	31.5 U
PCB-038 PCB-038 PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-049/04/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	11.3 U	47.6 U	4.55 U	2.35 U	2.05 U	4.47 U	6.99 U	29.3 U
PCB-039 PCB-039 PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	8360	6980	5630	7650	6220	5950	7760	4860
PCB-040/071 PCB-040/071 PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	115	48.7 U	4.79 U	25.2	20.9	20.4	7.35 U	275
PCB-041 PCB-041 PCB-042 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	1170	46.3 U	4.29 U	58.5	39.6	35.6	53.3	28.1 U
PCB-042 PCB-042 PCB-043 PCB-043 PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	104000	122000	659	1200	3620	3390	398	79900
PCB-043 PCB-043 PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	20400	19400	22	42.3	3.78 U	7.43 U	21.1	11200
PCB-044/047/065 PCB-044/047/065 PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	61900	59100	276	528	681	663	147	39600
PCB-045/051 PCB-045/051 PCB-046 PCB-046 PCB-048 PCB-048	9700	150 U	111	186	444	420	66.4	102 U
PCB-046 PCB-046 PCB-048 PCB-048	228000	356000	37300	50200	46400	43100	40800	213000
PCB-048 PCB-048	41800	34700	39	121	165	145	43.1	20400
	11300	3490	5.38 J	19.1	21.5	22.8 J	3.58 U	2400
	55000	47700	118	310	356	337	51.9	39900
	173000	335000	526	940	1040	953	325	211000
PCB-050/053 PCB-050/053	40200	35200	22.4	106	149	136	27.6	21300
PCB-052 PCB-052	307000	556000	417	1500	1900	1720	458	329000
PCB-054 PCB-054	302	255	3.47 U	1.02 U	1.87 J	2.23 U	4.95 U	259
PCB-055 PCB-055	1360	2610 J	6.33 U	4.48 U	5 U	10.2 U	6.99 U	73.6 U
PCB-056 PCB-056	79300	83500	414	693	894	883	274	56600
PCB-057 PCB-057	822	1560	8.77 J	15.3	33.4	35.2	11.6	1250
PCB-058 PCB-058	14.5 U	834	6.39 U	4.37 U	16.9	15	7.07 U	658
PCB-059/062/075 PCB-059/062/075	16700	29500	2750	3670	3360	3130	3070	18700
PCB-060 PCB-060	51000	124000	32600	37100	34900	33700	36500	87200
PCB-061/070/074/076 PCB-CE12	379000	838000	72500	80500	72100	71300	82500	584000
PCB-063 PCB-063	8780	24600	1360	1360	1300	1260	1040	18400
PCB-064 PCB-064	117000	193000	1160	1790	2450	2280	532	114000
PCB-066 PCB-066	206000	558000	128000	135000	122000	124000	147000	398000
PCB-067 PCB-067	4410	4490	85.8	94.8	112	107	40	3650
PCB-068 PCB-068	744	1680	181	227	192	184	191	1430
PCB-072 PCB-072		3880	72.6	57.2	117	116	42.2	3190
PCB-073 PCB-073	1550	10600	2.88 U	3.17 U	2.31 U	4.69 U	3.13 U	6200
PCB-077 PCB-077	13.7 U	16200	5980	8160	7250	6880	8710	13200
PCB-078 PCB-078 PCB-079 PCB-079		105 U 91.7 U	6.24 U 5.1 U	4.1 U 101	4.57 U 4.03 U	9.46 U 8.19 U	6.89 U 116	69.1 U 57.9 U

				1133UC AII	alytical Results				
	Location ID	PB063.1_A	PB063.2_A	PB063.2 A	PB069_A	PB069_A	PB069_A	PB084.1	PB084.1
	Sample ID		PB03-F-A-GKF-W-005-20110615	PB03-F-A-GKF-W-013-20110617	PB03-F-A-GKF-W-002-20110615	PB03-F-A-GKF-W-007-20110617	PB03-I-A-BCR-W-002-COMP-201106	PB04-F-A-GKF-W-005-20110627	PB04-F-A-GKF-W-013-20110627
	Sample Date	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/17/2011	6/19/2011	6/27/2011	6/27/2011
	Sample Type		N	N	N	N	N	N	N
	x x	3201509.99	3201561.28	3201561.28	3201132.36	3201132.36	3201132.36	3200727.85	3200727.85
	v	13831505.57	13831519.55	13831519.55	13830993.22	13830993.22	13830993.22	13829784.47	13829784.47
	Scientific_Name	Fundulus grandis	Fundulus grandis	Fundulus grandis	Fundulus grandis	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Fundulus grandis
	Size_Class	=<15	=<15	=<15	=<15	=<15	>2 to =<7.5	=<15	=<15
	Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-017	PCB-017	39700	37400	29600	47700	35200	162	23600	35400
PCB-018/030	PCB-018/030	96600	89200	63700	112000	81400	488	60200	88500
PCB-019	PCB-019	3530	3380	3520	4600	4330	38.5	2730	4160
PCB-020/028	PCB-020/028	313000	263000	224000	331000	241000	153000	113000	176000
PCB-021/033	PCB-021/033	38900	26300	23700	45300	32800	250	15500	27800
PCB-022	PCB-022	54100	44200	41500	55200	41700	344	24600	38200
PCB-023	PCB-022	129	88.6	88.2	140	107	2.31 U	43.9	65.7
PCB-024	PCB-023	599	471	427	694	562	2.62 U	323	522
PCB-024 PCB-025	PCB-024 PCB-025	9820	7980	6810	11300	8500	124	4820	7300
PCB-025 PCB-026/029	PCB-025	34300	28800	24500	37900	28400	398	14500	22500
PCB-026/029 PCB-027	PCB-026/029	4800	4240	3200	6100	4600	20.6	3000	4450
PCB-027 PCB-031	PCB-027	265000	229000	188000	284000	204000	2440	102000	157000
PCB-032	PCB-031	38600	36700	29200	43500	32100	211	21900	33900
PCB-034	PCB-032	857	670	590	1030	719	20.6	314	524
PCB-035	PCB-035	41.3 U	75.8 U	16.2 U	43.6 U	24.8 U	22.6 J	20.7	37
PCB-036	PCB-036	39.6 U	73.8 U	15.1 U	39.7 U	22.6 U	4 U	6.6 U	6.94 U
PCB-037	PCB-037	6220	5110	5350	7390	6360	12900	3120	6130
PCB-038	PCB-037	300	340	285	394	329	3.99 U	79.9	7.19 U
PCB-039	PCB-039	38.5 U	67.8 U	14.3 U	39.2 U	22.3 U	96.6	493	6.98 U
PCB-040/071	PCB-040/071	102000	115000	78800	164000	97900 J	1360	64700	102000
PCB-041	PCB-040/071	11500	13000	11100	18600	11500 J	49.3	10500	15100
PCB-042	PCB-041	47800	54800	40600	76800	45200 J	49.3	34900	50600
PCB-043	PCB-042	10900	14800	9170	21100	11800 J	416	6680	10500
PCB-044/047/065	PCB-044/047/065	285000	321000	215000	426000	245000 J	85700	175000	266000
PCB-044/047/003	PCB-044/047/003	23500	28200	19400	44200	23800 J	185	24800	36600
PCB-046	PCB-046	2600	2830	2140	5230	2780 J	29.1	3530	4940
PCB-048	PCB-048	47700	42900	29300	82800	46100 J	447	29400	48300
PCB-049/069	PCB-049/069	272000	305000	200000	409000	224000 J	1220	145000	225000
PCB-050/053	PCB-049/009	24100	32100	19700	46800	24200 J	178	25600	35900
PCB-052	PCB-052	438000	491000	319000	643000	346000 J	2260	263000	411000
PCB-054	PCB-052	302	241	244	320	290 J	4.86 U	174	253
PCB-055	PCB-055	3350	4690	1640	3190	2020 J	10.5 U	788	1260
PCB-056	PCB-056	75500	72000	59700	91300	63400 J	542	34000	51800
PCB-056 PCB-057	PCB-056	1440	1450	1150	1950	1250 J	30.7	602	915
PCB-057 PCB-058	PCB-057	852	772 J	665	1040	665 J	10.5 U	375	591
PCB-059/062/075	PCB-059/062/075	24600	26500	18600	37200	21800 J	6260	13800	21800
PCB-060	PCB-060	117000	108000	84100	129000	86200 J	53400	43500	68600
PCB-061/070/074/076	PCB-CE12	782000	747000	542000	945000	594000 J	138000	290000	471000
PCB-063	PCB-063	23600	22200	16800	29300	18500 J	2450	7810	12500
PCB-064	PCB-064	155000	169000	116000	228000	133000 J	2080	94000	147000
PCB-066	PCB-066	544000	526000	374000	625000	404000 J	259000	181000	295000
PCB-067	PCB-067	4440	4020	3310	6270	4070 J	166	1830	3040
PCB-068	PCB-068	1870	1650	1360	2300	1450 J	425	612	912
PCB-008 PCB-072	PCB-072	3780	3710	2900	5150	3120 J	160	1310	2040
PCB-072	PCB-072	56.2 U	81.2 U	13 U	22 U	29 UJ	6.03 U	5.96 U	8.84 U
PCB-073	PCB-073	14300	12700	12100	19100	14900	13400	7550	12400
PCB-077 PCB-078	PCB-077	62.6 U	110 U	22.8 U	69.8 U	33.9 UJ	10.5 U	20.5 U	12400 16 U
1 60 070	1 CB-076	02.0 0	110 0	1590	03.0 0	33.9 01	171	20.3 0	10 0

Appendix D
Tissue Analytical Results

Tissue Analytical Results							
	Location ID	PB095.2	PB101_D	PB101_D	PB101_D	PB102	
	Sample ID	PB04-I-A-BCR-W-001-COMP-201106	PB04-F-A-GKF-W-001-20110627	PB04-F-A-GKF-W-001-COMP-201106	PB04-F-A-GKF-W-009-20110627	PB04-F-A-PNF-W-001-COMP-201106	
	Sample Date	6/29/2011	6/27/2011	6/27/2011	6/27/2011	6/27/2011	
	Sample Type	N N	N	N	N	N N	
	х , , , ,	3200882	3201317.36	3201317.36	3201317.36	3201302.69	
	Υ	13829528	13828141.83	13828141.83	13828141.83	13828160.48	
	Scientific_Name	Callinectes sapidus	Fundulus grandis	Fundulus grandis	Fundulus grandis	Lagodon rhomboides	
	Size_Class	>2 to =<7.5	=<15	=<15	=<15	=<15	
	Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Migratory	
		Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	
Chemical	Tissue_Type CAS Number	Result	Result	Result	Result	Result	
PCB-017	PCB-017	39.1	26200	41500	23500	18500	
		33.1			57500		
PCB-018/030	PCB-018/030		66900	89300		71100	
PCB-019	PCB-019	8.96 U	2750	3330	1950	3110	
PCB-020/028	PCB-020/028	49800	143000	234000	138000	259000	
PCB-021/033	PCB-021/033	49.1	20600	25900	13700	9760	
PCB-022	PCB-022	96.3	30400	42800	27200	22500	
PCB-023	PCB-023	5 U	57.2	97.1	51.5	88.4	
PCB-024	PCB-024	5.19 U	364	538	336	687	
PCB-025	PCB-025	12.5 J	5610	9110	5570	3930	
PCB-026/029	PCB-026/029	17.5	17700	29400	17400	29700	
PCB-027	PCB-027	5.07 U	3270	4750	2740	1770	
PCB-031	PCB-031	286	126000	211000	122000	244000	
PCB-032	PCB-032	48.2	25000	37100	22800	8830	
PCB-034	PCB-034	4.85 U	418	675	399	511	
PCB-035	PCB-035	11.2 U	19.3 J	47.2	12.4 J	173 U	
PCB-036	PCB-036	10.4 U	8.94 U	15.2 U	8.78 U	166 U	
PCB-037	PCB-037	3410	3730	5650	2800	1940	
PCB-038	PCB-038	26.6	112	247	112	169 U	
PCB-039	PCB-039	22.5 J	597	14.6 U	544 J	159 U	
PCB-040/071	PCB-040/071	135	83400	117000	85100	29200	
PCB-041	PCB-041	13.5 U	12400	16900	12100	6550 U	
PCB-042	PCB-042	52.8 J	42700	67900	40700	50100	
PCB-042 PCB-043	PCB-042	12.5 U	8690	15600	9450	6730 U	
	PCB-043	28500	225000	332000	234000	320000	
PCB-044/047/065							
PCB-045/051	PCB-045/051	13 U	28900	35300	28900	25000	
PCB-046	PCB-046	15.4 U	3660	4020	2890	1770	
PCB-048	PCB-048	26.4	37500	59200	34400	14600	
PCB-049/069	PCB-049/069	88.5	195000	304000	211000	279000	
PCB-050/053	PCB-050/053	20.4 J	29600	37100	30800	8620	
PCB-052	PCB-052	119	345000	491000	382000	395000	
PCB-054	PCB-054	9.78 U	173	237	143	348	
PCB-055	PCB-055	19.7 U	1090	1260	1050	532 U	
PCB-056	PCB-056	28.2 J	42900	73400	43900	41900	
PCB-057	PCB-057	18.6 U	842	1350	960	895	
PCB-058	PCB-058	19.1 U	389	706	422	512 U	
PCB-059/062/075	PCB-059/062/075	2160	18600	26800	19100	22200	
PCB-060	PCB-060	19900	63800	83300	65500	80700	
PCB-061/070/074/076	PCB-CE12	42300	430000	648000	461000	629000	
PCB-063	PCB-063	368	12000	18500	12600	14700	
PCB-064	PCB-064	149	128000	168000	132000	174000	
PCB-066	PCB-066	71000	276000	433000	287000	428000	
PCB-067	PCB-067	17.2 U	2540	4530	2770	2240	
PCB-068	PCB-068	86.3	874	1520	1070	1270	
PCB-008	PCB-072	17.7 U	1960	3310	2130	2500	
PCB-072	PCB-072	19.9	6.63 U	26.8 U	20.1 U	3950 U	
PCB-073	PCB-073	3920	10300	13800	11500	1650	
		3920 17.9 U			11500 14.8 U	492 U	
PCB-078	PCB-078		16.5 U	27.8 U			
PCB-079	PCB-079	41.6	14.3 U	24.2 U	12.8 U	428 U	

					115542	Analytical Results				
	Location ID	Field QC	Field QC	PB002.1	PB003.1	PB005.1	PB006.1	PB008.2	PB010	PB010.1
	Sample ID	EB-110621-A01	EB-110624-A01	PB01-I-A-BCR-W-001-COMP-201106	PB01-F-A-STM-W-006-20110618	PB01-I-A-BCR-W-002-COMP-201106	PB01-I-B-WHS-W-002-COMP-201106	PB01-I-B-WHS-W-001-COMP-201106	PB01-F-A-GM-W-008-20110618	PB01-I-A-WHS-W-001-COMP-201106
	Sample Date	6/24/2011	6/24/2011	6/20/2011	6/18/2011	6/20/2011	6/20/2011	6/19/2011	6/18/2011	6/20/2011
	Sample Type	EB	EB	N	N	N N	N	N N	N	N N
	,			3202691	3202445.49	3202312	3202026.63	3202032	3201710.87	3201704.8
	Y			13836499	13836263.53	13836201	13836356.8	13836072	13836029.16	13836071.32
	Scientific_Name			Callinectes sapidus	Mugil cephalus	Callinectes sapidus	Penaeus setiferus	Penaeus setiferus	Brevoortia patronus	Penaeus setiferus
	Size_Class			>2 to =<7.5	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13	=<15	>2 to =<7.5
	Tissue_Category			Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type			Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080		34.4 U	5.21 U	8.19 UJ	4.9 U	14.6 UJ	21.6 U	12.1 UJ	46.2 U
PCB-081	PCB-081		32.5 U		78.9	115	233	135	147	217
PCB-082	PCB-082		406	5.06 U	1670 J	8.06 J	935 J	448	4070 J	905
PCB-083	PCB-083		327	5.17 U	760 J	5.92 U	1110 J	515 J	1390 J	1110
PCB-084	PCB-084		734	37.2	2890 J	30.1	727 J	344	5860 J	778
PCB-085/116/117	PCB-085/116/117		1410	4790	4490 J	6640	11800 J	7850	10700 J	10800
PCB-086/087/097/108/119/125	PCB-CE2		3880	1300	12300 J	1840	18100 J	10600	26200 J	15900
PCB-088/091	PCB-088/091		1060	81.5	3310 J	105	1950 J	1120	6710 J	1820
PCB-089	PCB-089		32.7 U	4.92 U	222 J	5.63 U	60.1 J	22.5	571 J	69.2
PCB-090/101/113	PCB-090/101/113		8980	598	20900 J	826	38900 J	29800	41100 J	37800
PCB-092	PCB-092		1870	59.5	3230 J	42.8	4550 J	2360	6900 J	4780
PCB-092 PCB-093/100	PCB-093/100		178	337	322 J	472	313 J	272	550 J	341
PCB-094	PCB-094		40.3 J	5.61	105 J	13.2	34.6 J	15 J	253 J	34.8
PCB-094	PCB-095		3020	133	9520 J	60.3	4810 J	2690	19000 J	4850
PCB-093	PCB-095		14 U	2.81 U	122 J	2.28 U	13.4 J	7.19 J	236 J	17
PCB-090 PCB-098/102	PCB-098/102		82.7	2.81 U	847 J	18.3	437 J	203	1500 J	388
PCB-099	PCB-099		6300	18500	14100 J	26500	27300 J	21000	28700 J	25300
	PCB-103			4.24 U			2/300 J 248 J	156		249
PCB-103			205 12.1 U		281 J	3.76 J			504 J 14.7 J	
PCB-104	PCB-104			2.01 U 8740	13.1 J 7360	1.46 U 11700	7.31 U 21200 J	4.19 U		9.93 U
PCB-105	PCB-105		1890 36.4 U					13400	16900	20000 J
PCB-106	PCB-106				13.6 J	6.32 U	15.3 UJ	41.1 U	32.6 J	21.9 UJ
PCB-107/124	PCB-107/124		213	22.7	725 J	39.7	1720 J	1260	1560 J	1620 J
PCB-109	PCB-109		1070	1080	1520 J	1690	3460 J	2400	3420 J	2990 J
PCB-110/115	PCB-110/115		5350	825	18200 J	1190	27600 J	17600	36600 J	24300 J
PCB-111	PCB-111		34.5	24.1	23 J	31.7 J	42.4 J	28.4	40.5 J	39.1 J
PCB-112	PCB-112		23 U		34.5 J	4.04 U	7.7 UJ	5.59 U	93.6 J	13.3 UJ
PCB-114	PCB-114		93.7 J	620	487	792	1470	978	1150	1330
PCB-118	PCB-118		7030	22700	17600 J	30200	48800 J	32700	37800 J	46200
PCB-120	PCB-120		189	157	148 J	205	220 J	173	260 J	218 J
PCB-121	PCB-121		22.8 U		11.1 J	16.7	7.57 UJ	5.5 U	16.4 J	13.1 UJ
PCB-122	PCB-122		38.7 U		261 J	6.98 U	629 J	418	569 J	573 J
PCB-123	PCB-123		95 J	476	440	623	1100 J	758	866	1120
PCB-126	PCB-126		45.1 U	68.8	41.8 J	95.6	133 J	69.6	104	147 J
PCB-127	PCB-127		34 U		27.3 J	6.37 U	15.3 UJ	41.1 U	51.8 J	21.9 UJ
PCB-128/166	PCB-128/166		1400	1290	1820	1400	2700 J	1800	3590	2740 J
PCB-129/138/163	PCB-129/138/163		12200	11000	13300	12100	19800 J	14000	25700	19200 J
PCB-130	PCB-130		780	189	949	268	1450 J	1010	1870	1320 J
PCB-131	PCB-131		52.9	7.7 J	110	7.32 U	24.1 J	69.2 U	238	21.2 UJ
PCB-132	PCB-132		1640	14.2	2740	14.6	885 J	414	5010	825 J
PCB-133	PCB-133		416	237	291	277	375 J	300	550	382 J
PCB-134	PCB-134		309	10.4 U	433	8.06 U	276 J	159	800	383 J
PCB-135/151	PCB-135/151		3040	50.9	3070	51.2	2140 J	1290	6180	2460 J
PCB-136	PCB-136		530	11.6	958	9.4	103 J	55.2	1640	114 J
PCB-137	PCB-137		261	511	660	647	1030 J	730	1290	959 J
PCB-139/140	PCB-139/140		205	273	285	369	304 J	229	514	344 J
PCB-141	PCB-141		783	8.79 U	1920	7.55	2980 J	2250	3700	2970 J
PCB-142	PCB-142		23.3 U		5.84 U	6.96 U	8.67 UJ	69.5 U	11.3 J	21.3 UJ
PCB-143	PCB-143		22.2 U	8.44 U	27.3	6.56 U	12 J	61.9 U	45.8	18.9 UJ

				rissue Ana	alytical Results				
	Location ID	PB010.3	PB013.1_A	PB013.3	PB013.3	PB015.1	PB015.2	PB015_A	PB015_A
	Sample ID	PB01-I-A-BCR-W-003-COMP-201106	_	PB01-F-A-GKF-W-004-COMP-201106	PB01-F-A-STM-W-003-20110616	PB01-F-A-GKF-W-002-COMP-201106	PB01-I-A-BCR-W-004-COMP-201106	_	PB01-I-A-BCR-W-006-COMP-20110
	Sample Date	6/19/2011	6/22/2011	6/21/2011	6/15/2011	6/15/2011	6/16/2011	6/25/2011	6/22/2011
	Sample Type		N	N	N	N	N	N	N N
	х х	3201743	3201306.47	3201433.88	3201433.88	3201175	3201180	3201168.73	3201168.73
	Y	13836152	13836191.23	13836075.12	13836075.12	13836251	13836183	13836254.66	13836254.66
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Fundulus grandis	Mugil cephalus	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Callinectes sapidus
	Size_Class	>2 to =<7.5	>2 to =<7.5	=<15	=<15	=<15	>2 to =<7.5	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Sedentary	Sedentary	Migratory	Sedentary	Sedentary	Sedentary	Sedentary
	Tissue_Type	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	8.01 U	4.42	28.4 U	31.5 U	17.5 U	9.81 U	43.8 U	17.7 U
PCB-081	PCB-081	253	218	573	239	492	307	681	223
PCB-082	PCB-082	41.8 J	5.89 J	5310	3120	4670	14.4	5420 J	21.2 UJ
PCB-083	PCB-083	7.82 U	2.38 U	4590	2880	3410	8.93 U	5630 J	23.6 UJ
PCB-084	PCB-084	94.6	42	8170	9070	7900	42.5	10800 J	52.2 J
PCB-085/116/117	PCB-085/116/117	17400	9620	45100	16300	38300	19200	62600 J	11700 J
PCB-086/087/097/108/119/125	PCB-083/110/117	4070	1940	81900	35200	71400	3760	110000 J	2470 J
PCB-086/087/097/108/119/125 PCB-088/091	PCB-088/091	391	1940	19400	8880	18300	165	28100 J	2470 J 117 J
PCB-088/091 PCB-089	PCB-088/091	7.44 U	2.56	19400	564	1210	8.48 U	28100 J 1520 J	20.3 UJ
PCB-090/101/113	PCB-099/101/113	4080	867	1410	63100	123000	2140	218000 J	928 J
PCB-090/101/113 PCB-092	PCB-090/101/113	205	80.9	28100	10500	24100	56.9	36800 J	88.3 J
		762			947	1640	839	2650 J	485 J
PCB-093/100	PCB-093/100		348	2090					
PCB-094	PCB-094	22.8 264	13.8	724	291	624	12.6 J 89.8	769 J	19.8 UJ
PCB-095	PCB-095		134	70800	28800	56100		87300 J	152 J
PCB-096	PCB-096	3.14 U	0.738 U	450	467	345	3.68 U	619 J	5.03 UJ
PCB-098/102	PCB-098/102	49.8	27.1	4380	2080	3910	29.5	6280 J	33 J
PCB-099	PCB-099	51400	27400	113000	41600	97800	59000	152000 J	37100 J
PCB-103	PCB-103	23.2	2.92	1600	767	1220	7.32 U	2130 J	18.4 UJ
PCB-104	PCB-104	2.06 U	0.93 U	27.3	34.5	26.1	2.43 U	43.6 J	6.78 UJ
PCB-105	PCB-105	29200	18500	70400	24500	62800	31100	101000	21600
PCB-106	PCB-106	9.73 U	3.82 U	49.9 U	24.9 U	9.84 U	12.7 U	20.2 UJ	20.9 UJ
PCB-107/124	PCB-107/124	189	50.7	4570	2440	4000	90.9	5680 J	42 J
PCB-109	PCB-109	4160	2230	13200	5140	11300	4220	17500 J	2360 J
PCB-110/115	PCB-110/115	4600	2010	115000	53700	103000	3370	156000 J	2140 J
PCB-111	PCB-111	47.7	26.3	103	53.8	87.3	51.6	160 J	39.8 J
PCB-112	PCB-112	5.34 U	1.47 U	5.78 U	16.9 U	5.95 U	6.1 U	14.1 UJ	14 UJ
PCB-114	PCB-114	2180	1340	5700	1650	4800	2300	7790	1430
PCB-118	PCB-118	68400	40600	151000	54000	126000	70900	216000	48500
PCB-120	PCB-120	328	170	581	335	524	348	867 J	292 J
PCB-121	PCB-121	22.8 J	1.46 U	5.67 U	21.3	5.84 U	29.1	13.1 UJ	14 UJ
PCB-122	PCB-122	49.5	7.88	559	696	588 J	14.1 U	516 J	24 UJ
PCB-123	PCB-123	1510	973	3850	1450	3250	1610	5560	1140
PCB-126	PCB-126	196	125	502	132	360 J	213	704	156
PCB-127	PCB-127	54.3 J	3.57 U	50.4 U	81.9 J	9.93 U	12.8 U	19.1 UJ	20.5 UJ
PCB-128/166	PCB-128/166	4190	1720	10600	5490	8820	3570	15900 J	2690
PCB-129/138/163	PCB-129/138/163	31900	14500	77700	38800	70300	28700	124000 J	23500
PCB-130	PCB-130	899	264	5270	2610	4430	640	8080 J	376
PCB-131	PCB-131	18.8	40.7 U	210	324	193	11.1 U	251 J	25.4 U
PCB-132	PCB-132	48.6	37.8 U	10800	5790	9520	23.5	14300 J	23.2 U
PCB-133	PCB-133	464	230	1560	786	1320	457	2580 J	393
PCB-134	PCB-134	13 J	39.6 U	1470	1070	1340	12.2 U	2000 J	27.3 U
PCB-135/151	PCB-135/151	179	49.4	16900	8120	14500	61.6 J	26300 J	78.9
PCB-136	PCB-136	30.5 J	9.72 J	2810	2590	2260	13.2	3860 J	8.79 J
PCB-137	PCB-137	2050	977	5270	2030	4610	1920	8440 J	1220
PCB-139/140	PCB-139/140	853	446	1920	750	1630	866	2920 J	514
PCB-141	PCB-141	182	33 U	11000	5590	9610	49 J	16700 J	11.2 J
PCB-142	PCB-142	11.6 U	37.3 U	9.36 U	20.5 U	10.6 U	13.1 J	13.6 UJ	22.8 U
PCB-143	PCB-143	11 U	34.6 U	141	20 U	122	9.95 U	198 J	21.2 U

Appendix D
Tissue Analytical Results

				HISSUC AII	alytical Results				
	Location ID	PB015 A	PB015_A	PB015_A	PB016.2	PB019.1	PB019.1	PB019.1	PB019.1
	Sample ID	PB01-I-A-BCR-W-007-COMP-201106	_	_	PB01-F-A-GKF-W-003-COMP-201106	PB01-F-A-SAS-W-005-20110622	PB01-I-B-BRS-W-012-20110621	PB01-I-B-BRS-W-014-20110621	PB01-I-B-BRS-W-015-20110621
	Sample Date	6/22/2011	6/22/2011	6/20/2011	6/15/2011	6/22/2011	6/21/2011	6/21/2011	6/21/2011
	Sample Type		N	N	N	N	N	N	N N
	х х	3201168.73	3201168.73	3201168.73	3201190	3201152.87	3201152.87	3201152.87	3201152.87
	Y	13836254.66	13836254.66	13836254.66	13836042	13835753.58	13835753.58	13835753.58	13835753.58
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Penaeus aztecus	Fundulus grandis	Cynoscion arenarius	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus
	Size_Class	>2 to =<7.5	>2 to =<7.5	>7.5 to =<13	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13
	Tissue_Category	Sedentary	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	6.32 UJ	8.98 UJ	29.5 U	14.3 U	14.6 UJ	197 U	49.8 U	72.5 U
PCB-081	PCB-081	238 J	334 J	345	257	46.2	504 J	413	412
PCB-082	PCB-082	11.6 J	26.6 J	1470 J	2040	4060 J	2810	2040 J	1940
PCB-083	PCB-083	5.46 UJ	6.45 UJ	1400 J	1980	12 UJ	223 U	57.4 UJ	107 U
PCB-084	PCB-084	40 J	34.3 J	972 J	4030	4810 J	1940	1600 J	1650
PCB-085/116/117	PCB-085/116/117	10700 J	15100 J	16100 J	22600	18100 J	26800	21900 J	19900
PCB-086/087/097/108/119/125	PCB-083/110/117	2000 J	2680 J	22600 J	38200	29200 J	38300	32500 J	27500
PCB-088/091	PCB-CE2 PCB-088/091	106 J	99.1 J	1800 J	10400	5130 J	3820	2740 J	2870
PCB-088/091 PCB-089	PCB-088/091 PCB-089	4.7 UJ	5.56 UJ	72.7 J	568	409 J	195 U	128 J	133 J
PCB-090/101/113	PCB-089 PCB-090/101/113	4.7 UJ 972 J	1230 J	72.7 J 50700 J	78700	409 J 46700 J	97400	71000 J	69400
PCB-090/101/113 PCB-092	PCB-090/101/113	972 J 64.9 J	1230 J 58.1 J	6090 J	14200	46700 J 7700 J	11400	8240 J	8210
PCB-093/100	PCB-092 PCB-093/100	310 J	371 J	6090 J 467 J	1060	810 J	11400	738 J	690
·									
PCB-094	PCB-094	7.96 J	5.42 UJ	35.7 J	310	78 J	202 U	78.2 J	87.2
PCB-095	PCB-095	106 J	79.2 J	5590 J	33100	12000 J	11600	8610 J	8980
PCB-096	PCB-096	1.18 UJ	1.31 UJ	28.4 J	213	327 J	127 U	66.4 J	71.4
PCB-098/102	PCB-098/102	20.6 J	20.3 J	511 J	2130	926 J	1220	873 J	866
PCB-099	PCB-099	29400 J	39300 J	38600 J	63200	44800 J	67400	52000 J	53400
PCB-103	PCB-103	4.26 UJ	5.04 UJ	364 J	825	591 J	824	552 J	590
PCB-104	PCB-104	2.28 UJ	2.94 UJ	5.45 UJ	14.8	13.1 J	126 U	39.5 U	55.4 U
PCB-105	PCB-105	22000 J	31100 J	29100	34100	25200 J	43400	39400 J	34200
PCB-106	PCB-106	6.43 UJ	6.31 UJ	12.2 UJ	6.11 U	11.4 UJ	171 U	55.2 UJ	92.3 U
PCB-107/124	PCB-107/124	55.9 J	75.5 J	2400 J	1980	1710 J	3820	3480 J	3230
PCB-109	PCB-109	2710 J	3750 J	5520 J	6620	5030 J	8110	7710 J	6460
PCB-110/115	PCB-110/115	1910 J	2390 J	24100 J	54800	44100 J	38300	35000 J	28700
PCB-111	PCB-111	38.6 J	46.5 J	52.9 J	55	50.7 J	133 U	88.3 J	93
PCB-112	PCB-112	3.25 UJ	26.4 J	7.34 UJ	4.15 U	7.18 UJ	134 U	34.5 UJ	64 U
PCB-114	PCB-114	1620 J	2140 J	2110	2750	1840 J	2970	2850 J	2370
PCB-118	PCB-118	51700 J	69900 J	65800	78500	55600 J	98100	87400 J	76800
PCB-120	PCB-120	225 J	278 J	313 J	325	317 J	463	454 J	407
PCB-121	PCB-121	13.4 J	15.9 J	7.09 UJ	4.08 U	17.4 J	135 U	34.8 UJ	64.6 U
PCB-122	PCB-122	7.36 UJ	7.23 UJ	854 J	294	137 J	1240	1210 J	1160
PCB-123	PCB-123	1180 J	1620 J	1770	1820	1040 J	2610	2190 J	2100
PCB-126	PCB-126	289 J	195 J	190 J	200	13.9 UJ	200 U	51.2 UJ	170 J
PCB-127	PCB-127	6.3 UJ	6.18 UJ	11.5 UJ	6.17 U	74.8 J	168 U	54.3 UJ	90.7 U
PCB-128/166	PCB-128/166	2460	3560	3860	5190	5120 J	7130	6560	6170
PCB-129/138/163	PCB-129/138/163	19000	26100	27700	42800	36400 J	57500	43300	40600
PCB-130	PCB-130	441	550	2090	2570	1400 J	3830	3340	3020
PCB-131	PCB-131	12.1	11.8 J	77.9	81.5	222 J	269 U	116	126
PCB-132	PCB-132	16	19	923	4960	3960 J	2000	1380	1360
PCB-133	PCB-133	336	414	599	851	628 J	1090	792	831
PCB-134	PCB-134	6.94 U	6.22 U	466	629	570 J	1010 J	43.1 U	537
PCB-135/151	PCB-135/151	50.7	56.5	3330	8960	4300 J	6410	4170	4310
PCB-136	PCB-136	9.51 J	9.28 J	152	1270	1080 J	508	257	301
PCB-137	PCB-137	1140	1550	1320	2630	2230 J	2760	2110	2070
PCB-139/140	PCB-139/140	398	512	509	930	858 J	1210	733	740
PCB-141	PCB-141	5.66 U	59.3	3390	5290	4220 J	7530	5590	5300
PCB-142	PCB-142	5.81 U	5.21 U	8.42 U	7.62 U	15.4 J	268 U	39.5 U	68.9 U
PCB-143	PCB-143	5.4 U	4.84 U	7.68 U	62.7	10.9 UJ	250 U	36.8 U	64.2 U

Tissue Analytical Results												
	Location ID	PB019.1	PB019.1	PB019.1	PB019.2	PB019_A	PB019_A	PB024.1	PB024.1			
	Sample ID		PB01-I-B-BRS-W-022-20110622	PB01-I-B-BRS-W-023-20110622		PB01-F-A-GKF-W-005-COMP-201106		PB01-F-A-GM-W-001-20110615	PB01-F-A-GM-W-002-20110615			
	Sample Date		6/22/2011	6/22/2011	6/28/2011	6/21/2011	6/23/2011	6/15/2011	6/15/2011			
	Sample Type		N	N	N	N	0,25,2011 N	N	N			
	y Sumple Type	3201152.87	3201152.87	3201152.87	3201194.8	3201314.96	3201314.96	3201090.96	3201090.96			
	^ v	13835753.58	13835753.58	13835753.58	13835735.1	13835750.59	13835750.59	13835160.37	13835160.37			
	Scientific_Name	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus	Crassostrea virginica	Fundulus grandis	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus			
	Size_Class	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15			
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Migratory	Migratory			
	Tissue_Type		Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body			
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result			
PCB-080	PCB-080	104 U	19.2 U	58.8 U	12.5 U	14.1 U	11.9	27 U	47.7 U			
PCB-081	PCB-081	570	459	406	288	556	266	196	207			
PCB-082	PCB-082	2660	2870	2590	5710 J	4720	6.88 UJ	6770	5780			
PCB-082	PCB-082	1870	13.6 U	37.4 U	2990 J	4380	3.75 UJ	2290	42.8 U			
PCB-084	PCB-083	2130	2280	2010	9590 J	7500	62.8 J	9920	9020			
PCB-085/116/117	PCB-085/116/117	30200	29500	27000	12800 J	40600	12400 J	16800	14600			
PCB-085/116/117 PCB-086/087/097/108/119/125	PCB-085/116/117	43300	42400	38700	38000 J	77100	2930 J	41700	36700			
PCB-088/091	PCB-CE2 PCB-088/091	43300 3570	3530	3280	9490 J	19200	2930 J 202 J	10900	10000			
<u> </u>	The state of the s	203 J	3530 178	3280 149			6.59 UJ	965	876			
PCB-089	PCB-089 PCB-090/101/113	105000	97800	86900	995 J	1210	6.59 UJ 2170 J		55100			
PCB-090/101/113		105000			56300 J	139000	21/0 J 155 J	61400				
PCB-092	PCB-092	11900	12000	10500 907	9130 J 885 J	25300		10500	9150 827			
PCB-093/100	PCB-093/100		1040			1620	491 J	878				
PCB-094	PCB-094	104 U	85.8	79.9	383 J	550	19.7 J	427	395			
PCB-095	PCB-095	11900	12000	10500	29500 J	61200	157 J	31700	29600			
PCB-096	PCB-096	91.1 J	76.8	63.2	479 J	339	1.29 UJ	412	388			
PCB-098/102	PCB-098/102	1140	1080	988	2640 J	4080	38.9 J	2320	2150			
PCB-099	PCB-099	68300	69300	63500	35900 J	107000	37500 J	41200	38000			
PCB-103	PCB-103	908	762	685	677 J	1460	11.2 J	713	669			
PCB-104	PCB-104	60.4 U	4.58 U	11.8 U	36.9 J	28.1 J	3.58 UJ	18.7	10.2 U			
PCB-105	PCB-105	45800	40100	38100	20300	71000	25500	25000	21600			
PCB-106	PCB-106	110 U	15.9 U	44.1 U	11 UJ	25.2 U	9.56 UJ	22.1 U	44.1 U			
PCB-107/124	PCB-107/124	4440	4130	3710	1850 J	4560	118 J	2500	2200			
PCB-109	PCB-109	9190	8750	8040	4790 J	13500	3230 J	5230	4600			
PCB-110/115	PCB-110/115	43300	42700	39400	47900 J	111000	2840 J	57000	50500			
PCB-111	PCB-111	111	86.4	75 J	45.6 J	120	70.4 J	49.9	58.2 J			
PCB-112	PCB-112	415 J	8.35 U	22.9 U	7.94 UJ	7.38 U	4.55 UJ	8.91 U	28.9 U			
PCB-114	PCB-114	3300	2860	2580	1290	5200	1780	1680	1480			
PCB-118	PCB-118	113000	91200	88000	56100	154000	57800	55200	45400			
PCB-120	PCB-120	565 69.6 U	514 31.8	466	253 J 7.66 UJ	618 7.24 U	362 J 28.7 J	311	296 28.2 U			
PCB-121	PCB-121			30.5				18.7	709			
PCB-122	PCB-122	1640	1390	1250	758 J	438 3820	13.5 J	800				
PCB-123 PCB-126	PCB-123 PCB-126	2690 309	2240 249 J	2120 256	1220 175	3820 473 J	1350 161	1330 162	1100 137 J			
PCB-126 PCB-127		309 109 U	249 J 106	101	175 10.4 UJ	4/3 J 25.5 U	9.36 UJ	78.1	42.6 U			
	PCB-127 PCB-128/166	9820 J	7540	6400	2400	10100	9.36 UJ 3540	5930	42.6 0			
PCB-128/166	PCB-128/166 PCB-129/138/163	66400 J	52200	44600	21000	81800	28900	41500	33300			
PCB-129/138/163 PCB-130	PCB-129/138/163 PCB-130	4700 J	3780	3150	1310	5330	793	2850	2280			
PCB-131	PCB-130	237 J	159	112	173	187	17.9	397	348			
PCB-131 PCB-132	PCB-131 PCB-132	237 J 2240 J	1850	1480	5800	10300	27.6	8980	7290			
PCB-132 PCB-133	PCB-132 PCB-133	1310 J	1080	854	452	10300	712	796	682			
PCB-133 PCB-134	PCB-133		730	854 696	973	1500			1230			
		1090 J 7650 J					11.2 U	1410	7460			
PCB-135/151	PCB-135/151 PCB-136		6630 374	5170 265	6390	16700	156	9270	7460 2170			
PCB-136		429 J			1960	2470	14.6	2690				
PCB-137	PCB-137	3390 J	2570	2190	138	5190	1630	2210	1710			
PCB-139/140	PCB-139/140	1320 J	985	771	334	1730	719	854	715			
PCB-141	PCB-141	9220 J	7340	6160	235	11300	127 J	6150	5060			
PCB-142	PCB-142	119 UJ	12.8 U	31.5 U	8.48 U	10.2 U	9.34 U	18.5 U	32.9 U			
PCB-143	PCB-143	111 UJ	12.2 U	29.8 U	7.73 U	12.2 J	8.68 U	18.1 U	32.2 U			

				Hissue Alli	alytical Results				
	Location ID	PB024.4	PB025_A	PB026.1	PB032_A	PB037.2	PB037.4	PB037.5	PB037.6
	Sample ID	PB01-I-A-BCR-W-005-COMP-201106	PB01-F-A-GM-W-004-20110615	PB01-F-A-STM-W-002-20110616	PB02-F-A-PNF-W-009-20110617	PB02-F-A-STM-W-006-20110616	PB02-I-B-BCR-W-001-COMP-201106	PB02-I-B-BRS-W-001-COMP-201106	PB02-I-A-BCR-W-002-COMP-201106
	Sample Date	6/20/2011	6/15/2011	6/15/2011	6/17/2011	6/16/2011	6/17/2011	6/17/2011	6/16/2011
	Sample Type		N	N N	N N	N	N	N	N
		3201319	3201214.42	3201153.65	3201407.03	3201346.97	3201504	3201380	3201402
	Υ	13835188	13835154.42	13835079.21	13834542.25	13834038.02	13833530	13834151	13833775
	Scientific_Name	Callinectes sapidus	Brevoortia patronus	Mugil cephalus	Lagodon rhomboides	Mugil cephalus	Callinectes sapidus	Penaeus aztecus	Callinectes sapidus
	Size_Class	· ·	=<15	=<15	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Migratory	Migratory	Migratory	Migratory	Migratory	Sedentary
	Tissue_Type	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	23.9 U	26.1 U	36.3 U	36.8 U	56.6 U	11 UJ	8.38 U	9.11 J
PCB-081	PCB-081	251	346	376	265	466	454 J	321	318
PCB-082	PCB-082	103 J	11800	3540	3630 J	7290	37.3 J	1120	20.4 J
PCB-083	PCB-083	27.7 U	3600	2140 J	1690 J	5010	7.69 UJ	1140	6.93 U
PCB-084	PCB-084	328	16700	12100	6200 J	17000	106 J	680	112
PCB-085/116/117	PCB-085/116/117	14700	27100	23600	24500 J	29900	25600 J	14700	16100
PCB-086/087/097/108/119/125	PCB-CE2	4480	68400	48500	46900 J	68500	4310 J	20100	3100
PCB-088/091	PCB-088/091	817	17600	10700	12100 J	17100	179 J	1320	330
PCB-089	PCB-089	26.4 U	1670	742	191 J	1430	6.63 UJ	62.9	6.58 U
PCB-090/101/113	PCB-090/101/113	5620	96300	85100	92600 J	113000	2060 J	43900	2580
PCB-092	PCB-092	740	16300	14300	11900 J	19600	135 J	5500	205
PCB-093/100	PCB-093/100	715	1370	1330	1190 J	1770	797 J	401	619
PCB-094	PCB-094	58.3 J	726	358	56.5 J	615	18.7 J	28.2	16.3
PCB-095	PCB-095	949	52200	35700	30100 J	53700	274 J	4290	358
PCB-096	PCB-096	9.71 U	697	676	228 J	940	2.2 UJ	18.7	3.3 U
PCB-098/102	PCB-098/102	224 J	3870	2630	431 J	4450	56.6 J	376	57.1
PCB-099	PCB-099	44600	64200	58900	64100 J	66600	66100 J	33900	44500
PCB-103	PCB-103	60.9	1050	957	911 J	1270	9.35 J	290	11.5
PCB-104	PCB-104	8.05 U	28.4 J	44.6	20.6 UJ	37.5	3.23 U	2.97 U	2.4 U
PCB-105	PCB-105	24700	39000	35600	35800 J	42800	51000 J	27200	29900
PCB-106	PCB-106	27.8 U	26.7 U	28.8 U	27.5 UJ	56.1 U	13.8 UJ	10.7 U	9.97 U
PCB-107/124	PCB-107/124	234	4030	3610	3010 J	4310	106 J	2310	121
PCB-109	PCB-109	3980	8080	7470	7030 J	8790	5200 J	4890	3760
PCB-110/115	PCB-110/115	5940	91800	71100	54500 J	95100	4870 J	19900	3830
PCB-111	PCB-111	109 J	75.4	63.2 J	69.4 J	73.6	56.5 J	39.7	41.9
PCB-112	PCB-112	18.9 U	19.6 U	16 U	26.8 UJ	28.4 U	4.58 UJ	4.36 U	4.73 U
PCB-114	PCB-114	1660	2710	2700	2660	3050	3780 J	1950	2170
PCB-118	PCB-118	55100	84900	79900	87100 J	93600	110000 J	59200	64200
PCB-120	PCB-120	421	426	385	425 J	442	372 J	213	267
PCB-121	PCB-121	61 J	28.7	25.3	30.4 J	36.5	23.1 J	4.21 U	14.4 J
PCB-122	PCB-122	30.8 U	1360	1010	580 J	1380	101 J	804	18.9
PCB-123	PCB-123	1200	2120	2050	2160	2540	2820 J	1480	1530
PCB-126	PCB-126	154 J	265	202	186 J	210	366 J	184 J	202
PCB-127	PCB-127	28 U	110	65.9	91.3 J	94	13.5 UJ	10.1 U	43.3
PCB-128/166	PCB-128/166	3200	8130	6950	5960	7880	5820 J	3090	3530
PCB-129/138/163	PCB-129/138/163	26700	56100	47000	42100	54100	44700 J	23600	25200
PCB-130	PCB-130	1260	3870	3000	2430	3780	660 J	1480	546
PCB-131	PCB-131	35.1 J	649	400	263	555	11.2 UJ	57.5	19.6
PCB-132	PCB-132	120	13100	5810	5800	9470	37 J	659	32.3
PCB-133	PCB-133	908	1050	847	601	998	672 J	438	379
PCB-134	PCB-134	40.8 U	2170	1400	672	1870	12.1 UJ	298	13.1 U
PCB-135/151	PCB-135/151	754	13300	9430	6100	12000	77.9 J	2390	115
PCB-136	PCB-136	88.5 J	3990	2980	1850	3940	21 J	102	26.5
PCB-137	PCB-137	1400	3050	2570	2240	3080	2810 J	1250	1730
PCB-139/140	PCB-139/140	812	1220	883	838	1090	923 J	417	694
PCB-141	PCB-141	416 J	8590	6930	5610	8460	9.83 UJ	3360	97.7
PCB-142	PCB-142	35.2 U	27.6	20.3 U	22.9 U	34.9 U	10.1 UJ	6.14 U	11.3 U
PCB-143	PCB-143	33.2 U	21.6 U	19.8 U	21.4 U	32.4 U	9.38 UJ	15.7	10.7 U
		ı		l		L	l .		

				Tissue Ana	alytical Results				
	Location ID	PB037_A	PB047.1_A	PB047.3	PB048.3	PB048.3	PB048.3	PB049_A	PB049_A
		PB02-I-A-BCR-W-004-COMP-201106	_	PB02-I-A-BCR-W-001-COMP-201106		PB02-F-A-GM-W-002-20110616		PB02-F-A-GKF-W-001-COMP-201106	<u> </u>
	Sample Date		6/15/2011	6/19/2011	6/16/2011	6/16/2011	6/16/2011	6/15/2011	6/22/2011
	Sample Type		N	N	N	N	N	N	N
	х х	3201325.97	3201461.79	3201645	3201519.15	3201519.15	3201519.15	3201468.39	3201468.39
	ν	13833885.45	13833007.68	13832925	13832869.89	13832869.89	13832869.89	13832722.8	13832722.8
	Scientific_Name	Callinectes sapidus	Lagodon rhomboides	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus	Mugil cephalus	Fundulus grandis	Callinectes sapidus
	Size_Class	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15	=<15	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Sedentary	Sedentary
	Tissue_Type	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	22 U	79 U	19.8 U	26.4 U	34 U	58.7 U	798 U	8.71 U
PCB-081	PCB-081	683	494	431	395	375	436	1080 U	240
PCB-082	PCB-082	15.7 U	9390	18.6 U	13000	12100	8750	6260	6.7 UJ
PCB-083	PCB-083	16.5 U	2380	19.1 U	3910	5320	3850	5210	7.26 UJ
PCB-084	PCB-084	100	11600	83.1 J	17800	17100	15700	10800	25.9 J
PCB-085/116/117	PCB-085/116/117	34900	59100	22900	30800	28800	28800	67700	9870 J
PCB-086/087/097/108/119/125	PCB-083/110/117	5210	111000	3580	76100	72200	67300	121000	2150 J
PCB-086/087/108/119/125 PCB-088/091	PCB-CE2	271	24100	191	18900	17700	15600	29800	78.8 J
PCB-088/091 PCB-089	PCB-088/091	15.2 U	252	191 18.1 U	1810	17700	15600	1830	6.69 UJ
PCB-089 PCB-090/101/113	PCB-089 PCB-090/101/113	2220	191000	18.1 0	106000	101000	98500	219000	797 J
PCB-090/101/113 PCB-092	PCB-090/101/113	112	21700	62.8 J	18300	17300	17000	44200	797 J 54 J
PCB-092 PCB-093/100	PCB-092 PCB-093/100	1050	2000	778	1470	1440	1660	2870	375 J
PCB-093/100 PCB-094	PCB-093/100 PCB-094		89.9 U			755	591	914	10.5 J
		28.2 322	54600	39.1	777				
PCB-095	PCB-095			146	55200	52300	48100	99000	79.2 J
PCB-096	PCB-096	11 U	389	10.3 U	731	697	869	539	2.62 UJ
PCB-098/102	PCB-098/102	83.9	490	48.2 J	4020	3970	4450	7030	16.2 J
PCB-099	PCB-099	95400	123000	60700	71400	60800	59000	168000	29300 J
PCB-103	PCB-103	13.1 U	1450	15.6 U	1100	1050	1030	2370	5.77 UJ
PCB-104	PCB-104	9.74 U	16.4 U	7.11 U	29.8 J	32.5	35.8	37.5 J	4.01 UJ
PCB-105	PCB-105	60300	86400	40500	47800	42800	39600	111000	21800
PCB-106	PCB-106	16.9 U	86.8 U	23.1 U	21.3 U	32.6 U	131 J	20.1 U	18.7 UJ
PCB-107/124	PCB-107/124	87.7	7080	87	4730	4180	3830	6100	40 J
PCB-109	PCB-109	6380	16200	4780	9350	8810	7920	16900	2430 J
PCB-110/115	PCB-110/115	5960	129000	4040	100000	96300	88000	165000	2020 J
PCB-111	PCB-111	50.8 J	58.1 U	48.8	75.4	71.8	45.9 J	136	46 J
PCB-112	PCB-112	10.6 U	64.9 U	13 U	14.3 U	473	406	8.16 U	44.4 J
PCB-114	PCB-114	4090	6680	3090	3390	3040	2950	8650	1630
PCB-118	PCB-118	119000	188000	89100	103000	90000	82100	219000	51200
PCB-120	PCB-120	311	637	278	466	421	299	726	246 J
PCB-121	PCB-121	10.4 U	61 U	12.9 U	28.1 J	30.7	35.1 U	8.01 U	16.2 J
PCB-122	PCB-122	17.2 U	881	25.5 U	1550	1610	1590	825	21.4 UJ
PCB-123	PCB-123	3450	4910	2090	2640	2130	2100	5480	1170
PCB-126	PCB-126	419	407	263	317	296	232	1220 J	138
PCB-127	PCB-127	15.5 U	191	67.5	136	116	54.5 U	20.3 U	17.5 UJ
PCB-128/166	PCB-128/166	5740	14300	3960	9390	8490	6320	14000	2140
PCB-129/138/163	PCB-129/138/163	38200	93700	27100	64000	59000	42000	109000	18700
PCB-130	PCB-130	497	4820	503	4300	4110	2810	6970	314
PCB-131	PCB-131	19.6	605	23.9 U	640	653	489	214	7.17 U
PCB-132	PCB-132	27.3	11600	23.1 U	13300	12900	8130	13000	17.1
PCB-133	PCB-133	569	920	428	1110	1080	684	1860	355
PCB-134	PCB-134	4.3 U	1020	26.3 U	2060 J	2160	1680	1590	6.96 U
PCB-135/151	PCB-135/151	39.2	10500	60.5	13600	12900	8960	22400	40
PCB-136	PCB-136	13 J	3080	11.3 U	3880	3790	3030	3070	5.62
PCB-137	PCB-137	3030	5780	2130	3560	3260	2510	7660	1050
PCB-139/140	PCB-139/140	1220	1820	827	1290	1260	875	2380	431
PCB-141	PCB-141	3.58 U	12500	22.3 U	9570	8990	6630	15600	61.3
PCB-142	PCB-142	4.05 U	51.8 U	15	27.2	30	30.5 U	9.37 U	6.56 U
PCB-143	PCB-143	3.75 U	48.1 U	21.4 U	15.3 U	109	117 J	179 J	6.08 U

Cocation Sample Sample Sample Sample Sample Sample Sample Sample Typ	PB02-F-A-SAS-W-002-20110616 6/16/2011 N 3201515.24 13832564.31 Cynoscion arenarius =<15 Migratory	PB052.1 PB02-F-A-STM-W-003-20110616 6/16/2011 N 3201515.24 13832564.31 Mugil cephalus =<15 Migratory Whole Body Result 71.9 U 815 11100 7100	PB053.2 PB02-F-A-GKF-W-025-20110621 6/21/2011 N 3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result 43.1 UJ 1080	PB053.2 PB02-F-A-GKF-W-029-20110622 6/22/2011 N 3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result 28.2 U	PB053_B PB02-F-A-GKF-W-013-20110618 6/18/2011 N 3201390.48 13832440.54 Fundulus grandis =<15 Sedentary Whole Body Result	PB056_A PB02-F-A-GKF-W-010-20110618 6/18/2011 N 3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body Result	PB056_A PB02-F-A-GKF-W-011-20110618 6/18/2011 N 3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body Result	PB057.2 PB03-F-A-GM-W-001-20110617 6/17/2011 N 3201603.32 13832054.52 Brevoortia patronus =<15 Migratory Whole Body
Sample Dat Sample Dat Sample Typ Scientific_Nam Size_Clas Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-088/091 PCB-088/091	6 6/16/2011 N 3201515.24 13832564.31 Cynoscion arenarius =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	6/16/2011 N 3201515.24 13832564.31 Mugil cephalus =<15 Migratory Whole Body Result 71.9 U 815	6/21/2011 N 3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result 43.1 UJ 1080	6/22/2011 N 3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result	6/18/2011 N 3201390.48 13832440.54 Fundulus grandis =<15 Sedentary Whole Body	6/18/2011 N 3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body	6/18/2011 N 3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body	6/17/2011 N 3201603.32 13832054.52 Brevoortia patronus =<15 Migratory Whole Body
Sample Typ Scientific_Nam Size_Clas Tissue_Tategor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	N 3201515.24 13832564.31 Cynoscion arenarius =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	N 3201515.24 13832564.31 Mugil cephalus =<15 Migratory Whole Body Result 71.9 U 815	N 3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result 43.1 UJ 1080	N 3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result	N 3201390.48 13832440.54 Fundulus grandis =<15 Sedentary Whole Body	N 3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body	N 3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body	N 3201603.32 13832054.52 Brevoortia patronus =<15 Migratory Whole Body
Scientific_Nam Size_Clas Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	3201515.24 13832564.31 Cynoscion arenarius =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	3201515.24 13832564.31 Mugil cephalus =<15 Migratory Whole Body Result 71.9 U 815	3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result 43.1 UJ 1080	3201266.16 13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result	3201390.48 13832440.54 Fundulus grandis =<15 Sedentary Whole Body	3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body	3201420.22 13832183.54 Fundulus grandis =<15 Sedentary Whole Body	3201603.32 13832054.52 Brevoortia patronus =<15 Migratory Whole Body
Scientific_Nam Size_Clas Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	3201515.24 13832564.31 Cynoscion arenarius =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	13832564.31 Mugil cephalus =<15 Migratory Whole Body Result 71.9 U 815	13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result 43.1 UJ 1080	13832433.69 Fundulus grandis =<15 Sedentary Whole Body Result	13832440.54 Fundulus grandis =<15 Sedentary Whole Body	13832183.54 Fundulus grandis =<15 Sedentary Whole Body	13832183.54 Fundulus grandis =<15 Sedentary Whole Body	13832054.52 Brevoortia patronus =<15 Migratory Whole Body
Size_Clas Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	Cynoscion arenarius =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	Mugil cephalus =<15 Migratory Whole Body Result 71.9 U 815	Fundulus grandis =<15 Sedentary Whole Body Result 43.1 UJ 1080	Fundulus grandis =<15 Sedentary Whole Body Result	Fundulus grandis =<15 Sedentary Whole Body	Fundulus grandis =<15 Sedentary Whole Body	Fundulus grandis =<15 Sedentary Whole Body	Brevoortia patronus =<15 Migratory Whole Body
Size_Clas Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	s =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	=<15 Migratory Whole Body Result 71.9 U 815 11100	=<15 Sedentary Whole Body Result 43.1 UJ 1080	=<15 Sedentary Whole Body Result	=<15 Sedentary Whole Body	=<15 Sedentary Whole Body	=<15 Sedentary Whole Body	=<15 Migratory Whole Body
Size_Clas Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	s =<15 Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	Migratory Whole Body Result 71.9 U 815 11100	Sedentary Whole Body Result 43.1 UJ 1080	Sedentary Whole Body Result	Sedentary Whole Body	=<15 Sedentary Whole Body	Sedentary Whole Body	Migratory Whole Body
Tissue_Categor Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	Migratory Whole Body Result 52.1 U 138 J 8130 3680 8550	Migratory Whole Body Result 71.9 U 815 11100	Sedentary Whole Body Result 43.1 UJ 1080	Sedentary Whole Body Result	Sedentary Whole Body	Sedentary Whole Body	Sedentary Whole Body	Migratory Whole Body
Tissue_Typ Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	Per Whole Body Result 52.1 U 138 J 8130 3680 8550	Whole Body Result 71.9 U 815 11100	Whole Body Result 43.1 UJ 1080	Whole Body Result	Whole Body	Whole Body	Whole Body	Whole Body
Chemical CAS Number PCB-080 PCB-080 PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	Result 52.1 U 138 J 8130 3680 8550	71.9 U 815 11100	Result 43.1 UJ 1080	Result	Result	Result	•	,
PCB-081 PCB-081 PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	138 J 8130 3680 8550	815 11100	1080	28.2 U			nesuit	Result
PCB-082 PCB-082 PCB-083 PCB-083 PCB-084 PCB-084/116/117 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	8130 3680 8550	11100			25.8 UJ	33.9 UJ	62.9 U	21.2 U
PCB-083 PCB-083 PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	3680 8550			865	1070	890	827	383
PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	8550	7100	14100 J	9410	10700 J	8000 J	8130 J	11200 J
PCB-084 PCB-084 PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091	8550		10000 J	6220	5420 J	5560 J	60.4 UJ	19.5 UJ
PCB-085/116/117 PCB-085/116/117 PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091		25200	19800 J	14900	14200 J	8960 J	11500 J	14800 J
PCB-086/087/097/108/119/125 PCB-CE2 PCB-088/091 PCB-088/091		47800	106000 J	87000	89200 J	68000 J	63500 J	27000 J
PCB-088/091 PCB-088/091	51900	103000	192000 J	152000	156000 J	18.5 UJ	124000 J	66800 J
	7970	23100	45500 J	36200	33300 J	23800 J	29700 J	15800 J
	836	2190	3390 J	2510	2390 J	1670 J	2080 J	1660 J
PCB-090/101/113 PCB-090/101/113	75200	158000	339000 J	268000	277000 J	205000 J	228000 J	94600 J
PCB-092 PCB-092	12900	26900	64100 J	52200	51300 J	37000 J	41700 J	16500 J
PCB-093/100 PCB-093/100	1130	2300	4630 J	3400	3230 J	2180 J	2760 J	1290 J
PCB-094 PCB-094	177	904	1600 J	1220	1170 J	756 J	1010 J	677 J
PCB-095 PCB-095	19400	71100	131000 J	115000	101000 J	67000 J	87900 J	44600 J
PCB-096 PCB-096	512	1250	841 J	732	640 J	382 J	603 J	639 J
PCB-098/102 PCB-098/102	1500	6050	10900 J	8550	8160 J	5360 J	7070 J	3810 J
PCB-099 PCB-099	66200	95600	229000 J	181000	211000 J	161000 J	169000 J	62500 J
PCB-103 PCB-103	790	1440	2780 J	2370	2280 J	1570 J	1970 J	921 J
PCB-104 PCB-104	25.4	52.8	49.1 J	36.8	45.2 J	79.4 J	58.1 J	24 Ј
PCB-105 PCB-105	48700	74600	143000 J	121000	142000 J	119000	106000 J	39400 J
PCB-106 PCB-106	50.6 U	269	28.7 UJ	21 U	101 J	19.3 UJ	39.6 UJ	16.7 UJ
PCB-107/124 PCB-107/124	3420	7250	12200 J	9350	10100 J	8070 J	7360 J	4150 J
PCB-109 PCB-109	8850	14100	31600 J	25900	26900 J	22500 J	19600 J	8530 J
PCB-110/115 PCB-110/115	82600	143000	257000 J	209000	205000 J	155000 J	164000 J	81700 J
PCB-111 PCB-111	63.1	93	193 J	147	179 J	138 J	120 J	59.2 J
PCB-112 PCB-112	360	732 J	25.3 UJ	12.3 U	849 J	16 UJ	38.1 UJ	11.7 UJ
PCB-114 PCB-114	3810	5780	11400 J	9400	12100 J	9790	8530	2900 J
PCB-118 PCB-118	107000	151000	282000 J	246000	279000 J	247000 J	211000 J	82500 J
PCB-120 PCB-120	409	450	1240 J	935	1080 J	895 J	770 J	366 J
PCB-121 PCB-121	22.3 U	38.7 U	63.5 J	46.1	51.1 J	41.4 J	37.3 UJ	22.1 J
PCB-122 PCB-122	568 J	2690	1530 J	1000	1540 J	1190 J	888 J	1420 J
PCB-123 PCB-123	2280	4150	8070 J	7040	8840 J	7060	6220	2120 J
PCB-126 PCB-126	202 J	423	992 J	833 J	946 J	823	731 J	281 J
PCB-127 PCB-127	136	152	374 J	272	397 J	361 J	287 J	135 J
PCB-128/166 PCB-128/166	9200	11300	23100 J	21700	21800 J	17100	15000	7060 J
PCB-129/138/163 PCB-129/138/163	59500	72100	162000 J	143000	143000 J	115000	105000	48800 J
PCB-130 PCB-130	2490	4860	10700 J	9360	9120 J	7360	6830	3340 J
PCB-131 PCB-131	401	708	510 J	348	354 J	266	256	512 J
PCB-132 PCB-132	7330	11300	24000 J	19500	17200 J	12700	13400	10100 J
PCB-133 PCB-133	916	1090	3060 J	2520	2440 J	1900	1780	863 J
PCB-134 PCB-134	1140	2580	2870 J	2530 J	2090 J	19.9 U	1610	1570 J
PCB-135/151 PCB-135/151	6590	12900	33800 J	28600	26000 J	17800	19600	10200 J
PCB-136 PCB-136	1690	4130	5120 J	4250	3640 J	2260	2870	2720 J
PCB-137 PCB-137	3770	4640	11500 J	9740	9980 J	7920	7550	2930 J
PCB-139/140 PCB-139/140	1260	1400	4200 J	3440	3180 J	2360	2380	1030 J
PCB-141 PCB-141	7460	11300	24300 J	20900	20400 J	15800	15400	7500 J
PCB-142 PCB-142	31.5 U	38.9 U	21.1 UJ	19.4 U	14.9 J	19.4 U	44.4 U	24.2 Ј
PCB-143 PCB-143	58.2	173	20.1 UJ	18.6 U	236 J	18.1 U	211	11.8 UJ

				Tissue Ana	lytical Results				
	Location ID	PB057.2	PB057.2	PB057.2	PB057_B	PB057_B	PB057_B	PB058.1	PB058.1
	Sample ID	PB03-F-A-STM-W-001-20110617	PB03-F-A-STM-W-001-COMP-201106	PB03-F-A-STM-W-002-20110617	PB03-F-A-GKF-W-001-20110615	PB03-F-A-GKF-W-006-20110617	PB03-F-A-PNF-W-001-COMP-201106	PB03-F-A-GM-W-003-20110618	PB03-F-A-GM-W-004-20110618
	Sample Date	6/17/2011	6/17/2011	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/18/2011	6/18/2011
	Sample Type	N	N	N	N	N	N	N	N
	х	3201603.32	3201603.32	3201603.32	3201644.06	3201644.06	3201644.06	3201682.3	3201682.3
	Υ	13832054.52	13832054.52	13832054.52	13832071.63	13832071.63	13832071.63	13832011.93	13832011.93
	Scientific_Name	Mugil cephalus	Mugil cephalus	Mugil cephalus	Fundulus grandis	Fundulus grandis	Lagodon rhomboides	Brevoortia patronus	Brevoortia patronus
	Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	- =<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	19.4 UJ	13.6 U	60.4 UJ	130 U	51.9 U	16.7 U	21.6 UJ	15 U
PCB-081	PCB-081	343	401	325	696	898	403	433	578
PCB-082	PCB-082	2650 J	4230	3180 J	9550	11500	4040	11900 J	21800
PCB-083	PCB-083	13.9 UJ	2780	39.6 UJ	9050	10100	1800 J	9.89 UJ	9030
PCB-084	PCB-084	6380 J	9740	7810 J	14600	17000	5580	16000 J	42800
PCB-085/116/117	PCB-085/116/117	16000 J	18200	16900 J	81000	87700	26100	27300 J	41400
PCB-086/087/097/108/119/125	PCB-CE2	32300 J	37500	36000 J	151000	161000	49100	68500 J	108000
PCB-088/091	PCB-088/091	6340 J	7690	7770 J	37900	39700	11300	16400 J	32900
PCB-089	PCB-089	502 J	743	635 J	2430	2830	148	1890 J	4120
PCB-090/101/113	PCB-090/101/113	57500 J	53500	60300 J	268000	278000	83500	94300 J	142000
PCB-092	PCB-092	9680 J	10400	10100 J	47800	51100	9310	16400 J	25500
PCB-093/100	PCB-093/100	808 J	994	893 J	3480	3440	888	1330 J	2420
PCB-094	PCB-094	218 J	261	260 J	1120	1310	32.6	791 J	1590
PCB-095	PCB-095	18800 J	26100	22700 J	116000	122000	24600	47800 J	109000
PCB-096	PCB-096	391 J	577	473 J	747	945	196	744 J	2290
PCB-098/102	PCB-098/102	1610 J	2030	2020 J	7960	9400	288	4440 J	8100
PCB-099	PCB-099	42000 J	36800	40300 J	182000	193000	59000	62100 J	87700
PCB-103	PCB-103	569 J	606	596 J	2280	2340	577	937 J	1560
PCB-104	PCB-104	37.7 J	27.6 J	30.2 J	39.5 J	46.9	1.59 U	23.4 J	49.8
PCB-105	PCB-105	27900 J	30700	25200 J	107000	118000	47100	39800 J	57500
PCB-106	PCB-106	15.6 UJ	7.02 U	33.9 UJ	96.7 U	55.8 U	4.53 U	20.4 UJ	16.1 U
PCB-107/124	PCB-107/124	2600 J	2450	2660 J	8050	8840	3150	4100 J	4620
PCB-109	PCB-109	5210 J	5250	5340 J	22400	25400	7610	8180 J	9590
PCB-110/115	PCB-110/115	45100 J	51500	47500 J	207000	217000	60400	83000 J	139000
PCB-111	PCB-111	42.4 J	31	43.4 J	132 J	138 J	40.1	57.6 J	63.8
PCB-112	PCB-112	8.37 UJ	2.7 U	23.6 UJ	78.8 U	24.4 U	3.04 U	5.91 UJ	15.6 U
PCB-114	PCB-114	2230	2320	1900 J	9070	9110	3890	2890 J	3950
PCB-118	PCB-118	56800 J	60000	53300 J	234000	257000	99200	78400 J	112000
PCB-120	PCB-120	263 J	173	232 J	809	837	245	343 J	357
PCB-121	PCB-121	16.9 J	2.68 U	22.5 UJ	80 U	46.7 J	3.02 U	19.3 J	39.7 J
PCB-122	PCB-122	778 J	910	856 J	850	1020	525 J	1490 J	1750
PCB-123	PCB-123	1710	1900	1460 J	6760	6630	2730	2130 J	2710
PCB-126	PCB-126	158 J	9.19 U	143 J	668	731	255	266 J	385
PCB-127	PCB-127	79.7 J	6.56 U	61.4 J	276	249 J	4.24 U	114 J	89.2
PCB-128/166	PCB-128/166	4170	4050	3930 J	18900	18800	5530	6610 J	8790
PCB-129/138/163	PCB-129/138/163	30900	27400	27000 J	121000	129000	41600	45700 J	57900
PCB-130	PCB-130	2010	1830	1750 J	7840	8130	1930	3100 J	4230
PCB-131	PCB-131	254	326	255 J	294	379	300	521 J	981
PCB-132	PCB-132	3460	4170	3280 J	16400	18400	5680	10000 J	19400
PCB-133	PCB-133	587	426	472 J	1910	2130	4.18 U	803 J	1050
PCB-134	PCB-134	949	850	895 J	2430	2780	224	1630 J	3680
PCB-135/151	PCB-135/151	6430	5840	4950 J	22400	26100	4270	10000 J	18100
PCB-136	PCB-136	1640	1960	1460 J	3580	4100	1260	2780 J	7150
PCB-137	PCB-137	1870	1760	1730 J	7870	8770	2610	2790 J	3690
PCB-139/140	PCB-139/140	569	664	549 J	2630	2910	922	988 J	1500
PCB-141	PCB-141	5000	4520	4060 J	17200	19000	5410	7080 J	10600
PCB-142	PCB-142	10.5 U	12.7	22.6 UJ	66 U	31.3 U	4.56 U	24.7 J	34.9 J
PCB-143	PCB-143	9.9 U	3.26 U	20.9 UJ	61.3 U	29.4 U	4.23 U	9.14 UJ	18.8 U
	1. 55 2.5	1 3.5 %	1 3.20 0	1 20.5 25	1 02.0 0	1		3.1. 00	10:0 0

				Tissue Ana	alytical Results				
	Location ID	PB058.1	PB059.1_A	PB059.1_A	PB059.3	PB059.4	PB059.4	PB059.5	PB063.1_A
	Sample ID		PB03-F-A-GKF-W-003-20110615	PB03-I-A-BCR-W-003-COMP-201106	PB03-I-B-BCR-W-002-20110627	PB03-I-B-BCR-W-003-20110628		PB03-I-A-BCR-W-001-COMP-201106	PB03-F-A-GKF-W-009-20110617
	Sample Date	6/18/2011	6/15/2011	6/19/2011	6/26/2011	6/27/2011	6/27/2011	6/27/2011	6/17/2011
	Sample Type	N	N	N N	N	N N	N	N	N
	х х	3201682.3	3201501.32	3201501.32	3201714.16	3201714.16	3201714.16	3201714	3201509.99
	Y	13832011.93	13831828.71	13831828.71	13831964.63	13831964.63	13831964.63	13831965	13831505.57
	Scientific_Name	Brevoortia patronus	Fundulus grandis	Callinectes sapidus	Callinectes sapidus	Callinectes sapidus		Callinectes sapidus	Fundulus grandis
	Size_Class	=<15	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13		>2 to =<7.5	=<15
	Tissue_Category	Migratory	Sedentary	Sedentary	Migratory	Migratory		Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body		Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	13 U	97.9 U	5.85 U	3.93 U	4.38 U	9.11 U	6.47 U	62.3 U
PCB-081	PCB-081	1220 J	861	340	437	400	368	456	631
PCB-082	PCB-082	21400	8640	22.8	52.8	39.1	40.2	28.5 J	6930 J
PCB-083	PCB-083	9520	7810	3.87 U	4.65 U	4.58 U	7.87 U	6.18 UJ	36.8 UJ
PCB-084	PCB-084	42000	13200	92.2	280	463	429	61.4 J	8840 J
PCB-085/116/117	PCB-085/116/117	41400	75900	15300	24200	23900	22600	17300 J	57200 J
PCB-086/087/097/108/119/125	PCB-CE2	109000	134000	2460	3960	3980	3810	2680 J	100000 J
PCB-088/091	PCB-088/091	33100	32000	273	419	735	687	113 J	21900 J
PCB-089	PCB-089	4040	2230	3.56 U	14.6	16.2	15.2	5.68 UJ	1590 J
PCB-090/101/113	PCB-090/101/113	141000	239000	1860	2700	3130	2930	1020 J	181000 J
PCB-092	PCB-092	25300	43000	137	314	519	484	80.1 J	31700 J
PCB-093/100	PCB-093/100	2460	3090	327	716	732	682	485 J	2020 J
PCB-094	PCB-094	1550	1070	18.2 J	38.3	109	104	16.7 J	713 J
PCB-095	PCB-095	109000	96200	259	550	1330	1220	168 J	63600 J
PCB-096	PCB-096	2290	703	1.52 U	4.57	3.74 J	4.64 U	1.8 UJ	424 J
PCB-098/102	PCB-098/102	7890	7340	69.7	136	248	241	5.42 UJ	5080 J
PCB-099	PCB-099	87800	167000	37300	60400	59000	55800	38900 J	131000 J
PCB-103	PCB-103	1620	1830	6.41	12.4	14.5	14.3	4.9 UJ	1450 J
PCB-104	PCB-104	51.1	20.3 U	1.88 U	1.5 U	6.34 J	3.31 U	2.43 UJ	23.8 UJ
PCB-105	PCB-105	57000	112000	29700	39200	36900	34100	37600	100000
PCB-106	PCB-106	15.5 U	66.9 U	5.69 U	4.98 U	5.04 U	8.63 U	4.83 UJ	36.6 UJ
PCB-107/124	PCB-107/124	4610	8340	84.8	110	154	147	48.1 J	6880 J
PCB-109	PCB-109	9680	22500	3470	4510	4170	3770	3990 J	20100 J
PCB-110/115	PCB-110/115	137000	184000	3310	4940	5750	5640	3380 J	135000 J
PCB-111	PCB-111	63 J	129	24.9	30.9	34.9	31.2	27 J	109 J
PCB-112	PCB-112	20.1 U	41.8 U	2.39 U	2.91 U	2.86 U	5.18 U	3.82 UJ	22.1 UJ
PCB-114	PCB-114	4120	8340	2290	2970	2840	2630	2780	8460
PCB-118	PCB-118	112000	227000	62100	81200	81900	75000	75000	213000 J
PCB-120	PCB-120	342	748	145	220	199	191	178 J	671 J
PCB-121	PCB-121	33.3 J	42.5 U	2.38 U	14.1 J	14.2 J	13.7 J	3.79 UJ	37.5 J
PCB-122	PCB-122	1720	1040	6.49 U	38.9 J	19.1	19.7 J	5.5 UJ	764 J
PCB-123	PCB-123	2680	6350	1750	2070	1970	1930	2100	6300
PCB-126	PCB-126	392	710	334	279	260	227	254	47.3 U
PCB-127	PCB-127	101	254	5.33 U	53.6	51.9 J	49.9	4.52 UJ	323 J
PCB-128/166	PCB-128/166	8830	17900	2510	4240	4110	3890	2570	17600 J
PCB-129/138/163	PCB-129/138/163	58700	113000	18500	27600	26100	25100	19900	126000 J
PCB-130	PCB-130	4270	7070	336	397	497	496	256	7740 J
PCB-131	PCB-131	949	259	18.8	16.5	23.7	11.7 U	8.86	303 J
PCB-132	PCB-132	19200	13500	21.8	47.5	37.8	38 J	34.8	13300 J
PCB-133	PCB-133	1040	1790	265	340	382	370	243	1930 J
PCB-134	PCB-134	3660	1560	4.57 U	15.9	11.9 U	13.1 U	4.02 U	1460 J
PCB-135/151	PCB-135/151	18000	18700	62.5	132	204	198	51.1	18100 J
PCB-136	PCB-136	7060	2750	12.8	34.5	49.2	46.3	10.3 J	2440 J
PCB-137	PCB-137	3740	7770	1520	2330	2210	2140	1730	9010 J
PCB-139/140	PCB-139/140	1480	2300	555	820	816	787	665	2500 J
PCB-141	PCB-141	10600	15800	3.81 U	105	109	116	3.35 U	17000 J
PCB-142	PCB-142	33.9	39.9 U	4.31 U	6.08 U	9.95 U	24 J	3.79 U	41.6 UJ
PCB-143	PCB-143	24.5 U	37.1 U	3.99 U	5.8	9.11 U	10.3 U	3.51 U	38.8 UJ

				1135UE AII	alytical Results				
	Location ID	PB063.1_A	PB063.2_A	PB063.2_A	PB069_A	PB069_A	PB069_A	PB084.1	PB084.1
	Sample ID	PB03-F-A-GKF-W-010-20110617	PB03-F-A-GKF-W-005-20110615	PB03-F-A-GKF-W-013-20110617	PB03-F-A-GKF-W-002-20110615	PB03-F-A-GKF-W-007-20110617	PB03-I-A-BCR-W-002-COMP-201106	PB04-F-A-GKF-W-005-20110627	PB04-F-A-GKF-W-013-20110627
	Sample Date	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/17/2011	6/19/2011	6/27/2011	6/27/2011
	Sample Type	N	N N	N N	N	N N	N N	N	N N
	. ,. x	3201509.99	3201561.28	3201561.28	3201132.36	3201132.36	3201132.36	3200727.85	3200727.85
	γ	13831505.57	13831519.55	13831519.55	13830993.22	13830993.22	13830993.22	13829784.47	13829784.47
	Scientific_Name	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Fundulus grandis				
	Size_Class	=<15	=<15	=<15	=<15	=<15	>2 to =<7.5	=<15	=<15
	Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body					
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-080	PCB-080	58.1 U	101 U	20.5 U	62.4 U	30.3 UJ	9.49 U	19.8 U	15.3 U
PCB-081	PCB-081	771	740	653	1010	713	679	377 J	554
PCB-082	PCB-082	7620 J	7900	7240 J	13100 J	8050 J	42 J	6270	8780
PCB-083	PCB-083	4450 J	7260	4620 J	10100 J	6170 J	3.15 UJ	6860	6620
PCB-084	PCB-084	10100 J	13000	9500 J	17700 J	10200 J	251 J	13400	19800
PCB-085/116/117	PCB-085/116/117	73800 J	74300	56500 J	94700 J	65300 J	27400 J	35400	58800
PCB-086/087/097/108/119/125	PCB-CE2	129000 J	131000	102000 J	179000 J	118000 J	4190 J	72400	115000
PCB-088/091	PCB-088/091	26400 J	32100	22000 J	43500 J	25100 J	520 J	21300	33500
PCB-089	PCB-089	1740 J	2160	1650 J	3100 J	1860 J	8.35 J	1730	2520
PCB-090/101/113	PCB-090/101/113	233000 J	236000	174000 J	320000 J	202000 J	3320 J	121000	196000
PCB-092	PCB-092	40600 J	41900	32300 J	57500 J	36300 J	331 J	21500	35300
PCB-092 PCB-093/100	PCB-093/100	2640 J	2860	2110 J	4090 J	2240 J	682 J	1740	2830
PCB-094	PCB-094	827 J	1050	745 J	1490 J	838 J	43.6 J	764	1130
PCB-095	PCB-095	76400 J	97800	65000 J	130000 J	74200 J	714 J	71400	115000
PCB-096	PCB-096	477 J	698	456 J	1100 J	563 J	2.96 J	833	1160
PCB-098/102	PCB-098/102	6070 J	7500	5180 J	10300 J	6320 J	181 J	4840	7760
PCB-099	PCB-099	171000 J	151000	129000 J	223000 J	147000 J	62300 J	78100	132000
PCB-103	PCB-103	1660 J	2040	1420 J	2840 J	1580 J	13.4 J	1080	1720
PCB-103	PCB-104	35.9 J	45.4 J	33.9 J	48.8 J	43.2 J	1.22 UJ	29.5 J	29.4
PCB-104	PCB-105	111000	107000	91800	141000 J	109000	52400	48500	78800
PCB-105	PCB-106	39 UJ	97.3 U	62.1 J	55.4 UJ	19.2 UJ	8.89 UJ	13 U	16.5 U
PCB-107/124	PCB-107/124	8470 J	7330	6420 J	10500 J	7570 J	122 J	3210	5230
PCB-107/124 PCB-109	PCB-107/124 PCB-109	24000 J	20800	17800 J	27500 J	20600 J	6540 J	9100	14700
PCB-109	PCB-110/115	175000 J	181000	132000 J	236000 J	158000 J	5300 J	99100	158000
PCB-1111	PCB-110/113	130 J	114 J	105 J	250000 J 179 J	120 J	44 J	50.9	83.8
PCB-111 PCB-112	PCB-111	793 J	1420	617 J	1080 J	663 J	1.95 UJ	9.93 U	17.9 U
PCB-112	PCB-112	9080	8700	7410	12200	8620	3690	3680	5990
PCB-114 PCB-118	PCB-114 PCB-118	217000 J	237000	188000 J	266000 J	217000 J	105000	107000	177000
PCB-110 PCB-120	PCB-118	830 J	696	645 J	1000 J	722 J	248 J	276	448
PCB-121	PCB-121	38.5 J	37.9 U	31.2 J	53.3 J	33.1 J	1.93 UJ	22.5 J	33.4 J
PCB-122	PCB-121	1480 J	1440	1020 J	1590 J	1100 J	1.93 UJ	407	659
PCB-122 PCB-123	PCB-123	6310	6100	4960	8480	6700	2890	2570	4180
PCB-126	PCB-126	682 J	684	577	968 J	765	378	320	523
PCB-127	PCB-127	328 J	263 J	252 J	357 J	295 J	8.32 UJ	85.2	140
PCB-127 PCB-128/166	PCB-128/166	18100	16500	13600	21400	18800 J	4300	7620	11900
PCB-129/188/163	PCB-128/100 PCB-129/138/163	122000	109000	92500	148000	125000 J	30300	50200	79000
PCB-129/138/163	PCB-129/138/103	7570	6800	5980	9180	7660 J	547	3420	5460
PCB-131	PCB-131	233	254	241	443	286 J	32.5	218	290
PCB-131	PCB-131	12900	13600	11300	19500	14700 J	40.9	9980	15000
PCB-132 PCB-133	PCB-132 PCB-133	1950	1750	1560	2360	1870 J	40.9	891	1400
PCB-133	PCB-133	1610	1680	1400	2660	1810 J	3.17 U	1340	1940
PCB-1354 PCB-135/151	PCB-134 PCB-135/151	19000	18100	15800	28600	20500 J	117	12300	19700
PCB-136	PCB-135/151 PCB-136	2360	2800	2260	4650	20500 J 2780 J	27	2700	3970
PCB-136 PCB-137	PCB-136 PCB-137	8510	7280	6350	10500	8930 J	2310	3480	5660
PCB-137 PCB-139/140	PCB-137 PCB-139/140	2500	2350	1990	3250	2500 J	952	1260	2110
PCB-139/140 PCB-141		16900	14900	1990	21400	2500 J 17300 J	952 84.5	8100	12900
PCB-141 PCB-142	PCB-141 PCB-142	23 U	60.3 U	12900 13.5 J	45 U	1/300 J 19.2 UJ	84.5 2.98 U	15.5 U	20.9 U
				13.5 J 177					
PCB-143	PCB-143	176	210	1//	277	206 J	3.22 J	14.6 U	171

Appendix D
Tissue Analytical Results

Tissue Analytical Results										
	Location ID	PB095.2	PB101_D	PB101_D	PB101_D	PB102				
	Sample ID	PB04-I-A-BCR-W-001-COMP-201106	PB04-F-A-GKF-W-001-20110627	PB04-F-A-GKF-W-001-COMP-201106	PB04-F-A-GKF-W-009-20110627	PB04-F-A-PNF-W-001-COMP-201106				
	Sample Date	6/29/2011	6/27/2011	6/27/2011	6/27/2011	6/27/2011				
	Sample Type	N N	N	N	N	N N				
	х х	3200882	3201317.36	3201317.36	3201317.36	3201302.69				
	Υ	13829528	13828141.83	13828141.83	13828141.83	13828160.48				
	Scientific_Name	Callinectes sapidus	Fundulus grandis	Fundulus grandis	Fundulus grandis	Lagodon rhomboides				
	Size_Class	>2 to =<7.5	=<15	=<15	=<15	=<15				
	Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Migratory				
		Whole Body	Whole Body	Whole Body	Whole Body	Whole Body				
Chemical	Tissue_Type CAS Number	Result	Result	Result	Result	Result				
PCB-080	PCB-080					465 U				
		17.3 U	15.9 U	26.3 U	14.2 U					
PCB-081	PCB-081	205	416 J	699	488	479 U				
PCB-082	PCB-082	14.4 U	8280	8790	7910	8860				
PCB-083	PCB-083	15 U	22.5 U	7170	18.9 U	3130				
PCB-084	PCB-084	26.3 J	16800	16700	16600	14900				
PCB-085/116/117	PCB-085/116/117	9780	55100	59200	60300	42900				
PCB-086/087/097/108/119/125	PCB-CE2	1380	108000	124000	116000	86200				
PCB-088/091	PCB-088/091	45.2 J	30100	35000	33700	24000				
PCB-089	PCB-089	14.1 U	2240	2710	2370	538				
PCB-090/101/113	PCB-090/101/113	441	186000	216000	204000	150000				
PCB-092	PCB-092	23.6 J	33700	40800	36800	18300				
PCB-093/100	PCB-093/100	297	2490	2830	2890	1760				
PCB-094	PCB-094	13.1 J	998	1230	1100	102 J				
PCB-095	PCB-095	73.9 J	100000	105000	116000	54400				
PCB-096	PCB-096	8.38 U	1000	908	1050	694				
PCB-098/102	PCB-098/102	14 J	6640	9220	7210	885				
PCB-099	PCB-099	27300	131000	151000	142000	102000				
PCB-103	PCB-103	12 U	1570	2110	1870	1190				
PCB-104	PCB-104	6.92 U	26.9	40.8	42.6	46.7 J				
PCB-105	PCB-105	16800	79500	95600	84200	64900				
PCB-106	PCB-106	18.1 U	15.7 U	27.6 U	15.8 U	49.7 U				
PCB-107/124	PCB-107/124	19.2 U	4960	6390	5850	4760				
PCB-109	PCB-107/124 PCB-109	1510	14600	15500	15700	9770				
PCB-109	PCB-109	1650	148000	158000	161000	115000				
·										
PCB-111	PCB-111	9.7 U	85.3	104	89.4	47.2				
PCB-112	PCB-112	10 U	14.6 U	17.5 U	12.3 U	16.5 U				
PCB-114	PCB-114	1260	5940	6940	6410	4760				
PCB-118	PCB-118	37400	172000	189000	188000	138000				
PCB-120	PCB-120	87.3	460	548	500	336				
PCB-121	PCB-121	9.96 U	38.3	36.8	41	16.1 U				
PCB-122	PCB-122	20.3 U	522	887	591	939				
PCB-123	PCB-123	1010	4180	4610	4540	3490				
PCB-126	PCB-126	121	551	639	588	321 J				
PCB-127	PCB-127	27.5	147	25.3 U	154	45.4 U				
PCB-128/166	PCB-128/166	1250	12900	12100	13100	6870				
PCB-129/138/163	PCB-129/138/163	8940	86000	87600	86500	50700				
PCB-130	PCB-130	115	5830	5890	5890	2210				
PCB-131	PCB-131	18.8 U	24.3 U	379	284	426				
PCB-132	PCB-132	18.1 U	15800	15400	16300	8320				
PCB-133	PCB-133	183	1530	1690	1520	399				
PCB-134	PCB-134	19.6 U	1940	2030	2140	402				
PCB-135/151	PCB-135/151	13.2 U	20900	20000	22300	5610				
PCB-136	PCB-136	7.64 U	4060	3900	4480	2070				
PCB-137	PCB-137	918	6010	6180	6050	3210				
PCB-139/140	PCB-139/140	352	2200	2370	2270	1130				
PCB-141	PCB-139/140	16.8 U	13800	13500	14300	6740				
PCB-141	PCB-141 PCB-142	17.7 U	280	23 U	22.2 U	16.6 U				
					175					
PCB-143	PCB-143	16.9 U	193 J	235	1/5	15.4 U				

Appendix D
Tissue Analytical Results

Sample S	PB010.1 PB01-I-A-WHS-W-001-COMP-2011 6/20/2011 N 3201704.8 13836071.32 Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570 2050 J
Semple to \$4.3004_100 \$2.3004_2000_2000_2000_2000_2000_2000_2000_	618 PB01-I-A-WHS-W-001-COMP-2011 6/20/2011 N 3201704.8 13836071.32 Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
Separation Sep	N 3201704.8 13836071.32 Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
N	N 3201704.8 13836071.32 Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
Part	3201704.8 13836071.32 Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
	13836071.32 Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
Second Common Second Commo	Penaeus setiferus >2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
Size_Class	>2 to =<7.5 Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
	Migratory Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
Times Type Result Resu	Whole Body Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
Chemical CAS Number Result Resu	Result 409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
PCS-544	409 J 5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
PCP-145	5.42 UJ 3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
PCR-146	3770 J 6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
	6470 J 44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
PC8-148	44.2 J 5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
PCB-150	5.46 UJ 5.19 UJ 14900 J 435 J 96.3 2570
PCB-152	5.19 UJ 14900 J 435 J 96.3 2570
PCB-153/168	14900 J 435 J 96.3 2570
PCB.154 PCB.155 — 638 554 363 642 419 J 360 674 PCB.155 — 608 J 77 57.7 90.3 93.3 60.2 112 PCB.156/157 — 686 J 1390 1300 1610 2620 1840 2800 PCB.158 — 696 1100 1220 1380 2100 J 1520 2440 PCB.159 — 20.8 U 4.81 U 3.76 U 3.56 U 10.5 UJ 4.52 U 206 PCB.160 — 15.3 U 7.32 U 4.38 U 5.7 U 6.41 UJ 5.14 U 6.63 U PCB.161 — 20.8 U 4.81 U 3.76 U 3.56 U 10.5 UJ 4.52 U 206 206 PCB.162 — 15.3 U 7.32 U 4.38 U 5.7 U 6.41 UJ 5.14 U 6.63 U 6.07 U 6.07 U 6.07 U 6.07 U 6.07 U 6.07 U <	435 J 96.3 2570
PCB-155 PCB-155 - 60.8 J 77 57.7 90.3 93.3 60.2 112 PCB-156/157 PCB-156/157 - 686 J 1350 1300 1610 2620 1840 2800 PCB-158 PCB-158 PCB-158 - 486 J 1100 1220 1350 22100 J 1520 2440 PCB-159 PCB-159 - 20.8 U 4.81 U 3.76 U 3.56 U 10.5 UJ 452 U 206 PCB-160 PCB-161 - 15.3 U 7.32 U 4.88 U 5.7 U 6.41 UJ 51.4 U 6.63 U PCB-161 PCB-162 - 61.5 63.2 51.3 82.8 16.1 U 49.5 U 6.07 U 6.07 U 9.08 6.07 U 6.07 U 9.08 6.07 U 9.08 6.07 U 9.08 6.07 U 9.08 9.08 6.07 U 9.08 9.00 9.00 4.00 U 4.00 U 8.00 U <th< td=""><td>96.3 2570</td></th<>	96.3 2570
PCB-156/157 PCB-156/157 — 686 J 1350 1300 1610 2620 1840 2800 PCB-158 — 496 1100 1290 1350 2100 J 1520 2440 PCB-159 PCB-159 — 208 U 481 U 3.76 U 3.56 U 100 J 190 J 452 U 206 PCB-160 — 15.3 U 7.32 U 4.38 U 5.7 U 6.41 UJ 51.4 U 6.63 U PCB-161 — 2910 6.09 U 4.02 U 4.74 U 6.18 UJ 4.95 U 6.67 U PCB-162 PCB-162 — 61.5 G 6.2 S 51.3 S 82.8 S 163 J 52 J 106 6.67 U PCB-162 PCB-164 — 362 L 12.6 J 845 S 31.1 I 1030 J 726 I 1660 I PCB-164 PCB-164 — 362 L 12.6 J 845 S 31.1 J 1030 J 726 I 1660 I PCB-165<	2570
PCB-158 PCB-158 — 496 1100 1290 1350 2100 J 1520 2440 PCB-159 PCB-159 — 20.8 U 4.81 U 3.76 U 3.56 U 10.5 U 4.52 U 206 PCB-160 PCB-161 — 15.3 U 7.32 U 4.88 U 5.7 U 6.41 UU 51.4 U 6.63 U PCB-161 PCB-161 — 2910 6.09 U 4.02 U 4.74 U 6.18 UJ 4.95 U 6.07 U PCB-162 PCB-164 — 61.5 S 63.2 S 51.3 S 82.8 I 163 J 52 J 106 PCB-162 PCB-164 — 362 S 122 J 8.45 S 31.1 I 1000 J 726 I 1660 O PCB-165 PCB-166 — 17.3 U 10.7 I 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-166 — 437 G 601 513 G 693 G 908 G 676 1090 G PCB-167	
PCB-159 PCB-159 - 20.8 U 4.81 U 3.76 U 3.56 U 10.5 UJ 4.52 U 206 PCB-160 PCB-160 - 15.3 U 7.32 U 4.38 U 5.7 U 6.41 UJ 51.4 U 6.63 U PCB-161 PCB-161 - 2910 6.09 U 4.02 U 4.74 U 6.18 UJ 49.5 U 6.07 U PCB-162 PCB-161 - 2910 6.09 U 4.02 U 4.74 U 6.18 UJ 49.5 U 6.07 U PCB-162 PCB-163 - 61.5 68.2 51.3 82.8 163 J 52 J 106 PCB-164 PCB-165 - 362 12.6 J 845 31.1 1000 J 726 1660 PCB-165 PCB-165 - 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-168 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12	
PCB-160 PCB-160 - 15.3 U 7.32 U 4.38 U 5.7 U 6.41 UJ 51.4 U 6.63 U PCB-161 PCB-161 - 2910 6.09 U 4.02 U 4.74 U 6.18 UJ 49.5 U 6.07 U PCB-162 PCB-163 - 61.5 63.2 51.3 82.8 163 J 52 J 106 PCB-164 PCB-164 - 362 12.6 J 845 31.1 1030 J 726 1660 PCB-165 PCB-166 - 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-168 - 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-167 - 437 U 601 513 693 908 676 1090 PCB-169 PCB-167 - 437 U 601 513 693 908 676 1090 PCB-170 PCB-170 <td></td>	
PCB-161 PCB-161 - 2910 6.09 U 4.02 U 4.74 U 6.18 UJ 49.5 U 6.07 U PCB-162 PCB-162 - 61.5 63.2 51.3 82.8 163 J 52 J 106 PCB-164 PCB-164 - 362 12.6 J 845 31.1 1030 J 726 1660 PCB-165 PCB-165 - 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-167 - 437 601 513 693 908 676 1090 PCB-169 PCB-169 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 U PCB-170 PCB-169 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 U PCB-170 PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-171/173	11.9 UJ
PCB-162 PCB-162 61.5 63.2 51.3 82.8 163 J 52 J 106 PCB-164 PCB-164 362 12.6 J 845 31.1 1030 J 726 1660 PCB-165 PCB-166 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-167 437 601 513 693 908 676 1090 PCB-169 PCB-169 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 PCB-170 PCB-169 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 PCB-170 PCB-170 1340 1020 2100 1040 2450 1930 4930 PCB-171/173 550 525 689 536 743 615 1610 PCB-172 PCB-172<	15.7 UJ
PCB-164 PCB-164 - 362 12.6 J 845 31.1 1030 J 726 1660 PCB-165 PCB-165 - 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-167 - 437 601 513 693 908 676 1090 PCB-169 PCB-169 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-171/173 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-171/173 PCB-173 - 550 525 689 536 743 615 1610 PCB-172 PCB-173 - 335<	15.1 UJ
PCB-165 PCB-165 - 17.3 U 10.7 12.2 J 11 J 6.56 UJ 52.6 U 18 PCB-167 PCB-167 - 437 601 513 693 908 676 1090 PCB-169 PCB-169 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-173 PCB-171/173 - 550 525 689 536 743 615 1610 PCB-172 PCB-173 - 550 525 689 536 743 615 1610 PCB-174 PCB-175 - 335 250 418 298 562 510 1020 PCB-175 PCB-176 - 127 78 135 102 131 110 318 PCB-176 PCB-177 -	172 J
PCB-167 PCB-167 - 437 601 513 693 908 676 1090 PCB-169 PCB-169 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-171/173 PCB-171/173 - 550 525 689 536 743 615 1610 PCB-172 PCB-172 - 335 250 418 298 582 510 1020 PCB-174 PCB-174 - 1050 14.9 1700 22.7 J 1390 984 4030 PCB-175 PCB-176 - 127 78 135 102 131 110 318 PCB-176 - 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-178 - 1490 333	1020 J
PCB-169 PCB-169 - 16.4 U 8.26 J 16.1 J 4.37 U 17.7 J 15.1 9.12 PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-171/173 PCB-171/173 - 550 525 689 536 743 615 1610 PCB-172 - 335 250 418 298 582 510 1020 PCB-174 - 1050 14.9 1700 22.7 J 1390 984 4030 PCB-175 - 127 78 135 102 131 110 318 PCB-176 - 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-177 - 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 - 1050 476 615 519 673 <t< td=""><td>16.1 UJ</td></t<>	16.1 UJ
PCB-170 PCB-170 - 1340 1020 2100 1040 2450 1930 4930 PCB-171/173 PCB-171/173 - 550 525 689 536 743 615 1610 PCB-172 PCB-172 - 335 250 418 298 582 510 1020 PCB-174 PCB-174 - 1050 14.9 1700 22.7 J 1390 984 4030 PCB-175 PCB-175 - 127 78 135 102 131 110 318 PCB-176 PCB-176 - 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-178 - 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 - 1050 476 615 519 673 581 1430 PCB-180/193 PCB-180/193 - 3980 4340 </td <td>892 J</td>	892 J
PCB-171/173 PCB-171/173 - 550 525 689 536 743 615 1610 PCB-172 PCB-172 - 335 250 418 298 582 510 1020 PCB-174 PCB-174 - 1050 14.9 1700 22.7 J 1390 984 4030 PCB-175 PCB-175 - 127 78 135 102 131 110 318 PCB-176 PCB-176 - 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-177 - 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 - 1050 476 615 519 673 581 1430 PCB-179 - 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 - 3980 4340 5220	23.1 J
PCB-172 335 250 418 298 582 510 1020 PCB-174 1050 14.9 1700 22.7 J 1390 984 4030 PCB-175 PCB-175 127 78 135 102 131 110 318 PCB-176 PCB-176 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-177 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 1050 476 615 519 673 581 1430 PCB-179 PCB-179 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	2740
PCB-174 PCB-174 1050 14.9 1700 22.7 J 1390 984 4030 PCB-175 PCB-175 127 78 135 102 131 110 318 PCB-176 PCB-176 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-177 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 1050 476 615 519 673 581 1430 PCB-179 PCB-179 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	854
PCB-175 PCB-175 127 78 135 102 131 110 318 PCB-176 PCB-176 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-178 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 1050 476 615 519 673 581 1430 PCB-179 PCB-179 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	598
PCB-176 203 5.9 J 284 13.5 J 62.1 35.8 623 PCB-177 PCB-178 1490 333 1270 444 1450 1130 2820 PCB-178 PCB-178 1050 476 615 519 673 581 1430 PCB-179 PCB-179 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	1500
PCB-177	160
PCB-178 PCB-178 1050 476 615 519 673 581 1430 PCB-179 PCB-179 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	70.9
PCB-179 PCB-179 927 4.4 U 840 6.5 214 178 1780 PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	1610
PCB-180/193 PCB-180/193 3980 4340 5220 4180 6700 5670 12500	779
	289
	7190
PCB-181 PCB-181 11.4 U 23.7 34.8 31.9 42.7 40.1 76.8 PCB-182 PCB-183 PCB-183 <td>52.6</td>	52.6
PCB-182 PCB-182 44 24.7 18.2 29.2 28.8 23.9 61.8	42.8 J
PCB-183/185 PCB-183/185 1650 1460 1860 1420 1880 1620 4170 PCB-184 PCB-184 28.2 40.2 39.4 59.6 32.6 32.1 78.8	1900 35.8
	35.8 4.47 U
PCB-187	6130
PCB-188 PCB-188 119 J 49.2 28.5 40.3 30.8 22.2 51.8	32.9
PCB-189 PCB-189 62.1 J 78.6 78.3 87.1 130 93 215 PCB-100 PCB-100 243 251 420 250 635 513 1020	138
PCB-190 PCB-190 243 251 430 259 635 512 1030	670
PCB-191 PCB-191 50.6 J 71.3 107 74.7 136 116 253 PCB-192 PCB-192 9.27 U 3.86 U 2.82 J 3.74 U 6 4.43 U 8.55 J	156
	7.27 U
PCB-194 PCB-194 1090 539 871 566 1020 707 2740	919
PCB-195 PCB-195 - 291 154 370 176 352 227 999	
PCB-196 PCB-196 - 622 446 567 442 530 362 1550	718 U
PCB-197/200 PCB-197/200 - 200 141 285 177 173 130 727	538
PCB-198/199 PCB-198/199 - 1790 752 1280 906 1230 994 3550	538 177
PCB-201	538 177 1110
PCB-202 PCB-202 1160 687 753 608 614 526 1430	538 177

				1135UE Alle	alytical Results				
	Location ID	PB010.3	PB013.1_A	PB013.3	PB013.3	PB015.1	PB015.2	PB015_A	PB015_A
			_	PB01-F-A-GKF-W-004-COMP-201106		PB01-F-A-GKF-W-002-COMP-201106		_	_
	Sample Date		6/22/2011	6/21/2011	6/15/2011	6/15/2011	6/16/2011	6/25/2011	6/22/2011
	Sample Type		N N	N	N	N	N	N	N
	х х	3201743	3201306.47	3201433.88	3201433.88	3201175	3201180	3201168.73	3201168.73
	v	13836152	13836191.23	13836075.12	13836075.12	13836251	13836183	13836254.66	13836254.66
	Scientific_Name		Callinectes sapidus	Fundulus grandis	Mugil cephalus	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Callinectes sapidus
	Size_Class	>2 to =<7.5	>2 to =<7.5	=<15	=<15	=<15	>2 to =<7.5	=<15	>2 to =<7.5
	Tissue_Category	Sedentary	Sedentary	Sedentary	Migratory	Sedentary	Sedentary	Sedentary	Sedentary
	Tissue_Category	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-144	PCB-144	76.1	16.3	2980	1280	2420	31.2	4480 J	24.7
PCB-144 PCB-145	PCB-144 PCB-145	76.1 3.27 U	0.52 U	17.2	12.5	13.6 J	4.68 U	20 J	4.07 U
PCB-146	PCB-146	6710	2720	15100	25.4 J	12400	6460	23200 J	4620
PCB-147/149	PCB-147/149	1870	417	37600	18000	33100	1080	54800 J	609
PCB-148	PCB-148	115	50.9	180	91.5	150	127	291 J	77
PCB-150	PCB-150	3.17 U	0.48 U	112	100	83.5	4.53 U	172 J	3.93 U
PCB-152	PCB-152	3.08 U	0.469 U	41.3	32.1	36.4	4.4 U	55.6 J	3.92 U
PCB-153/168	PCB-153/168	40500	16300	76500	36900	67400	37300	119000 J	27500
PCB-154	PCB-154	1120	442	1680	770	1410	1150	2700 J	691
PCB-155	PCB-155	96.1	58.5	168	141	129	128	224	151
PCB-156/157	PCB-156/157	4460	2060	9930	3830	8240	3990	13700	3040
PCB-158	PCB-158	4190	1760	7980	3800	7040	3690	12500 J	2770
PCB-159	PCB-159	7.12 U	7.27	7.11 U	15.7 U	5.71 U	13.1	7.66 UJ	14 U
PCB-160	PCB-160	9.51 U	24.4 U	6.76 U	14.2 U	7.66 U	8.64 U	9.8 UJ	16.2 U
PCB-161	PCB-161	7.91 U	22.9 U	6.09 U	5730	6.9 U	7.19 U	8.82 UJ	15.7 U
PCB-162	PCB-162	155	87	623	163 J	545	171	1050 J	155
PCB-164	PCB-164	177	23.8 U	3500	1910	3380	43.8	5520 J	16.4 U
PCB-165	PCB-165	18.2	26.1 U	49.8	40.5	37.5	18.4	76.6 J	17.6 U
PCB-167	PCB-167	1850	788	3910	1480	3290	1570	5590	1290
PCB-169	PCB-169	8.25 U	1.74 U	75.5	10.1 U	71.7	10.1 U	222 J	29.5
PCB-170	PCB-170	3760	1450	12800	7580	9910	2830	18500	3120
PCB-171/173	PCB-171/173	1550	674	4330	2160	3330	1290	6030	1240
PCB-172	PCB-172	809	356	2450	1530	2020	691	3560	748
PCB-174	PCB-174	108	25.1	7590	3470	6070	38.4	9830	32.5
PCB-175	PCB-175	251	103	759	389	590	222	1110	179
PCB-176	PCB-176	44.8 J	5.35 J	1000	811	831	30.3 J	1340	14.7 J
PCB-177	PCB-177	1280	418	6780	3690	5110	947	9630	762
PCB-178	PCB-178	883	467	3540	1720	2720	822	5090	854
PCB-179	PCB-179	17.3	5.7	3270	2540	2670	6.7	4640	10.6 J
PCB-180/193	PCB-180/193	13100	5530	33100	19100	25500	11000	45500	10800
PCB-181	PCB-181	86.6	39.7	221	106	169	83.8	307	64.2
PCB-182	PCB-182	70.2	36.3	155	88.1	110 J	59.1 J	202 J	61
PCB-183/185	PCB-183/185	4300	1820	10800	6010	8030	3680	15100	2870
PCB-184	PCB-184	63.6	31.8	152	87.6	109	81.4	198	65.2
PCB-186	PCB-186	3.28 U	0.463 U	4.49 J	3.25 U	3.1	3.9 U	4.2 J	6.43 U
PCB-187	PCB-187	7370	2980	18400	11500	14400	6300	25600	6200
PCB-188	PCB-188	64.2	34.7	105	69.1	78.4	63.9	146	60.7
PCB-189	PCB-189	226	84.7	460	251 J	384	210	644	197
PCB-190	PCB-190	836	337	2000	1540	1680	684	2800	796
PCB-191	PCB-191	229	90.9	616	366	487	192	887	212
PCB-192	PCB-192	5.23 U	1.75 J	6.53	10.2 U	6.39 J	6.51 U	7.54 J	9.58 U
PCB-194	PCB-194	1660	550	4440	3310	3660	1440	5680	1560
PCB-195	PCB-195	539	169	2280	1220	1670	423	3160	437
PCB-195	PCB-193	1310	509	3640	1800	2820	1170	5630	1070
PCB-190 PCB-197/200	PCB-190 PCB-197/200	283	126	1380	695	960	294	1850	241
PCB-197/200 PCB-198/199	PCB-197/200 PCB-198/199	283	947	7980	3720	5850	294	11900	1860
PCB-198/199 PCB-201	PCB-198/199 PCB-201	391	171	1610	604	1090	389	2300	306
LCD-701	LCD-201	221	1/1	1010	004	1090	202	2300	300

				Tissue An	alytical Results				
	Location ID	PB015_A	PB015_A	PB015_A	PB016.2	PB019.1	PB019.1	PB019.1	PB019.1
		=	6 PB01-I-A-BCR-W-009-COMP-201106	_			PB01-I-B-BRS-W-012-20110621	PB01-I-B-BRS-W-014-20110621	PB01-I-B-BRS-W-015-20110621
	Sample Date	6/22/2011	6/22/2011	6/20/2011	6/15/2011	6/22/2011	6/21/2011	6/21/2011	6/21/2011
	Sample Type	0, ==, ==== N	N	N	N	N	N	N	0,, N
	y x	3201168.73	3201168.73	3201168.73	3201190	3201152.87	3201152.87	3201152.87	3201152.87
	Ŷ	13836254.66	13836254.66	13836254.66	13836042	13835753.58	13835753.58	13835753.58	13835753.58
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Penaeus aztecus	Fundulus grandis	Cynoscion arenarius	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus
	Size_Class	>2 to =<7.5	>2 to =<7.5	>7.5 to =<13	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13
	Tissue_Category	Sedentary	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-144	PCB-144	20.9	27.8	728	1520	1040 J	1450 J	876	1010
PCB-145	PCB-145	0.874 U	1.13 U	2.47 U	9.24	5.28 J	114 U	20.7 U	40 U
PCB-146	PCB-146	3780	4940	5460	7850	25.6 J	9570	6990	6900
PCB-147/149	PCB-147/149	636	819	8440	18400	7780 J	16900	11800	11300
PCB-148	PCB-148	51.7	64.3	65.7	97.9	73 J	208	100	124 J
PCB-150	PCB-150	0.845 U	1.09 U	15.7	56.5	60.2 J	110 U	20 U	38.6 U
PCB-150	PCB-152	0.843 U	1.09 U	4.42	19.6	18.6 J	100 U	19.7 U	37.9 U
PCB-152 PCB-153/168	PCB-153/168	20500	27800	24200	43400	38500 J	47100	34300	34200
PCB-153/108	PCB-153/108	476	603	688	939	877 J	1290	881	869
PCB-154 PCB-155	PCB-155	81.6	104	137	91	110	192 J	189 J	179
PCB-133 PCB-156/157	PCB-133	2750	3710	3630	4760	3950	5280	4880	4360
PCB-150/157 PCB-158	PCB-138/137 PCB-158	2320	3130	2930	4380	3760 J	5900	4880	4560
PCB-159	PCB-159	3.88 U	5.56 U	17.2 U	6.1 U	8.52 UJ	158 U	31.4 U	160 J
PCB-139	PCB-159	4.11 U	3.69 U	6.08 U	5.5 U	8.41 UJ	193 U	28.4 U	49.7 U
PCB-160 PCB-161	PCB-161	4.11 U	3.59 U	5.77 U	4.95 U	3540 J	193 U	27.5 U	49.7 U
PCB-101 PCB-162	PCB-162	137	186	16.5 U	224	165 J	303	27.3 U	255 J
PCB-162	PCB-164	4.19 U	50.3	1460	1900	1360 J	2550	2070	1880
PCB-165	PCB-104 PCB-165	13.7	13.3	18.8	27.9	17.6 J	206 U	30.4 U	53 U
PCB-163	PCB-167	1090	15.5	1400	1960	17.0 J 1390 J	1990	1800	1600
PCB-167 PCB-169	PCB-167 PCB-169	18.5	5.13 U	1400 14.3 U	38.7	7.04 UJ	288 U	25 U	58.6 U
PCB-109	PCB-109	2180 J	3500	4050	6020	6920 J	5600	6150	5140
PCB-170 PCB-171/173	PCB-170	886 J	1330	1130	2090	2200 J	1610	1560	1480
PCB-171/173	PCB-171/173	589 J	881	921	1200	1320 J	1290	1230	1080
PCB-172	PCB-172	20.8 J	41.1 J	1670	3100	2110 J	2340	2210	1940
PCB-174 PCB-175	PCB-174	141 J	192	220	378	341 J	372 J	301	287
PCB-173	PCB-175	22.9 J	17.6	53.2 J	463	416 J	222 J	177	196
PCB-170	PCB-170	672 J	1090	2680	3200	1470 J	3370	3500	3350
PCB-177	PCB-177	666 J	891	1100	1770	1460 J	1660	1470	1390
PCB-178	PCB-178	4.85 J	5.69	261	1510	1030 J	528	407	426
PCB-180/193	PCB-180/193	7760 J	12100	10400	16200	19000 J	14000	14400	12400
PCB-181	PCB-181	47.5 J	72.4	70	103	124 J	87.4 U	97.5	45.7 U
PCB-182	PCB-182	44.5 J	57.4	46.6	77.6	82.9 J	82.8 U	51.5	32.5 U
PCB-183/185	PCB-183/185	2150 J	3120	3010	5550	6270 J	4600	4440	4230
PCB-183/183	PCB-183/183	32.9 J	42	45.2	71.7	95.5 J	91 J	70	72.8
PCB-184 PCB-186	PCB-184 PCB-186	0.671 UJ	1.08 U	2.06 U	1.7 U	1.27 UJ	67.9 U	13.9 U	26.7 U
PCB-180 PCB-187	PCB-186	5710 J	7710	9650	9090	5520 J	11400	13300	10100
PCB-187	PCB-187	45.4	52.6	48.7	46.8	58.5	102 U	68.5 J	83.4 J
PCB-1889	PCB-189	45.4 125 J	195	203	212	248 J	296	256	259
PCB-189 PCB-190	PCB-189	571 J	852	981	1020	1430 J	1300	1400	1260
PCB-190 PCB-191	PCB-190 PCB-191	149 J	217	220	317	359 J	329	290	253
PCB-191	PCB-191 PCB-192	3.19 UJ	3.05 U	7.85 J	2.56 U	10.7 J	73.9 U	18 U	38.7 U
PCB-192 PCB-194	PCB-192 PCB-194	895	1550	1640	2.36 0	4330	2410	2270	2090
PCB-194 PCB-195	PCB-194 PCB-195	300	536	545	943	1420	689	764	694
PCB-195 PCB-196	PCB-195 PCB-196	832	1250	830	1550	2300	1040	1100	1010
	PCB-196 PCB-197/200	171	227	232	503	736	369	344	318
PCB-197/200		1/1	2480	1660	3190	3780	2260	2450	2420
DCD 109/100						3/00		/4311	Z4ZU
PCB-198/199 PCB-201	PCB-198/199 PCB-201	260	337	225	604	744	324	330	303

				Tissue An	alytical Results				
	Location ID	PB019.1	PB019.1	PB019.1	PB019.2	PB019_A	PB019_A	PB024.1	PB024.1
	Sample ID		PB01-I-B-BRS-W-022-20110622	PB01-I-B-BRS-W-023-20110622		PB01-F-A-GKF-W-005-COMP-201106		PB01-F-A-GM-W-001-20110615	PB01-F-A-GM-W-002-20110615
	Sample Date	6/21/2011	6/22/2011	6/22/2011	6/28/2011	6/21/2011	6/23/2011	6/15/2011	6/15/2011
	Sample Type	N	N	N	N	N N	N	N	N
	Х	3201152.87	3201152.87	3201152.87	3201194.8	3201314.96	3201314.96	3201090.96	3201090.96
	Υ	13835753.58	13835753.58	13835753.58	13835735.1	13835750.59	13835750.59	13835160.37	13835160.37
	Scientific_Name	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus	Crassostrea virginica	Fundulus grandis	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus
	Size_Class	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-144	PCB-144	1590 J	1350	1070	502	2720	49.7	1390	1130
PCB-145	PCB-145	68.4 UJ	1.24 U	5.15 U	9.4	12.4 U	1.85 U	7.21	7.6 J
PCB-146	PCB-146	11200 J	36.8	7600	4160	15600	7460	31.9	27 U
PCB-147/149	PCB-147/149	20500 J	16000	12800	16300	34300	1360	24500	20700
PCB-148	PCB-148	183 J	140	103	61.9	235	155	96.8	85.7
PCB-150	PCB-150	66.1 UJ	27.3	22.1	65.7	98.6	1.79 U	101	84.4
PCB-152	PCB-152	64.8 UJ	13.3	11 J	24.3	33.5	1.78 U	26.6	25.1
PCB-153/168	PCB-153/168	59900 J	44300	37700	22300	81300	34100	38500	31700
PCB-154	PCB-154	1650 J	1300	932	599	1670	1190	852	719
PCB-155	PCB-155	252	155	187	126	190	397	109	104
PCB-156/157	PCB-156/157	6230	5180	4680	1650	9230	3260	3970	3360
PCB-158	PCB-158	6700 J	5480	4620	253	8020	3460	3960	3190
PCB-159	PCB-159	200 J	10.1 U	24.4 U	5.79 U	32.5 U	19.9 U	16.1 U	23.5 U
PCB-160	PCB-160	85.9 UJ	9.69 U	23.7 U	6.13 U	7.35 U	6.62 U	12.8 U	22.9 U
PCB-161	PCB-161	83 UJ	8240	22.6 U	5.81 U	6.62 U	6.44 U	6150	5020
PCB-162	PCB-162	326 J	237 J	192 J	205	680	127	172	148 J
PCB-164	PCB-164	3190 J	2590	2180	1150	3870	118	2500	2000
PCB-165	PCB-165	91.8 UJ	12	35.8	16.5 J	64.5	32.8	49	40.2
PCB-167	PCB-167	2160	1900	1810	762	3800	1370	1540	1290
PCB-169	PCB-169	247 UJ	6.79 U	13.9 U	6.06 U	91	34.2 U	14 U	16 U
PCB-170	PCB-170	7270	6980	7290	565	13800	2940 J	7330	6120
PCB-171/173	PCB-171/173	2010	2110	1970	540	4300	1230 J	2400	1980
PCB-172	PCB-172	1630	1500	1430	101	2690	902 J	1590	1390
PCB-174	PCB-174	3030	2990	2650	208	7810	68 J	6240	5090
PCB-175	PCB-175	429	413	341	78.8	784	235 J	423	346
PCB-176	PCB-176	261	252	207	316	981	16.7 J	883	750
PCB-177	PCB-177	4470	4680	4360	2090	6690	908 J	4350	3560
PCB-178	PCB-178	1970	2080	1780	804	3630	1060 J	1880	1540
PCB-179	PCB-179	580	618	461	1290	3120	11.3 J	2600	2140
PCB-180/193	PCB-180/193	17700	16900	16800	628	34000	11100 J	19100	16200
PCB-181	PCB-181	148	143	110	7.68	221	97.8 J	119	103
PCB-182	PCB-182	114 J	97.3	5.26 U	2.62 U	192	114 J	86.7	51.4
PCB-183/185	PCB-183/185	5430	5660	4960	1160	10700	3060 J	6210	5030
PCB-184	PCB-184	33.8 U	82.1	76.7	23.8	182	154 J	85	76.9
PCB-186	PCB-186	36.6 U	1.51 U	4.42 U	1.9 U	6.49 J	2.36 UJ	3.53 J	6.2 U
PCB-187	PCB-187	13900	12900	12900	6630	20900	7880 J	11600	9950
PCB-188	PCB-188	89.9	64.4	66.3	41.6	129	151	58.8	54.2
PCB-189	PCB-189	350	274	265	53	464	189 J	284	258
PCB-190	PCB-190	1650	1750	1810	87	2210	745 J	1500	1290
PCB-191	PCB-191	387	364	351	8.88 J	677	225 J	352	311
PCB-192	PCB-192	40.2 U	5.97 U	15.8 U	2.97 U	15.5	14.4 J	10.5	12.4 U
PCB-194	PCB-194	2850	2230	2460	70.6	4790	1210	4140	3780
PCB-195	PCB-195	880	783	803	26.9	2300	502	1410	1260
PCB-196	PCB-196	1240	1280	1240	15.5	4040	1590	2070	1900
PCB-197/200	PCB-197/200	418	390	367	41.8	1380	665	872	781
PCB-198/199	PCB-198/199	2750	2940	2780	167	8140	3160	4780	4200
PCB-201	PCB-201	398	402	355	110	1660	740	715	624
PCB-202	PCB-202	1370	1050	1130	917	3860	2090	1600	1390

Appendix D
Tissue Analytical Results

				1133UE AII	alytical Results				
	Location ID	PB024.4	PB025_A	PB026.1	PB032_A	PB037.2	PB037.4	PB037.5	PB037.6
		PB01-I-A-BCR-W-005-COMP-201106	_	PB01-F-A-STM-W-002-20110616	PB02-F-A-PNF-W-009-20110617	PB02-F-A-STM-W-006-20110616		PB02-I-B-BRS-W-001-COMP-201106	
	Sample Date		6/15/2011	6/15/2011	6/17/2011	6/16/2011	6/17/2011	6/17/2011	6/16/2011
	Sample Type		N N	N	N	N	N N	N	N
	х х	3201319	3201214.42	3201153.65	3201407.03	3201346.97	3201504	3201380	3201402
	· · · · · · · · · · · · · · · · · · ·	13835188	13835154.42	13835079.21	13834542.25	13834038.02	13833530	13834151	13833775
	Scientific_Name	Callinectes sapidus	Brevoortia patronus	Mugil cephalus	Lagodon rhomboides	Mugil cephalus	Callinectes sapidus	Penaeus aztecus	Callinectes sapidus
	Size_Class	>2 to =<7.5	=<15	=<15	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Migratory	Migratory	Migratory	Migratory	Migratory	Sedentary
	Tissue_Type	· ·	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-144	PCB-144	134	2020	1590	1390	1960	27.8 J	581	49.1
PCB-145	PCB-145	10.3 U	10.8 J	14.9 J	13.2 U	20.3 J	3.66 UJ	1.04 J	2.72 U
PCB-146	PCB-146	8870	41.4	29	7420	35.8	7640 J	4170	5000
PCB-147/149	PCB-147/149	4010	34800	18700	21100	27700	883 J	6090	1180
PCB-148	PCB-148	301	132	93.8	127	121	90 J	50.6	83.9
PCB-150	PCB-150	9.96 U	135	106	93.9	118	3.54 UJ	8.27	2.63 U
PCB-152	PCB-150	9.50 U	42.1	44.9	12.7 U	64.1	3.53 UJ	4.34 J	2.56 U
PCB-132 PCB-153/168	PCB-152 PCB-153/168	36400	50000	42300	44000	48900	46500 J	18700	30800
PCB-154	PCB-153/168 PCB-154	1680	1140	905	871	953	40300 J 1100 J	457	904
PCB-155	PCB-154 PCB-155	279	120	93.3	199	116	88.9	61.5	54.4
PCB-155 PCB-156/157	PCB-155 PCB-156/157	3330	5700	5090	4660	5860	6940	3210	3810
PCB-158	PCB-150/157	3270	5500	4750	4170	5360	5560 J	2680	3510
PCB-159	PCB-158 PCB-159	23 U	19.2 U	18.2 U	181 J	28.6 U	14.2 UJ	4.22 U	9.29 J
PCB-160	PCB-160	28.8 U	15.4 U	18.2 U	17.1 U	28.0 U	7.15 UJ	4.43 U	9.27 U
PCB-161	PCB-161	28.8 U	7980	6280	17.1 U	7460	6.96 UJ	4.43 U	7.71 U
PCB-162	PCB-161	126 J	227	204	165	216	163 J	135	157
PCB-164	PCB-162	261	3410	2050	2090	2800	56.8 J	1130	84.7
PCB-165	PCB-165	46.3	63.3	45 J	36 J	63.1	20.1 J	11.5	12.8 J
PCB-167	PCB-167	1360	2170	1940	1800	2190	2510	11.5	1470
PCB-169	PCB-169	25.1 U	14.3 U	12.2 U	41.2	2190 21 U	12.2 UJ	3.76 U	27.4
PCB-170	PCB-109	3240	9670	8700	7070	8920	5920	2950	27.4
PCB-171/173	PCB-171/173	1630	3210	2440	2270	2670	2080	862	1140
PCB-1712	PCB-171/173	1000	2050	1650	1210	1750	1430	605	605
PCB-172	PCB-172	454	8350	3240	3310	5450	40.7	1330	50.5
PCB-174	PCB-174	370	544	428	428	474	311	159	175
PCB-175	PCB-176	112	1210	823	640	1050	10.9 J	65	20.4 J
PCB-177	PCB-170	1960	5750	3710	2670	4540	1130	1770	715
PCB-177	PCB-177	1810	2510	1820	1190	2040	1450	742	715
PCB-179	PCB-178	118	3620	2630	1540	3250	8.78	180	9.66
PCB-180/193	PCB-180/193	12800	24400	21300	17900	21800	21000	7890	9800
PCB-180/193	PCB-180/193	1200	165	145	129	147	118	55.5	75
PCB-182	PCB-181	165	111	86.7	90.2	84.2	88.3	32.3	32.7
PCB-182 PCB-183/185	PCB-182 PCB-183/185	4390	8040	6620	6090	7010	5180	2320	3040
PCB-184	PCB-183/183	183	92.3	58.3	90.2	73.8	36.9	21.1	32.7
PCB-186	PCB-184	6.76 U	2.9 U	3.66 U	9.6 U	7.16 U	4.25 U	1.19 U	1.83 U
PCB-187	PCB-187	9740	14900	12100	12000	12600	12600	5640	5250
PCB-188	PCB-187	167	71	58.7	97.9	64.4	49.5	27.5	35.4 J
PCB-189	PCB-189	238	377	283	248	307	321	143	175
PCB-190	PCB-190	840	1920	1750	1580	1790	1420	754	669
PCB-191	PCB-190	249	461	399	391	419	400	150	175
PCB-192	PCB-192	17.7 U	13.7 J	8.58 U	16.4	13.5 U	4.85 U	4.15 J	3.75 U
PCB-194	PCB-192	17.7 0	4950	3560	3080	3700	2740	1060	1260
PCB-194	PCB-194	603	1850	1300	1250	1350	714	351	373
PCB-195	PCB-195	1560	2670	1920	1910	1970	1990	564	952
PCB-190 PCB-197/200	PCB-197/200	637	1070	603	725	731	250	150	156
PCB-197/200 PCB-198/199	PCB-198/199	3320	6190	3920	3200	4210	3150	1190	1610
PCB-198/199 PCB-201	PCB-198/199 PCB-201	803	904	565	707	642	405	166	245
. 50 201	, CD 201	003]] 303	1 ,0,	072	771	100	273

Tissue Analytical Results										
	Location ID	PB037_A	PB047.1_A	PB047.3	PB048.3	PB048.3	PB048.3	PB049_A	PB049_A	
	Sample ID	PB02-I-A-BCR-W-004-COMP-201106	PB02-F-A-PNF-W-002-20110615	PB02-I-A-BCR-W-001-COMP-201106	PB02-F-A-GM-W-001-20110616	PB02-F-A-GM-W-002-20110616	PB02-F-A-STM-W-001-20110616	PB02-F-A-GKF-W-001-COMP-201106	PB02-I-A-BCR-W-003-COMP-201106	
	Sample Date	6/19/2011	6/15/2011	6/19/2011	6/16/2011	6/16/2011	6/16/2011	6/15/2011	6/22/2011	
	Sample Type	N	N	N	N	N	N	N	N	
	х	3201325.97	3201461.79	3201645	3201519.15	3201519.15	3201519.15	3201468.39	3201468.39	
	Υ	13833885.45	13833007.68	13832925	13832869.89	13832869.89	13832869.89	13832722.8	13832722.8	
	Scientific_Name	Callinectes sapidus	Lagodon rhomboides	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus	Mugil cephalus	Fundulus grandis	Callinectes sapidus	
	Size_Class	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15	=<15	=<15	>2 to =<7.5	
	Tissue_Category	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Sedentary	Sedentary	
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result	
PCB-144	PCB-144	27.5	2780	39.1 J	2140	2050	1560	4050	16.3	
PCB-145	PCB-145	0.66 U	9.24 U	11.8 U	11.8	12.6 J	18.8 J	23.1 J	1.61 U	
PCB-146	PCB-146	6060	40.6	4950	41.9	9550	6150	18300	3820	
PCB-147/149	PCB-147/149	739	33700	741	35100	34500	21000	44500	386	
PCB-148	PCB-148	64.5	136	90.6	127	122	72.9	192	64.8	
PCB-150	PCB-150	0.61 U	113	11.4 U	134	132	83.8	123	1.49 U	
PCB-152	PCB-152	0.596 U	18 J	11.1 U	41.9	44.9 J	51.1	56.2	1.45 U	
PCB-153/168	PCB-153/168	35100	82700	31200	56500	52200	35600	96400	23300	
PCB-154	PCB-154	698	1130	852	1140	1090	678	1760	577	
PCB-155	PCB-155	45.6	116	68.4 J	130	121	43.3 J	149	61.7	
PCB-156/157	PCB-156/157	6300	11100	4510	6900	6080	4740	13900	2840	
PCB-158	PCB-158	5060	10000	3920	6380	5780	4200	11400	2300	
PCB-159	PCB-159	12.1	32.9 U	16.8 U	8.9 U	15.7 U	23.9 U	25.2 U	8.5 U	
PCB-160	PCB-160	2.65 U	35.6 U	18.6 U	10.9 U	14.8 U	21 U	6.77 U	4.29 U	
PCB-161	PCB-161	2.48 U	12100	15.5 U	8680	14.4 U	20.4 U	6.09 U	4.02 U	
PCB-162	PCB-162	207	363	170	314	280	200	787	119	
PCB-164	PCB-164	2.58 U	4070	43.3	3770	3540	2230	4940	31.4	
PCB-165	PCB-165	15.8	42.3 J	16.6 J	58.7	36.7 J	38.8	50.4	15 J	
PCB-167	PCB-167	2200	4160	1570	2640	2320	1630	5360	1110	
PCB-169	PCB-169	3.67 U	46.5	18.7 U	51.9	11.3 U	19 U	95.8	7.08 U	
PCB-170	PCB-170	4040	15500	2610	11400	10100	5800	17500	1950	
PCB-171/173 PCB-172	PCB-171/173 PCB-172	1520 908	4240 2640	1110 617	3640 2340	3170 2150	1810 1120	5630 3020	805 538	
PCB-172 PCB-174	PCB-172 PCB-174	23.1	6020	20.9 U	9010	9010	3990	9710	21.9 J	
PCB-174 PCB-175	PCB-174 PCB-175	23.1	735	190	596	567	303	993	133	
PCB-173	PCB-176	71.9 J	1060	30.1	1260	1160	687	1270	6.32	
PCB-170	PCB-177	507	4340	556	6320	5610	2970	8550	524	
PCB-178	PCB-178	1030	1980	751	2690	2480	1290	4460	699	
PCB-179	PCB-179	4.53	2430	7.26 U	3690	3480	2150	4040	6.17 J	
PCB-180/193	PCB-180/193	12600	38100	9680	28800	25000	13900	41900	7920	
PCB-180/133	PCB-181	94.5	249	72.9	180	170	107	308	48.9	
PCB-182	PCB-182	45.8	8.05 U	59	117	111	57.9	183	43.5	
PCB-183/185	PCB-183/185	3830	11300	3090	9130	8250	4670	13900	2090	
PCB-184	PCB-184	32	72.2	90.3	96	88.6	29.7	152	31	
PCB-186	PCB-186	0.5 U	6.92 U	8.06 U	2.55 U	3.25 U	5.8 U	4.94 J	1.37 U	
PCB-187	PCB-187	6320	22200	5020	18300	15900	8130	22600	4750	
PCB-188	PCB-188	29.5	84.4	79.1	78.8	73.2	32.7	94.8	55.3	
PCB-189	PCB-189	214	464	188	423	375	196	524	127	
PCB-190	PCB-190	869	3350	643	2330	2070	1150	2710	513	
PCB-191	PCB-191	224	721	166	538	480	266	769	143	
PCB-192	PCB-192	1.33 U	19 U	16.7 U	11.9	14.7	13.1 U	6.41 U	2.56 U	
PCB-194	PCB-194	1480	7580	1210	6010	5360	2300	4650	856	
PCB-195	PCB-195	460	2610	332	2070	1870	870	2670	268	
PCB-196	PCB-196	1160	3930	978	2940	2720	1280	4760	741	
PCB-197/200	PCB-197/200	186	931	450	1100	1060	428	1440	160	
PCB-198/199	PCB-198/199	2220	6460	1640	6690	6040	2690	9700	1560	
PCB-201	PCB-201	311	1020	381	922	888	378	1830	247	
PCB-202	PCB-202	557	1710	623	2050	1860	755	2570	493	

				Tissue An	alytical Results				
	Location ID	PB052.1	PB052.1	PB053.2	PB053.2	PB053_B	PB056_A	PB056_A	PB057.2
	Sample ID	PB02-F-A-SAS-W-002-20110616	PB02-F-A-STM-W-003-20110616	PB02-F-A-GKF-W-025-20110621	PB02-F-A-GKF-W-029-20110622	PB02-F-A-GKF-W-013-20110618	PB02-F-A-GKF-W-010-20110618	PB02-F-A-GKF-W-011-20110618	PB03-F-A-GM-W-001-20110617
	Sample Date	6/16/2011	6/16/2011	6/21/2011	6/22/2011	6/18/2011	6/18/2011	6/18/2011	6/17/2011
	Sample Type	N	N	N	N	N	N	N	N
	Х	3201515.24	3201515.24	3201266.16	3201266.16	3201390.48	3201420.22	3201420.22	3201603.32
	Υ	13832564.31	13832564.31	13832433.69	13832433.69	13832440.54	13832183.54	13832183.54	13832054.52
	Scientific_Name	Cynoscion arenarius	Mugil cephalus	Fundulus grandis	Brevoortia patronus				
	Size Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
	Tissue_Category	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Migratory
	Tissue_Type	Whole Body							
Chemical	CAS Number	Result							
PCB-144	PCB-144	1530	2370	6220 J	5250	4980 J	3420	3710	1730 J
PCB-145	PCB-145	11.9 J	32.8	28.6 J	24.5 J	32 J	16.3	22.4	9.35 J
PCB-146	PCB-146	42.9 J	10100	94.4 J	76.1	21500 J	16900	15500	25.6 J
PCB-147/149	PCB-147/149	11600	30300	77300 J	62600	54600 J	40100	41800	25900 J
PCB-148	PCB-148	87.1	94.4	309 J	236	234 J	170	182	94.6 J
PCB-150	PCB-150	67.5	105	193 J	154	140 J	98.2	111	93.6 J
PCB-152	PCB-152	29.8 J	78	87 J	74.9	67.6 J	41.4	58	33.7 J
PCB-153/168	PCB-153/168	55900	60100	161000 J	132000	133000 J	106000	98100	44200 J
PCB-154	PCB-154	996	897	3010 J	2280	2320 J	1670	1610	864 J
PCB-155	PCB-155	125	62.6	202	166	182 J	172	117	77.3
PCB-156/157	PCB-156/157	7260	8830	18800 J	15800	18700 J	15400	12900	5610
PCB-158	PCB-158	6350	7490	17400 J	15100	15700 J	12700	11100	4930 J
PCB-159	PCB-159	19.5 U	22.5 U	12.9 UJ	12.5 U	559 J	483	401	7.22 UJ
PCB-160	PCB-160	21.6 U	26.7 U	13.8 UJ	12.8 U	9.23 UJ	14.5 U	33.3 U	9.12 UJ
PCB-161	PCB-161	5020	26 U	23800 J	19500	8.46 UJ	13.3 U	30.5 U	6840 J
PCB-162	PCB-162	344	302	743 J	684	755 J	1070	640 J	213 J
PCB-164	PCB-164	2470	3500	7530 J	6500	6520 J	5210	4810	2820 J
PCB-165	PCB-165	28.6 J	58.2	101 J	66.9	75.2 J	56.7	57.5 J	47.9 J
PCB-167	PCB-167	2450	3180	7390 J	6150	7230 J	5980	5270	2000 J
PCB-169	PCB-169	13.5 U	15.6 U	13.5 UJ	11.6 U	24 J	8.52 U	21.3 U	5.81 UJ
PCB-170	PCB-170	12000	11900	21100	19400	22000 J	20100	15000	8340 J
PCB-171/173	PCB-171/173	3460	3160	7170	6280	6750 J	5590	4550	2580 J
PCB-172	PCB-172	2220	2240	4380	3730	4030 J	3450	2720	1810 J
PCB-174	PCB-174	3900	5300	13600	10900	11100 J	9280	7750	6740 J
PCB-175	PCB-175	500	505	1370	1090	1280 J	1030	857	473 J
PCB-176	PCB-176	622	1040	2020	1550	1520 J	1180	1130	884 J
PCB-177	PCB-177	2590	4980	10900	9300	9520 J	7940	6320	4250 J
PCB-178	PCB-178	2090	2150	5710	4690	5340 J	4230	3540	2000 J
PCB-179	PCB-179	1740	3230	5610	4500	4340 J	3070	3160	2590 J
PCB-180/193	PCB-180/193	31100	28500	58000	49800	55800 J	50500	38300	21800 J
PCB-181	PCB-181	194	194	429	353	394 J	323	249	137 J
PCB-182	PCB-182	109	100 J	266	222	218 J	180	105 J	92.3 J
PCB-183/185	PCB-183/185	8880	8140	20900	17200	18600 J	15200	12400	7120 J
PCB-184	PCB-184	94.8	39	203	151	162 J	128	112	66.1 J
PCB-186	PCB-186	3.95 U	5.71 J	2.67 U	6.19 J	4.42 J	3.67 U	12.3 U	2.92 J
PCB-187	PCB-187	9980	16000	30900	28400	33100 J	31300	21400	12300 J
PCB-188	PCB-188	75.8	49.9	134	118	125	117	89	51.8
PCB-189	PCB-189	428	396	780	616	745 J	618	542	304 J
PCB-190	PCB-190	2450	2340	3440	3160	3720 J	3330	2520	1670 J
PCB-191	PCB-191	566	542	1110	979	1140 J	1020	760	411 J
PCB-192	PCB-192	17.2 J	12.9 U	6.82 U	6.76 U	9.58 J	8.31 J	15.8 U	7.96 J
PCB-194	PCB-194	7610	5000	7800	7470	8450 J	7940	5830	4460
PCB-195	PCB-195	2430	1720	3890	3640	3550 J	3190	2340	1430
PCB-196	PCB-196	3480	2480	7130	5910	6270 J	5590	4180	2340
PCB-197/200	PCB-197/200	1000	705	2250	1690	1720 J	1450	729	837
PCB-198/199	PCB-198/199	6330	5240	15100	12100	12800 J	11500	8390	5160
PCB-201	PCB-201	997	671	2650	1990	2060 J	1670	1320	741
PCB-202	PCB-202	2180	1410	3820	3180	3570	3270	2410	1440
			1	l .	l .	l .			

Part					Tissue Ana	lytical Results				
		Location ID	PB057.2	PB057.2	PB057.2	PB057_B	PB057_B	PB057_B	PB058.1	PB058.1
		Sample ID	PB03-F-A-STM-W-001-20110617	PB03-F-A-STM-W-001-COMP-201106	PB03-F-A-STM-W-002-20110617	PB03-F-A-GKF-W-001-20110615	PB03-F-A-GKF-W-006-20110617	PB03-F-A-PNF-W-001-COMP-201106	PB03-F-A-GM-W-003-20110618	PB03-F-A-GM-W-004-20110618
		Sample Date	6/17/2011	6/17/2011	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/18/2011	6/18/2011
Page		Sample Type	. N	N	N	N	N	N	N	N
		х	3201603.32	3201603.32	3201603.32	3201644.06	3201644.06	3201644.06	3201682.3	3201682.3
Page		Υ	13832054.52	13832054.52	13832054.52	13832071.63	13832071.63	13832071.63	13832011.93	13832011.93
Property Property		Scientific_Name	Mugil cephalus	Mugil cephalus	Mugil cephalus	Fundulus grandis	Fundulus grandis	Lagodon rhomboides	Brevoortia patronus	Brevoortia patronus
Ober Manual Tomography Work body Mode body Mode body Mode body Work body		Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
Central Chanter Chanter Standam Standam <t< th=""><th></th><th>Tissue_Category</th><th>Migratory</th><th>Migratory</th><th>Migratory</th><th>Sedentary</th><th>Sedentary</th><th>Migratory</th><th>Migratory</th><th>Migratory</th></t<>		Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Migratory	Migratory	Migratory
March Marc		Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
March Marc	Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Personal Personal	PCB-144	PCB-144	1150	1060			4710	1290	1700 J	2820
Part	PCB-145	PCB-145	9.75 J	15.4	11.9 J	22.7	26.6	2.67 J	12.2 J	28.1
PS-946 PS-946 PS-94	PCB-146	PCB-146	18.7 J	4410	17.7 UJ	17900	60.4	6750	23.9 J	10200
Post	PCB-147/149	PCB-147/149	12700	11100	10300 J	48600	58100	14400	25100 J	46300
Post Post	PCB-148	PCB-148	73.2	41.4	48.8 J	176	186	52.5	91.8 J	134
Per	PCB-150	PCB-150			48.6 J		138	45.6	92.3 J	176
Profession Pro	PCB-152	PCB-152	28.5	33	28.4 J	61.7	79.1		39.6 J	85.6
No. No. No. No. No. No. No. No. No. No.	-	PCB-153/168	30900	22400	24900 J	105000	114000	35800	40900 J	50000
Part	PCB-154	PCB-154	734	428	459 J	1700	1940	474	810 J	1200
March Marc	PCB-155	PCB-155	35.8	30.5	34.5 J	163	133	37.9	74.5	85.4
Process Proc	PCB-156/157	PCB-156/157	3710	3580	3250 J	14100	14500	5510	5130 J	7070
Property Property	PCB-158	PCB-158	3190	2860	2840 J	13000	13300	4310	4600 J	6040
Probate Prob	PCB-159	PCB-159	7.98 U	8.33 U	21 UJ	49.2 U	16.3 U	3.52 U	12.1 UJ	298
Probabil Probabil	PCB-160	PCB-160	7.56 U	2.3 U	16.2 UJ	49 U	22 U	2.98 U	7.08 UJ	15.3 U
PEPAR	PCB-161	PCB-161	4530	2.15 U	3640 J	46.9 U	17100	2.8 U	6370 J	13.2 U
PC-1655 PC-1	PCB-162	PCB-162	130	176	112 J	592 J	599 J	218	240 J	313 J
Page Page	PCB-164	PCB-164	1390	1160	1150 J	5460	5710	1690	2680 J	3730
March Marc	PCB-165	PCB-165	36.4	11	24.9 J	50.6 U	55.2	13.8	44.3 J	31.6 J
PE-1970	PCB-167	PCB-167	1360	1250	1180 J	5560	5660	2020	1830 J	2470
Peb-11/173	PCB-169	PCB-169	7.22 U	8.12 U	17 UJ	32.5 U	12.9 U	3.3 U	9.15 UJ	13.5 U
Fig. 1.2	PCB-170	PCB-170	5310 J	3870	4460 J	21800	17100	5910	7060	7600 J
PC-174	PCB-171/173	PCB-171/173	1560 J	1150	1200 J	5700	5020	1820	2260	2820 J
PG-175 PG-175 PG-175 PG-175 PG-175 PG-176 PG-176 PG-176 PG-176 PG-176 PG-176 PG-177 PG-176 PG-177 P	PCB-172	PCB-172	1100 J	735	894 J	3770	3030	996	1550	1670 J
Page Page	PCB-174	PCB-174	2410 J	1620	1670 J	9950	9170	2260	5840	8060 J
PCB-177 PCB-177 PCB-177 PCB-177 PCB-178 PCB-	PCB-175	PCB-175	304 J	211	229 J	989	893	305	408	469 J
PC-179 PC-180 PC-180 PC-180 PC-181 P	PCB-176	PCB-176	580 J	430	394 J	1270	1290	389	814	1240 J
PC-1379 PC-1379 PC-130713 PC-13071	PCB-177	PCB-177	2420 J	1640	1730 J	8600	7500	1510	3820	4870 J
PCB-180/193 PCB-180/193 14000 J 8480 11600 J 50200 42400 1500 18100 19300 J PCB-181 PCB-181 88.1 J 31.2 U 75.6 J 324 288 94.1 125 146 J PCB-182 PCB-182 37.4 J 27.7 47.8 J 122 4.31 U 48.6 7.9 48.9 J PCB-182 PCB-182 37.4 J 27.7 47.8 J 122 4.31 U 48.6 7.9 48.9 J PCB-183/185 PCB-184 4530 J 2670 3700 J 14800 13700 4330 6130 6130 6880 J PCB-184 PCB-186 22.1 J 20.2 27.7 J 119 111 21.5 61.0 78.5 J 45.4 U PCB-186 PCB-186 23.8 U 28.1 3.88 U 9.20 U 3.03 U 0.038 U 2.57 J 4.54 U PCB-187 PCB-186 23.8 U 2.88 U 2.10 U 3.00 U 3.00 U 3.00 U 3.00 U	PCB-178	PCB-178								2230 J
PCB-181 PCB-181 83.1 J 31.2 U 75.6 J 324 268 94.1 125 146 J PCB-182 PCB-182 37.4 J 27 47.8 J 122 4.31 U 48.6 79 48.9 J PCB-183/185 PCB-183/185 4530 J 2670 3700 J 14800 13700 4330 6130 6880 J PCB-184 PCB-184 24.1 J 20.2 23.7 J 119 111 21.5 61 79.5 J PCB-186 PCB-186 2.8 U J 20.2 23.7 J 119 111 21.5 61 79.5 J PCB-186 PCB-186 3.28 U J 2.81 3.88 U J 9.26 U 3.63 U 0.818 U 2.57 J 4.54 U PCB-187 PCB-187 7970 J 4880 U 3.88 U J 9.26 U 3.63 U 0.818 U 2.57 J 4.54 U PCB-188 2.89 U J 19 U 2.78 110 U 9.08 3.43 U 4.56 U 2.77 U 3.73 U	PCB-179	PCB-179	1700 J	1330	1240 J			958	2370	3870 J
PCB-182 PCB-182 37.4 J 27 47.8 J 122 4.31 U 48.6 79 48.9 J PCB-183/185 PCB-183/185 4530 J 2670 3700 J 14800 13700 4330 6330 6630 6680 J PCB-184										
PCB-183/185 PCB-183/185 4530 J 2670 3700 J 14800 13700 4330 6130 6880 J PCB-184 PCB-184 PCB-184 24.1 J 20.2 23.7 J 119 111 21.5 61 79.5 J PCB-186 PCB-186 3.28 UJ 2810 3.88 UJ 9.26 U 3.63 U 0.018 U 2.57 J 4.54 U PCB-187 PCB-187 7970 J 4980 6180 J 30000 24800 8060 10600 11300 J PCB-188 PCB-188 28.9 19 27.8 110 90.8 34.3 45.6 56.1 J PCB-189 PCB-189 221 J 145 19 J 364 54 20 20 22 373 PCB-190 PCB-191 267 J 152 19 J 369 280 1180 1400 3170 364 J PCB-191 PCB-191 267 J 152 221 J 1010 825 266 347 364										
PCB-184 PCB-184 24.1 J 20.2 23.7 J 119 111 21.5 61 79.5 J PCB-186 PCB-186 3.28 UJ 2.81 3.68 UJ 9.26 U 3.63 U 0.818 U 2.57 J 4.54 U PCB-187 PCB-187 PCB-187 7970 J 4980 6180 J 30000 24800 8060 10600 11300 J PCB-188 PCB-187 7970 J 4980 6180 J 30000 24800 8060 10600 11300 J PCB-189 PCB-188 2.81 J 19 27.8 110 9.08 34.3 45.6 56.1 J PCB-189 PCB-189 221 J 145 159 J 534 542 202 272 373 PCB-190 PCB-190 PCB-191 257 J 152 221 J 1010 825 266 347 364 J PCB-191 PCB-191 257 J 881 J 221 J 1010 825 266 347 364 J <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>										
PCB-186 CB-186 3.28 UJ 2.81 3.68 UJ 9.26 U 3.63 U 0.818 U 2.57 J 4.54 U PCB-187 PCB-187 7970 J 4990 6180 J 30000 24800 8060 10600 11300 J PCB-188 PCB-188 2.89 19 2.78 110 9.8 3.43 4.5 56.1 J PCB-198 PCB-198 2.21 J 145 159 J 534 542 20.2 272 373 PCB-190 PCB-199 1.090 J 731 92 J 3650 2880 1180 1400 1530 J PCB-191 PCB-191 2.67 J 152 J 1510 J 1010 J 825 J 266 347 J 364 J PCB-192 PCB-192 4.25 U 2.81 U 8.70 U 8.97 U 5.44 U 6.9 8.60 U PCB-192 PCB-192 4.65 U 8.81 U 2.46 U 8.97 U 5.44 U 6.9 8.60 U PCB-193 PCB-195 8	PCB-183/185	-								
PCB-187 PCB-187 7970 J 4980 6180 J 30000 24800 8060 10600 11300 J PCB-188 PCB-188 28.9 19 27.8 110 90.8 34.3 45.6 56.1 J PCB-189 PCB-189 PCB-190 145 159 J 534 542 202 272 373 PCB-190 PCB-190 1090 J 731 912 J 3660 2880 1180 1400 1530 J PCB-191 PCB-191 267 J 152 912 J 3660 2880 1180 1400 1530 J PCB-192 PCB-194 267 J 152 211 J 1010 855 266 140 409 364 J PCB-192 PCB-192 4.25 UJ 25 U 881 UJ 24.6 U 8.97 U 5.44 6.9 4.6 9 8.56 U PCB-194 PCB-194 2320 1200 8700 7130 2310 3650 3700 PCB-195	PCB-184									
PCB-188 PCB-188 28.9 19 27.8 110 90.8 34.3 45.6 56.1 J PCB-189 PCB-189 PCB-189 221 J 145 159 J 534 542 202 272 373 PCB-190 PCB-190 1090 J 731 912 J 3650 2880 1180 1400 1530 J PCB-191 PCB-191 267 J 152 221 J 1010 825 266 37 364 J PCB-192 PCB-192 4.25 UJ 25 U 8.81 UJ 24.6 U 8.97 U 5.44 6.9 8.56 U PCB-194 PCB-194 2320 1200 2160 8700 7130 2310 3650 370 PCB-195 RS4 465 693 3330 2850 897 1300 1410 PCB-197/200 PCB-197/200 1340 714 1140 5380 4670 1380 286 287 1300 1310 2310 2310 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>										
PCB-189 PCB-189 221 J 145 159 J 534 542 202 272 373 PCB-190 PCB-190 1090 J 731 912 J 3650 2880 1180 1400 1530 J PCB-191 PCB-191 267 J 152 221 J 1010 825 266 347 364 J PCB-192 PCB-192 4.25 UJ 25 U 8.81 UJ 24.6 U 8.97 U 5.44 6.9 8.56 U PCB-193 PCB-194 2320 1200 2160 8700 7130 2310 3650 3700 PCB-195 834 465 693 3330 2850 897 1300 1410 PCB-196 1410 714 1140 538 4670 1380 280 280 231 PCB-197/200 PCB-197/200 395 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>										
PCB-190 PCB-190 1090 J 731 912 J 3650 2880 1180 1400 1530 J PCB-191 PCB-191 267 J 152 221 J 1010 825 266 347 364 J PCB-192 PCB-192 4.25 UJ 25 U 8.81 UJ 24.6 U 8.97 U 5.44 6.9 8.56 U PCB-194 PCB-194 2320 1200 2160 8700 7130 2310 3650 370 PCB-195 834 465 693 3330 2850 897 1300 1410 PCB-196 CB-196 1410 714 1140 5380 4670 1380 2080 2130 PCB-197/200 PCB-197/200 756 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 1380 418 683 775										
PCB-191 PCB-191 152 221 J 1010 825 266 347 364 J PCB-192 PCB-192 4.25 UJ 25 U 8.81 UJ 24.6 U 8.97 U 5.44 6.9 8.56 U PCB-194 PCB-194 2320 1200 2160 8700 7130 2310 3650 3700 PCB-195 834 465 693 3330 2850 897 1300 1410 PCB-196 PCB-196 1410 714 1140 5380 4670 1380 2080 2130 PCB-197/200 PCB-197/200 95 35 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 2150 4640 5000 PCB-201 PCB-201 390 236 320 1420 1380 418 683 775										
PCB-192 4.25 UJ 25 U 8.81 UJ 24.6 U 8.97 U 5.44 6.9 8.56 U PCB-194 PCB-194 2320 1200 2160 8700 7130 2310 3650 3700 PCB-195 PCB-195 834 465 693 3330 2850 897 1300 1410 PCB-196 PCB-196 1410 714 1140 5380 4670 1380 2080 2130 PCB-197/200 PCB-197/200 PCB-197/200 9CB-197/200 395 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 418 683 775										
PCB-194 PCB-194 2320 1200 2160 8700 7130 2310 3650 3700 PCB-195 PCB-195 834 465 693 3330 2850 897 1300 1410 PCB-196 PCB-196 1410 714 1140 5380 4670 1380 2080 2130 PCB-197/200 PCB-197/200 395 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 2150 4640 5000 PCB-201 PCB-201 390 236 320 1420 1380 418 683 775		_								
PCB-195 R34 465 693 3330 2850 897 1300 1410 PCB-196 PCB-196 1410 714 1140 5380 4670 1380 2080 2130 PCB-197/200 PCB-197/200 395 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 2150 4640 5000 PCB-201 PCB-201 390 236 320 1420 1380 418 683 775										
PCB-196 PCB-196 1410 714 1140 5380 4670 1380 2080 2130 PCB-197/200 PCB-197/200 395 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 2150 4640 5000 PCB-201 PCB-201 390 236 320 1420 1380 418 683 775										
PCB-197/200 PCB-197/200 395 256 311 1230 1170 354 782 531 PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 2150 4640 5000 PCB-201 PCB-201 390 236 320 1420 1380 418 683 775										
PCB-198/199 PCB-198/199 2640 1560 2220 10500 9310 2150 4640 5000 PCB-201 PCB-201 390 236 320 1420 1380 418 683 775										
PCB-201 PCB-201 390 236 320 1420 1380 418 683 775										
	PCB-198/199	· ·								
PCB-202 PCB-202 707 501 635 2940 2510 613 1310 1550										
	PCB-202	PCB-202	707	501	635	2940	2510	613	1310	1550

				rissue Ana	alytical Results				
	Location ID	PB058.1	PB059.1_A	PB059.1_A	PB059.3	PB059.4	PB059.4	PB059.5	PB063.1_A
	Sample ID		PB03-F-A-GKF-W-003-20110615	PB03-I-A-BCR-W-003-COMP-201106		PB03-I-B-BCR-W-003-20110628		PB03-I-A-BCR-W-001-COMP-201106	PB03-F-A-GKF-W-009-20110617
	Sample Date	6/18/2011	6/15/2011	6/19/2011	6/26/2011	6/27/2011	6/27/2011	6/27/2011	6/17/2011
	Sample Type		N	N	N	N	N	N	N
	x x	3201682.3	3201501.32	3201501.32	3201714.16	3201714.16	3201714.16	3201714	3201509.99
	v	13832011.93	13831828.71	13831828.71	13831964.63	13831964.63	13831964.63	13831965	13831505.57
	Scientific_Name	Brevoortia patronus	Fundulus grandis	Callinectes sapidus	Callinectes sapidus	Callinectes sapidus	13831304.03	Callinectes sapidus	Fundulus grandis
	Size_Class	=<15	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13		>2 to =<7.5	=<15
	Tissue_Category	Migratory	Sedentary	Sedentary	Migratory	Migratory		Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body		Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-144	PCB-144	2770	3480	37.6	41.4	70.2	70.8	18.3	3570 J
PCB-145	PCB-144 PCB-145	25.1	20.2	1.21 U	1.95 U	2.14 U	3.48 U	2.82 U	22 J
PCB-146	PCB-143	10200	32.7 U	3160	4370	4240	4070	3030	67 J
PCB-147/149	PCB-146 PCB-147/149	45700	39200	808	868	1370	1330	381	42400 J
	·	128			56.4		56.2		
PCB-148	PCB-148		140	33.5		55.3		31.2	150 J
PCB-150	PCB-150	177	84.9	1.12 U	1.87 U	2.05 U	3.38 U	2.61 U	93.3 J
PCB-152	PCB-152	84.8 J	57.3	1.1 U	0.974 J	2.52 J	2.08 J	2.55 U	48.8 J
PCB-153/168	PCB-153/168	50200	97800	17100	27300	26100	25300	20300	113000 J
PCB-154	PCB-154	1180	1390	324	586	573	536	367	1510 J
PCB-155	PCB-155	92.1	117	25.7	34.3	34.7	31.8	22	120
PCB-156/157	PCB-156/157	7110	14000	3050	4130	3980	3700	3790	13900
PCB-158	PCB-158	6190	12300	2510	3940	3720	3600	2920	13500 J
PCB-159	PCB-159	323	32 U	7.33 U	9.21 J	4.18 U	6.29 U	5.3 U	25.3 UJ
PCB-160	PCB-160	19.9 U	29.7 U	2.82 U	4.72 U	7.72 U	8.42 U	2.48 U	30 UJ
PCB-161	PCB-161	17.2 U	14200	2.64 U	4.03 U	6.58 U	7.25 U	2.33 U	15800 J
PCB-162	PCB-162	222	565	106	132	106	107	123	550 J
PCB-164	PCB-164	3740	4840	2.75 U	46.1	101	104	2.42 U	5590 J
PCB-165	PCB-165	30.9	30.6 U	8.05 J	10.7	11.5 J	11.2	4.73 J	73.9 J
PCB-167	PCB-167	2560	5180	1090	1510	1410	1340	1250	5080
PCB-169	PCB-169	15.4 U	19.4 U	6.93 U	4.51 U	4.54 U	6.32 U	4.72 U	85.3 UJ
PCB-170	PCB-170	8220 J	19700	1950	2350 J	2290 J	2120	1650	15300
PCB-171/173	PCB-171/173	2930 J	4820	739	979 J	949 J	898	673	4190
PCB-172	PCB-172	1760 J	3360	441	506 J	482 J	464	405	3030
PCB-174	PCB-174	8410 J	7770	28.3	47.7 J	58.7 J	55.4 J	20.7 J	7020
PCB-175	PCB-175	508 J	807	126	142 J	146 J	139	101	750
PCB-176	PCB-176	1260 J	953	10.7	26 J	1.72 U	3.45 U	1.58 U	833
PCB-177	PCB-177	5090 J	6880	349	360 J	435 J	394	260	5580
PCB-178	PCB-178	2320 J	3440	497	579 J	650 J	609	401	3040
PCB-179	PCB-179	3970 J	2790	5.95 J	10.9 J	7.27 J	8.85	7.61	2460
PCB-180/193	PCB-180/193	20600 J	46200	6380	8060 J	7450 J	6850	6040	41100
PCB-181	PCB-181	152 J	272	48.1	67.8 J	62 J	58	48.6	259
PCB-182	PCB-182	90.2 J	166	31.1	42 J	39.8 J	39.3	33.3	136 J
PCB-183/185	PCB-183/185	7150 J	12400	1850	2570 J	2540 J	2400	1770	11900
PCB-184	PCB-184	81.7 J	90.9	18.5	29.7 J	28.4 J	24.1	16.8	87.9
PCB-186	PCB-186	5.53 U	5.7 U	0.675 U	2.46 U	1.82 U	3.59 U	1.79 U	8.15 U
PCB-187	PCB-187	11900 J	26700	3400	3990 J	3670 J	3410	3340	23100
PCB-188	PCB-188	53.5 J	80.9	23.1	25.4 J	20.7 J	20.5 J	14.7	76.7
PCB-189	PCB-189	386	478	94.7	133	133	122	108	476
PCB-190	PCB-190	1620 J	3300	415	536 J	481 J	443	423	2700
PCB-191	PCB-191	380 J	929	109	151 J	139 J	132	111	813
PCB-192	PCB-192	8.2 U	18.3 U	76.3 U	2.73 U	2.94 U	5.28 U	1.96 U	10.6 U
PCB-194	PCB-194	4020	7440	677	840	824	773	607	6480
PCB-195	PCB-195	1510	2760	183	257	270	254	135	2630
PCB-196	PCB-196	2320	4600	535	773	760	682	537	3870
PCB-197/200	PCB-197/200	994	935	85.8	136	138	126	79	911
PCB-198/199	PCB-198/199	5460	8730	974	1360	1320	1160	901	7480
PCB-201	PCB-201	832	1140	158	213	208	194	123	1070
PCB-202	PCB-202	1650	2240	312	339	329	300	211	2360

					Tissue Ana	alytical Results				
Part		Location ID	PB063.1_A	PB063.2_A	PB063.2_A	PB069_A	PB069_A	PB069_A	PB084.1	PB084.1
Part		Sample ID	PB03-F-A-GKF-W-010-20110617	PB03-F-A-GKF-W-005-20110615	PB03-F-A-GKF-W-013-20110617	PB03-F-A-GKF-W-002-20110615	PB03-F-A-GKF-W-007-20110617	PB03-I-A-BCR-W-002-COMP-201106	PB04-F-A-GKF-W-005-20110627	PB04-F-A-GKF-W-013-20110627
		Sample Date	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/17/2011	6/19/2011	6/27/2011	6/27/2011
1899 1899		Sample Type	N	N	N	N	N	N	N	N
Profession Pro		х	3201509.99	3201561.28	3201561.28	3201132.36	3201132.36	3201132.36	3200727.85	3200727.85
Page		Υ	13831505.57	13831519.55	13831519.55	13830993.22	13830993.22	13830993.22	13829784.47	13829784.47
Tree Tree Part		Scientific_Name	Fundulus grandis	Fundulus grandis	Fundulus grandis	Fundulus grandis	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Fundulus grandis
Classical (State) Tissoc-lying (State) Mode Rody (State) White Rody (State) White Rody (Rod) (Rod) White Rody (Rod) (Ro		Size_Class	=<15	=<15	=<15	=<15	=<15	>2 to =<7.5	=<15	=<15
formes C5 Stations Fault 1000 1000 1500		Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary
Pose Pose		Tissue_Type	Whole Body	Whole Body	Whole Body					
Profession	Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
Profession	PCB-144	PCB-144	3650	3300	3080	5510	4010 J	73.4	2100	3450
Principle Prin	PCB-145	PCB-145	19.4	22.3 J	20	36.6 J	23.1 J	3.53 U	17.8	28.4 J
March Marc	PCB-146	PCB-146	18300	15900	13500	22200	17800 J	5080	8850	13900
March	PCB-147/149	PCB-147/149	40700	39800	34600	60500	44500 J	1260	26100	41300
Probability	PCB-148	PCB-148	153	142	135	237	158 J	46.6	83	135
Process	PCB-150	PCB-150	86.8	95	85.3	170	103 J	3.26 U	73.2	97.6
Part Part	PCB-152	PCB-152	52.9	63.7	42.6	99.1	58.3 J	3.19 U	45	76.2
March Marc	PCB-153/168	PCB-153/168	110000	93400	83300	132000	106000 J	27800	45900	71600
PASSIGNET PASSIGNET 14500 14500 14500 14500 14500 14701 14501 14501 14501 1500	PCB-154	PCB-154	1580	1350	1340	2230	1610 J	497	799	1240
March Marc	PCB-155	PCB-155	116	112	110	145	97.4	34.1	52.7	78.5
K6.169 K6.19 K6.19 K6.10 A8.1 K6.11 A8.1 K6.11 A8.1 K6.12 A8.2	PCB-156/157	PCB-156/157	13500	12900	11500	18500	13500	4720	6250	9560
Page Page	PCB-158	PCB-158	13100	11400	9980	15900	13500 J	3920	5620	8850
	PCB-159	PCB-159	16.1 U	36 U	343	561	424 J	8.69 J	160	10.6 U
Profession	PCB-160	PCB-160	17.1 U	41.4 U	7.25 U	32.6 U	13.9 UJ	1.95 U	12.6 U	16.9 U
PC-164 PC-164 PC-164 PC-165 P	PCB-161	PCB-161	16.4 U	40.4 U	6.64 U	31 U	13.2 UJ	1.83 U	10.7 U	14.3 U
Part Part	PCB-162	PCB-162	576	590	479	756	637 J	165	269 J	284
PASS PASS	PCB-164	PCB-164	5280	4750	4160	6860	5500 J	1.9 U	2490	3920
Pach Pach	PCB-165	PCB-165	54.5	53.6 J	47.4	62.4 J	58.3 J	12.8 J	22.2	41.6
	PCB-167	PCB-167	5380	5070	4530	6800	5120	1680	2460	3720
PG-17/17/27	PCB-169	PCB-169	11.9 U	25.7 U	65.8	R	25 UJ	1.92 U	11.8 U	11.5 U
PCH-172	PCB-170	PCB-170	19300	16800	13900	19200	15100	2810	6310 J	9460 J
	PCB-171/173	PCB-171/173	5060	4540	4140	5550	4130	1110	2020 J	3090 J
	PCB-172	PCB-172	3480	2760	2430	3700	2770	652	1190 J	1750 J
Page Page	PCB-174	PCB-174	7870	7690	6790	10400	7380	39.5	3960 J	5740 J
PCB-177 PCB-177 PCB-177 PCB-177 PCB-178 3780 3120 3240 4540 3330 836 1710 2270 2270 2270 2270 2470	PCB-175	PCB-175	913	757	765	1080	805	186	369 J	581 J
PCB-178 PCB-178 2780 3120 3240 4540 3330 836 1710 J 2570 J PCB-179 PCB-179 2500 2700 2620 4140 2610 9.39 1730 J 2710 J PCB-180/193 A6200 33500 33500 43900 37300 8620 16000 J 23800 J PCB-181 PCB-181 286 255 228 325 239 64.8 102 J 164 J PCB-182 PCB-182 187 144 J 128 196 141 40 50.1 J 95.9 J PCB-183/195 PCB-183/195 13900 11400 11000 15400 11200 2710 2530 46.6 J 88.7 J PCB-184 PCB-184 100 8.2 90.5 140 77.9 25.3 46.6 J 88.7 J PCB-186 PCB-186 3.36 6.19 J 3.09 J 9.55 J PCB-187 PCB-188 88.6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-189 PCB-189 48.4 4.59 4.25 4.55 4.41 4.5 4.40 4.31 PCB-190 PCB-190 91.30 2590 2230 3140 2460 561 310 233 318 PCB-191 PCB-192 PCB-192 PCB-192 PCB-192 PCB-193 PCB-193 PCB-193 PCB-194 PCB-194 PCB-195 PCB-195 PCB-196 PCB-197 PCB-176	PCB-176	1000	929	923	1450	924	16.8	557 J	820 J	
PCB-179 PCB-179 2500 2700 2620 4140 2610 9.39 1730 J 2710 J PCB-180/133 PCB-180/133 48200 39500 38500 49300 37300 8620 16000 J 28800 J PCB-181 PCB-181 286 255 228 325 239 64.8 10.2 J 1164 J PCB-182 PCB-183/185 187 144 J 128 196 141 40 50.1 J 55.9 J PCB-183/185 1870 1140 1100 15400 11200 2710 5210 J 800 J PCB-184 PCB-183/185 13900 11400 11000 15400 11200 2710 J 5210 J 800 J PCB-186 PCB-187 CCB-186 3.36 6.19 U 3.09 U 9.45 U 4.11 U 0.847 U 4.24 U 3.91 U PCB-187 PCB-187 26600 2.210 U 2.0000 2.8500 2.2500 5150 U 8820 J 13000 J	PCB-177	PCB-177	6920	6190	5830	7730	5640	439	3050 J	4470 J
PCB-180/193	PCB-178	PCB-178	3780	3120	3240	4540	3330	836	1710 J	2570 J
PCB-181 PCB 181 286 255 228 325 239 64.8 102 J 164 J PCB-182 PCB-182 187 144 J 128 196 141 40 50.1 J 95.9 J PCB-183/185 PCB-183/185 1390 11400 11000 1540 1120 2710 5210 J 8000 J PCB-184 PCB-187 100 82 90.5 140 77.9 25.3 64.6 J 88.7 J PCB-186 PCB-186 3.36 6.19 U 3.09 U 9.45 U 4.11 U 0.847 U 4.24 U 3.91 U PCB-187 PCB-187 26600 24100 20600 28500 222600 5190 8820 J 3.91 U PCB-188 B.8.6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-192 PCB-188 88.6 47.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-199 PCB-189	PCB-179	PCB-179	2900	2700	2620	4140	2610	9.39	1730 J	2710 J
PCB-182 PCB-182 187 144 J 128 196 141 40 50.1 J 95.9 J PCB-183/185 PCB-183/185 13900 11400 11000 15400 11200 2710 55.1 J 80.00 J PCB-184 PCB-184 100 8.2 90.5 140 77.9 25.3 64.6 J 88.7 J PCB-186 PCB-186 3.36 6.19 U 3.09 U 94.5 U 4.11 U 0.847 U 4.24 U 3.91 U PCB-187 PCB-187 26600 24100 20600 228500 22600 5190 8820 J 13000 J PCB-188 PCB-188 8.8 6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-189 PCB-189 484 45.9 42.5 69.4 49.9 140 23.3 33.8 PCB-190 PCB-190 3130 2590 2230 3140 2460 2660 2660 2660 2660 PCB-191 PCB-191 94.7 78.3 70.3 88.4 74.7 16.3 27.7 J 45.5 J PCB-192 PCB-194 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-196 PCB-196 5020 4100 3850 5720 4060 70.9 1860 27.30 J PCB-197/000 PCB-197/00 PCB-197/00 1070 901 1040 1400 94.6 1100 94.6 119 511 711 J PCB-201 PCB-201 1340 1110 1290 1630 1630 1600 221 605 918 J PCB-201 PCB-201 1340 1110 1290 1630 1630 1600 221 605 918 J PCB-197 PCB-201 PCB-201 1340 1100 1200 1630 1630 1600 1200 1210 605 918 J PCB-197 PCB-201 PCB-201 1340 1100 1200 1630 1630 1630 1600 221 605 918 J PCB-197 PCB-197 PCB-201 1340 1100 1200 1630 1630 1630 1600 1200 1210 1200 1210 1200	PCB-180/193	PCB-180/193	48200	39500	35800	49300	37300	8620	16000 J	23800 J
PCB-183/185 PCB-183/185 13900 11400 11000 15400 11200 2710 5210 8000 1	PCB-181	PCB-181	286	255	228	325	239	64.8	102 J	164 J
PCB-184 PCB-184 100 82 90.5 140 77.9 25.3 64.6 J 88.7 J PCB-186 PCB-186 3.36 6.19 U 3.09 U 9.45 U 4.11 U 0.847 U 4.24 U 3.91 U PCB-187 PCB-187 26600 24100 20600 28500 22600 5190 8820 J 1300 J PCB-188 PCB-188 88.6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-189 PCB-189 484 459 425 694 499 140 233 338 PCB-190 PCB-190 3130 2590 2230 3140 2460 561 1010 J 1440 J PCB-191 PCB-192 9.21 U 2.06 U 536 13 U 5.87 U 163 297 J 456 J PCB-192 9.21 U 2.06 U 536 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-193 PCB-194 <t< td=""><td>PCB-182</td><td>PCB-182</td><td>187</td><td>144 J</td><td>128</td><td>196</td><td></td><td>40</td><td>50.1 J</td><td>95.9 J</td></t<>	PCB-182	PCB-182	187	144 J	128	196		40	50.1 J	95.9 J
PCB-186 PCB-186 3.36 6.19 U 3.09 U 9.45 U 4.11 U 0.847 U 4.24 U 3.91 U PCB-187 PCB-187 26600 24100 20600 28500 22600 5190 8820 J 13000 J PCB-188 PCB-188 8.8.6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-189 484 459 425 694 499 140 233 338 PCB-190 PCB-190 3130 2590 2230 3140 2460 561 1010 J 1440 J PCB-191 PCB-192 9.21 U 22.0 U 5.36 13 U 5.87 U 1.63 U 297 J 456 J PCB-192 P.CB-192 9.21 U 20.6 U 5.36 U 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 2780	PCB-183/185	PCB-183/185	13900	11400	11000	15400	11200	2710	5210 J	8000 J
PCB-187 PCB-188 2600 2410 26600 28500 22600 5190 8820 J 13000 J PCB-188 PCB-188 88.6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-189 PCB-189 484 459 425 694 499 140 233 338 PCB-190 PCB-191 947 783 703 984 747 163 29.1 140 J 140 J PCB-191 PCB-192 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J	PCB-184	PCB-184	100		90.5	140	77.9	25.3	64.6 J	88.7 J
PCB-188 PCB-188 88.6 77.8 87.2 94.5 73.9 25.4 34.9 J 46.9 J PCB-189 PCB-189 484 459 425 694 499 140 233 338 PCB-190 PCB-190 3130 2590 2230 3140 2460 561 1010 J 1440 J PCB-191 947 783 703 984 747 163 297 J 456 J PCB-192 PCB-192 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 1070 901 3100 1400 946 119 511 711 J PCB-198/199 PCB-197 7680 <t< td=""><td>PCB-186</td><td>PCB-186</td><td>3.36</td><td>6.19 U</td><td>3.09 U</td><td>9.45 U</td><td>4.11 U</td><td>0.847 U</td><td>4.24 U</td><td>3.91 U</td></t<>	PCB-186	PCB-186	3.36	6.19 U	3.09 U	9.45 U	4.11 U	0.847 U	4.24 U	3.91 U
PCB-189 484 459 425 694 499 140 233 338 PCB-190 PCB-190 3130 2590 2230 3140 2460 561 1010 J 1440 J PCB-191 PCB-191 947 783 703 984 747 163 297 J 456 J PCB-192 PCB-192 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-195 2780 2570 2140 3200 2230 290 1110 1610 J PCB-196 PCB-196 5020 4100 3850 5720 4060 799 1860 2730 J PCB-197/200 PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199	PCB-187	PCB-187	26600	24100	20600	28500	22600	5190	8820 J	13000 J
PCB-190 PCB-190 3130 2590 2230 3140 2460 561 1010 J 1440 J PCB-191 PCB-191 947 783 703 984 747 163 297 J 456 J PCB-192 PCB-192 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-195 2780 2570 2140 3200 2230 290 1110 1610 J PCB-196 PCB-198 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 1080 8080 1400 3590 5290 J PCB-2	PCB-188	PCB-188	88.6	77.8	87.2	94.5	73.9	25.4	34.9 J	46.9 J
PCB-191 947 783 703 984 747 163 297 J 456 J PCB-192 PCB-192 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-195 2780 2570 2140 3200 2230 290 1110 1610 J PCB-196 PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 PCB-201 1340 1110 1290 1630 1160 121 605 918 J	PCB-189	PCB-189	484	459	425	694	499	140	233	338
PCB-192 9.21 U 20.6 U 5.36 13 U 5.87 U 1.26 U 6.43 U 7.65 U PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-195 2780 2570 2140 3200 2230 290 1110 1610 J PCB-196 PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-190	PCB-190	3130	2590	2230	3140	2460	561	1010 J	1440 J
PCB-194 7150 7050 4970 7860 5520 990 2570 3620 J PCB-195 PCB-195 2780 2570 2140 3200 2230 290 1110 1610 J PCB-196 PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-191	PCB-191	947	783	703	984	747	163	297 J	456 J
PCB-195 2780 2570 2140 3200 2230 290 1110 1610 J PCB-196 PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-192	PCB-192	9.21 U	20.6 U	5.36	13 U	5.87 U	1.26 U	6.43 U	7.65 U
PCB-196 5020 4100 3850 5720 4060 709 1860 2730 J PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-194	PCB-194	7150	7050	4970	7860	5520	990	2570	3620 J
PCB-197/200 PCB-197/200 1070 901 1040 1400 946 119 511 711 J PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-195	PCB-195	2780	2570	2140	3200	2230	290	1110	1610 J
PCB-198/199 PCB-198/199 9670 7680 8160 10800 8080 1400 3590 5290 J PCB-201 PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-196	PCB-196	5020	4100	3850	5720	4060	709	1860	2730 J
PCB-201 PCB-201 1340 1110 1290 1630 1160 221 605 918 J	PCB-197/200	PCB-197/200	1070	901	1040	1400	946	119	511	711 J
	PCB-198/199	PCB-198/199	9670	7680	8160	10800	8080	1400	3590	5290 J
PCB-202 PCB-202 2490 2300 2210 2880 2120 464 1030 1540 J	PCB-201	PCB-201	1340	1110	1290	1630	1160	221	605	918 J
	PCB-202	PCB-202	2490	2300	2210	2880	2120	464	1030	1540 J

Appendix D
Tissue Analytical Results

	Tissue Analytical Results								
Sumple Type Park		Location ID	PB095.2	PB101_D	PB101_D	PB101_D	PB102		
Second		Sample ID	PB04-I-A-BCR-W-001-COMP-201106	PB04-F-A-GKF-W-001-20110627	PB04-F-A-GKF-W-001-COMP-201106	PB04-F-A-GKF-W-009-20110627	PB04-F-A-PNF-W-001-COMP-201106		
Second		Sample Date	6/29/2011	6/27/2011	6/27/2011	6/27/2011	6/27/2011		
		•	· ·		1				
Part		х х							
		 V							
Sec. Page		Scientific Name							
		_	·	_	_	•			
Total Total (Section) Cast Number Benul Secul Secul Member Benul R3 344 R3 144 9.72 U 38.60 373.0 393.3 335.0 R3 344 R3 140 8.70 U 3.60 373.0 393.3 335.0 R3 444 R3 140 155.0 155.00 150.0 150.0 150.0 250.0 </th <th></th> <th>_</th> <th></th> <th></th> <th></th> <th></th> <th></th>		_							
Chemical Colo Nombre Result Res			, in the second		· ·	•			
No. No.	Chamical		·	•	·	•			
PAPER									
PC-1-16									
PC 1477/26F									
PC-146									
PK-120	<u> </u>	-							
PR 132									
RPS-153/186 RPS-155 RPS-1									
PC-1545									
PCB-155 PCB-156 PCB-156 13.2 10.7 38.9 1.23 4.12 1.25	<u> </u>		9790						
PASS PASS	PCB-154	PCB-154	211	1370	1500	1430	563		
PATE PATE	PCB-155	PCB-155	13.2	107	98.9	123	41.2		
PA-199	PCB-156/157	PCB-156/157	1630	10200	12200	11000	6420		
PER-1640	PCB-158	PCB-158	1470	9560	9570	9560	5290		
PG-1616 PG-1616 PG-1616 PG-162 PG-1626 PG-16	PCB-159	PCB-159	10.3 U	263	26.4 U	282	4.28 U		
	PCB-160	PCB-160	14 U	18.1 U	17.5 U	18 U	10.9 U		
PCB-164 PCB-165 13.1 U 4190 4270 4370 1510 PCB-165 PCB-165 13.8 U 42 44.6 47 11.6 U PCB-167 PCB-167 630 3960 4400 4290 4290 PCB-168 PCB-169 12 U 17.1 U 30.8 U 19.3 U 3.84 U PCB-170 PCB-170 630 16000 11000 11000 11000 11000 PCB-170 PCB-170 630 16000 13000 11000 11000 1300 U PCB-171/73 PCB-172 287 3401 3750 3600 3600 1830 PCB-171/73 PCB-172 207 1300 2370 2170 2270 PCB-174 PCB-174 112 U 6480 7780 3780 3600 2480 PCB-175 PCB-176 6550 U 9681 1070 3680 3680 3880 PCB-177 PCB-178 6550 U 9681 1070 3680 3680 3880 PCB-177 PCB-178 6550 U 9681 1070 3680 3680 3880 PCB-178 PCB-178 6550 U 9681 1070 3680 3680 3880 PCB-179 PCB-178 6550 U 9681 1070 3680 3890	PCB-161	PCB-161	12.6 U	15.3 U	15.5 U	15.3 U	10.2 U		
PCB-156 PCB-156 PCB-156 13.8 U 419 4270 4370 4370 1310 PCB-156 PCB-156 13.8 U 42.1 44.6 J 47 11.6 U PCB-157 PCB-157 PCB-157 PCB-157 PCB-157 PCB-157 PCB-157 PCB-157 PCB-157 PCB-159 PCB-150 PCB-157 PCB-171/173 PCB-177 PCB-178 PCB-162	PCB-162	48.3	380	456	432	225			
PCB-155 PCB-157	PCB-164	PCB-164		4190	4270	4370	1910		
PCB-167 PCB-167 610 3960 4400 4290 2260 PCB-169 PCB-169 12 U 17.1 U 33.8 U 133 U 3.8 U PCB-170 610 10600 I 11000 J 11200 J 650 PCB-171/173 287 3410 J 3750 J 3830 J 1830 PCB-172/173 287 3410 J 3750 J 3830 J 1830 PCB-174 PCB-172/173 287 3410 J 3750 J 3830 J 1830 PCB-176 PCB-177 207 1980 J 2770 J 2170 J 2270 J PCB-176 PCB-177 51.7 640 J 7734 J 687 J 2850 J 287 PCB-176 PCB-178 55.7 640 J 734 J 687 J 287 PCB-177 PCB-178 659 U 908 J 1070 J 1530 J 434 PCB-187 PCB-187 657 U 980 J 2570 J 5570 J 5550 J 1140 PCB-1880/93									
PCB-169 PCB-169 12 U 17.1 U 30.8 U 193 U 3.8 U PCB-170 610 100001 110001 112001 6550 PCB-171/173 PCB-171/173 287 34101 37501 36301 1130 PCB-172 PCB-172 207 19801 23701 21701 927 PCB-174 PCB-174 112 U 64801 77801 71601 2250 PCB-175 PCB-176 6.53 U 9081 10701 10001 2450 PCB-176 PCB-177 947 50701 58701 5001 144 PCB-177 PCB-177 947 50701 58701 55501 144 PCB-178 PCB-178 228 29801 32501 31001 777 PCB-179 PCB-187 6271 28001 33001 10001 777 PCB-181 PCB-181 2286 1221 2071 1941 102 PCB-181 PCB-180									
PCB-170 PCB-170 610 10600 J 11000 J 11200 J 6550 PCB-171/173 PCB-171/173 287 3410 J 3750 J 3630 J 1830 PCB-172 PCB-172 207 1980 J 22370 J 2170 J 927 PCB-174 PCB-175 51.7 6408 J 7780 J 150 J 2450 PCB-175 51.7 640 J 7784 J 687 J 2287 PCB-176 PCB-176 6.59 U 908 J 1070 J 1030 J 434 PCB-177 PCB-176 6.59 U 908 J 1070 J 1030 J 434 PCB-177 PCB-178 94.7 5070 J 5870 J 5550 J 1440 PCB-187 PCB-178 233 2980 J 3320 J 3300 J 727 PCB-180 J PCB-181 283 2980 J 3420 J 3300 J 1050 PCB-180 J PCB-181 283 2980 J 3420 J 3300 J 1050 PCB-180									
PCB-171/173 PCB-172 287 3410 J 3750 J 3630 J 1830 PCB-172 PCB-172 207 1980 J 2370 J 2170 J 927 PCB-174 PCB-174 112 U 6480 J 7780 J 7160 J 2450 PCB-175 PCB-176 PCB-177 51.7 640 J 774 J 687 J 287 PCB-176 PCB-176 6.59 U 908 J 1070 J 1030 J 434 PCB-177 PCB-177 94.7 5070 J 5870 J 300 J 727 PCB-178 PCB-179 6.27 U 2800 J 3250 J 3300 J 1060 PCB-178 PCB-179 6.27 U 2800 J 3420 J 3300 J 1060 PCB-189 PCB-189 2270 O 2880 J 3420 J 3300 J 1060 PCB-181 PCB-181 28.6 182 J 207 J 194 J 102 PCB-182 PCB-183 28.6 182 J 207 J 194 J 102									
PCB-172 PCB-174 PCB-174 112 U 6480 J 7780 J 7150 J 927 PCB-174 PCB-174 11.2 U 6480 J 7780 J 7160 J 2450 PCB-175 PCB-176 51.7 640 J 734 J 687 J 287 PCB-176 PCB-176 6.59 U 908 J 1070 J 1030 J 434 PCB-177 PCB-177 94.7 5070 J 5570 J 5550 J 1440 PCB-178 PCB-179 94.7 5070 J 5570 J 5550 J 1440 PCB-179 PCB-179 94.7 5070 J 3550 J 3100 J 727 PCB-179 PCB-179 6.27 U 2800 J 3420 J 3300 J 1060 PCB-189 PCB-189 J 2700 26800 J 30400 J 22800 J 15000 PCB-1818 PCB-1817 286 182 J 207 J 194 J 102 PCB-182 8.5 U 110 J 126 J 117 J 49.8									
PCB-174 PCB-175 PCB-175 S1.7 640 J 7780 J 7160 J 2450 PCB-175 PCB-175 S1.7 640 J 734 J 687 J 287 PCB-176 PCB-176 PCB-176 PCB-177 94.7 5070 J 5870 J 5550 J 1440 PCB-178 PCB-178 283 2980 J 3250 J 3100 J 727 PCB-179 PCB-179 6.27 U 2880 J 3420 J 3300 J 1060 PCB-180/193 PCB-181 28.6 182 J 207 J 194 J 102 PCB-181 PCB-181 28.6 182 J 207 J 194 J 102 PCB-182 PCB-181 28.6 182 J 207 J 194 J 102 PCB-183 /185 PCB-181 28.6 182 J 207 J 194 J 102 PCB-184 PCB-184 8.5 U 110 J 126 J 117 J 49.8 PCB-183 /185 PCB-183 /185 861 8850 J 11400 J		-							
PCB-175 PCB-176 PCB-176 65.9 U 988 J 1070 J 1030 J 434 PCB-176 PCB-177 94.7 5070 J 5870 J 5550 J 1440 PCB-178 PCB-178 283 2980 J 3250 J 3100 J 727 PCB-179 FCB-179 6.27 U 2800 J 3420 J 3300 J 1060 PCB-180/193 PCB-180/193 2700 26800 J 3420 J 3300 J 1060 PCB-180/193 PCB-180/193 2700 26800 J 36000 J 28500 J 15000 PCB-181 2.86 182 J 207 J 194 J 102 PCB-182 PCB-181 2.86 182 J 207 J 194 J 102 PCB-182 PCB-183 A 85 U 110 J 126 J 117 J 49.8 PCB-182 PCB-183 A 11.7 J 124 J 114 J 147 J 25.6 PCB-184 11.7 J 124 J 114 J 147 J 125 J									
PCB-176 PCB-177 94.7 5070 J 5870 J 5550 J 1434 PCB-178 PCB-177 94.7 5070 J 5870 J 5550 J 1440 PCB-178 PCB-178 283 2980 J 3250 J 3100 J 727 PCB-179 PCB-179 6.27 U 2800 J 3420 J 3300 J 1060 PCB-180/193 PCB-181 28.6 182 J 207 J 194 J 102 PCB-181 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-183/185 PCB-183/185 861 8850 J 11400 J 9640 J 4360 PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-185 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.165 U PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-186 6.76 U 4.13 U									
PCB-177 PCB-177 94.7 5070 J 5870 J 5550 J 1440 PCB-178 PCB-178 283 2980 J 3250 J 3100 J 727 PCB-179 PCB-179 6.27 U 2800 J 3420 J 3300 J 1060 PCB-180/193 PCB-180/193 2700 26800 J 3000 J 28500 J 15000 PCB-181 PCB-181 28.6 182 J 207 J 194 J 102 PCB-182 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-183/185 PCB-183/185 861 8850 J 11400 J 9640 J 4360 PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-186 PCB-186 6.76 U 4.13 U 8.99 UJ 4.67 U 1.65 U PCB-187 PCB-188 11 155.8 J 67.1 J 55.6 J 31.1 PCB-188 PCB-189 45.1 394 492 452 214 PCB-191									
PCB-178 PCB-179 CB-179 6.27 U 2800 J 3250 J 3100 J 727 PCB-189/193 PCB-180/193 2700 26800 J 30400 J 3300 J 1660 PCB-180/193 PCB-1818/185 2700 26800 J 30400 J 28500 J 15000 PCB-181 PCB-1811 28.6 182 J 207 J 194 J 102 PCB-182 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-183/185 PCB-184 11.7 J 124 J 1140 J 9640 J 4360 PCB-184 PCB-186 6.76 U 4.13 U 8.93 U 4.67 U 1.65 U PCB-186 PCB-186 6.76 U 4.13 U 8.93 U 4.67 U 1.65 U PCB-187 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-199 PCB-199 152 1680 J 1800 J 1780 J <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>									
PCB-179 PCB-179 6.27 U 2800 J 3420 J 3300 J 1060 PCB-180/193 PCB-180/193 2700 26800 J 30400 J 28500 J 15000 PCB-181 PCB-181 28.6 182 J 207 J 194 J 102 PCB-182 PCB-182 8.5 U 110 J 125 J 117 J 49.8 PCB-183/185 PCB-183/185 861 8850 J 11400 J 9640 J 4360 PCB-184 PCB-187 361 8850 J 11400 J 9640 J 4360 PCB-186 PCB-187 861 8850 J 11400 J 9640 J 4360 PCB-186 PCB-186 6.76 U 4.13 U 8.93 U 4.67 U 1.65 U PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-199 PCB-189									
PCB-180/193 PCB-180/193 2700 26800 J 30400 J 28500 J 15000 PCB-181 PCB-181 28.6 182 J 207 J 194 J 102 PCB-182 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-183/185 PCB-183/185 86 I 8850 J 11400 J 9640 J 4560 PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-186 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.65 U PCB-187 PCB-188 11 1800 J 15700 J 15600 J 7640 PCB-188 11 1800 J 15700 J 15600 J 7640 PCB-188 11 15.8 J 67.1 J 55.6 J 31.1 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 51.1 J 51.2 J 569 J <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>									
PCB-181 PCB-182 28.6 182 J 207 J 194 J 102 PCB-182 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-183/185 PCB-183/185 861 8850 J 11400 J 9640 J 4360 PCB-184 PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.65 U PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 51.1 J 51.2 J 569 J 540 J 282 PCB-192 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 282 4230 4500 4560 1790	PCB-179	PCB-179		2800 J		3300 J			
PCB-182 PCB-182 8.5 U 110 J 126 J 117 J 49.8 PCB-183/185 PCB-183/185 861 8850 J 11400 J 9640 J 4360 PCB-184 PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-186 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.65 U PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-198 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 8.91 U 8.62 U 162 UJ 7.58 U 5.31 PCB-194 282 4230 4500 4560 1790 PCB-195 73.2 J 1860 2170 19	PCB-180/193	PCB-180/193	2700	26800 J	30400 J	28500 J	15000		
PCB-183/185 PCB-183/185 861 8850 J 11400 J 9640 J 4360 PCB-184 PCB-184 11.7 J 124 J 114 J 147 J 25.6 PCB-186 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.65 U PCB-187 PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-194 282 4230 4500 4560 1790 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 26	PCB-181	PCB-181	28.6	182 J	207 J	194 J	102		
PCB-184 PCB-186 11.7 J 124 J 114 J 147 J 25.6 PCB-186 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.65 U PCB-187 PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-193 891 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-198/199 505 6260 7880 6570 2490 PCB-2	PCB-182	PCB-182	8.5 U	110 J	126 J	117 J	49.8		
PCB-186 PCB-186 6.76 U 4.13 U 8.93 UJ 4.67 U 1.65 U PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-193 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-198/199 505 6260 7880 6570 2490 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490	PCB-183/185	PCB-183/185	861	8850 J	11400 J	9640 J	4360		
PCB-187 1340 14800 J 15700 J 15600 J 7640 PCB-188 PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-192 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201	PCB-184	PCB-184	11.7 J	124 J	114 J	147 J	25.6		
PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-193 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433	PCB-186	PCB-186	6.76 U	4.13 U	8.93 UJ	4.67 U	1.65 U		
PCB-188 11 55.8 J 67.1 J 55.6 J 31.1 PCB-189 PCB-189 45.1 394 492 452 214 PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-193 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433	PCB-187	PCB-187	1340	14800 J	15700 J	15600 J	7640		
PCB-189 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-192 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-190 PCB-190 152 1680 J 1800 J 1780 J 1290 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-192 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-191 PCB-191 51.1 J 512 J 569 J 540 J 282 PCB-192 PCB-192 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-192 PCB-192 8.91 U 8.62 U 16.2 UJ 7.58 U 5.31 PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-194 PCB-194 282 4230 4500 4560 1790 PCB-195 PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-195 73.2 J 1860 2170 1940 987 PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-196 PCB-196 261 3070 3760 3250 1600 PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-197/200 PCB-197/200 37.6 808 1180 885 390 PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-198/199 PCB-198/199 505 6260 7880 6570 2490 PCB-201 PCB-201 72.2 1040 1390 1110 433									
PCB-201 PCB-201 72.2 1040 1390 1110 433		-							
PCB-202 PCB-202 148 1780 2120 1930 477									
	PCB-202	PCB-202	148	1780	2120	1930	477		

					Tissue	Analytical Results				
	Location ID	Field QC	Field QC	PB002.1	PB003.1	PB005.1	PB006.1	PB008.2	PB010	PB010.1
	Sample ID	EB-110621-A01	EB-110624-A01	PB01-I-A-BCR-W-001-COMP-201106	PB01-F-A-STM-W-006-20110618	PB01-I-A-BCR-W-002-COMP-201106	PB01-I-B-WHS-W-002-COMP-201106	PB01-I-B-WHS-W-001-COMP-201106	PB01-F-A-GM-W-008-20110618	PB01-I-A-WHS-W-001-COMP-201100
	Sample Date	6/24/2011	6/24/2011	6/20/2011	6/18/2011	6/20/2011	6/20/2011	6/19/2011	6/18/2011	6/20/2011
	Sample Type	EB	EB	N N	N	N N	N	N	N	N N
	y y	25		3202691	3202445.49	3202312	3202026.63	3202032	3201710.87	3201704.8
	v			13836499	13836263.53	13836201	13836356.8	13836072	13836029.16	13836071.32
	Scientific_Name			Callinectes sapidus	Mugil cephalus	Callinectes sapidus	Penaeus setiferus	Penaeus setiferus	Brevoortia patronus	Penaeus setiferus
	_			·	• .	·			·	
	Size_Class			>2 to =<7.5	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13	=<15	>2 to =<7.5
	Tissue_Category			Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
Chemical	Tissue_Type CAS Number	Result	Result	Whole Body Result	Whole Body Result	Whole Body Result	Whole Body Result	Whole Body Result	Whole Body Result	Whole Body Result
PCB-203	PCB-203		823	441	721	464	704	550	1970	598
PCB-204	PCB-203		7.62	12.9	19.9	18.7	18.2	11.5 J	50.4	11.5 U
PCB-205	PCB-205		50.9	20.9 J	77.7	27.3	55.4	38.8	210	517 U
PCB-206	PCB-206		1690	503	848	593	623	473	2910	654
PCB-207	PCB-207		301	443	579	589	406	276	2300	440
PCB-208	PCB-208		1080	647	897	811	607	460	3330	600
PCB-209	PCB-209		2190	14200	41000	18300	16900	13000	107000	17300
Total Decachlorobiphenyl homologs (U = 1/2)	tSum_DecPCB_N		2190	14200	41000	18300	16900	13000	107000	17300
Total Dichlorobiphenyl homologs (U = 1/2)	tSum_DiPCB_N		164.63 J	464.32 J	5661.9 J	509.89 J	3773.2 J	1958.2 J	4070.8 J	2716.6 J
Total Heptachlorobiphenyl homologs (U = 1/2)	tSum_HepPCB_N		19841.61 J	12146.74 J	19540.87 J	12006.29 J	23448.98	18096.45	45398.18 J	24954.87 J
Total Hexachlorobiphenyl homologs (U = 1/2)	tSum_HexPCB_N		51122.17 J	36659.94 J	56270.44 J	41035 J	64768.2 J	47612.03 J	106457.76 J	64616.74 J
Total Monochlorobiphenyl homologs (U = 1/2)	tSum_MonPCB_N		6.05 UJ	1.86 U	146.81	6.33 U	177.5 J	45.05 J	125.07 J	146.1 J
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum_NonPCB_N		3071	1593	2324	1993	1636	1209	8540	1694
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N		6472.52	3359.8 J	5174.6	3581	4830.6	3642.6 J	13794.4	4647.3
Total PCB Congener (U = 1/2)	tPCBCong_N		166151.68 J	228676.53 J	596176.03 J	324017.32 J	918637.82 J	547939.37 J	1259030.88 J	837229.96 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N		163961.68 J	214476.53 J	555176.03 J	305717.32 J	901737.82 J	534939.37 J	1152030.88 J	819929.96 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98_N		17.5 J	97.57 J	103.7 J	147.29	312.2 J	177.4	197.09 J	277.3 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N		0.2 J	0.72 J	0.6 J	1.01	1.7 J	1	1.24 J	1.7 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	tPCBCongCPM_N		2.8 J	8.35 J	5.7 J	11.28	16.7 J	9.3	12.88 J	18.1 J
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum_PenPCB_N		44587.6 J	60624.15 J	121974.4 J	83162.72 J	217689.09 J	147236.43 J	253722.6 J	203719.17 J
Total Tetrachlorobiphenyl homologs (U = 1/2)	tSum_TetPCB_N		33512.9 J	78996.76 J	266157.14 J	129201.4 J	446355.2 J	244379.72	564484.6 J	406720.8
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N		5181.37 J	20629.13 J	77925.88 J	34223.48	139059.04 J	70759.94	155437.47 J	110714.38
Dioxin Furans (ng/kg)										
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0		0.301 U	15	747	9.24	21.6	31.1	738	25.2 J
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)			7.23	9.24	69.6	10.2	8.74	9.66	96.9	10.9
1 1 1	67562-39-4		0.103 U	1.63 J	22.6	1.64 J	2.73	4.1	29.7	3.08 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)			0.738 J	1.43 U	6.64	1.95 U	1.57 J	1.62 J	9.26	1 U
	55673-89-7		0.738 J 0.173 U	0.291 U	2.89 J	0.348 U	0.649 U	0.783 U	5.05	0.705 U
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9		0.112 U	0.532 U	1.88 J	0.819 U	1.87 J	1.38 J	2.23 J	1.66 J
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6		0.112 U	0.829 U	0.227 U	1.11 U	0.505 U	0.381 U	0.308 U	0.526 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9		0.101 U	0.829 U	0.909 J	0.365 U	0.303 U 0.772 J	0.569 J	1.11 J	0.320 U
	57653-85-7		0.112 U		0.909 J 0.236 U	0.303 U 1.1 U	0.772 J 0.505 U	0.309 J 0.425 U	0.547 J	0.493 U 0.567 U
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9			0.783 U	0.236 U	0.209 U	0.505 U	0.423 U	0.324 U	0.567 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3		0.168 U	0.83 U	0.235 U	1.14 U	0.513 U	0.41 U	0.324 U	0.556 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6		0.357 U	1.01 U	1.73 J	1.46 U	4.55	3.63 J	3.41	4.65
	40321-76-4		0.405 J	0.252 U	0.174 U	0.379 U	0.253 U	0.337 U	0.673 J	0.48 U
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5		0.134 U	0.136 U	0.739 J	0.174 U	0.426 J	0.514 U	0.728 J	0.622 U
	57117-31-4		0.484 U	1.14 U	2.06 J	1.5 J	5.05	3.5	3.1	4.07
<u> </u>	51207-31-9		3.73	6.8	59.2	7.89	15.5	12	9.79	13.7
	1746-01-6		0.579	3.68	15.5	3.67	3.64 J	4.34	7.15	3.36
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4		0.841 U	7.66 J	7.53 J	10.1	23.5	17.3 J	8.51	21.8 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9		0.405 J	0.252 U	0.174 U	0.379 U	2.74 J	1.75 J	0.673 J	1.5 J
	tDioxFurB_N		5.016 J	11.366 J	77.754 J	13.512 J	25.179 J	20.709 J	21.947 J	22.169 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurF_N		1.371 J	4.745 J	20.438 J	5.429 J	7.792 J	7.445 J	10.98 J	6.98 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	tDioxFurM_N	-	1.497 J	4.879 J	23.151 J	5.4 J	7.431 J	7.257 J	11.034 J	6.715 J

			Tissue Ana	alytical Results				
Location	ID PB010.3	PB013.1_A	PB013.3	PB013.3	PB015.1	PB015.2	PB015_A	PB015_A
Sample	ID PB01-I-A-BCR-W-003-COMP-201106	_	PB01-F-A-GKF-W-004-COMP-201106	PB01-F-A-STM-W-003-20110616	PB01-F-A-GKF-W-002-COMP-20110	6 PB01-I-A-BCR-W-004-COMP-20110	_	- 06 PB01-I-A-BCR-W-006-COMP-201106
Sample Da	te 6/19/2011	6/22/2011	6/21/2011	6/15/2011	6/15/2011	6/16/2011	6/25/2011	6/22/2011
Sample Ty		N	N N	N	N	N	N	N
	X 3201743	3201306.47	3201433.88	3201433.88	3201175	3201180	3201168.73	3201168.73
	Y 13836152	13836191.23	13836075.12	13836075.12	13836251	13836183	13836254.66	13836254.66
Scientific Nar		Callinectes sapidus	Fundulus grandis	Mugil cephalus	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Callinectes sapidus
Size_Cla	· ·	>2 to =<7.5	=<15	=<15	=<15	>2 to =<7.5	=<15	>2 to =<7.5
Tissue_Catego		Sedentary	Sedentary	Migratory	Sedentary	Sedentary	Sedentary	Sedentary
Tissue_Ty		Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203 PCB-203	1300	431	3550	2270	2740	1220	4710	1070
PCB-204 PCB-204	33.7	15.9	116	51.8	73.4	33.6	147	26.3
PCB-205 PCB-205	61.7	16.6	232	221	181	50.9	293	47.8
PCB-206 PCB-206	1220	492	3280	2630	2470	1190	3890	927
PCB-207 PCB-207	1090	399	4990	1960	2990	1060	5670	856
PCB-208 PCB-208	1270	559	5760	2530	3720	1350	7080	1100
PCB-209 PCB-209	25800	12200	115000	92100	70800	31100	124000	26000
Total Decachlorobiphenyl homologs (U = 1/2) tSum_DecPCB_N	1 25800	12200	115000	92100	70800	31100	124000	26000
Total Dichlorobiphenyl homologs (U = 1/2) tSum DiPCB N	1297.88	1525.67 J	12709.06 J	6329.8 J	12808.11 J	1657.66 J	14345.09 J	1381.61 J
Total Heptachlorobiphenyl homologs (U = 1/2) tSum_HepPCB_1		14491.232 J	108539.02 J	63014.52 J	84207.89 J	29236.81 J	151125.74 J	28294.91 J
Total Hexachlorobiphenyl homologs (U = 1/2) tSum HexPCB N		43064.795 J	306112.46	148296.3 J	268417.13 J	92122.29 J	469150.14 J	69854.95 J
Total Monochlorobiphenyl homologs (U = 1/2) tSum MonPCB		3.322 J	248 J	150.3 J	297.4 J	5.82 U	235.32 J	5.93 U
Total Nonachlorobiphenyl homologs (U = 1/2) tSum NonPCB I		1450	14030	7120	9180	3600	16640	2883
Total Octachlorobiphenyl homologs (U = 1/2) tSum_OctPCB_N		3281.5	27828	15481.8	21014.4	7885.5	39370	7368.1
Total PCB Congener (U = 1/2) tPCBCong_N	680864.4 J	457647.511 J	3781237.62 J	1793318.68 J	3359203.87 J	763874.14 J	5163120.59 J	560775.99 J
Total PCB Congener minus 209 (U = 1/2) tPCBCong209_N	655064.4 J	445447.511 J	3666237.62 J	1701218.68 J	3288403.87 J	732774.14 J	5039120.59 J	534775.99 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2) tPCBCngCPB98_		260.92	602.3	234.7 J	526.2 J	362	728.24894 J	294.5
Total PCB Congener TEQ 1998 (Fish) (U = 1/2) tPCBCngCPF98 L		1.5	5	1.6 J	4 J	2.4	6.77302 J	1.8
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2) tPCBCongCPM_1		14.97	61	16.4 J	45.4 J	25.5	88.92282 J	19.4
Total Pentachlorobiphenyl homologs (U = 1/2) tSum_PenPCB_N		106633.234 J	896572.17	368821.4 J	770680.18 J	199620.57 J	1242471.9 J	130929.55 J
Total Tetrachlorobiphenyl homologs (U = 1/2) tSum_TetPCB_N	250552.35 J	193569.33	1789095.8	853044.2	1694080.5 J	307173.83 J	2444027.5 J	222955.13 J
Total Trichlorobiphenyl homologs (U = 1/2) tSum_TriPCB_N	64119.63	81428.44	511103.2	238960.3 J	427718.2	91471.43 J	661754.9	71100.21 J
Dioxin Furans (ng/kg)	04113.03	01420.44	311103.2	230300.3 3	427710.2	31471.43 3	001734.3	71100.21 7
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) 39001-02-0	15.3	9.46	7.5	671	3.34 J	22	3.42 J	22.8
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD) 3268-87-9	46.4	14.5	5.13	40.6	9.73	50.6	6.88	7.65
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) 67562-39-4	2.09 J	2.51	1.72 J	18.2	0.87 J	7.16	1.24 J	5.44
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD) 35822-46-9	2.71 U	1.96 J	1.26 J	4.44 J	1.04 J	5.08	1.4 J	1.38 J
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) 55673-89-7	0.462 U	0.206 U	0.208 U	3.28	0.292 U	0.447 J	0.417 U	0.393 U
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) 70648-26-9	1.38 U	1.51 J	0.469 J	1.35 J	0.292 U	2.48	0.789 J	2.3 J
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD) 39227-28-6	1.38 U	0.156 U	0.403 J 0.13 U	0.271 U	0.173 U	0.354 U	0.348 J	0.275 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) 57117-44-9	0.428 U	0.633 J	0.41 J	0.271 U	0.202 J	0.873 U	0.461 J	1.13 J
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD) 57653-85-7	1.17 U	0.178 U	0.41 J	0.289 U	0.202 J 0.182 U	0.436 J	0.569 J	0.5 J
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) 72918-21-9	0.271 U	0.169 U	0.174 U	0.553 U	0.259 U	0.332 U	0.134 U	0.44 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD) 19408-74-3	1.22 U	0.169 U	0.14 U	0.285 U	0.18 U	0.368 U	0.185 J	0.288 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) 57117-41-6	2.61	2.5	1.77 J	2.15 J	1.27 J	2.65	2.88	6.65
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD) 40321-76-4	0.488 U	0.14 U	0.886 J	0.501 J	0.137 U	0.323 J	2.37 J	0.411 J
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) 60851-34-5	0.488 U	0.294 J	0.128 U	0.523 U	0.137 U	0.383 U	0.273 U	0.431 J
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) 57117-31-4	2.57 J	2.35 J	2.66	2.97	2.21 J	2.64	3.46	6.19
2,3,7,8-Tetrachlorodibenzofuran (TCDF) 51207-31-9	10.7	10.4	11.9	15.9	9.96	9.13 J	13.5	33
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) 1746-01-6	3.59	2.42	5.82	7.33	5.42	4.02	8.53	4
Total Pentachlorodibenzofuran (PeCDF) 30402-15-4	3.59 14 J	2.42 14.4 J	5.62 5.5 J	7.55 8.51 J	4.39 J	16.7 J	9.49 J	27.4 J
Total Pentachlorodibenzo-p-dioxin (PeCDD) 36088-22-9	0.488 U	2.15 J	0.886 J	0.501 J	0.137 U	2.33 J	3.35 J	27.4 J
	0.488 U 17.608 J	2.15 J 15.786 J	0.886 J 21.58 J	0.501 J 27.479 J	17.894 J	2.33 J 16.825 J	28.352 J	44.761 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2) tDioxFurB_N					7.294 J			
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2) tDioxFurF_N	6.245 J	4.633 J	8.88 J	10.83 J		6.763 J	13.793 J	10.033 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2) tDioxFurM_N	6.104 J	4.64 J	8.932 J	11.142 J	7.318 J	6.683 J	13.662 J	10.333 J

				Tissue Ana	alytical Results				
	Location ID	PB015_A	PB015_A	PB015_A	PB016.2	PB019.1	PB019.1	PB019.1	PB019.1
	Sample ID	PB01-I-A-BCR-W-007-COMP-201106	PB01-I-A-BCR-W-009-COMP-201106	PB01-I-B-BRS-W-001-COMP-201106	PB01-F-A-GKF-W-003-COMP-201106	PB01-F-A-SAS-W-005-20110622	PB01-I-B-BRS-W-012-20110621	PB01-I-B-BRS-W-014-20110621	PB01-I-B-BRS-W-015-20110621
	Sample Date	6/22/2011	6/22/2011	6/20/2011	6/15/2011	6/22/2011	6/21/2011	6/21/2011	6/21/2011
	Sample Type	N	N	N	N	N	N	N	N
		3201168.73	3201168.73	3201168.73	3201190	3201152.87	3201152.87	3201152.87	3201152.87
	Υ	13836254.66	13836254.66	13836254.66	13836042	13835753.58	13835753.58	13835753.58	13835753.58
	Scientific_Name	Callinectes sapidus	Callinectes sapidus	Penaeus aztecus	Fundulus grandis	Cynoscion arenarius	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus
	Size_Class	>2 to =<7.5	>2 to =<7.5	>7.5 to =<13	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13
	Tissue_Category	Sedentary	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203	PCB-203	827	1240	931	1480	3030	1350	1390	1220
PCB-204	PCB-204	20.3	25.5	24.5	43.7	63.6	62.7 U	44.3	36.4 J
PCB-205	PCB-205	36.5	48.6	89.4	96.4	207 U	140	108 J	108 J
PCB-206	PCB-206	663	921	1010	1370	4250	1350	1270	1220
PCB-207	PCB-207	554	802	657	1910	3670	889	857	844
PCB-208	PCB-208	733	990	873	2180	4500	1200	1180	1030
PCB-209	PCB-209	15800	22000	26800	44400	112000	30300	29200	29300
Total Decachlorobiphenyl homologs (U = 1/2)	tSum_DecPCB_N	15800	22000	26800	44400	112000	30300	29200	29300
Total Dichlorobiphenyl homologs (U = 1/2)	tSum_DiPCB_N	1499.97 J	2657.3 J	4353.8 J	5656.24 J	4016.84 J	8913.6 J	7238 J	6032.9
	tSum_Bir CB_N	21819.781 J	32368.85 J	36768.58 J	52421.23	50466.24 J	49215 J	51394.4 J	44019 J
	tSum HexPCB N	55759.991 J	75818.36 J	93633.73	161910.13	123693.11 J	184786 J	137480.3 J	133081.5 J
1 / 0 (/ / /	tSum MonPCB N	4.06 U	4.08 U	92.26 J	140.3	267.1 J	172.7	180.8 J	157.1 J
. , ,	tSum_NonPCB_N	1950	2713	2540	5460	12420	3439	3307	3094
. , ,	tSum_NonFCB_N	5588.8	8391.1	7015.9	11480.1	17937.1	9623.4	9890.3 J	9146.4 J
Total PCB Congener (U = 1/2)	tPCBCong_N	487928.797 J	674944.25 J	1167288.08 J	1964621.79 J	1448892.45 J	2092123 J	1730559.8 J	1552684.1 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	472128.797 J	652944.24 J	1140488.08 J	1920221.8 J	1336892.45 J	2061822.9 J	1730339.8 J 1701359.8 J	1523384.1 J
	tPCBCngCPB98_N	312.4 J	434.33 J	433.2 J	270.7	77.5 J	586.73896 J	559.5 J	504.6 J
	tPCBCngCPF98_N	2.5 J	434.33 J 2.45 J	2.4 J	2.2	0.64 J	2.57243 J	2 J	2.5 J
	tPCBCongCPM_N	32.5 J	23.74 J	23.2 J	25.4	3.63 J	20.15058 J	8.2 J	22.5 J
		125501.57 J	170559.12 J	279032.49 J	455025.05	310793.24 J	464624	383011.3 J	352214.6 J
	tSum_PenPCB_N	190602.49 J	260096.18 J	538309.6	1015345.92	656383.8 J	1078482.5 J	858453.7	764013.2 J
	tSum_TetPCB_N	69400.62 J	100334.51 J	178741.77 J	212782.83 J	160915.02	262566.8 J	250404 J	211625.6 J
	tSum_TriPCB_N	69400.62 J	100334.51 J	1/8/41.// J	212/02.03 J	160915.02	202500.8 J	250404 J	211623.0 J
Dioxin Furans (ng/kg) 1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	20001 02 0	12.0	15.2	124	9.66	5.39			
	39001-02-0	12.8	15.2	124					
	3268-87-9	9.09 2.39	8.36 2.71	10.7 5.65	3.18 J 0.762 J	2.81 J			
	67562-39-4		_			0.469 J			
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)		1.25 J	1.37 J	0.799 U	0.581 J	0.662 J	-		
	55673-89-7	0.48 U	0.401 U	1.11 U	0.209 U	0.24 U	-		
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9	1.23 J	1.56 J	2.41 J	0.14 U	0.221 J			
	39227-28-6	0.347 U	0.535 J	0.55 U	0.157 U	0.136 U	-		
	57117-44-9	0.665 J	0.916 J	0.978 J	0.137 U	0.155 J			
	57653-85-7	0.366 U	0.376 U	0.567 U	0.167 U	0.145 U			
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9	0.196 U	0.414 U	0.936 U	0.209 U	0.165 U			
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3	0.363 U	0.378 U	0.567 U	0.165 U	0.143 U			
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6	2.75	4.11	5.89	0.786 J	0.871 J			
	40321-76-4	0.42 J	0.291 U	0.395 U	0.116 U	0.108 U			
	60851-34-5	0.189 U	0.329 U	0.712 U	0.148 U	0.147 U			
	57117-31-4	2.71	4.18	6.77	1.31 J	0.216 J			
	51207-31-9	12.4	17.8	21.1	6.08	0.877			
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1746-01-6	3.23	4.75	4.54	3.46	0.849			
, ,	30402-15-4	15 J	21.9 J	33 J	2.58 J	1.25 J			
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9	3.78 J	3.63 J	5.04 J	0.116 U	0.108 U			
	tDioxFurB_N	19.302 J	27.652 J	33.739 J	11.042 J	2.155 J			
. , , , , , ,	tDioxFurF_N	6.091 J	8.67 J	10.112 J	4.599 J	1.194 J			
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	tDioxFurM_N	6.093 J	8.479 J	9.667 J	4.617 J	1.171 J			

				lissue An	alytical Results				
	Location ID	PB019.1	PB019.1	PB019.1	PB019.2	PB019 A	PB019_A	PB024.1	PB024.1
	Sample ID		PB01-I-B-BRS-W-022-20110622	PB01-I-B-BRS-W-023-20110622		6 PB01-F-A-GKF-W-005-COMP-201106	- I		PB01-F-A-GM-W-002-20110615
	Sample Date	6/21/2011	6/22/2011	6/22/2011	6/28/2011	6/21/2011	6/23/2011	6/15/2011	6/15/2011
	Sample Type	N	N	N	N	N	N N	N	N
	y x	3201152.87	3201152.87	3201152.87	3201194.8	3201314.96	3201314.96	3201090.96	3201090.96
	v	13835753.58	13835753.58	13835753.58	13835735.1	13835750.59	13835750.59	13835160.37	13835160.37
	Scientific_Name	Penaeus aztecus	Penaeus aztecus	Penaeus aztecus	Crassostrea virginica	Fundulus grandis	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus
	Size_Class	>7.5 to =<13	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15
	Tissue_Category	Migratory		Migratory	Sedentary	Sedentary	Sedentary	Migratory	Migratory
	Tissue_Category	Whole Body	Migratory Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203	PCB-203	1580	1620	1580	99.2	3640	1480	2670	2330
PCB-204	PCB-204	33.4 U	38	36	3.12 J	180	140	58	50.5
PCB-205	PCB-205	134 J	116	109	7.42 J	262	92.8	280	248
PCB-205	PCB-206	1590	1320	1370	34.2	4510	2450	4060	3760
		1090	868	948	32.7	6450	2260	3100	3150
PCB-207 PCB-208	PCB-207 PCB-208	1390	1050	1200	32.7 45.9	7020	3130	4290	4080
PCB-208 PCB-209	PCB-208 PCB-209	44800	29900	31300			80800	152000	138000
		44800	29900	31300	911	172000	80800		138000
Total Decachlorobiphenyl homologs (U = 1/2)	tSum_DecPCB_N				911	172000		152000	
Total Dichlorobiphenyl homologs (U = 1/2)	tSum_DiPCB_N	7431.8 J	7361.52	8065.8 J	6302.64 J	11955.7 J	1784.88 J	6048.3 J	5811.1 J
Total Heptachlorobiphenyl homologs (U = 1/2)	tSum_HepPCB_N	61474.2 J	59861.54	57840.74	14636.5 J	113501.99 J	31102.38 J	67105.53 J	56249.8
Total Hexachlorobiphenyl homologs (U = 1/2)	tSum_HexPCB_N	222355.5 J	172225.81 J	145866.42 J	89369.8 J	310336.34 J	89767.5 J	167032.26	136616.3 J
Total Monochlorobiphenyl homologs (U = 1/2)	tSum_MonPCB_N	115 J	506.2 J	723.1 J	72.73	315.4 J	4.02 U	174.5	137.72
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum_NonPCB_N	4070	3238	3518	112.8	17980	7840	11450	10990
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N	11636.7 J	10849	10860	1458.54 J	30252	11669.8	18595	16563.5
Total PCB Congener (U = 1/2)	tPCBCong_N	2192371.5 J	2031988.89 J	1964859.31 J	1429439.04 J	3824700.99 J	695219.28 J	1896422.09 J	1735872.21 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	2147571.5 J	2002088.89 J	1933559.31 J	1428528.04 J	3652700.99 J	614419.28 J	1744422.09 J	1597872.21 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98_N	694.7385 J	567.57 J	544.7	361.21	577.2 J	331.4 J	267.5	241.5 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N	3.903825 J	3.18 J	3.1	2.05	4.8 J	2 J	1.8	1.6 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2		41.1819 J	30.43 J	31	20.74	58.6 J	20 J	19.6	16.6 J
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum_PenPCB_N	509308.6 J	469281.01 J	432038.7 J	350206 J	860048.76 J	149413.98 J	388288.9	341433.6 J
Total Tetrachlorobiphenyl homologs (U = 1/2)	tSum_TetPCB_N	1017143.8 J	1003156.4	989340.5 J	731573.8	1882811.6	237348.65 J	841480.9	798314.6
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N	314036 J	275609.3 J	285306 J	234795.18 J	425499.16	85487.44 J	244246.6 J	231755.6 J
Dioxin Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0				10.9	6.12	49.2	1350	1500
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD	,				6.89	2.56 J	6.16	253	293
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	67562-39-4				0.794 J	0.836 J	7.68	49	54.8
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCD					0.988 J	0.564 J	0.808 J	25.9	30.6
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	55673-89-7				0.454 U	0.199 U	0.421 U	9.15	13.4
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9				0.257 U	0.394 J	1.79 J	5.14	6.41
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6				0.328 U	0.15 U	0.301 U	0.331 U	0.362 J
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9				0.251 U	0.118 U	0.934 J	2.46 J	2.94
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	57653-85-7				0.359 U	0.152 U	0.329 U	1.35 J	1.59 J
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9				0.425 U	0.177 U	0.395 U	1.05 J	1.39 J
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3				0.349 U	0.154 U	0.32 U	0.875 J	0.981 J
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6				1.05 J	1.22 J	3.22	5.34	5.86
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	40321-76-4				0.325 U	0.495 J	1.12 J	0.897 J	0.931 J
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5				0.361 U	0.165 U	0.362 U	1.81 J	2.2 J
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	57117-31-4				2.62	2.48	3.53	5.49	6.39
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	51207-31-9				14.5	10.4	13.6	14.8	14.4
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1746-01-6				2.36	5.32	3.42	9.31	10.6
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4				12.4 J	5.02 J	21.8 J	18.1 J	21 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9				3.27 J	0.495 J	4.16 J	2.58 J	3.48 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	tDioxFurB_N				19.853 J	18.902 J	22.413 J	32.954 J	35.225 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurF_N				4.773 J	7.748 J	7.62 J	15.878 J	18.132 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	tDioxFurM_N				4.932 J	7.738 J	7.517 J	16.101 J	18.177 J

				Tissue An	alytical Results				
	Location ID	PB024.4	PB025_A	PB026.1	PB032 A	PB037.2	PB037.4	PB037.5	PB037.6
	Sample ID	PB01-I-A-BCR-W-005-COMP-201106	PB01-F-A-GM-W-004-20110615	PB01-F-A-STM-W-002-20110616	PB02-F-A-PNF-W-009-20110617	PB02-F-A-STM-W-006-20110616	PB02-I-B-BCR-W-001-COMP-201106	PB02-I-B-BRS-W-001-COMP-201106	PB02-I-A-BCR-W-002-COMP-201106
	Sample Date	6/20/2011	6/15/2011	6/15/2011	6/17/2011	6/16/2011	6/17/2011	6/17/2011	6/16/2011
	Sample Type		N	N	N	N	N	N	N
		3201319	3201214.42	3201153.65	3201407.03	3201346.97	3201504	3201380	3201402
	Υ	13835188	13835154.42	13835079.21	13834542.25	13834038.02	13833530	13834151	13833775
	Scientific_Name		Brevoortia patronus	Mugil cephalus	Lagodon rhomboides	Mugil cephalus	Callinectes sapidus	Penaeus aztecus	Callinectes sapidus
	Size_Class		=<15	=<15	=<15	=<15	>7.5 to =<13	>7.5 to =<13	>2 to =<7.5
	Tissue_Category	Sedentary	Migratory	Migratory	Migratory	Migratory	Migratory	Migratory	Sedentary
	Tissue_Type		Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203	PCB-203	1550	3380	2360	2410	2440	1850	704	980
PCB-204	PCB-204	157	66.8	35.9	71.7	45.3	23.4	14	13.3 J
PCB-205	PCB-205	141	325	216	269	230	57.1 U	54.3	32.4 J
PCB-206	PCB-206	3440	4410	2040 J	3450	2260	988	623	709
PCB-207	PCB-207	3240	3260	1240	2720	1460	827	343	585
PCB-208	PCB-208	4920	4540	1620 J	2830	2020	1010	483	694
PCB-209	PCB-209	175000	143000	44600	92300	46900	22200	14200	14900
Total Decachlorobiphenyl homologs (U = 1/2)	tSum DecPCB N	175000	143000	44600	92300	46900	22200	14200	14900
Total Dichlorobiphenyl homologs (U = 1/2)	tSum DiPCB N	2599.9 J	9101.47 J	7557.6 J	3536 J	11446.8 J	1788.19 J	4412.14	1620.78
Total Heptachlorobiphenyl homologs (U = 1/2)	tSum_HepPCB_N	39598.23	87466.45 J	68247.82	58965.5	74449.73	53599.63 J	25701.15 J	25438.15 J
Total Hexachlorobiphenyl homologs (U = 1/2)	tSum HexPCB N	95880.92 J	228773.4 J	172478.2 J	162859.6 J	211230.8 J	127352.43 J	75598.83 J	79552.68 J
Total Monochlorobiphenyl homologs (U = 1/2)	tSum MonPCB N	7.83 U	199.3	183.7 J	150.81 J	247.18 J	3.17 U	107.19	2.25 U
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum NonPCB N	11600	12210	4900 J	9000	5740	2825	1449	1988
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N	13691	23255.8	15699.9	15052.7	16748.3	11921.9	4747.3	6059.7 J
Total PCB Congener (U = 1/2)	tPCBCong_N	864807.31 J	2862773.47 J	2310204.97 J	2113550.66 J	3115985.51 J	935995.24 J	996652.52 J	626864.72 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	689807.31 J	2719773.47 J	2265604.97 J	2021250.66 J	3069085.51 J	913795.24 J	982452.52 J	611964.73 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98 N	307.6 J	426.7	321.5	89.6 J	453.3	509.3 J	369.86 J	340.8
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N	1.9 J	2.9	2.4	1.8 J	2.8	3.8 J	2.18 J	2.3
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)		19 J	31.6	24.8	24 J	26.7	43.1 J	22.01 J	24.8
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum PenPCB N	167042.88 J	618539.7 J	505521.4 J	490407 J	660680.8	278403.36 J	246013.67 J	174124.76 J
Total Tetrachlorobiphenyl homologs (U = $1/2$)	tSum_TetPCB_N	270156.8 J	1352624.2	1153568.1	1059613	1639047.1 J	354626.08 J	465520.07 J	249875.45 J
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N	89226.24 J	387603.2 J	337448.2	221666 J	449494.8 J	83274.04 J	158903.19 J	73302.11 J
Dioxin Furans (ng/kg)	todin_iiii cb_iv	05220.24 3	367003.2 3	337440.2	221000 3	445454.0 3	03274.04 3	150505.15 7	75502.11 7
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0	184	944	327	169	108	7.57	21.4	8.29
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDI)		13.5 J	207	41.7	15	26	9.16	12.2	20.7
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	67562-39-4	13.7	35.1	9.7	5.41	3.84 J	3.4	2.77 J	1.71 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDI)		2.22 U	21.2	4.42	1.24 J	2.78 J	2.1 J	2.38 J	4.48
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	55673-89-7	1.39 J	7.21	2.87	0.573 U	0.599 U	0.378 U	0.75 U	0.191 U
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9	2.05 J	4.21	1.69 J	0.573 U	0.399 U 0.872 J	3.11	2.28 J	1.91 U
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6	0.816 U	0.343 J	0.148 U	0.02 J 0.235 U	0.872 J 0.343 U	0.409 U	0.424 U	0.471 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9	1.12 U	2.18 J	0.679 J	0.233 U 0.276 J	0.476 J	1.2 J	0.424 U	0.471 U
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	57653-85-7	0.834 U	1.19 J	0.378 J	0.242 U	0.362 U	0.667 J	0.458 U	0.764 U
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9	0.834 U	1.01 J	0.254 U	0.419 U	0.302 U 0.411 U	0.397 U	0.438 U	0.433 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3	0.25 U	0.733 J	0.254 U	0.419 U	0.411 U 0.359 U	0.429 U	0.773 U	0.476 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6	2.38	5.58	3.45	0.242 0 1.44 J	0.339 U 2.71 J	5.02	5.53	4.39
1,2,3,7,8-Pentachiorodibenzo-p-dioxin (PeCDD)	40321-76-4	0.267 U	1.08 J	0.382 J	0.262 J	0.455 J	0.535 J	0.416 U	0.133 U
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5	0.634 U	1.75 J	0.566 J	0.202 J 0.364 U	0.453 J 0.369 U	0.333 J 0.311 U	0.410 U	0.133 U 0.251 U
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	57117-31-4	2.45	5.78	3.71	1.56 J	3.21	4.81	5.71	3.66
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	51207-31-9	9.58	15	23	10.7	28.2	18.5	18.2	12.8
2,3,7,8-Tetrachiorodibenzo-p-dioxin (TCDD)	1746-01-6	9.58 2.81 J	9.79	7.19	3.46 J	9.62	5.24	3.79	3.27
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4	2.81 J 18.6 J	9.79 14.6 J	7.19 9.51 J	3.46 J 4.73 J	9.62 8.2 J	21.3 J	28.5 J	3.27 21 J
<u> </u>		18.6 J		9.51 J 0.782 J	0.262 J	0.455 J	21.3 J 2.38 J	28.5 J 4.03 J	
Total Piovin/Furan TEO 1008 (Avian) (LL = 1/2)	36088-22-9	1.6 J 15.758 J	1.93 J 33.785 J	0.782 J 35.115 J	0.262 J 16.351 J	0.455 J 42.016 J	30.131 J	4.03 J 28.93 J	2.52 J 20.451 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	tDioxFurB_N								
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurF_N	5.458 J	16.454 J	11.264 J	5.376 J	13.546 J	9.973 J	8.584 J	6.326 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	tDioxFurM_N	5.361 J	16.393 J	11.728 J	5.592 J	14.276 J	9.856 J	8.225 J	6.14 J

			Tissue Ana	llytical Results				
Location ID	PB037_A	PB047.1_A	PB047.3	PB048.3	PB048.3	PB048.3	PB049_A	PB049 A
Sample ID	- PB02-I-A-BCR-W-004-COMP-201106	_	PB02-I-A-BCR-W-001-COMP-201106	PB02-F-A-GM-W-001-20110616	PB02-F-A-GM-W-002-20110616	PB02-F-A-STM-W-001-20110616	PB02-F-A-GKF-W-001-COMP-201106	PB02-I-A-BCR-W-003-COMP-201106
Sample Date	6/19/2011	6/15/2011	6/19/2011	6/16/2011	6/16/2011	6/16/2011	6/15/2011	6/22/2011
Sample Type	N	N	N N	N	N	N	N	N
x	3201325.97	3201461.79	3201645	3201519.15	3201519.15	3201519.15	3201468.39	3201468.39
y	13833885.45	13833007.68	13832925	13832869.89	13832869.89	13832869.89	13832722.8	13832722.8
Scientific Name	Callinectes sapidus	Lagodon rhomboides	Callinectes sapidus	Brevoortia patronus	Brevoortia patronus	Mugil cephalus	Fundulus grandis	Callinectes sapidus
Size_Class	>2 to =<7.5	=<15	>2 to =<7.5	=<15	=<15	=<15	=<15	>2 to =<7.5
Tissue_Category	Sedentary	Migratory	Sedentary	Migratory	Migratory	Migratory	Sedentary	Sedentary
	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Tissue_Type Chemical CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203 PCB-203	1130	5140	1000	3730	3520	1560	3800	764
PCB-204 PCB-204	16.6	51.6	50.5 J	58.7	54.7	17.9	110	22.1
PCB-205 PCB-205	31.6	389	37.5	357	317	132	320 U	26
PCB-206 PCB-206	807	4340	814	4680	4190	1140	2930	735
PCB-200 PCB-207	628	2750	1970	3270	3060	635	3860	517
PCB-208 PCB-208	789	3160	1050	4480	4060	850	4560	742
PCB-209 Total Describes his head because (14, 4/2) Total Describes his head because (14, 4/2)	13000	67300	18800	125000	113000	18600	80600	13500
Total Decachlorobiphenyl homologs (U = 1/2) tSum_DecPCB_N	13000	67300	18800	125000	113000	18600	80600	13500
Total Dichlorobiphenyl homologs (U = 1/2) tSum_DiPCB_N	4095.23	3723.3 J	1920.3	12059.98	10768.48	8480 J	14399.75 J	2300.57
Total Heptachlorobiphenyl homologs (U = 1/2) tSum_HepPCB_N	32602.245 J	115502.58	24978.86	100951.98	90261.13	48668.75	138321.94 J	20407.06 J
Total Hexachlorobiphenyl homologs (U = 1/2) tSum_HexPCB_N	105899.063 J	323349.22 J	83310.6 J	250905.95 J	236619.3 J	162589.1 J	400033.22 J	57782.83 J
Total Monochlorobiphenyl homologs (U = 1/2) tSum_MonPCB_N	1.75 U	150.9 J	6.24 U	263.1 J	218.58 J	122.56 J	360.7 J	2.926 J
Total Nonachlorobiphenyl homologs (U = 1/2) tSum_NonPCB_N	2224	10250	3834	12430	11310	2625	11350	1994
Total Octachlorobiphenyl homologs (U = 1/2) tSum_OctPCB_N	7552.2	29821.6	6702 J	25927.7	23689.7	10410.9	31690	5137.1
Total PCB Congener (U = 1/2) tPCBCong_N	1264098.298 J	3972325 J	825405.09 J	2424485.76 J	2275858.27 J	2745075.55 J	5144331.74 J	520316.246 J
Total PCB Congener minus 209 (U = 1/2) tPCBCong209_N	1251098.298 J	3905025 J	806605.09 J	2299485.76 J	2162858.28 J	2726475.56 J	5063731.74 J	506816.246 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2) tPCBCngCPB98_N	703.52	145	469.1	476.1	445.2	433.9	771.8 J	285.97
Total PCB Congener TEQ 1998 (Fish) (U = 1/2) tPCBCngCPF98_N	4.58	3.9	3	3.4	3.1	2.8	9.4 J	1.7
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2) tPCBCongCPM_N	49.2	51.4	31.7	39.2	35	28.3	137.1 J	16.86
Total Pentachlorobiphenyl homologs (U = 1/2) tSum_PenPCB_N	339821.52 J	1053025.5	234814.15 J	694848.1 J	644629.3	602024.5 J	1321625.78 J	123582.52 J
Total Tetrachlorobiphenyl homologs (U = 1/2) tSum_TetPCB_N	556153.18	2052944.2 J	360217.6 J	768205.6	721529.09	1502390.9 J	2602385.3	209317.02 J
Total Trichlorobiphenyl homologs (U = 1/2) tSum_TriPCB_N	202748.78 J	316257.6 J	90818.84 J	433893.4	423832.7	389163.8	543565.1	86292.23 J
Dioxin Furans (ng/kg)								
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) 39001-02-0	2.97 J	9	4.51 J	98.7	96.5	119	0.279 U	9.53
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD) 3268-87-9	6.05	12.1	11.9	76.7	76.7	75.8	1.93 J	5.24
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) 67562-39-4	1.51 J	0.821 J	1.27 J	6.03	6.08	5.72	0.107 U	1.86 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD) 35822-46-9	1.82 J	1.23 J	1.3 U	9.4	8.36	7.91	0.252 U	0.974 J
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) 55673-89-7	0.306 U	0.385 U	0.315 U	1.27 J	1.34 J	0.702 U	0.173 U	0.334 U
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) 70648-26-9	2.36 J	0.492 J	1.16 U	2.31 J	2.33 J	2.17 J	0.206 J	1.23 J
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD) 39227-28-6	0.819 J	0.209 U	0.668 U	0.321 U	0.403 U	0.513 U	0.129 U	0.337 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) 57117-44-9	1.21 J	0.251 J	0.496 U	1.06 J	1.18 J	0.99 J	0.167 J	0.603 J
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD) 57653-85-7	0.287 U	0.226 U	0.664 U	1.16 J	1.05 J	0.591 U	0.229 J	0.359 U
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) 72918-21-9	0.339 U	0.235 U	0.21 U	0.492 U	0.679 U	0.51 U	0.147 U	0.32 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD) 19408-74-3	0.425 J	0.221 U	0.687 U	0.348 U	0.421 U	0.561 U	0.136 U	0.354 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) 57117-41-6	4.9	2.38 J	2.95	4.94	4.79	3.48	0.130 U	2.81
	0.709 J	0.652 J	0.525 J	0.95 J	0.797 J	0.322 U	0.425 J	0.225 U
	0.709 J 0.299 U	0.032 J 0.239 U	0.525 J 0.166 U	0.854 J	0.797 J 0.671 U	0.322 U 0.493 U	0.425 J 0.112 U	0.304 U
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) 60851-34-5								
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) 57117-31-4	4.66	3.19	2.57	5.23	4.54	4.15	1.9 J	2.69
2,3,7,8-Tetrachlorodibenzofuran (TCDF) 51207-31-9	15.2	11.4	11.5	13.4	15.2	28.7	8.81	12 J
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) 1746-01-6	3.06	7.96	2.47	11	10.6	5.37	3.86	2.7
Total Pentachlorodibenzofuran (PeCDF) 30402-15-4	25.6	6.65 J	15.6 J	12.6 J	11 J	10.5	3.11 J	14.1 J
Total Pentachlorodibenzo-p-dioxin (PeCDD) 36088-22-9	6.05 J	0.652 J	1.8 J	0.95 J	0.797 J	0.322 U	0.425 J	1.26 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2) tDioxFurB_N	24.612 J	23.569 J	17.533 J	31.658 J	32.176 J	39.227 J	15.152 J	18.049 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2) tDioxFurF_N	7.927 J	11.062 J	5.294 J	16.123 J	15.298 J	9.803 J	5.809 J	5.223 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2) tDioxFurM_N	7.399 J	10.94 J	5.233 J	15.823 J	15.197 J	10.398 J	5.853 J	5.205 J

Tissue Analytical Results									
	Location ID	PB052.1	PB052.1	PB053.2	PB053.2	PB053 B	PB056_A	PB056_A	PB057.2
	Sample ID	PB02-F-A-SAS-W-002-20110616	PB02-F-A-STM-W-003-20110616	PB02-F-A-GKF-W-025-20110621	PB02-F-A-GKF-W-029-20110622	PB02-F-A-GKF-W-013-20110618	PB02-F-A-GKF-W-010-20110618	PB02-F-A-GKF-W-011-20110618	PB03-F-A-GM-W-001-20110617
	Sample Date	6/16/2011	6/16/2011	6/21/2011	6/22/2011	6/18/2011	6/18/2011	6/18/2011	6/17/2011
	Sample Type	N	N	N	N	N	N	N	N
	y x	3201515.24	3201515.24	3201266.16	3201266.16	3201390.48	3201420.22	3201420.22	3201603.32
	Ŷ	13832564.31	13832564.31	13832433.69	13832433.69	13832440.54	13832183.54	13832183.54	13832054.52
	Scientific_Name	Cynoscion arenarius	Mugil cephalus	Fundulus grandis	Brevoortia patronus				
	Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
	Tissue_Category	Migratory	Migratory	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Migratory
	Tissue_Type	Whole Body							
Chemical	CAS Number	Result							
PCB-203	PCB-203	4610	3140	7680	6050	6730 J	5890	4420	3020
PCB-204	PCB-204	74.8	28.9 J	138	105	106 J	102	73.6	41.7
PCB-205	PCB-205	380	250	345	283	331 J	289	232	240
PCB-206	PCB-206	6590	2160	4410	3750	4180	3990	3030	2920
PCB-207	PCB-207	5400 6720	1200 1500	5350 6450	4880 5620	4650 5970	4880	3340 3840	2010 2840
PCB-208	PCB-208						6190		
PCB-209	PCB-209	175000	30900	127000	107000 J	112000 J	116000	72000	73500
Total Dishlarshiphenyl homologs (U = 1/2)	tSum_DecPCB_N	175000	30900	127000	107000 J	112000 J	116000	72000	73500
Total Dichlorobiphenyl homologs (U = 1/2)	tSum_DiPCB_N	4904.9	16565.6 J	29765.61 J	25007.79	24812.5 J	20990.5 J	23556 J	9695.4 J
Total Heptachlorobiphenyl homologs (U = 1/2)	tSum_HepPCB_N	83018.77 J	90818.06 J	188026.74	162448.57 J	179898 J	158596.14 J	121518 J	73630.08 J
Total Hexachlorobiphenyl homologs (U = 1/2)	tSum_HexPCB_N	196161.1 J	266045.8	629408.7 J	532587.9 J	526082.54 J	410000.46	388340.7 J	192155.23 J
Total Monochlorobiphenyl homologs (U = 1/2)	tSum_MonPCB_N	67.08	304.39	377.4	358.8 J	377.84 J	379.9 J	363.6 J	189.9 J
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum_NonPCB_N	18710	4860	16210	14250	14800	15060	10210	7770
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N	29091.8	20644.9 J	50803	42418	45587 J	40891	29924.6	19709.7
Total PCB Congener (U = 1/2)	tPCBCong_N	2511956.11 J	4538570.8 J	8847995.31 J	7110917.26 J	6767235.33 J	5242306.61 J	5593763.6 J	2890241.46 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	2336956.11 J	4507670.8 J	8720995.31 J	7003917.26 J	6655235.33 J	5126306.61 J	5521763.6 J	2816741.46 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98_N	188.1 J	789.3	1152.5 J	987 J	1091.86215 J	943.33 J	865.8 J	440.06 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N	2.2 J	5.1	9.7 J	8.2 J	9.349275 J	8.09 J	7.2 J	3.01 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	tPCBCongCPM_N	25.9 J	51.5	115.7 J	97.5 J	111.43945 J	96.35 J	85.6 J	33.08 J
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum_PenPCB_N	548834.9 J	987795.2 J	1924379.6 J	1571688.6 J	1656118.3 J	1168212.7 J	1304574.8 J	586428.2 J
Total Tetrachlorobiphenyl homologs (U = 1/2)	tSum_TetPCB_N	1159897.1 J	2416992.6 J	4655465.5 J	3715735.1	3325852.4 J	2620869.3 J	3063492.9 J	1512803.4
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N	296270.5 J	703644.2	1226558.7 J	939422.6	881706.7 J	691306.7 J	579783	414359.5
Dioxin Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0	3.2 J	15.2	0.23 U	0.89 J	1.43 J	1.76 J	0.988 J	57.5
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	3268-87-9	3.08 J	21.7	5.04	5.13	6.83	6.95	4.03 J	57.7
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	67562-39-4	0.441 J	1.31 J	0.132 U	0.326 J	0.217 U	2.08 J	0.353 J	4.46
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	35822-46-9	0.549 J	2.38	0.621 J	0.881 J	1.25 J	0.796 J	0.601 J	6.42
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	55673-89-7	0.294 U	0.211 U	0.223 U	0.19 U	0.362 U	0.404 U	0.283 U	0.697 J
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9	0.403 J	0.794 J	0.365 J	0.397 J	0.19 U	0.33 J	0.198 U	1.58 J
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6	0.121 U	0.136 U	0.155 U	0.104 U	0.29 U	0.214 U	0.23 U	0.189 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9	0.243 J	0.34 J	0.345 J	0.25 J	0.183 U	0.522 J	0.196 U	0.712 J
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	57653-85-7	0.285 J	0.325 J	0.158 U	0.459 J	0.316 U	0.225 U	0.233 U	0.587 J
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9	0.181 U	0.145 U	0.249 U	0.198 U	0.304 U	0.421 U	0.314 U	0.3 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3	0.129 U	0.146 U	0.159 U	0.197 U	0.308 U	0.224 U	0.235 U	0.198 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6	1.08 J	2.13 J	2.25 J	2.26 J	1.72 J	1.72 J	1.22 J	3.11
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	40321-76-4	0.162 U	0.43 J	1.03 J	1.21 J	2.22 J	0.685 J	0.63 J	0.726 J
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5	0.181 J	0.223 J	0.258 U	0.187 U	0.247 U	0.44 U	0.316 U	0.552 J
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	57117-31-4	0.356 J	2.88	4.55 J	3.87	3.9 J	3.3	2.56	3.43
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	51207-31-9	1.05 U	18.8	15.6	13.2	12.7	12.1	9.91	10.1
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1746-01-6	0.8 J	3.65	8.22	6.18	7.43	6.64	4.81	5.97
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4	1.92 J	6.94	8.09 J	7.04 J	7.43 7.23 J	10.8 J	4.71 J	8.67 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9	0.162 U	0.43 J	1.03 J	1.21 J	2.66 J	0.685 J	0.63 J	0.949 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	tDioxFurB_N	1.981 J	26.15 J	29.737 J	24.793 J	26.497 J	23.067 J	18.108 J	20.926 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurB_N	1.961 J	6.768 J	12.557 J	10.219 J	26.497 J 12.448 J	9.874 J	7.394 J	9.495 J
. , , , , ,	tDioxFurF_N	1.272 J 1.219 J	7.126 J	12.372 J	10.219 J 10.099 J	12.448 J 12.251 J	9.874 J 9.771 J	7.394 J 7.334 J	9.495 J 9.356 J
rotal Dioxili/Furali TEQ 2005 (Mammal) (U = 1/2)	thioxentivi_N	1.219 J	7.120 J	12.3/2 J	10.099 1	12.251 J	9.//1 J	7.334 J	9.350 J

Tissue Analytical Results									
	Location ID	PB057.2	PB057.2	PB057.2	PB057_B	PB057 B	PB057_B	PB058.1	PB058.1
	Sample ID		PB03-F-A-STM-W-001-COMP-201106		PB03-F-A-GKF-W-001-20110615	PB03-F-A-GKF-W-006-20110617	PB03-F-A-PNF-W-001-COMP-201106	PB03-F-A-GM-W-003-20110618	PB03-F-A-GM-W-004-20110618
	Sample Date	6/17/2011	6/17/2011	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/18/2011	6/18/2011
	Sample Type	N	N	N	N	N	N	N	N
	х	3201603.32	3201603.32	3201603.32	3201644.06	3201644.06	3201644.06	3201682.3	3201682.3
	Ŷ	13832054.52	13832054.52	13832054.52	13832071.63	13832071.63	13832071.63	13832011.93	13832011.93
	Scientific_Name	Mugil cephalus	Mugil cephalus	Mugil cephalus	Fundulus grandis	Fundulus grandis	Lagodon rhomboides	Brevoortia patronus	Brevoortia patronus
	Size_Class	=<15	=<15	=<15	=<15	=<15	=<15	=<15	=<15
	Tissue_Category	Migratory	Migratory	Migratory	Sedentary	Sedentary	Migratory	Migratory	Migratory
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203	PCB-203	1700	821	1490	5620	4860	1710	2700	2710
PCB-204	PCB-204	17	11	16.4	73.7	64.7	18.7	39.7	42.5
PCB-205	PCB-205	132	84.6	113	283	248	158	215	267
PCB-206	PCB-206	1080	736	1080	3350	3430	1550	2790	3320
PCB-207	PCB-207	586	445	592	3990	3810	966	1920	2280
PCB-208	PCB-208	739	586	768	5390	4290	1060	2640	3110
PCB-209	PCB-209	15600	12300	18500	94200	83900	21500	71800	83800
Total Decachlorobiphenyl homologs (U = 1/2)	tSum_DecPCB_N	15600	12300	18500	94200	83900	21500	71800	83800
, , , , , , , , , , , , , , , , , , , ,	tSum_DecPCB_N	4950.3 J	6646.06	7032.5 J	94200 21726.7 J	23867	3417.7 J	11648.8 J	16831.96
Total Dichlorobiphenyl homologs (U = 1/2) Total Heptachlorobiphenyl homologs (U = 1/2)		4950.3 J 44949.26 J	29164.11	35706.14 J	160714.93	137128.26	44589.349	63061.07 J	72913.05 J
	tSum_HepPCB_N	117921.73 J	102427.08	99345.8 J	439167.2 J	13/128.26 476852.3 J	44589.349 142368.5 J	181784.04 J	268118 J
Total Hexachlorobiphenyl homologs (U = 1/2) Total Monochlorobiphenyl homologs (U = 1/2)	tSum_HexPCB_N		102427.08 156.76 J	99345.8 J 114.91 J	439167.2 J 315.5 J	389.1 J	142368.5 J 117.65 J	230.1	349.1 J
1 7 0 1 7	tSum_MonPCB_N	93.8 J							
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum_NonPCB_N	2405	1767	2440	12730	11530	3576	7350	8710
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N	10545	5848.6	9098.4	39476.7	34192.7	10008.7	17399.7	18115.5
Total PCB Congener (U = 1/2)	tPCBCong_N	1504838.03 J	1735514.48 J	1870516.01 J	4600511.58 J	7061455.81 J	2206629.039 J	3070347.7 J	4005386.97 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	1489238.03 J	1723214.48 J	1852016.01 J	4506311.58 J	6977555.81 J	2185129.039 J	2998547.7 J	3921586.97 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98_N	273.09 J	324.86	267.9 J	826.9	959.8	102.59	497.01 J	714.3
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N	1.87 J	1.28	1.7 J	6.9	7.7	2.34	3.04 J	4.4
	tPCBCongCPM_N	19.27 J	4.26	17.7 J	80.2	87.5	30.55	31.62 J	45.7
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum_PenPCB_N	338692.63 J	367426.67 J	348722.8 J	1532099.3 J	1636453.7 J	501612.91 J	591663.4 J	972330.4 J
Total Tetrachlorobiphenyl homologs (U = 1/2)	tSum_TetPCB_N	752136.5 J	924411.3	1043528.6 J	1533578.6	3689904.7	1218044.4 J	1648366.8 J	1975824.46
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N	217543.8 J	285366.89	306026.8 J	766502.8	967238.1	261393.88	477043.9	588394.6
Dioxin Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0	48.6	25.8	44.4	1.06 J	1.47 J	1.69 J	83.7	122
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)		52.4	33.4	63.5	5.77	7.4	3.46 J	136	233
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	67562-39-4	3.34	2.36 J	4.12	0.463 J	0.315 J	0.255 J	8.89	11.4
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)		5.76	3.69	7.12	0.707 J	0.747 J	0.253 U	14.7	28
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	55673-89-7	0.679 U	0.382 U	0.721 J	0.142 U	0.248 U	0.207 U	1.27 J	2.13 J
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9	1.17 J	0.767 J	1.33 J	0.241 J	0.231 J	0.11 U	2.6	3.44
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6	0.461 U	0.214 U	0.262 U	0.0871 U	0.134 U	0.165 U	0.292 U	2.83 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9	0.542 J	0.373 J	0.815 J	0.149 J	0.12 J	0.102 U	1.31 J	2.01 J
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	57653-85-7	0.493 U	0.289 J	0.462 J	0.204 J	0.248 J	0.184 U	0.962 J	2.76 U
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9	0.681 U	0.331 U	0.365 U	0.198 U	0.11 U	0.156 U	0.443 U	0.464 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3	0.484 U	0.227 U	0.286 U	0.0913 U	0.144 U	0.178 U	0.294 U	2.89 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6	1.43 J	1.16 J	1.74 J	1.01 J	1.12 J	0.678 J	3.29	4.87
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	40321-76-4	0.449 J	0.163 U	0.473 J	0.752 J	0.536 J	0.267 J	0.784 J	0.687 U
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5	0.617 U	0.273 U	0.56 J	0.172 U	0.118 J	0.145 U	0.782 J	1.38 J
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	57117-31-4	1.83 J	1.37 J	2.38 J	2.58	2.59	0.916 J	3.92	5.25
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	51207-31-9	10	8.12	10.5	10.9	10.9	4.27	11.2	16.8
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1746-01-6	2.02	1.43	2.81	6.57	5.64	2.29	5.86 J	8.18
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4	5.97 J	3.57 J	6.91 J	4.53 J	4.68 J	1.59 J	11.7 J	17.8 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9	0.449 J	0.163 U	0.473 J	0.752 J	0.536 J	0.267 J	1.74 J	0.687 U
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	tDioxFurB_N	14.769 J	11.316 J	16.718 J	20.976 J	19.849 J	7.855 J	22.754 J	32.195 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurF_N	4.364 J	2.897 J	5.512 J	9.296 J	8.167 J	3.335 J	10.042 J	13.873 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	tDioxFurM_N	4.494 J	3.045 J	5.614 J	9.3176 J	8.182 J	3.338 J	9.97 J	13.577 J

Tissue Analytical Results									
	Location ID	PB058.1	PB059.1_A	PB059.1_A	PB059.3	PB059.4	PB059.4	PB059.5	PB063.1_A
	Sample ID	PB03-F-A-GM-W-005-20110618	PB03-F-A-GKF-W-003-20110615	PB03-I-A-BCR-W-003-COMP-201106	PB03-I-B-BCR-W-002-20110627	PB03-I-B-BCR-W-003-20110628	PB03-I-B-BCR-W-003-20110628DUP	PB03-I-A-BCR-W-001-COMP-201106	PB03-F-A-GKF-W-009-20110617
	Sample Date	6/18/2011	6/15/2011	6/19/2011	6/26/2011	6/27/2011	6/27/2011	6/27/2011	6/17/2011
	Sample Type	N	N	N	N	N	N	N	N
	х х	3201682.3	3201501.32	3201501.32	3201714.16	3201714.16	3201714.16	3201714	3201509.99
	Υ	13832011.93	13831828.71	13831828.71	13831964.63	13831964.63	13831964.63	13831965	13831505.57
	Scientific_Name	Brevoortia patronus	Fundulus grandis	Callinectes sapidus	Callinectes sapidus	Callinectes sapidus		Callinectes sapidus	Fundulus grandis
	Size_Class	=<15	=<15	>2 to =<7.5	>7.5 to =<13	>7.5 to =<13		>2 to =<7.5	=<15
	Tissue_Category	Migratory	Sedentary	Sedentary	Migratory	Migratory		Sedentary	Sedentary
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body		Whole Body	Whole Body
Chemical	CAS Number	Result	Result	Result	Result	Result	Result	Result	Result
PCB-203	PCB-203	2870	4780	517	727	712	672	486	4340
PCB-204	PCB-204	46.2 J	50.5	7.81	15.3	14.6	10.7	7.67 J	56.2
PCB-205	PCB-205	283	221	16.5	19.7 J	17.6	19.1	10.7	201
PCB-206	PCB-206	3670	2950	417	435	405	403	234	2810
PCB-207	PCB-207	2420	3040	263	408	433	413	230	3130
PCB-208	PCB-208	3350	3880	397	458	475	442	262	3830
PCB-209	PCB-209	92100	76300	5250	8140	9540	8910	4790	74800
Total Decachlorobiphenyl homologs (U = 1/2)	tSum_DecPCB_N	92100	76300	5250	8140	9540	8910	4790	74800
Total Dichlorobiphenyl homologs (U = 1/2)	tSum_DiPCB_N	17103.07	29896.1 J	1813.96 J	2069.37 J	2110.92	1897.32 J	1708.73 J	21805.3 J
Total Heptachlorobiphenyl homologs (U = 1/2)	tSum_HepPCB_N	76888.26 J	141148.8	16554.938 J	20608.1 J	19585.11 J	18193.31 J	15426.38 J	122860.98 J
Total Hexachlorobiphenyl homologs (U = 1/2)	tSum_HexPCB_N	268334.3 J	395884.1	52076.04 J	79130.864 J	76342.01 J	73499.5 J	57726.24 J	432968.5 J
Total Monochlorobiphenyl homologs (U = 1/2)	tSum_MonPCB_N	312.7 J	444	3.11 J	9.52 J	6.153 J	3.04 U	2.75 U	309.9 J
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum_NonPCB_N	9440	9870	1077	1301	1313	1258	726	9770
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N	19985.2 J	32896.5	3466.11	4680 J	4593.2	4190.8	3097.37 J	29398.2
Total PCB Congener (U = 1/2)	tPCBCong_N	4003442.03 J	6501404.6 J	623702.118 J	752900.594 J	722043.627 J	694017.63 J	695657.24 J	4685390.62 J
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	3911342.03 J	6425104.6 J	618452.118 J	744760.594 J	712503.627 J	685107.63 J	690867.24 J	4610590.62 J
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98_N	789.2 J	982.9	370.56	485.08	433.73	408.33	511.7	740 J
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N	4.7 J	7.5	2.94	3.09	2.87	2.6	2.98	3.5 J
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2	tPCBCongCPM_N	46.6 J	84.4	37.21	32.85	30.79	27.16	30.16	15.6 J
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum_PenPCB_N	969180.2 J	1404509.7	161350.56 J	229441.19 J	228495.22 J	212821.01 J	186234.23 J	1105174.8 J
Total Tetrachlorobiphenyl homologs (U = 1/2)	tSum_TetPCB_N	1934015.4 J	3462382.3 J	284966.08 J	324368.23	299915.71 J	295170.65 J	322840.44	2275250.4
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N	616083 J	948073.1	97144.32 J	83152.32	80142.318 J	78072.95 J	103103.76 J	613052.4
Dioxin Furans (ng/kg)	l								
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0	134	0.49 U	1.71 J				2.52 J	0.204 U
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDI	0) 3268-87-9	215	7.3 J	3.97 J				13.3	3.15 J
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	67562-39-4	12.9	0.272 U	0.49 J				1.44 J	0.522 J
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCD		25	0.647 J	0.977 J				2.31 J	0.401 J
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	55673-89-7	1.72 J	0.467 U	0.202 U				0.359 U	0.217 U
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9	3.7 J	0.205 U	0.668 J				1.12 J	0.095 U
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6	2.29 U	0.24 U	0.18 U				0.275 U	0.154 U
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9	2 J	0.19 U	0.352 J				0.628 J	0.0981 U
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	57653-85-7	2.35 U	0.246 U	0.408 J				0.431 J	0.167 U
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9	0.533 U	0.312 U	0.2 U				0.421 U	0.158 U
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3	2.39 U	0.247 U	0.19 U				0.292 U	0.163 U
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6	4.44	0.827 J	1.16 J				2.27 J	0.624 J
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	40321-76-4	0.618 U	0.613 J	0.309 J				0.593 J	1.31 J
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5	1.18 J	0.267 U	0.141 U				0.382 U	0.149 U
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	57117-31-4	5.93 J	2.22 J	1.28 J				1.89 J	1.76 J
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	51207-31-9	17	10.7	4.45				9.23	7.52
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1746-01-6	8.13 J	4.86	1.09 J				2.25 J	3.84
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4	18.9 J	3.86 J	6.46 J				12 J	3.93 J
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9	0.618 U	0.613 J	1.86 J				3.78 J	1.31 J
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	tDioxFurB_N	32.922 J	18.549 J	7.39 J				14.451 J	14.5373 J
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurF_N	13.993 J	7.276 J	2.496 J				4.673 J	6.5094 J
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)		13.618 J	7.332 J	2.459 J				4.732 J	6.5092 J
	_	[l .	1		1	1	j	

Tissue Analytical Results									
Location ID	PB063.1_A	PB063.2_A	PB063.2_A	PB069_A	PB069 A	PB069_A	PB084.1	PB084.1	
	PB03-F-A-GKF-W-010-20110617	PB03-F-A-GKF-W-005-20110615	PB03-F-A-GKF-W-013-20110617	PB03-F-A-GKF-W-002-20110615	PB03-F-A-GKF-W-007-20110617	PB03-I-A-BCR-W-002-COMP-201106	PB04-F-A-GKF-W-005-20110627	PB04-F-A-GKF-W-013-20110627	
Sample Date	6/17/2011	6/15/2011	6/17/2011	6/15/2011	6/17/2011	6/19/2011	6/27/2011	6/27/2011	
Sample Type	N	N	N	N	N	N N	N	N	
у х	3201509.99	3201561.28	3201561.28	3201132.36	3201132.36	3201132.36	3200727.85	3200727.85	
	13831505.57	13831519.55	13831519.55	13830993.22	13830993.22	13830993.22	13829784.47	13829784.47	
Scientific Name	Fundulus grandis	Callinectes sapidus	Fundulus grandis	Fundulus grandis					
Size_Class	=<15	=<15	=<15	=<15	=<15	>2 to =<7.5	=<15	=<15	
Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	Sedentary	
Tissue_Type	Whole Body	Whole Body	Whole Body						
Chemical CAS Number	Result	Result	Result	Result	Result	Result	Result	Result	
PCB-203 PCB-203	5210	4060	4080	5900	4280	666	1910	2790 J	
PCB-204 PCB-204	63.2	53.8	64.3	74.7	51	11.2	25 J	36.3 J	
PCB-205 PCB-205	215	196	183	302 J	190	21.3	103	147	
PCB-206 PCB-206	2790	2740	2630	3320	2340	469	1260	1800	
PCB-207 PCB-207	3000	3060	2780	3370	2360	397	1430	1990	
PCB-208 PCB-208	3780	3680	3640	4270	2990	544	1630	2270	
PCB-208 PCB-209 PCB-209	75100 J	74600	69500	82200	55400	7980	33400	44200	
	75100 J 75100 J	74600	69500	82200	55400	7980	33400	44200	
	26948.8 J	20284.5 J		34020.4 J	27457.5 J	3766.99 J	12526.84	20469.57	
Total Dichlorobiphenyl homologs (U = 1/2) tSum_DiPCB_N Total Heptachlorobiphenyl homologs (U = 1/2) tSum_HepPCB_N	145153.57	20284.5 J 124890.2 J	18115.41 J 111936.6	34020.4 J 156574.73	118031.79	23639.244	52722.93 J	78626.28 J	
Total Hexachlorobiphenyl homologs (U = 1/2) tSum_HexPCB_N	426399.9	384274.5 J	333396.54 J	544841.4 J	435476.8 J	84502.58 J	202833.5 J	317819.4 J	
Total Monochlorobiphenyl homologs (U = 1/2) tSum_MonPCB_N	338.32	265.9 J	313.94	598.7 J	496.3 J	5.53 J	269.63 J	409.7 J	
Total Nonachlorobiphenyl homologs (U = 1/2) tSum_NonPCB_N	9570	9480	9050	10960	7690	1410	4320	6060	
Total Octachlorobiphenyl homologs (U = 1/2) tSum_OctPCB_N	35008.2	30020.8	27987.3	39766.7 J	28637	4891.5	13314 J	19392.3 J	
Total PCB Congener (U = 1/2) tPCBCong_N	6000677.74 J	5938890.2 J	4473197.15 J	7758932.13 J	5075277.69 J	1142429.629 J	2915816.39 J	4586762.27 J	
Total PCB Congener minus 209 (U = 1/2) tPCBCong209_N	5925577.73 J	5864290.2 J	4403697.16 J	7676732.12 J	5019877.69 J	1134449.629 J	2882416.4 J	4542562.27 J	
Total PCB Congener TEQ 1998 (Avian) (U = 1/2) tPCBCngCPB98_N	876 J	792.8	741.1 J	1172.78974 J	908.2 J	782.88	454.2 J	739	
Total PCB Congener TEQ 1998 (Fish) (U = 1/2) tPCBCngCPF98_N	7 J	6.9	6 J	9.52337 J	7.5 J	4.42	3.4 J	5.5	
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2) tPCBCongCPM_N	80.9 J	81.6	70.3 J	112.62322 J	89.4 J	44.49	38.2 J	62.3	
Total Pentachlorobiphenyl homologs (U = 1/2) tSum_PenPCB_N	1340980.9 J	1381228 J	1065454.2 J	1818503.8 J	1236961.9 J	276629.09 J	732979.56 J	1183753.8 J	
Total Tetrachlorobiphenyl homologs (U = 1/2) tSum_TetPCB_N	3019963.4	3123259.1 J	2181050.1	4065557.1	2429584.6 J	569039.04	1461402.13 J	2295631.97	
Total Trichlorobiphenyl homologs (U = 1/2) tSum_TriPCB_N	921214.7	790587.2	656393	1005909.3	735541.8	170565.66 J	402047.8	620399.26	
Dioxin Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF) 39001-02-0	0.467 J	0.302 U	0.434 U	1 J	0.925 J	1.8 J	24.7	6.67	
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD) 3268-87-9	3.23 J	5.57	6.27	4.83	10.8	4.8	48.2	23.8	
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF) 67562-39-4	0.426 J	0.315 J	0.159 U	0.291 J	1.83 J	0.944 J	1.98 J	3.59	
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD) 35822-46-9	0.443 J	0.726 J	0.566 J	0.417 J	1.23 J	1.54 J	4.71 U	2.42 U	
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF) 55673-89-7	0.285 U	0.21 U	0.272 U	0.244 U	0.28 U	0.205 U	1.14 U	0.534 U	
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF) 70648-26-9	0.165 J	0.212 J	0.143 U	0.147 J	0.183 J	0.678 J	0.534 U	0.272 U	
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD) 39227-28-6	0.183 U	0.133 U	0.206 U	0.162 U	0.156 U	0.18 U	3.05 U	1.77 U	
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF) 57117-44-9	0.156 J	0.167 J	0.147 U	0.126 J	0.529 J	0.429 J	0.529 U	0.271 U	
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD) 57653-85-7	0.193 U	0.231 J	0.217 U	0.24 J	0.177 U	0.481 J	3.06 U	1.85 U	
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF) 72918-21-9	0.225 U	0.144 U	0.241 U	0.189 U	0.239 U	0.226 U	0.76 U	0.401 U	
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD) 19408-74-3	0.191 U	0.137 U	0.215 U	0.164 U	0.17 U	0.186 U	3.16 U	1.87 U	
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF) 57117-41-6	0.813 J	0.827 J	0.846 J	0.964 J	0.581 J	1.15 J	0.776 J	0.654 J	
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD) 40321-76-4	0.702 J	0.608 J	0.599 J	1.46 J	1.34 J	0.416 J	1.22 U	0.581 U	
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF) 60851-34-5	0.174 U	0.135 U	0.212 U	0.186 U	0.226 U	0.207 U	0.666 U	0.327 U	
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF) 57117-31-4	2.24 J	2.35 J	2.2 J	2.48	1.67 J	1.57 J	1.66 J	1.31 J	
2,3,7,8-Tetrachlorodibenzofuran (TCDF) 51207-31-9	9.42	9.36	8.31	10.4	8.16	6	8.86	8.32	
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) 1746-01-6	4.23	4.58	4.06	5	3.34	1.69 J	3	2.85	
Total Pentachlorodibenzofuran (PeCDF) 30402-15-4	4.14 J	3.96 J	3.76 J	4.5 J	5.96 J	8.06 J	2.43 J	2.38 J	
Total Pentachlorodibenzo-p-dioxin (PeCDD) 36088-22-9	0.702 J	0.608 J	0.599 J	1.46 J	1.34 J	4.18 J	1.22 U	0.581 U	
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2) tDioxFurB_N	16.747 J		15.311 J	19.502 J	14.698 J	9.955 J	14.617 J	13.089 J	
Total Dioxili/Furall TEQ 1996 (Aviall) (0 = 1/2) [tDioxFurb IV]	16.747 J	17.051 J	13.311 1	19.302 1	14.050 3		14.017 3		
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	6.67 J	6.966 J	6.311 J	8.363 J	6.109 J	3.444 J	5.875 J	4.812 J	

	Tissue Analytical Results								
	Location ID	PB095.2	PB101_D	PB101_D	PB101_D	PB102			
	Sample ID	PB04-I-A-BCR-W-001-COMP-201106	PB04-F-A-GKF-W-001-20110627	PB04-F-A-GKF-W-001-COMP-201106	PB04-F-A-GKF-W-009-20110627	PB04-F-A-PNF-W-001-COMP-201106			
	Sample Date	6/29/2011	6/27/2011	6/27/2011	6/27/2011	6/27/2011			
	Sample Type	N	N	N	N	N			
	х х	3200882	3201317.36	3201317.36	3201317.36	3201302.69			
	Υ	13829528	13828141.83	13828141.83	13828141.83	13828160.48			
	Scientific_Name		Fundulus grandis	Fundulus grandis	Fundulus grandis	Lagodon rhomboides			
	Size_Class	>2 to =<7.5	=<15	=<15	=<15	=<15			
	Tissue_Category	Sedentary	Sedentary	Sedentary	Sedentary	Migratory			
	Tissue_Type	Whole Body	Whole Body	Whole Body	Whole Body	Whole Body			
Chemical	CAS Number	Result	Result	Result	Result	Result			
PCB-203	PCB-203	250	3150	3950	3470	1860			
PCB-204	PCB-204	7.38 U	43	72.4 J	47.8	18.3			
PCB-205	PCB-205	12.4 U	158	226	180	149			
PCB-206	PCB-206	111	2010	3020	2480	1240			
PCB-207	PCB-207	95.8	2100	3100	2510	776			
PCB-208	PCB-208	119	2500	3810	3000	728			
PCB-208	PCB-208	1640	47500	64600	59600	14300			
Total Decachlorobiphenyl homologs (U = 1/2)		1640	47500	64600	59600	14300			
1 , 0 , , ,	tSum_DecPCB_N			22549.28 J	9430.48				
Total Dichlorobiphenyl homologs (U = 1/2)	tSum_DiPCB_N	1208.4 J 6758.01 J	14274.78 88382.18 J	22549.28 J 100131.66 J	9430.48 94858.73 J	13722.05 J 44705.63			
Total Heptachlorobiphenyl homologs (U = 1/2)	tSum_HepPCB_N								
Total Hexachlorobiphenyl homologs (U = 1/2)	tSum_HexPCB_N	27359.26 J	341351 J	351778.9 J	349287.2	177881.54			
Total Monochlorobiphenyl homologs (U = 1/2)	tSum_MonPCB_N	6.69 U	244.54 J	499.7 J	174.76 J	460.43			
Total Nonachlorobiphenyl homologs (U = 1/2)	tSum_NonPCB_N	325.8	6610	9930	7990	2744			
Total Octachlorobiphenyl homologs (U = 1/2)	tSum_OctPCB_N	1638.89 J	22399	27248.4 J	23942.8	10194.3			
Total PCB Congener (U = 1/2)	tPCBCong_N	361364.24 J	4123009.04 J	5547280.5 J	4322176.71 J	4333632.6 J			
Total PCB Congener minus 209 (U = 1/2)	tPCBCong209_N	359724.24 J	4075509.04 J	5482680.5 J	4262576.71 J	4319332.6 J			
Total PCB Congener TEQ 1998 (Avian) (U = 1/2)	tPCBCngCPB98_N	231	623.1 J	837.3	694.7	147.6 J			
Total PCB Congener TEQ 1998 (Fish) (U = 1/2)	tPCBCngCPF98_N	1.4	5.4 J	6.5	5.8	2.99 J			
Total PCB Congener TEQ 2005 (Mammal) (U = 1/2)	tPCBCongCPM_N	14.5	64.8 J	75.3	69.4	39 J			
Total Pentachlorobiphenyl homologs (U = 1/2)	tSum_PenPCB_N	99338.93 J	1114954.9	1260998.8	1217529.5	852292.8 J			
Total Tetrachlorobiphenyl homologs (U = 1/2)	tSum_TetPCB_N	169215.94 J	2001760.66 J	2958484.5	2113004	2532762			
Total Trichlorobiphenyl homologs (U = 1/2)	tSum_TriPCB_N	53869.91 J	485531.97 J	751059.2	446359.29 J	684569.9			
Dioxin Furans (ng/kg)									
1,2,3,4,6,7,8,9-Octachlorodibenzofuran (OCDF)	39001-02-0	2.23 J	4.46 J	0.477 U	2.92 J	1.56 J			
1,2,3,4,6,7,8,9-Octachlorodibenzo-p-dioxin (OCDD)	3268-87-9	12.7	36	5.74	25.6	3.82 J			
1,2,3,4,6,7,8-Heptachlorodibenzofuran (HpCDF)	67562-39-4	0.637 J	1.1 J	0.126 U	0.42 U	0.14 U			
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin (HpCDD)	35822-46-9	1.02 U	5.55 J	0.69 J	2.69 U	0.357 U			
1,2,3,4,7,8,9-Heptachlorodibenzofuran (HpCDF)	55673-89-7	0.225 U	0.794 U	0.227 U	0.693 U	0.242 U			
1,2,3,4,7,8-Hexachlorodibenzofuran (HxCDF)	70648-26-9	0.417 U	0.577 J	0.119 U	0.262 U	0.405 J			
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	39227-28-6	0.561 U	1.92 U	0.177 U	2.01 U	0.135 U			
1,2,3,6,7,8-Hexachlorodibenzofuran (HxCDF)	57117-44-9	0.156 U	0.322 U	0.117 U	0.251 U	0.153 U			
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin (HxCDD)	57653-85-7	0.575 U	2.02 U	0.19 U	1.97 U	0.141 U			
1,2,3,7,8,9-Hexachlorodibenzofuran (HxCDF)	72918-21-9	0.121 U	0.477 U	0.188 U	0.382 U	0.229 U			
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin (HxCDD)	19408-74-3	0.586 U	2.04 U	0.187 U	2.05 U	0.14 U			
1,2,3,7,8-Pentachlorodibenzofuran (PeCDF)	57117-41-6	0.334 U	0.651 J	0.896 J	0.559 J	1.5 J			
1,2,3,7,8-Pentachlorodibenzo-p-dioxin (PeCDD)	40321-76-4	0.15 U	0.724 U	0.67 J	0.643 U	0.152 U			
2,3,4,6,7,8-Hexachlorodibenzofuran (HxCDF)	60851-34-5	0.118 U	0.368 U	0.164 U	0.343 U	0.172 U			
2,3,4,7,8-Pentachlorodibenzofuran (PeCDF)	57117-31-4	0.554 U	1.47 J	2.33 J	1.79 J	1.68 J			
2,3,7,8-Tetrachlorodibenzofuran (TCDF)	51207-31-9	1.65	6.21	8.24	7.56	10.7			
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)	1746-01-6	0.398 J	3.28	3.99	7.50 3.98 J	2.64			
Total Pentachlorodibenzofuran (PeCDF)	30402-15-4	0.398 J 4.5 J	3.28 2.92 J	3.99 J	3.98 J	2.04 3.58 J			
· ,									
Total Pentachlorodibenzo-p-dioxin (PeCDD)	36088-22-9	0.268 J	0.724 U	0.67 J	0.643 U	0.152 U			
Total Dioxin/Furan TEQ 1998 (Avian) (U = 1/2)	tDioxFurB_N	2.513 J	11.688 J	15.367 J	13.942 J	15.328 J			
Total Dioxin/Furan TEQ 1998 (Fish) (U = 1/2)	tDioxFurF_N	0.899 J	5.361 J	6.36 J	6.197 J	4.272 J			
Total Dioxin/Furan TEQ 2005 (Mammal) (U = 1/2)	tDioxFurM_N	0.87 J	5.221 J	6.277 J	6.002 J	4.429 J			

Notes:

J = Estimated value
U = Compound analyzed, but not detected above detection limit
UJ = Compound analyzed, but not detected above estimated detection limit
R = Rejected

-- Results not reported or not applicable

pct = percent mg/kg = milligrams per kilogram μg/kg = micrograms per kilogram ng/kg = nanograms per kilogram

EB - Equipment Blank
N = Normal Field Sample

TEQ values as of 2005, World Health Organization.